

# Do methylenetetrahydrofolate dehydrogenase, cyclohydrolase, and formyltetrahydrofolate synthetase 1 polymorphisms modify changes in intelligence of school-age children in areas of endemic fluorosis?

Zichen Feng<sup>1</sup>, Ning An<sup>1</sup>, Fangfang Yu<sup>1</sup>, Jun Ma<sup>2</sup>, Na Li<sup>3</sup>, Yuhui Du<sup>1</sup>, Meng Guo<sup>1</sup>, Kaihong Xu<sup>1</sup>, Xiangbo Hou<sup>1</sup>, Zhiyuan Li<sup>1</sup>, Guoyu Zhou<sup>1</sup>, Yue Ba<sup>1</sup>

<sup>1</sup>Department of Occupational and Environmental Health, School of Public Health, Zhengzhou University, Zhengzhou, Henan 450001, China;

<sup>2</sup>Department of Endemic Disease, Kaifeng Center for Disease Control and Prevention, Kaifeng, Henan 475004, China;

<sup>3</sup>Dietary Center, Zhengzhou Health Vocational College, Zhengzhou, Henan 450122, China.

## Abstract

**Background:** Excessive exposure to fluoride can reduce intelligence. Methylenetetrahydrofolate dehydrogenase, cyclohydrolase, and formyltetrahydrofolate synthetase 1 (*MTHFD1*) polymorphisms have important roles in neurodevelopment. However, the association of *MTHFD1* polymorphisms with children's intelligence changes in endemic fluorosis areas has been rarely explored.

**Methods:** A cross-sectional study was conducted in four randomly selected primary schools in Tongxu County, Henan Province, from April to May in 2017. A total of 694 children aged 8 to 12 years were included in the study with the recruitment by the cluster sampling method. Urinary fluoride (UF) and urinary creatinine were separately determined using the fluoride ion-selective electrode and creatinine assay kit. Children were classified as the high fluoride group and control group according to the median of urinary creatinine-adjusted urinary fluoride (UF<sub>Cr</sub>) level. Four loci of *MTHFD1* were genotyped, and the Combined Raven's Test was used to evaluate children's intelligence quotient (IQ). Generalized linear model and multinomial logistic regression model were performed to analyze the associations between children's UF<sub>Cr</sub> level, *MTHFD1* polymorphisms, and intelligence. The general linear model was used to explore the effects of gene-environment and gene-gene interaction on intelligence.

**Results:** In the high fluoride group, children's IQ scores decreased by 2.502 when the UF<sub>Cr</sub> level increased by 1.0 mg/L ( $\beta = -2.502$ , 95% confidence interval [CI]: -4.411, -0.593), and the possibility for having "excellent" intelligence decreased by 46.3% (odds ratio = 0.537, 95% CI: 0.290, 0.994). Children with the GG genotype showed increased IQ scores than those with the AA genotype of rs11627387 locus in the high fluoride group ( $P < 0.05$ ). Interactions between fluoride exposure and *MTHFD1* polymorphisms on intelligence were observed (Pinteraction < 0.05).

**Conclusion:** Our findings suggest that excessive fluoride exposure may have adverse effects on children's intelligence, and changes in children's intelligence may be associated with the interaction between fluoride and *MTHFD1* polymorphisms.

**Keywords:** Fluoride; Intelligence; Interaction; *MTHFD1* gene

## Introduction

Low-dose fluoride can prevent dental caries and is beneficial for bone growth.<sup>[1,2]</sup> However, the health damage caused by excessive fluoride should not be ignored.<sup>[3]</sup> Dental and skeletal fluorosis are the most specific diseases caused by chronic intake of excessive fluoride.<sup>[4,5]</sup> Besides, fluoride exposure is related to the dysfunction of the reproductive system, as well as the liver and brain damage.<sup>[6-8]</sup>

Increasing studies have suggested that excessive fluoride is associated with disorders in cognition, learning, and memory ability.<sup>[9-11]</sup> Among them, a birth cohort study

conducted in Canada suggested that increased fluoride concentrations in drinking water were associated with intellectual impairment in children.<sup>[12]</sup> Another study reported a negative correlation between fluoride exposure and intelligence quotient (IQ) in children,<sup>[13]</sup> which was similar to our previous study.<sup>[14]</sup> In addition, an animal study demonstrated that fluoride exposure during development could induce cognitive deficits in mice.<sup>[15]</sup> However, there was no evidence that reductions in IQ scores were caused by excessive fluoride in a prospective study conducted in a New Zealand community.<sup>[16]</sup> Similarly, a cross-sectional study conducted in China mentioned that there was no statistical difference in IQ

Access this article online	
Quick Response Code:	Website: www.cmj.org
	DOI: 10.1097/CM9.0000000000002062

**Correspondence to:** Yue Ba, Department of Occupational and Environmental Health, School of Public Health, Zhengzhou University, Zhengzhou, Henan 450001, China  
E-Mail: byyue@zzu.edu.cn

Copyright © 2022 The Chinese Medical Association, produced by Wolters Kluwer, Inc. under the CC-BY-NC-ND license. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Chinese Medical Journal 2022;135(15)

Received: 25-11-2021; Online: 18-07-2022 Edited by: Jing Ni

scores between children in different fluorosis areas and children in the control group (CG).<sup>[17]</sup> Behind the different levels of fluoride exposure and assessment methods, different ethnic and genetic backgrounds may also explain the inconsistent results.

Individual susceptibility (eg, gene polymorphisms) is an important factor that affects the sensitivity of body health to environmental factors.<sup>[18,19]</sup> Several studies have reported that genetic polymorphisms can modify the harmful effects of fluoride, such as dental and skeletal fluorosis.<sup>[20,21]</sup> Studies have also shown that gene polymorphisms may modulate sensitivity to the effects of fluoride exposure on intelligence. For example, polymorphisms of the catechol-O-methyltransferase gene and dopamine receptor-2 gene have been reported to modify the adverse effects of fluoride on IQ scores.<sup>[22,23]</sup> Methylenetetrahydrofolate dehydrogenase, cyclohydro-lase, and formyltetrahydrofolate synthetase 1 (*MTHFD1*) gene encode a nicotinamide adenine dinucleotide phosphate-dependent trifunctional enzyme, which provides the one-carbon derivatives of tetrahydrofolate in three sequential reactions and is involved in the folate pathway.<sup>[24]</sup> Comprehension of the influential factors of neural development is incomplete, but folate deficiency has been implicated consistently in neuropathological lesions.<sup>[25]</sup> Previous studies have pointed out that mouse with *MTHFD1* mutations seemingly showed impaired functions of one-carbon metabolism and higher plasma homocysteine levels,<sup>[26,27]</sup> which have been further linked to cognitive impairment and Alzheimer's disease.<sup>[28,29]</sup> In addition, certain loci polymorphisms of *MTHFD1* were associated with neurological diseases in an epidemiological study.<sup>[30]</sup> All these evidences suggest that *MTHFD1* mutations may relate to neurodevelopment. However, few studies have focused on the effects of *MTHFD1* polymorphisms on children's intelligence.

Present study focused on four loci of *MTHFD1* which are related to neurodevelopment: rs11627387, rs1076991, rs2236224, and rs2236225.<sup>[31-33]</sup> Studies have suggested that variations of these four loci can modify the impact of certain factors on health.<sup>[34-36]</sup> However, whether *MTHFD1* polymorphisms are involved in the effect of fluoride exposure on children's intelligence remains unclear.

Based on the above analyses, we conducted a cross-sectional study in an endemic drinking water-borne fluorosis area in Tongxu County, Kaifeng (Henan Province, China). We aimed at evaluating the effects of polymorphisms of *MTHFD1* loci (rs11627387, rs1076991, rs2236224, and rs2236225) and excessive exposure to fluoride on children's intelligence and exploring the potential effect of their interaction on children's intelligence, which further provides a new clue for the study of fluoride neurotoxicity.

## Methods

### Ethical approval

The study protocol (No. ZZUIRB2017-018) was approved by the Ethics Review Board of Zhengzhou University (Zhengzhou, China). Children and their

guardians were fully aware of the aim and process of our research and provided written informed consent.

### Study design and population

As described thoroughly in our previous study,<sup>[14]</sup> we conducted a cross-sectional study in Tongxu County, Henan Province, from April to May in 2017. Four primary schools were randomly selected. We excluded children who were non-local residents, on calcium supplements, had diseases based on calcium or phosphorus metabolism, had digestive diseases, and had thyroid diseases. Subsequently, 694 school-age children aged 8 to 12 years from grades 2 to 6 in four selected schools were recruited by cluster sampling. All children lived on campus and they had similar living conditions, living habits, and dietary structure.

A questionnaire was designed in advance and included information on sociodemographic data, medical history, maternal pregnancy, information on birth, and other information (eg, exercise). The height and weight of children were measured twice and the mean value was taken, which were accurate to 0.1 cm and 0.1 kg, respectively. Then, the body mass index (BMI) was calculated. In addition, fluoride-free containers were used to collect mid-flow morning urine and whole blood from the cubital vein after an overnight fasting. Samples of urine and blood were stored, respectively, at  $-20^{\circ}\text{C}$  and  $-80^{\circ}\text{C}$  for subsequent measurements.

### Exposure assessment

Exposure assessment has been described in detail in our previous study.<sup>[14]</sup> In accordance with the standard detailed by the health industry of China (WS/T 892015), a method based on a fluoride ion-selective electrode (Shanghai Exactitude Instruments, Shanghai, China) was conducted to determine the urinary fluoride (UF) level of children. A creatinine assay kit (Jiancheng Bioengineering Institute, Nanjing, China) was used to determine the concentration of urinary creatinine ( $U_{Cr}$ ). Each determination of levels of UF and  $U_{Cr}$  was undertaken twice and averaged for data analyses. We calculated the urinary creatinine-adjusted urinary fluoride ( $UF_{Cr}$ ) level to correct the influence of urine dilution on the UF level using the following equation:  $UF_{Cr}(\text{mg/L}) = (UF[\text{mg/L}]/U_{Cr}[\text{mg/L}]) \times U_{Cr\text{ mean}}(\text{mg/L})$ , where  $U_{Cr\text{ mean}}$  denotes the mean  $U_{Cr}$  concentration of the total population.<sup>[37]</sup> Then, according to the median value of  $UF_{Cr}$ , children were separated into the high fluoride group (HFG,  $UF_{Cr} > 1.33 \text{ mg/L}$ ) and CG,  $UF_{Cr} \leq 1.33 \text{ mg/L}$ .

### Intelligence assessment

The IQ was assessed using the second revision of the Combined Raven's Test—the Rural in China (CRTRC2).<sup>[38]</sup> Each student completed the paper independently with the supervision of trained investigators. Answer sheets were scored in accordance with the standard of the Combined Raven's Test. The intelligence levels were defined by the IQ scores and classified as “retarded” ( $\leq 69$ ); “marginal” (70–79); “dull normal” (80–89); “normal” (90–109); “highnormal” (110–119); “superior” (120–129) and “excellent” ( $\geq 130$ ). Only 11 children had retarded,

marginal, or dull normal intelligence (IQ score < 90); so, 683 children were finally included and were further separated into four groups (normal, high normal, superior, and excellent) according to their IQ scores in this study.

### Genotyping of gene polymorphisms

Genomic DNA was extracted from whole-blood samples by a genomic DNA miniprep kit (LifeFeng Biotechnology, Shanghai, China). Four single-nucleotide polymorphism (SNP) loci with minor allele frequency > 0.1 were retrieved from Haploview ([www.broadinstitute.org/haploview/haploview](http://www.broadinstitute.org/haploview/haploview)). All of these loci polymorphisms have been reported to be related to neurodevelopmental defects.<sup>[31-33]</sup> rs11627387, rs1076991, rs2236224, and rs2236225 loci were located in intron 26, 2 KB upstream, intron 21, and exon 20, respectively, in *MTHFD1*. All polymorphisms of the four loci were genotyped through a custom-by-design 48-Plex SNPscan™ Kit (catalog number, G0104; Genesky Biotechnologies, Shanghai, China). The kit was developed according to the patented SNP genotyping technology of Genesky Biotechnologies, which was based on double ligation and multiplex fluorescence polymerase chain reaction (PCR).<sup>[39]</sup> Briefly, DNA samples (100–200 ng) were denatured at 98°C for 5 min and then mixed with the premix containing ligase and the probe. The ligation reaction was carried out in a thermal cycler (ABI2720; Applied Biosystems, Foster City, CA, USA). Then, two fluorescent PCRs were undertaken for each ligation product. PCR products were separated and detected by capillary electrophoresis in a sequencer (ABI3730XL; Applied Biosystems). Genotyping was completed according to the obtained information for the labeling dye color and fragment size of allele-specific ligation PCR products. About 4% of the genotyping was done repeatedly, and the consistency rate was > 96%.

### Statistical analyses

A database was set up by Epidata 3.0 (Epidata Association, Odense, Denmark) in which two operators independently imported all data. Mean ± standard deviation and number (%) are presented for continuous and categorical variables, respectively.

Differences in continuous data between the two groups were compared using the Student's *t*-test or Mann-Whitney U-test. The distribution of categorical variables was compared using the Chi-squared test. Potential confounders (children's age, gender, BMI, age at which pregnancy occurred, gestational weeks, birth weight, birth modes, paternal and maternal education level) were chosen as adjustment variables based on existing literature and population characteristics of this study.<sup>[13]</sup> The generalized linear model (GLM) was used to analyze the association between children's UF<sub>Cr</sub> level, *MTHFD1* polymorphisms, and IQ scores. The multinomial logistic regression model was applied to analyze the relationship between children's UF<sub>Cr</sub> level, *MTHFD1* polymorphisms, and intelligence levels. And normal intelligence children were reference in the analyses of the intelligence levels. For exploration of the relationship between the UF<sub>Cr</sub> level and children's intelligence, the UF<sub>Cr</sub> level was separated into categorical variables according to tertiles, and the median

of each segment was regarded as a continuous variable to estimate the linear trend of children's intelligence. The general linear model was used to explore models of possible gene-environment and gene-gene interaction on intelligence. Data were processed using SPSS 21.0 (IBM, Armonk, NY, USA). Plots were drawn by GraphPad Prism 8.0.1. *P* < 0.05 was considered significant.

## Results

### General characteristics of participants

Since 11 children had an IQ score < 90, a total of 683 eligible children aged 8 to 12 years were included in this study and were further classified as the CG (*n* = 342) and HFG (*n* = 341) according to the median of children's UF<sub>Cr</sub> level (1.33 mg/L). The distribution of children's age was consistent in the CG and HFG (10.05 ± 1.24 and 10.08 ± 1.23 years, respectively). The concentration of UF<sub>Cr</sub> in the HFG (2.15 ± 0.91 mg/L) was significantly higher than that in the CG (0.83 ± 0.30 mg/L) (*P* < 0.001), whereas the distribution of other sociodemographic characteristics (except U<sub>Cr</sub> and UF levels) presented no significant differences between the two groups (*P* > 0.05 for all) [Table 1].

### Association between children's UF<sub>Cr</sub> level and intelligence

The GLM and multinomial logistic regression model were employed to evaluate if there were associations between the UF<sub>Cr</sub> level and children's IQ scores or intelligence levels, respectively [Figure 1]. For each increase of 1.0 mg/L in the UF<sub>Cr</sub> level, children's IQ scores decreased by 2.502 ( $\beta = -2.502$ , 95% confidence interval [CI]: -4.411, -0.593, *P* = 0.010), and the possibility of developing "excellent" intelligence decreased by 46.3% with reference to the normal intelligence children in the HFG (odds ratio [OR] = 0.537, 95% CI: 0.290, 0.994, *P* = 0.048). After stratifying children according to the tertiles of children's UF<sub>Cr</sub> concentration in different groups, the trend test showed no significance (*P*<sub>trend</sub> > 0.05 for all).

### Association between *MTHFD1* polymorphisms and intelligence


The genotype distributions of rs11627387, rs1076991, rs2236224, and rs2236225 loci in *MTHFD1* were in accordance with the Hardy-Weinberg equilibrium (*P* > 0.05 for all loci) [Supplementary Table 1, <http://links.lww.com/CM9/A987>], which indicated that the investigated participants were representative of the population. We also estimated the difference in the distribution of genotypes/allele of rs11627387, rs1076991, rs2236224, and rs2236225 between the CG and HFG, but significant differences were not found (*P* > 0.05 for all) [Supplementary Table 2, <http://links.lww.com/CM9/A987>].

Furthermore, the relationship between *MTHFD1* polymorphisms and intelligence was assessed by the GLM and multinomial logistic regression model. In the general population and HFG, children with the GG genotype of rs11627387 showed increased IQ score relative to those with the AA genotype of rs11627387 ( $\beta = 3.574$ , 95% CI:

**Table 1: Demographic data of the study population.**

Variables	Total (n = 683)	CG (n = 342)	HFG (n = 341)	Statistics	P values
Age (years)	10.07 ± 1.24	10.05 ± 1.24	10.08 ± 1.23	0.375*	0.708
Gender				0.333 <sup>†</sup>	0.564
Boys	324 (47.44)	166 (48.54)	158 (46.33)		
Girls	359 (52.56)	176 (51.46)	183 (53.67)		
BMI (kg/m <sup>2</sup> )	17.50 ± 2.96	17.68 ± 2.96	17.33 ± 2.95	-1.618*	0.106
U <sub>Cr</sub> (mg/L)	1089 ± 607	1277 ± 606	901 ± 548	-8.261*	<0.001
UF (mg/L)	1.27 ± 0.79	0.98 ± 0.62	1.56 ± 0.82	9.811*	<0.001
UF <sub>Cr</sub> (mg/L)	1.49 ± 0.95	0.83 ± 0.30	2.15 ± 0.91	22.616*	<0.001
Age at which pregnancy occurred (years)	25.82 ± 4.28	25.95 ± 4.52	25.70 ± 4.01	-0.750*	0.453
Gestational weeks (weeks)	36.87 ± 4.84	36.53 ± 5.31	37.22 ± 4.30	1.742*	0.081
Birth weight (kg)	3.34 ± 0.52	3.31 ± 0.55	3.36 ± 0.48	0.987*	0.324
Birth modes				0.280 <sup>†</sup>	0.869
Natural birth	467 (68.37)	232 (67.84)	235 (68.91)		
Cesarean delivery	209 (30.60)	107 (31.29)	102 (29.91)		
Rest	7 (1.02)	3 (0.88)	4 (1.17)		
Paternal education				0.497 <sup>†</sup>	0.780
Primary school and below	57 (8.35)	31 (9.06)	26 (7.62)		
Middle school	463 (67.79)	229 (66.96)	234 (68.62)		
High school and above	163 (23.87)	82 (23.98)	81 (23.75)		
Maternal education				0.457 <sup>†</sup>	0.796
Primary school and below	87 (12.74)	46 (13.45)	41 (12.02)		
Middle school	453 (66.33)	223 (65.20)	230 (67.45)		
High school and above	143 (20.94)	73 (21.35)	70 (20.53)		
IQ scores	122.05 ± 11.88	121.50 ± 12.14	122.61 ± 11.61	1.059*	0.290
Intelligence levels				2.162 <sup>†</sup>	0.539
90–109	112 (16.40)	60 (17.54)	52 (15.25)		
110–119	173 (25.33)	85 (24.85)	88 (25.81)		
120–129	216 (31.63)	113 (33.04)	103 (30.21)		
≥130	182 (26.65)	84 (24.56)	98 (28.74)		

Data are presented as mean ± SD or n (%). \* Mann–Whitney U-test. <sup>†</sup>  $\chi^2$  test. BMI: Body mass index; CG: Control group; HFG: High fluoride group; IQ: Intelligence quotient; SD: Standard deviation; U<sub>Cr</sub>: Urinary creatinine; UF: Urinary fluoride; UF<sub>Cr</sub>: Urinary creatinine-adjusted urinary fluoride.



**Figure 1:** Relationship between fluoride exposure and children's intelligence. Analyses were adjusted for age, gender, BMI, age at which pregnancy occurred, gestational weeks, birth modes, birth weight, and paternal and maternal education level. Normal intelligence children were reference to the analyses of the intelligence levels. BMI: Body mass index; CG: Control group; CI: Confidence interval; HFG: High fluoride group; IQ: Intelligence quotient; OR: Odds ratio; UF<sub>Cr</sub>: Urinary creatinine-adjusted urinary fluoride.

0.274, 6.874,  $P=0.034$  for the general population,  $\beta=4.723$ , 95% CI: 0.277, 9.168,  $P=0.037$  for the HFG) [Figure 2]. There was an increment in IQ score in children carrying the *G* allele than in those with the *A* allele of rs11627387 in the HFG, and the association was borderline significant ( $P=0.059$ ) [Figure 2]. In addition, the possibility of developing “high normal” intelligence was lower in the HFG in children with the *AG* genotype when compared with those carrying the *AA* genotype of rs11627387 (OR = 0.212, 95% CI: 0.045, 0.997,  $P=0.049$ ) [Figure 3]. With respect to rs2236225 locus, participants with the *AA* genotype seemingly showed a lower possibility of developing “high normal” intelligence when compared with children with the *GG* genotype in the HFG, and the association was borderline significant ( $P=0.056$ ) [Figure 3]. However, a statistical significance was not found in the association between *MTHFD1* polymorphism and children’s intelligence levels in the total group and CG ( $P > 0.050$ ) [Supplementary Tables 3 and 4, <http://links.lww.com/CM9/A987>].


**Effects of gene-gene and gene-environment interaction on children’s intelligence**

The general linear model was used to explore gene-environment and gene-gene interactions. Loci rs11627387,

rs1076991, and rs2236225 may have interactive effects on the IQ scores according to analyses of gene-gene interaction [Figure 4], where the association showed marginal significance ( $F=1.726$ ,  $P=0.059$ ). In addition, the interaction between rs11627387, rs1076991, rs2236224, and the  $UF_{Cr}$  level might affect children’s IQ scores ( $F=1.669$ ,  $P=0.021$ ), as well as the interactive effects of loci rs11627387, rs1076991, rs2236225, and the  $UF_{Cr}$  level ( $F=1.764$ ,  $P=0.012$ ). An effect of the interaction between these four loci and the  $UF_{Cr}$  level on the IQ scores of children was also found ( $F=1.614$ ,  $P=0.012$ ). However, statistical significances were not found in other models [Supplementary Table 5, <http://links.lww.com/CM9/A987>].

**Discussion**


We evaluated the interaction between fluoride exposure and *MTHFD1* polymorphisms on the intelligence of children living in areas of endemic fluorosis. We found that the decline in the IQ scores in children was associated with excessive exposure to fluoride and that changes in children’s intelligence might be modified by rs11627387 locus polymorphisms of *MTHFD1* to some extent. Moreover, the interactive effects of the four loci of *MTHFD1* and fluoride exposure on children’s IQ scores showed different models in our study.



**Figure 2:** Regression analyses of *MTHFD1* polymorphisms and the IQ scores. Analyses were adjusted for age, gender, BMI, age at which pregnancy occurred, gestational weeks, birth modes, birth weight, and paternal and maternal education level. Twenty-two participants were not genotyped successfully for rs11627387 and rs2236225 respectively, and 23 participants were not genotyped successfully for rs1076991 and rs2236224 respectively. BMI: Body mass index; CG: Control group; CI: Confidence interval; HFG: High fluoride group; IQ: Intelligence quotient; *MTHFD1*: Methylenetetrahydrofolate dehydrogenase, cyclohydrolase, and formyltetrahydrofolate synthetase 1.

Loci	Genotypes / Allele	Intelligence levels					
		High normal (n = 88)		Superior (n = 103)		Excellent (n = 98)	
		OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
<b>HFG (n = 341)</b> rs11627387	AA	Reference		Reference		Reference	
	AG		0.049		0.357		0.697
	GG		0.449		0.848		0.573
	AG+GG		0.109		0.543		0.949
	A	Reference		Reference		Reference	
rs1076991	G		0.666		0.524		0.328
	TT	Reference		Reference		Reference	
	TC		0.500		0.792		0.814
	CC		0.850		0.832		0.233
	TC+CC		0.711		0.799		0.460
rs2236224	T	Reference		Reference		Reference	
	C		0.835		0.858		0.277
	GG	Reference		Reference		Reference	
	GA		0.140		0.457		1.000
	AA		0.164		0.610		0.549
rs2236225	GA+AA		0.069		0.276		0.771
	G	Reference		Reference		Reference	
	A		0.063		0.247		0.687
	GG	Reference		Reference		Reference	
	GA		0.721		0.559		0.926
	AA		0.056		0.266		0.531
	GA+AA		0.312		0.325		0.961
	G	Reference		Reference		Reference	
	A		0.100		0.167		0.820

**Figure 3:** Regression analyses of *MTHFD1* polymorphisms and intelligence levels in the HFG. Analyses were adjusted for age, gender, BMI, age at which pregnancy occurred, gestational weeks, birth modes, birth weight, and paternal and maternal education level. Normal intelligence children were referenced (n = 52). BMI: Body mass index; CI: Confidence interval; HFG: High fluoride group; *MTHFD1*: Methylenetetrahydrofolate dehydrogenase, cyclohydrolase, and formyltetrahydrofolate synthetase 1; OR: Odds ratio.



**Figure 4:** Interaction analyses of fluoride exposure and *MTHFD1* polymorphisms. Analyses were adjusted for age, gender, BMI, age at which pregnancy occurred, gestational weeks, birth modes, birth weight, and paternal and maternal education level. Dashed line denotes the P value of 0.05 for significance. BMI: Body mass index; Locus 1: rs11627387; Locus 2: rs1076991; Locus 3: rs2236224; Locus 4: rs2236225; *MTHFD1*: Methylenetetrahydrofolate dehydrogenase, cyclohydrolase, and formyltetrahydrofolate synthetase 1; UF<sub>Cr</sub>: Urinary creatinine-adjusted urinary fluoride.

The health and function of the nervous system can be affected by exposure to various environmental factors.<sup>[40,41]</sup>

In terms of fluoride, drinking water containing a high concentration of fluoride is the main way to the exposure of superabundant fluoride.<sup>[42]</sup> Fluoride entering the body can be distributed widely throughout the body after absorption, and most of it is deposited in bone and teeth.<sup>[43]</sup> The absorption rate of fluoride in children is about 80% to 90%, much higher than that in adults.<sup>[44]</sup> In addition, fluoride can penetrate the blood-brain barrier into brain tissue and seems to accumulate in the areas of the brain responsible for learning and memory functions.<sup>[45,46]</sup> But the evidence regarding a link between fluoride exposure and intelligence impairment is not definitive. Most of the scholars conceive that fluoride in brain tissue can damage nerve functions and even lead to intellectual loss.<sup>[47-49]</sup> For example, animal studies found that, upon exposure to increasing concentrations of sodium fluoride in drinking water, the learning abilities of mice were impaired.<sup>[10,50]</sup> However, the results of population-based epidemiological studies are not completely consistent. A cohort study from Canada revealed a positive association between fluoride exposure during pregnancy and intelligence decline in offspring.<sup>[51]</sup>

Our current study support this finding, that is, an inverse association between excessive exposure to fluoride and children's IQ scores in the HFG, which recollects that excessive fluoride exposure may have negative influence on normal development of children's intelligence. Whereas, a prospective study conducted in New Zealand did not find a positive association between lower intelligence and a higher fluoride level.<sup>[16]</sup> A cross-sectional study conducted in China did not observe the significant difference in IQ scores between children in fluorosis areas and children in the CG, either.<sup>[17]</sup> Differences in study design, levels of fluoride exposure, exposure patterns, ethnicity, and assessment methods can explain these inconsistencies to a certain extent. On the other hand, genetic susceptibility (eg, genetic polymorphisms) is also one of the important reasons that is worthy of further discussion.

We further discussed the association between *MTHFD1* polymorphisms and children's intelligence according the study design. We found that, compared with children carrying the AA genotype of the rs11627387 locus, children carrying the GG genotype might have increased IQ score in the HFG, whereas the possibility for having "high normal" intelligence was lower for children carrying the AG genotype. It can be suggested that polymorphisms of the rs11627387 locus of *MTHFD1* may have effects on changes in children's intelligence. Specifically, the GG genotype of rs11627387 may retard the intelligence decline caused by excessive fluoride exposure in schoolage children, as carrying the GG genotype of rs11627387 may exert a positive effect on IQ scores compared to carrying the AA genotype when UFCr >1.33 mg/L in children. Although few studies have focused on the relationship between *MTHFD1* polymorphisms and intelligence, others are still evaluating the effects of *MTHFD1* polymorphisms on neural development based on animals and humans. An animal study reported that mice with loss of one allele of *MTHFD1* via a gene-trap mutation showed impaired learning ability.<sup>[52]</sup> Another study revealed that reduced gene expression of the alpha seven nicotinic cholinergic receptor was observed in a mouse model which simulated polymorphisms at the rs2236225 locus of *MTHFD1* in humans.<sup>[53]</sup> Also, agonists of the alpha seven nicotinic cholinergic receptor gene could be used to treat neurocognitive dysfunction in schizophrenia.<sup>[54]</sup> In addition, population-based studies also pointed out that polymorphisms of rs11627387, rs1076991, rs2236224, and rs2236225 loci are involved in neural tube defects.<sup>[31,33]</sup> These studies demonstrate that *MTHFD1* polymorphisms can have effects on normal neurocognitive functions and may modify the susceptibility of neurological diseases. Overall, our results support the correlation between *MTHFD1* polymorphisms and intellectual changes in school-age children, which is worthy of confirmation in further studies.

Intelligence is not only modified by genetic factors such as gene polymorphisms but also related to gene-environment interaction. For example, Zhao *et al*<sup>[55]</sup> conducted a cross-sectional study in Tianjin and reported that ankyrin repeat and kinase domain 1 (ANKK1), catechol-O-methyltransferase (COMT), monoamine oxidase A

(MAOA) gene polymorphisms may have interactive effects with UF on children's intelligence. In our study, the effects of gene-environment and gene-gene interaction on children's IQ were explored. We did not observe an interaction between a single locus of *MTHFD1* and fluoride exposure on children's intelligence but found the different models of interaction between multiple loci of *MTHFD1* and fluoride exposure. These results suggest that phenotypes can be affected by the interaction of environmental factors with multiple loci of one gene and even multiple genes<sup>[56]</sup>; the modification effect of one locus polymorphism is minor. All these findings suggest that *MTHFD1* polymorphisms may be involved in the effects of fluoride exposure on intelligence in school-age children, and we further provide a novel clue for the study of neurotoxicological mechanisms of fluoride.

There are several advantages in this study. First, the study was a population-based epidemiological study which explored the impact of fluoride and gene polymorphisms on children's intelligence, where the results suggest that people living in fluorosis areas should pay attention to the harmful effects of excessive fluoride intake on school-age children's intelligence. Second, this study was conducted in the middle of the semester which avoided the impact of students' psychological stress caused by the beginning of the semester or the final exam on intelligence tests. Third, as described in the study design in our previous publication,<sup>[14]</sup> all the children lived on campus and they had similar living conditions, living habits, and dietary structure, which minimized the bias. Finally, the investigated areas in Tongxu County, Henan Province, are relatively underdeveloped, with no industrial fluorine and other pollutants such as lead and mercury, etc, that may affect intelligence.

Our study also had some limitations. First, we did not adjust the children's diet, although children included in the study were all boarding school students, and the dietary structure was relatively consistent; the differences in dietary habits and physical fitness of children may lead to differences in nutrients intake, which may further affect mental development. Therefore, children's diet will be considered in our subsequent studies. Second, the study is a cross-sectional study with only one sampling; so, the causal relationship is weak. So, long-term large-scale epidemiological or cohort studies should be conducted to provide more evidence. Third, although we adjusted for confounding factors such as children's age, gender, BMI, maternal age at which pregnancy occurred, gestational weeks, birth weight, birth modes, and paternal and maternal education level, there may still be some other confounding factors (such as folic acid). However, bounded to the eugenics policy, almost all the mothers of the investigated students had taken folic acid in the early stage of pregnancy<sup>[57]</sup>; so, the bias may be reduced. Studies involving larger populations and more questionnaire information are in progress.

## Conclusions

We evaluated the effects of polymorphisms of *MTHFD1* and fluoride exposure on children's intelligence in endemic

fluorosis areas. Excessive fluoride exposure may have adverse effects on children's intelligence, and changes in children's intelligence may be associated with the interaction between fluoride and *MTHFD1* polymorphisms.

### Acknowledgements

The authors would like to express their sincere thanks to all the staff and volunteers involved in the study. The authors also thank Francis-Kojo Afrim for his English editing help.

### Funding

This study was supported by the National Natural Science Foundation of China (Nos. 81972981, 82003401, and 81673116) and Key Projects of Colleges and Universities of Henan Education Department (21A330006).

### Conflicts of interest

None.

### References

- Whelton HP, Spencer AJ, Do LG, Rugg-Gunn AJ. Fluoride revolution and dental caries: evolution of policies for global use. *J Dent Res* 2019;98:837–846. doi: 10.1177/0022034519843495.
- Riedel C, Zimmermann EA, Zustin J, Niecke M, Amling M, Grynepas M, *et al.* The incorporation of fluoride and strontium in hydroxyapatite affects the composition, structure, and mechanical properties of human cortical bone. *J Biomed Mater Res A* 2017;105:433–442. doi: 10.1002/jbm.a.35917.
- Guth S, Huser S, Roth A, Degen G, Diel P, Edlund K, *et al.* Toxicity of fluoride: critical evaluation of evidence for human developmental neurotoxicity in epidemiological studies, animal experiments and in vitro analyses. *Arch Toxicol* 2020;94:1375–1415. doi: 10.1007/s00204-020-02725-2.
- Wei W, Pang S, Sun D. The pathogenesis of endemic fluorosis: research progress in the last 5 years. *J Cell Mol Med* 2019;23:2333–2342. doi: 10.1111/jcmm.14185.
- Yuan L, Fei W, Jia F, Jun-Ping L, Qi L, Fang-Ru N, *et al.* Health risk in children to fluoride exposure in a typical endemic fluorosis area on Loess Plateau, North China, in the last decade. *Chemosphere* 2020;243:125451. doi: 10.1016/j.chemosphere.2019.125451.
- Wang HW, Liu J, Wei SS, Zhao WP, Zhu SQ, Zhou BH. Mitochondrial respiratory chain damage and mitochondrial fusion disorder are involved in liver dysfunction of fluoride-induced mice. *Chemosphere* 2020;241:125099. doi: 10.1016/j.chemosphere.2019.125099.
- Jaiswal P, Mandal M, Mishra A. Effect of hesperidin on fluoride-induced neurobehavioral and biochemical changes in rats. *J Biochem Mol Toxicol* 2020;34:e22575. doi: 10.1002/jbt.22575.
- Ma Q, Huang H, Sun L, Zhou T, Zhu J, Cheng X, *et al.* Gene-environment interaction: does fluoride influence the reproductive hormones in male farmers modified by ER $\alpha$  gene polymorphisms? *Chemosphere* 2017;188:525–531. doi: 10.1016/j.chemosphere.2017.08.166.
- Grandjean P. Developmental fluoride neurotoxicity: an updated review. *Environ Health* 2019;18:110. doi: 10.1186/s12940-019-0551-x.
- Yuan J, Li Q, Niu R, Wang J. Fluoride exposure decreased learning ability and the expressions of the insulin receptor in male mouse hippocampus and olfactory bulb. *Chemosphere* 2019;224:71–76. doi: 10.1016/j.chemosphere.2019.02.113.
- Xin J, Wang H, Sun N, Bughio S, Zeng D, Li L, *et al.* Probiotic alleviate fluoride-induced memory impairment by reconstructing gut microbiota in mice. *Ecotoxicol Environ Saf* 2021;215:112108. doi: 10.1016/j.ecoenv.2021.112108.
- Till C, Green R, Flora D, Hornung R, Martinez-Mier EA, Blazer M, *et al.* Fluoride exposure from infant formula and child IQ in a Canadian birth cohort. *Environ Int* 2020;134:105315. doi: 10.1016/j.envint.2019.105315.
- Wang M, Liu L, Li H, Li Y, Liu H, Hou C, *et al.* Thyroid function, intelligence, and low-moderate fluoride exposure among Chinese school-age children. *Environ Int* 2020;134:105229. doi: 10.1016/j.envint.2019.105229.
- Xu K, An N, Huang H, Duan L, Ma J, Ding J, *et al.* Fluoride exposure and intelligence in school-age children: evidence from different windows of exposure susceptibility. *BMC Public Health* 2020;20:1657. doi: 10.1186/s12889-020-09765-4.
- Ge Y, Chen L, Yin Z, Song X, Ruan T, Hua L, *et al.* Fluoride-induced alterations of synapse-related proteins in the cerebral cortex of ICR offspring mouse brain. *Chemosphere* 2018;201:874–883. doi: 10.1016/j.chemosphere.2018.02.167.
- Broadbent JM, Thomson WM, Ramrakha S, Moffitt TE, Zeng J, Page LAF, *et al.* Community water fluoridation and intelligence: prospective study in New Zealand. *Am J Public Health* 2015;105:72–76. doi: 10.2105/AJPH.2013.301857.
- Li FH, Chen X, Huang RJ, Xie YP. Intelligence impact of children with endemic fluorosis caused by fluoride from coal burning (in Chinese). *J Environ Health* 2009;26:338–340. doi: 10.16241/j.cnki.1001-5914.2009.04.040.
- Kang X, Guo T, Liu L, Ding SZ, Lei C, Luo H. Association between PTC1 gene polymorphisms and chronic obstructive pulmonary disease susceptibility in a Chinese Han population: a case-control study. *Chin Med J* 2020;133:2071–2077. doi: 10.1097/CM9.0000000000000858.
- Wen D, Zhou XL, Du X, Dong JZ, Ma CS. Association of interleukin-18 gene polymorphisms with Takayasu arteritis in a Chinese Han population. *Chin Med J* 2020;133:2315–2320. doi: 10.1097/CM9.0000000000001047.
- Abbasoglu Z, Dalledone M, Wambier LM, Pecharki G, Baratto-Filho F, Andrades KMR, *et al.* Single nucleotide polymorphism rs4284505 in microRNA17 and risk of dental fluorosis. *Acta Odontol Scand* 2020;78:463–466. doi: 10.1080/00016357.2020.1786600.
- Chu Y, Liu Y, Guo N, Lou Q, Wang L, Huang W, *et al.* Association between ALOX15 gene polymorphism and brick-tea type skeletal fluorosis in Tibetans, Kazaks and Han, China. *Int J Environ Health Res* 2021;31:421–432. doi: 10.1080/09603123.2019.1666972.
- Zhang S, Zhang X, Liu H, Qu W, Guan Z, Zeng Q, *et al.* Modifying effect of COMT gene polymorphism and a predictive role for proteomics analysis in children's intelligence in endemic fluorosis area in Tianjin, China. *Toxicol Sci* 2015;144:238–245. doi: 10.1093/toxsci/kfu311.
- Cui Y, Zhang B, Ma J, Wang Y, Zhao L, Hou C, *et al.* Dopamine receptor D2 gene polymorphism, urine fluoride, and intelligence impairment of children in China: a school-based cross-sectional study. *Ecotoxicol Environ Saf* 2018;165:270–277. doi: 10.1016/j.ecoenv.2018.09.018.
- Sutherland HG, Hermile H, Sanche R, Menon S, Lea RA, Haupt LM, *et al.* Association study of MTHFD1 coding polymorphisms R134K and R653Q with migraine susceptibility. *Headache* 2014;54:1506–1514. doi: 10.1111/head.12428.
- Greene NDE, Copp AJ. Neural tube defects. *Annu Rev Neurosci* 2014;37:221–242. doi: 10.1146/annurev-neuro-062012-170354.
- Field MS, Shields KS, Abarinov EV, Malysheva OV, Allen RH, Stabler SP, *et al.* Reduced MTHFD1 activity in male mice perturbs folate- and choline-dependent one-carbon metabolism as well as transsulfuration. *J Nutr* 2013;143:41–45. doi: 10.3945/jn.112.169821.
- MacFarlane AJ, Perry CA, Girnary HH, Gao D, Allen RH, Stabler SP, *et al.* Mthfd1 is an essential gene in mice and alters biomarkers of impaired one-carbon metabolism. *J Biol Chem* 2009;284:1533–1539. doi: 10.1074/jbc.M808281200.
- Smith AD, Refsum H, Homocysteine . B vitamins, and cognitive impairment. *Annu Rev Nutr* 2016;36:211–239. doi: 10.1146/annurev-nutr-071715-050947.
- Dayon L, Guiraud SP, Corthesy J, Da Silva L, Migliavacca E, Tautvydaite D, *et al.* One-carbon metabolism, cognitive impairment and CSF measures of Alzheimer pathology: homocysteine and beyond. *Alzheimers Res Ther* 2017;9:43. doi: 10.1186/s13195-017-0270-x.
- Wu J, Bao Y, Lu X, Wu L, Zhang T, Guo J, *et al.* Polymorphisms in MTHFD1 gene and susceptibility to neural tube defects: a casecontrol study in a Chinese Han population with relatively low folate levels. *Med Sci Monit* 2015;21:2630–2637. doi: 10.12659/MSM.895155.



31. Etheredge AJ, Finnell RH, Carmichael SL, Lammer EJ, Zhu H, Mitchell LE, *et al.* Maternal and infant gene-folate interactions and the risk of neural tube defects. *Am J Med Genet A* 2012;158A:2439–2446. doi: 10.1002/ajmg.a.35552.
32. Meng J, Han L, Zhuang B. Association between MTHFD1 polymorphisms and neural tube defect susceptibility. *J Neurol Sci* 2015;348:188–194. doi: 10.1016/j.jns.2014.12.001.
33. Carroll N, Pangilinan F, Molloy AM, Troendle J, Mills JL, Kirke PN, *et al.* Analysis of the MTHFD1 promoter and risk of neural tube defects. *Hum Genet* 2009;125:247–256. doi: 10.1007/s00439-008-0616-3.
34. Liu AY, Scherer D, Poole E, Potter JD, Curtin K, Makar K, *et al.* Gene-diet-interactions in folate-mediated one-carbon metabolism modify colon cancer risk. *Mol Nutr Food Res* 2013;57:721–734. doi: 10.1002/mnfr.201200180.
35. Shaw GM, Lu W, Zhu H, Yang W, Briggs FBS, Carmichael SL, *et al.* 118 SNPs of folate-related genes and risks of spina bifida and conotruncal heart defects. *BMC Med Genet* 2009;10:49. doi: 10.1186/1471-2350-10-49.
36. Zhu H, Yang W, Lu W, Etheredge AJ, Lammer EJ, Finnell RH, *et al.* Gene variants in the folate-mediated one-carbon metabolism (FOCM) pathway as risk factors for conotruncal heart defects. *Am J Med Genet A* 2012;158A:1124–1134. doi: 10.1002/ajmg.a.35313.
37. Thomas DB, Basu N, Martinez-Mier EA, Sanchez BN, Zhang Z, Liu Y, *et al.* Urinary and plasma fluoride levels in pregnant women from Mexico City. *Environ Res* 2016;150:489–495. doi: 10.1016/j.envres.2016.06.046.
38. Liu HL, Lam LT, Zeng Q, Han SQ, Fu G, Hou CC. Effects of drinking water with high iodine concentration on the intelligence of children in Tianjin, China. *J Public Health (Oxf)* 2009;31:32–38. doi: 10.1093/pubmed/fdn097.
39. Chen X, Li S, Yang Y, Yang X, Liu Y, Liu Y, *et al.* Genome-wide association study validation identifies novel loci for atherosclerotic cardiovascular disease. *J Thromb Haemost* 2012;10:1508–1514. doi: 10.1111/j.1538-7836.2012.04815.x.
40. Liu WT, Yan XX, Cheng DZ, Zhang HZ, Ding N, Xu KM, *et al.* Oxcarbazepine monotherapy in children with benign epilepsy with centrotemporal spikes improves quality of life. *Chin Med J* 2020;133:1649–1654. doi: 10.1097/CM9.0000000000000925.
41. Tian DY, Wang J, Sun BL, Wang Z, Xu W, Chen Y, *et al.* Spicy food consumption is associated with cognition and cerebrospinal fluid biomarkers of Alzheimer disease. *Chin Med J* 2020;134:173–177. doi: 10.1097/CM9.0000000000001318.
42. Li P, Qian H, Wu J, Chen J, Zhang Y, Zhang H. Occurrence and hydrogeochemistry of fluoride in alluvial aquifer of Weihe River, China. *Environ Earth Sci* 2013;71:3133–3145. doi: 10.1007/s12665-013-2691-6.
43. Chakraborti D, Rahman MM, Chatterjee A, Das D, Das B, Nayak B, *et al.* Fate of over 480 million inhabitants living in arsenic and fluoride endemic Indian districts: magnitude, health, socio-economic effects and mitigation approaches. *J Trace Elem Med Biol* 2016;38:33–45. doi: 10.1016/j.jtemb.2016.05.001.
44. O'Mullane DM, Baez RJ, Jones S, Lennon MA, Petersen PE, Rugg-Gunn AJ, *et al.* Fluoride and oral health. *Commun Dent Health* 2016;33:69–99. <https://eprints.whiterose.ac.uk/101379/>.
45. Gori S, Inno A, Lunardi G, Gorgoni G, Malfatti V, Severi F, *et al.* 18F-Sodium fluoride PET-CT for the assessment of brain metastasis from lung adenocarcinoma. *J Thorac Oncol* 2015;10:e67–e68. doi: 10.1097/JTO.0000000000000523.
46. Pereira M, Dombrowski PA, Lusso EM, Chioca LR, Da Cunha C, Andreatini R. Memory impairment induced by sodium fluoride is associated with changes in brain monoamine levels. *Neurotox Res* 2011;19:55–62. doi: 10.1007/s12640-009-9139-5.
47. Zhou G, Tang S, Yang L, Niu Q, Chen J, Xia T, *et al.* Effects of long-term fluoride exposure on cognitive ability and the underlying mechanisms: role of autophagy and its association with apoptosis. *Toxicol Appl Pharmacol* 2019;378:114608. doi: 10.1016/j.taap.2019.114608.
48. Ding Y, Gao Y, Sun H, Han H, Wang W, Ji X, *et al.* The relationships between low levels of urine fluoride on children's intelligence, dental fluorosis in endemic fluorosis areas in Hulunbuir, Inner Mongolia, China. *J Hazard Mater* 2011;186:1942–1946. doi: 10.1016/j.jhazmat.2010.12.097.
49. Sun Z, Zhang Y, Xue X, Niu R, Wang J. Maternal fluoride exposure during gestation and lactation decreased learning and memory ability, and glutamate receptor mRNA expressions of mouse pups. *Hum Exp Toxicol* 2018;37:87–93. doi: 10.1177/0960327117693067.
50. Wang J, Zhang Y, Guo Z, Li R, Xue X, Sun Z, *et al.* Effects of perinatal fluoride exposure on the expressions of miR-124 and miR-132 in hippocampus of mouse pups. *Chemosphere* 2018;197:117–122. doi: 10.1016/j.chemosphere.2018.01.029.
51. Green R, Lanphear B, Hornung R, Flora D, Martinez-Mier EA, Neufeld R, *et al.* Association between maternal fluoride exposure during pregnancy and IQ scores in offspring in Canada. *JAMA Pediatr* 2019;173:940–948. doi: 10.1001/jamapediatrics.2019.1729.
52. Pjetri E, Zeisel SH. Deletion of one allele of Mthfd1 (methylene-tetrahydrofolate dehydrogenase 1) impairs learning in mice. *Behav Brain Res* 2017;332:71–74. doi: 10.1016/j.bbr.2017.05.051.
53. Ash JA, Jiang X, Malysheva OV, Fiorenza CG, Bisogni AJ, Levitsky DA, *et al.* Dietary and genetic manipulations of folate metabolism differentially affect neocortical functions in mice. *Neurotoxicol Teratol* 2013;38:79–91. doi: 10.1016/j.ntt.2013.05.002.
54. Freedman R. (7-nicotinic acetylcholine receptor agonists for cognitive enhancement in schizophrenia. *Annu Rev Med* 2014;65:245–261. doi: 10.1146/annurev-med-092112-142937.
55. Zhao L, Yu C, Lv J, Cui Y, Wang Y, Hou C, *et al.* Fluoride exposure, dopamine relative gene polymorphism and intelligence: a cross-sectional study in China. *Ecotoxicol Environ Saf* 2020;209:111826. doi: 10.1016/j.ecoenv.2020.111826.
56. Rapin I, Tuchman RF. What is new in autism? *Curr Opin Neurol* 2008;21:143–149. doi: 10.1097/WCO.0b013e3282f49579.
57. Wang A, Duan L, Huang H, Ma J, Zhang Y, Ma Q, *et al.* Association between fluoride exposure and behavioural outcomes of school-age children: a pilot study in China. *Int J Environ Health Res* 2020;1–10. doi: 10.1080/09603123.2020.1747601.

---

**How to cite this article:** Feng Z, An N, Yu F, Ma J, Li N, Du Y, Guo M, Xu K, Hou X, Li Z, Zhou G, Ba Y. Do methylenetetrahydrofolate dehydrogenase, cyclohydrolase, and formyltetrahydrofolate synthetase 1 polymorphisms modify changes in intelligence of school-age children in areas of endemic fluorosis?. *Chin Med J* 2022;135:1846–1854. doi: 10.1097/CM9.0000000000002062