

Development and Application of Global Health Events-Mental Stress Scale for Assessment of Medical Staff's Acute Mental Stress Responses

Xin-Ying Cai¹, Shao-Yan Zheng², Zhen-Su Lin², Su-Zhi Chen², Wei-Yi Zhu², Jia-Jia Huang², Ze-Li Zheng², Yu-Hua Zhou²

¹Clinical Research Center, Shantou Central Hospital, Shantou City, People's Republic of China; ²Nursing Department, Shantou Central Hospital, Shantou City, People's Republic of China

Correspondence: Yu-Hua Zhou, Nursing Department, Shantou Central Hospital, Wai-ma Road 114, Shantou City, People's Republic of China, Tel +86 0754-88903584, Fax +86 0754-88548117, Email zhouyuhua1023@163.com

Background: Medical workers have been increasingly involved in emergent public health events, which can lead to severe stress. However, no standardized, officially recognized, unified tool exists for mental distress measurement in medical workers who experienced the public health events.

Purpose: In the present study, we propose the Global Health Events-Mental Stress Scale (GHE-MSS), as a revised version of the Impact of Event Scale-Revision (IES-R), for assessment of medical workers' acute mental stress responses within one month and their chronic mental stress responses within six months after major health events.

Patients and methods: The IES-R was slightly modified, developed, and its reliability and validity were tested using the Delphi survey, primary survey with 115 participants, formal survey with 300 participants, and clinical evaluation with 566 participants.

Results: Exploratory factor analysis and confirmatory factor analysis confirmed a promising validity of the scale. The values of Cronbach's alpha coefficient, the Spearman-Brown coefficient, and the retested Cronbach's alpha coefficient of the scale applied for the clinical evaluation were 0.88, 0.87, and 0.98, respectively, which confirmed a good internal consistency and stability. The results of the goodness-of-fit test indicated a good adaptation of the model. A correlation analysis was conducted to assess the correlation between the GHE-MSS and the PCL-C, which had a correlation coefficient of 0.68 ($P < 0.01$).

Conclusion: GHE-MSS can be applied with a promising reliability and validity for the assessment of the acute mental stress response of medical workers experiencing public health events. This method can also be used for the screening of mental stress-associated disorders.

Keywords: IES-R, global health events, mental stress scale, medical workers, mental disorders, mental stress responses

Introduction

A large number of people worldwide are annually infected and killed by epidemic diseases.¹⁻⁵ Due to the tremendous pressure of the epidemic, medical workers fighting the epidemic are prone to negative impacts of physical deterioration and fatigue, including physical exhaustion and mental disorders. On the psychological level, such medical workers face traumatic reactions, interpersonal difficulties, communication problems, loss of faith in fairness and kindness, and severe mental disorders (eg, depression). These challenges further reveal the link among interpersonal difficulties, traumatic reactions, and negative clinical outcome such as suicidal behavior.⁶ In a previous study, measures of affective temperament-types were independently and more strongly associated with lifetime suicide attempts than was diagnosis of a major affective disorder or other variables.⁷

The development of public health strategies largely depends on social and behavioral changes. Therefore, it is crucial to evaluate and compare the behavioral and stress responses to public health events.⁸ Although several investigations have been concentrated on the psychological anxiety and sleep quality assessment for medical workers,⁹⁻¹⁵ currently, no standardized, officially recognized, unified scale exists for the measurement of medical workers' mental stress responses

after experiencing public health events. Most people experience distress after their exposure to an extreme event. In case of good psychosocial resilience and access to social support, the distress of such people may be relatively transient, and adaptation processes can soon be initiated.¹⁶ However, in a minority of patients, a mental health disorder is developed from one to six months after a traumatic event; nevertheless, treatment after one year of such an encounter can often provide satisfactory therapeutic results.¹⁷ Therefore, the development of a mental stress scale for medical workers, especially after experiencing public health events, is essential for early diagnosis and timely treatment of mental disorders.¹⁸

The Impact of Event Scale-Revision (IES-R)¹⁹ is a tool utilized to evaluate post-traumatic stress disorder (PTSD). The original IES-R requires subjects to respond to their psychological conditions in a specific stressful environment in the past week. In the present study, we assumed that after experiencing public health events, medical workers would prefer the use of self-test scales specifically developed for the medical population rather than ones employed for assessment in the general population. Here, we propose GHE-MSS as an addition to the existing established measures of medical workers' acute mental disorder rather than a replacement.

Methods

Ethics Statement

The present study was approved by the Ethics Committee of Shantou Central Hospital (Shantou, China; Approval No. 2020-Ke-Yan (011)). This research was conducted in compliance with the Declaration of Helsinki. Informed consent was obtained from all the participants.

Development of the Scale

Design of the Survey

Modifications to the original IES-R questionnaire were performed accordingly. We modified the original questionnaire as follows: we changed “Below is a list of difficulties that people sometimes experience after stressful events” to “The following questions relates to thoughts, feelings, and behaviors after participating in the treatment of public epidemic diseases”. In addition, we replaced “it” in the items 1, 3, 5, 6, 7, 8, 9, 11, 12, 13, 16, 17, 19, 20, and 22 with “the treatment experience”. We also added “After providing treatment” to items 2, 4, 10, 15, 18, and 21, and to change “that time” to “the treatment time” in item 14.

The Delphi Survey

Two rounds of the Delphi survey were undertaken to evaluate 22 items and validate the content. The first step of the Delphi survey was to determine the number of experts based on the specific research content. Usually, 17–30 experts are selected for exploring underlying information or assumptions. However, 4–16 experts could be selected for consultations to obtain satisfactory answers for an ordinary topic, such as slight modifications of an existing psychological scale.²⁰ In the present study, a total number of ten experts were selected by the convenience sampling method: three psychologists, five neuropsychologists, and two psychiatrists invited from a psychological services clinic, a neurology clinic, and a mental health center of two tertiary hospitals. In addition, experts could access the questionnaire through software, namely “questionnaire star and email”. The questionnaires that were not filled out in accordance with the instructions were treated as invalid and were not included in the data processing. The data collected in the first round, were directly displayed in the second round after statistical analysis. The second round of expert consultation questionnaires was issued, and data were collected for subsequent analysis. Finally, the experts made judgments based on the importance of each indicator. The newly developed scale, GHE-MSS, is a revised version of IES-R.¹⁹ The revised Delphi survey was previously applied to the Chinese version of the IES-R.²¹ The content of Chinese version of GHE-MSS was further verified by back-translation until the meaning of each item was matched with the original item of GHE-MSS. Finally, the Chinese and English GHE-MSS versions were released. The agreement of each item was evaluated using the five-point Likert's scale (five = strongly agree, four = agree, three = neutral, two = disagree, and one = strongly disagree). The quality control of the Delphi survey was based on the following criteria: (1) The return rate of the questionnaire used to measure the positive degree marks given by experts on this questionnaire;²² (2) The authoritative coefficient (Cr) was

composed of the basis of expert judgment (Ca) and the expert familiarity with the questions (Cs). The formula applied was $Cr = (Ca + Cs)/2$. The values of Ca and Cs were obtained through expert self-evaluation.²³ The mean and coefficient of variation were used to describe experts' opinions. The screening and revision of the items were carried out on the basis of mean > 3.5 and coefficient of variation < 0.25 .²⁴ The Kendall coefficient was implemented to indicate the level of consensus among experts' opinions, where Kendall coefficient > 0.7 indicated credibility.²⁵ The authority degree of the experts was expressed by the authority coefficient, and an authority coefficient > 0.7 showed high authority of the experts.

Testing of the Proposed Scale

Participants were asked to complete an online socio-demographic questionnaire (eg, data on sex, identity, and technical titles) via WeChat and Wen-juan Star software. All personal information including demographic was anonymized to maintain and protect confidentiality. Then, the subjects were asked to fill the questionnaire after familiarizing with the introductory part attached to the form to better understand the aim of the research. Next, the participants were prompted to answer questions and to complete the GHE-MSS.

Here, the target subjects were medical workers who had participated in the frontline treatment of COVID-19 since its outbreak in December 2019. The inclusion criteria applied were as follows: (1) medical workers' age ≥ 18 years; (2) medical workers with valid practicing certificates; (3) participation in the prevention, diagnosis, and treatment of COVID-19 for at least two weeks; and (4) voluntary participation in this research. Hospital administrative workers, professional researchers, non-clinical nurses, and non-doctors were excluded from the study cohort. The sample size in the primary survey was set to be at least five-fold higher than the number of the entries, and the sample size in the formal survey was set to be at least ten-fold higher than the number of the entries²⁶ (Figure 1).

1. The IES-R¹⁹ is a 22-item self-report measure that assesses subjective distress caused by traumatic events, but it is not appropriate for PTSD diagnosis. Items are rated on a five-point scale, ranging from zero ("not at all") to four ("extremely"). The IES-R yields a total score (ranging from zero to 88), and intrusion, avoidance, and hyperarousal subscale scores can also be calculated. Scores exceeding a value of 24 are quite meaningful. Those with such high scores, who do not have full PTSD, would have partial PTSD or at least some of the symptoms.²⁷
2. PTSD Checklist-civilian Version (PCL-C). The PCL is a standardized self-report rating scale for PTSD comprising 17 items that correspond to the key symptoms of PTSD. Currently, two versions of the PCL exist: 1) PCL-M that is specific to PTSD caused by military experience and 2) PCL-C that is applied to any traumatic events. PCL-C was utilized in this study based on the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV). Scores for each question ranged from zero (not at all) to five (extremely). A total symptom severity score (range=17–85) can be obtained by summing the scores for each of the 17 items.




Figure 1 Flow chart of the study process.

The best PCL-C cut-off score is used to screen PTSD.^{28–30} This approach for diagnosing PTSD has shown reasonable sensitivity (0.60) and a high specificity (0.99).³¹ The PCL-C was used for the external validity of GHE-MSS in the present study.

Primary Survey

Primary survey was conducted based on the results of the Delphi survey. For this purpose, 115 medical workers from the Emergency Department, Intensive Care Unit (ICU), and the Infection Department of a tertiary hospital in Guangdong Province (China) were enrolled in the primary survey conducted in March 2020. Principal component analysis was employed to evaluate the primary survey results. The reliability of the primary survey was analyzed by Cronbach's alpha coefficient.³² The common-method variance was utilized to investigate the validity of the primary survey.

Formal Survey

The reliability and validity of the formal survey were assessed via the participation of 300 medical workers from the Emergency Departments, ICUs, Infection Departments, Respiratory Departments, and the Clinical Laboratories of five tertiary and secondary hospitals located in Guangdong Province (China). The formal survey was conducted in April 2020. The Cronbach's alpha coefficient was employed to evaluate internal consistency, and the reliability and validity of the formal survey were confirmed. Bartlett's test and the Kaiser-Meyer-Olkin (KMO) measure were used for the determination of the exploratory factor analysis adequacy. The KMO statistics ranged from zero to one, with values closer to one denoting greater adequacy of the factor analysis (KMO \geq 0.6 (low adequacy), KMO \geq 0.7 (medium adequacy), KMO \geq 0.8 (high adequacy), and KMO \geq 0.9 (very high adequacy)).^{33,34}

Clinical Evaluation

Here, 566 medical workers from the Emergency Departments, ICUs, Infection Departments, Respiratory Departments, and the Clinical Laboratories of five tertiary and secondary hospitals in Guangdong Province were surveyed in September 2020. We used the PCL-C to assess the consistency and validity of PTSD occurrence. The Pearson's correlation coefficient (PCC) was employed to calculate the correlation coefficient between each item score and the total score. Confirmatory factor analysis (CFA) was next used to establish the validity of the scale structure.³⁵ A standard load factor $>$ 0.7 was considered to indicate a strong correlation, whereas a factor $<$ 0.4 represented a weak correlation between the item and the factor. Criterion-related validity was utilized to test the correlation between scores of the new scale and the standard scale. The correlation coefficient was within the range 0.40–0.80; the validity was considered to be confirmed at $P < 0.05$.

Statistical Analysis

Data were statistically analyzed with SPSSAU 2016–2021 (Qing-Si Technology Ltd, Beijing, China) and GraphPad Prism 5.0 (GraphPad Prism Software Inc., San Diego, CA, USA) software. CFA was conducted using SPSSAU. Kline's references³⁶ were utilized to analyze univariate normality (skew index \leq 3.0; kurtosis index \leq 10.0). Good model fit cut-off values adopted were those proposed by Hair et al³⁷ and Schumacker and Lomax³⁸ (samples with $N >$ 250 and number of observed variables between 12 and 30): Chi-square test of the model fit (χ^2/df) values less than 5 indicated a good model fit; comparative fit index (CFI) values close to 0.90 or 0.95 represented a good model fit; goodness-of-fit index (GFI) values close to 0.90 or 0.95 indicated a good model fit; standardized root mean square residual (SRMR) and root mean square error of approximation (RMSEA) values below 0.05 revealed good fit, and values between 0.05 to 0.08 with a CFI of 0.92 or higher indicated close model fit.³⁹ Convergent validity was established by the average variance extracted (AVE \geq 0.50)³⁷ and composite reliability (CR \geq 0.70).⁴⁰ In addition, we calculated the heterotrait-monotrait (HTMT) ratio of correlation through a multitrait-multimethod matrix among the GHE-MSS factors by using a value lower than 0.85 to indicate adequate discriminant validity.^{40,41}

Results

The Delphi Survey

Approximately 50% of the study participants were aged >40 years. Besides, 80% of the participants had a Bachelor's degree or above, and 40% of them had work experience >15 years. Experts' research fields were psychotherapy education or research, clinical psychotherapy, and psychological care (Table 1). The authority coefficient was within the range 0.63–0.93, and the overall authority coefficient was above 0.76, indicating that the experts had high authority. Two rounds of the Delphi survey were carried out for each group of experts. Ten questionnaires were sent out in each of the both rounds, which were then recovered. The recovery rate of the two rounds of the Delphi was 100%, indicating that the experts were highly concerned about the questionnaire. The Kendall coefficient in the first round of the Delphi survey was equal to 0.47 ($P < 0.01$). The mean number of entries in the first Delphi round was 4.73, with a coefficient of variation of 0.11. After the modification of experts' opinions, the mean number of the entries in the second round of the Delphi was 4.85, with a coefficient of variation of 0.08. The Kendall coefficient in the second round was equal to 0.74 ($P < 0.01$). Therefore, experts' opinions were relatively consistent and the survey results were accordingly reliable (Table 2).

Proposed Scale Testing

The number of participants in the primary survey was 115 (Table 3). Based on the results of the primary survey, a Cronbach's alpha coefficient of 0.98 was determined. The range of common-method variance was 0.71–0.86. The sample size of the formal survey was 300 (Table 3). KMO value was 0.97, and the Bartlett's test of sphericity showed that degree-of-freedom value was equal to 231 ($P < 0.01$). The percentages of the variance of the three measures were 40.71%, 23.77%, and 10.73%, respectively, with cumulative percentages of variance after the rotation of 75.21%, which showed a good validity. The Cronbach's alpha coefficient value in the formal survey was 0.98, which indicated that the proposed scale had a promising internal consistency.

Clinical Evaluation

Reliability, Validity, and Relevance

In the current survey, 566 subjects participated in the clinical evaluation (Table 3). Based on the analysis of univariate normality, the score of each scale basically showed normal distribution (Supplementary Table 1). The Cronbach's alpha

Table 1 Experts' Demographic Characteristics

Characteristics	Frequency (N)	Proportion (%)
Age (years)		
30–40	5	50
40–50	2	20
≥50	3	30
Highest educational degree		
College	2	20
Undergraduate	6	60
Master's and doctorate	2	20
Title		
Junior professional title	3	30
Senior professional title	3	30
Associate senior professional title	4	40
Work experience (years)		
11–15	6	60
15–20	1	10
≥20	3	30
Research field		
Psychotherapy education or research	1	10
Clinical psychotherapy	7	70
Psychological care	2	20

Table 2 Recovery of the Questionnaire and Suggestions Offered by Delphi Consultation

	First Round	Second Round
Number of questionnaires distributed	10	10
Number of questionnaires collected	10	10
Number of experts who expressed their opinions	3	1
Effective questionnaires	10	10
Recovery rate (%)	100	100
Authority coefficient	0.76	0.76
Mean (coefficient of variation)	4.73 (0.11)	4.85 (0.08)
Kendall coefficient	0.47	0.74
P-value	0.00**	0.00**

Notes: **Significant difference compared between-groups ($p < 0.01$).

Table 3 Medical Workers' Demographic Characteristics in the Three Main Stages of the Survey

Characteristics	Primary Survey (N = 115)		Formal Survey (N = 300)		Clinical Evaluation (N = 566)	
Gender						
Male	28	24.35%	51	17.00%	103	18.20%
Female	87	75.65%	249	83.00%	463	81.80%
Identity						
Doctor	16	13.91%	36	12.00%	70	12.37%
Nurse	99	86.09%	264	88.00%	496	87.63%
Technical titles						
Junior	84	73.04%	206	68.67%	417	73.67%
Intermediate	20	17.39%	74	24.67%	118	20.85%
Senior	11	9.57%	20	6.67%	31	5.48%
Department						
Emergency department	12	10.43%	52	17.33%	158	27.92%
Fever clinic	9	7.83%	27	9.00%	27	4.77%
Respiratory department	14	12.17%	76	25.33%	106	18.73%
Infections department	23	20.00%	50	16.67%	91	16.08%
ICU	41	35.65%	64	21.33%	82	14.49%
Clinical laboratory	8	6.96%	17	5.67%	17	3.00%
Others	8	6.96%	14	4.67%	85	15.02%

Abbreviation: ICU, Intensive Care Unit.

and the Spearman-Brown coefficients of the GHE-MSS were 0.88 and 0.87, respectively, indicating a credible internal consistency of the scale. The results of the analysis of the retest reliability revealed that the reliability of the three measures and the total scale were both above 0.9, confirming a high stability. The correlation coefficient between the relative item and the score of the avoidance measure was within the range 0.76–0.83 ($P < 0.01$). The Pearson's correlation coefficients of the dimensions of the PTSD's hyperarousal and intrusion symptoms were 0.78–0.84 and 0.81–0.90, correspondingly ($P < 0.01$) (Table 4).

Correlation analysis was conducted to assess the correlation between the GHE-MSS and the PCL-C, for which a correlation coefficient of 0.68 ($P < 0.01$) was determined, indicating that the correlation validity of the scale was relatively high (Figure 2). CFA showed that the standard load factor was within the range 0.73–0.89 (Table 5). The fitting index met the requirements for the model fitting adequacy comparative fit index (CFI) = 0.947; the standardized root mean square residual (SRMR) = 0.038; the root mean square error of approximation (RMSEA) = 0.067, representing a reasonable goodness-of-fit (Table 6). Good convergent and discriminant validity indices were established and evaluated (Supplementary Tables 2 and 3).

Table 4 Reliability, Validity, and Relevance Analyzed by Clinical Evaluation of GHE-MSS

	Clinical Evaluation of GHE-MSS (N = 566)			
	Cronbach's Alpha Coefficient	Spearman-Brown Coefficient	Cronbach's Alpha Coefficient (Test-Retest, N = 161)	Pearson's Correlation Coefficient
INT	0.95	0.94	0.99	0.81–0.90**
AVD	0.91	0.86	0.95	0.76–0.83**
HYP	0.89	0.88	0.98	0.78–0.84**

Notes: **Significant difference compared between-groups ($p < 0.01$).

Abbreviations: GHE- MSS, Global Health Events-Mental Stress Scale; INT, intrusion; HYP, hyperarousal; AVD, avoidance.


Assessment Results

The clinical evaluation of the scale was performed six months after conducting the formal survey. We found that the PTSD incidence was from 30.00% to 19.79% lower than that in the formal survey (Figure 3). Furthermore, the incidence of PTSD in PCL-C was 8.13%. Therefore, the incidence of PTSD in GHE-MSS was markedly higher than that in PCL-C (Figure 4).

Discussion

GHE-MSS was developed that included three measures: avoidance, PTSD's hyperarousal, and PTSD's intrusion symptoms. Each measure contained eight, six, and eight items, accounting for a total number of 22 items. Each of the items included in the questionnaire was measured by the five-point Likert's scale. The GHE-MSS was composed of the Delphi consultation, primary survey, formal survey, and clinical evaluation. After clinical evaluation, the Cronbach's alpha coefficient (0.88) showed a good internal consistency. In addition, the items of the scale were slightly modified based on the IES-R via two rounds of Delphi consultation, which enhanced the content validity. Exploratory factor analysis and CFA were implemented to verify all the indicators of the scale, which confirmed that the scale had a promising validity. GHE-MSS was developed to accurately and effectively evaluate the occurrence of mental stress responses in medical workers who experienced public health events. To date, research has been concentrated mainly on indicators, such as acute anxiety, depression, and acute mental stress. In spite of the importance of public health events, few investigations have been performed to determine and assess the incidence of chronic stress among medical workers. Hence, the present study presents a revised version of the IES-R for assessment of chronic stress incidence among medical workers. The proposed scale has the potential to evaluate medical workers' mental stress responses after stressful public health events within six months.

Mental stress response is delayed and thus often fails to attract quickly people's attention. Therefore, a targeted scale should be developed for early screening. Early intervention in people experiencing a major trauma was highly, meaningful in several cases.^{42,43} The initial IES is a 15-item self-reported scale assessing subjective distress caused by

**Figure 2** Correlations between GHE-MSS and PCL-C.

Abbreviations: GHE-MSS, Global Health Events-Mental Stress Scale; PCL-C, PTSD Checklist-Civilian Version.

Table 5 Factor Loadings of the Scale Items (Pattern Matrix)

	Item	HYP	AVD	INT
5	I avoided letting myself get upset when I thought about the treatment experience or was reminded of it.		0.77	
7	I felt as if the treatment experience had not happened or was not real.		0.79	
8	I stayed away from reminders of the treatment experience.		0.76	
11	I tried not to think about the treatment experience.		0.75	
12	I was aware that I still had a lot of feelings about the treatment experience, but I did not deal with them.		0.75	
13	My feelings about the treatment experience were kind of numb.		0.78	
17	I tried to remove the treatment experience from my memory.		0.74	
22	I tried not to talk about the treatment experience.		0.74	
4	After providing treatment, I felt irritable and angry.	0.73		
10	After providing treatment, I was jumpy and easily startled.	0.78		
15	After providing treatment, I had trouble falling asleep.	0.74		
18	After providing treatment, I had trouble concentrating.	0.80		
19	Reminders of the treatment experience caused me to have physical reactions, such as sweating, trouble breathing, nausea, or a pounding heart.	0.77		
21	After providing treatment, I felt watchful and on-guard.	0.79		
1	Any reminder brought back feelings about the treatment experience.			0.86
2	After providing treatment, I had trouble staying asleep.			0.84
3	Other things kept making me think about the treatment experience.			0.85
6	I thought about the treatment experience when I did not mean to.			0.89
9	Pictures about the treatment experience popped into my mind			0.87
14	I found myself acting or feeling like I was back at the treatment time.			0.84
16	I had waves of strong feelings about the treatment experience.			0.79
20	I had dreams about the treatment experience.			0.84
% explained variance		24.11	23.54	21.17

Abbreviations: INT, intrusion; HYP, hyperarousal; AVD, avoidance.

Table 6 Confirmatory Factor Analysis of the Oblique Bi-Factorial Model

	X ²	df	P	X ² /df	CFI	GFI	SRMR	RMSEA
Model	729.016	206	0.000**	3.539	0.947	0.886	0.038	0.067

Notes: **Significant difference compared between-groups ($p < 0.01$).

Abbreviations: X²/df, Chi-square test of model fit; CFI, comparative fit index; GFI, goodness-of-fit index; SRMR, standardized root mean square residual; RMSEA, root mean square error of approximation.

traumatic events. The IES has been revised to IES-R via the addition of seven items related to excessive arousal symptoms of PTSD. Earlier studies showed that a high IES score within one week after a traumatic event was associated with the appearance of PTSD symptoms six months later, with a sensitivity of 92%.^{44,45} Therefore, the development of the scale in the present study included a primary survey, a formal survey in the early stage of the COVID-19 outbreak, and a clinical evaluation of the stress response six months later. The PCL-C was employed as the standard for accuracy and validity assessment. The results showed that the Cronbach's alpha coefficient and the Spearman-Brown coefficient of the scale both had values above 0.85. The CFA revealed that the modified model fitted well, suggesting that the scale has a promising reliability and stability. Each measure was characterized by good internal consistency and stability. However, the incidence of PTSD in the GHE-MSS in the clinical evaluation was significantly higher than that in the PCL-C, which may be due to the differences in the reliability and validity of the mentioned scales. As previously reported, IES-R is not appropriate for PTSD diagnosis.¹⁹ Previous evidence showed that PCL-C had a reasonable sensitivity and a higher specificity for PTSD diagnosis.³¹ Generally, a positive correlation exists in the incidence of PTSD between GHE-MSS and PCL-C, confirming that GHE-MSS can be applied to assess the mental status of medical workers experiencing public




Figure 3 Estimation of the incidence of PTSD using the GHE-MSS in the stages of scale testing and clinical evaluation.

Abbreviations: GHE-MSS, Global Health Events-Mental Stress Scale; PTSD, post-traumatic stress disorder.




Figure 4 The rate of PTSD symptoms assessed by GHE-MSS and PCL-C in the stage of clinical evaluation.

Abbreviations: GHE-MSS, Global Health Events-Mental Stress Scale; PTSD, post-traumatic stress disorder; PCL-C, The PTSD Checklist-Civilian Version; ICU, Intensive Care Unit.

health events. In the present study, GHE-MSS was developed on the basis of IES-R. We found that the GHE-MSS was more promising for the assessment of acute and chronic stress mental responses of medical workers involved in public health events rather than the diagnosis of PTSD symptoms.

In addition to the assessment of the acute stress mental response, the early identification of subgroups of the population of medical workers exposed to a higher risk of mental stress could provide guidance on the timely application of therapeutic measures after major adverse health events. In this regard, our study has several strengths. To the best of our knowledge, this is the first self-rating scale for the assessment of the mental response of medical workers experiencing major adverse health events, specifically an infectious disease outbreak. Nevertheless, further research is still urgently needed. As known, infectious disease outbreaks have a profound psychological impact on healthcare workers. In this respect, Tam et al established that healthcare workers experienced acute stress reactions during the Severe Acute Respiratory Syndrome (SARS) outbreak in 2003.^{46,47} Recently, several reports have confirmed the substantial psychological stress as well as physical symptoms experienced by health care professionals dealing with the COVID-19 outbreak.^{48,49} In addition, this study was commenced during the initial wave of COVID-19 pandemic, emphasizing the acute mental response of medical workers. With the long-expected entry into the post-COVID-19 era, local and national mental health institutions have conducted evaluations and provided support for the general public,

including health care professionals, such as online mental health services.⁵⁰ Therefore, increasingly more healthcare facilities and governmental efforts are likely to focus on the psychological aspect of severe infectious disease outbreaks such as that of COVID-19. It is hence necessary that medical workers exposed to significant work pressure should perform appropriate mental disorder self-tests. Furthermore, this study assessed mental stress in both tertiary and secondary hospitals in different areas, which is often a missing element of research exploring stress in healthcare workers.

This study provides initial evidence for the validity and reliability of scores derived for GHE-MSS. Earlier findings support the view that GHE-MSS can be used to measure mental distress, especially in medical workers experiencing public health events. For example, Han et al⁵¹ reported that 34.6% of the medical staff in China had depressive symptoms from February 17 to 24, 2020. Another study also showed the combined prevalence of anxiety, depression, and insomnia of Hubei medical staff, which reached as high as 38%.⁵² Additionally, Kheradmand et al reported that healthcare workers in Tehran developed mild (41.4%) to moderate (31.5%) impact mental issues during the initial phase of the pandemic.⁵³ The results of these earlier studies are similar to our findings on the incidence of PTSD among medical workers in the early stages of COVID-19. Hopefully, GHE-MSS can be assessed and validated in other countries to further deepen this evidence. We plan to do a large-sample, cross-sectional study by applying this scale in national hospitals through cooperation with the related government departments and psychological support centers. In this future study, we intend collect and assess more clinical evidence on the application of GHE-MSS. Further studies could also facilitate the development of effective strategies for the prevention of mental disorders, which could not only allow early intervention and identification of high-risk medical individuals, but also provide the supportive environment for medical workers seeking help.

Limitations

The present study contains several limitations. First, this is a cross-sectional study, which might have limited the possibility for establishing temporality between exposure and health outcome, increasing the risk for reporting bias. For instance, reporting bias can occur when the same person reports both exposure and health effects as in the case of questionnaire studies and when there is a public awareness of the problem.⁵⁴ In such a situation, a higher risk might exist for systematic over-reporting of mental disorder-associated problems in diseased subjects when compared with healthy ones and/or a risk for over-reporting of symptoms in subjects after experiencing public health events when compared with people without such experience. Such reporting bias does not occur to the same extent in longitudinal studies, and, therefore, further research is required. Second, the developed scale was more appropriate for the assessment of mental stress, particularly associated with public health events, whereas the participants were medical workers who participated in COVID-19 treatment in China. We failed to include medical workers who participated in other health events (eg, influenza A infection outbreaks). Therefore, this scale still needs to be validated in research on other infectious diseases. In addition, the Chinese version of GHE-MSS was verified by back-translation with the original item of GHE-MSS. The targeted population of our study consisted of Chinese medical workers, which may decrease the generalizability of our results because of the differences between languages and the inhomogeneous distribution of mental response across the globe. It is unknown whether the results of this study are applicable to other medical populations in China or other countries. Hopefully, this self-rating scale can be assessed and validated in other countries in the future. The results of the convenience sampling cannot be generalized to the target population because of the potential bias of the sampling technique. Young medical workers and people experiencing heavy mental stress that are familiar with social media were more likely to participate in our research. However, senior doctors and nurses were reluctant to participate in this study. Therefore, the degree of mental stress reaction in our study might have been overestimated. Meanwhile, we could not assess how many people had been invited to participate in the research, and the response rate could not be, therefore, calculated. Our study included medical workers from only five tertiary and secondary hospitals in Guangdong Province. Hence, the source and size of the samples were limited. Thus, further research with a larger sample size is warranted to eliminate the aforementioned shortcomings and to confirm our findings.

Conclusion

In conclusion, here, we propose a revised version of the IES-R, called GHE-MSS. Our findings revealed that the proposed scale has a good internal consistency and validity. GHE-MSS may facilitate the screening of mental stress

responses in medical workers in the early stage of their occurrence after experiencing public health events. Therefore, the identification of subgroups of medical workers with higher susceptibility to PTSD may be of significance. The proposed scale is also promising for developing targeted therapies for individuals who are at high-risk of mental stress exposure due to their participation in the treatment of pandemics-affected subjects.

Acknowledgments

This study was supported by the Nursing Department of Shantou Central Hospital. We would like to express our gratitude to Ms. Li-Xian Yang from the Second People's Hospital of Shantou, Ms. Qiu-Lian Lin from the Third People's Hospital of Shantou, Ms. Li-Fang Lin from Chao-yang District People's Hospital of Shantou, and Dr. Qing-Hua Huang from Nan-ao County People's Hospital of Shantou for their contributions to the study. We are also thankful to Professor Kitty K. Wu for approving the use of the CIES-R. Special thanks are due to Professor Marmar Charles for approving the use of IES-R. This study was financially supported by the Shantou Science and Technology Plan Medical and Health Category Project (Grant No. Shan-Fu-Ke [2020]23-03).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Qiu J. One world, one health: combating infectious diseases in the age of globalization. *Natl Sci Rev*. 2017;4(3):493–499. doi:10.1093/nsr/nwx047
2. Sullivan SG, Arriola CS, Bocacao J, Burgos P, Wood T. Heterogeneity in influenza seasonality and vaccine effectiveness in Australia, Chile, New Zealand and South Africa: early estimates of the 2019 influenza season. *Eurosurveillance*. 2019;24(45). doi:10.2807/1560-7917.ES.2019.24.45.1900645
3. Shi W, Ke C, Fang S, Li J, Wang D. Co-circulation and persistence of multiple A/H3N2 influenza variants in China. *Emerg Microbes Infect*. 2019;8(1):1157–1167. doi:10.1080/22221751.2019.1648183
4. Vallès X, Stenseth NC, Demeure C, Horby P, Baril L. Human plague: an old scourge that needs new answers. *PLoS Negl Trop Dis*. 2020;14(8):e0008251. doi:10.1371/journal.pntd.0008251
5. Randremanana R, Raberahona M, Randria M, et al. An open-label, randomized, non-inferiority trial of the efficacy and safety of ciprofloxacin versus streptomycin + ciprofloxacin in the treatment of bubonic plague (IMASOY): study protocol for a randomized control trial. *Trials*. 2020;21(1):722. doi:10.1186/s13063-020-04642-2
6. Solano P, Ustulin M, Pizzorno E, et al. A google-based approach for monitoring suicide risk. *Psychiatry Res*. 2016;246:581–586. doi:10.1016/j.psychres.2016.10.030
7. Baldessarini R, Innamorati M, Erbuto D, et al. Differential associations of affective temperaments and diagnosis of major affective disorders with suicidal behavior. *J Affect Disord*. 2017;210:19–21. doi:10.1016/j.jad.2016.12.003
8. Philippe FL, Houle I. Cognitive integration of personal or public events affects mental health: examining memory networks in a case of natural flooding disaster. *J Pers*. 2019;88:861–873. doi:10.1111/jopy.12531
9. Torales J, O'Higgins M, Castaldelli-Maia JM, Ventriglio A. The outbreak of COVID-19 coronavirus and its impact on global mental health. *Int J Soc Psychiatry*. 2020;66(4):317–320. doi:10.1177/0020764020915212
10. Wang C, Pan R, Wan X, Tan Y, Ho RC. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health*. 2020;17(5):1729. doi:10.3390/ijerph17051729
11. Qiu J, Shen B, Zhao M, Wang Z, Xu Y. A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *Gen Psychiatry*. 2020;33(2):3. doi:10.1136/gpsych-2020-100213
12. Sanguino CG, Ausín B, Castellanos MA, Saiz J, Muoz M. Mental health consequences during the initial stage of the 2020 coronavirus pandemic (COVID-19) in Spain. *Brain Behav Immun*. 2020;87:172–176.
13. Zhang Y, Ma ZF. Impact of the COVID-19 pandemic on mental health and quality of life among local residents in Liaoning Province, China: a cross-sectional study. *Int J Environ Res Public Health*. 2020;17(7):2381.
14. Mazza C, Ricci E, Biondi S, et al. A nationwide survey of psychological distress among Italian people during the COVID-19 pandemic: immediate psychological responses and associated factors. *Int J Environ Res Public Health*. 2020;17(9):3165. doi:10.3390/ijerph17093165
15. Cw A, Rp A, Xw A, et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun*. 2020;87:40–48. doi:10.1016/j.bbi.2020.04.028
16. Lock S, Rubin GJ, Murray V, Rogers MB, Williams R. Secondary stressors and extreme events and disasters: a systematic review of primary research from 2010–2011. *PLoS Curr*. 2012;4. doi:10.1371/currents.dis.a9b76fed1b2dd5c5bfcfc13c87a2f24f
17. Deng M, Medicine P. New progress of clinical research to posttraumatic stress disorder (DSM-5 update). *Chin J Health Psychol*. 2016;1:7.
18. Widiger T, McCabe G. The alternative model of personality disorders (AMPD) from the perspective of the five-factor model. *Psychopathology*. 2020;53(3–4):1–8. doi:10.1159/000507378
19. Weiss D, Marmar C. *Impact of Event Scale - Revised (IES-R) - PTSD*. National Center for PTSD; 1997.
20. Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs*. 2000;32(4):1008.
21. Wu KK, Chan KS. The development of the Chinese version of impact of event scale – revised (CIES-R). *Soc Psychiatry Psychiatr Epidemiol*. 2003;38(2):94–98. doi:10.1007/s00127-003-0611-x

22. Li Y, Ehiri J, Hu D, et al. Framework of behavioral indicators for outcome evaluation of TB health promotion: a Delphi study of TB suspects and Tb patients. *BMC Infect Dis.* 2014;14(1):268. doi:10.1186/1471-2334-14-268
23. Xing Q, Zhang M, Zhao F, Zhou Y, Yuan L. The development of a standardized framework for primary nurse specialists in diabetes care in China: a Delphi study. *J Nurs Res.* 2019;27(6):1. doi:10.1097/jnr.0000000000000330
24. Hou YC, Wang M, Li XQ, Wen S, Xu YH. [Construction of surveillance and early-warning-index-system on tuberculosis in China]. *Zhonghua liu xing bing xue za zhi.* 2012;33(5):505–508. Chinese. doi:10.1016/j.tmaid.2012.03.001
25. Zhai XD, Wang CX, Ma YJ, et al. Using the modified Delphi method to research the influencing factors of long-term health-related quality of life in patients with unruptured intracranial aneurysms after endovascular treatment. *Chin Neurosurg J.* 2020;6(1):7. doi:10.1186/s41016-020-00186-1
26. Carey RG, Seibert JH. A patient survey system to measure quality improvement: questionnaire reliability and validity. *Med Care.* 1993;31(9):834–845. doi:10.1097/00005650-199309000-00008
27. Asukai N, Kato H, Kawamura N, et al. Reliability and validity of the Japanese-language version of the impact of event scale-revised (IES-R-J): four studies of different traumatic events. *J Nerv Ment.* 2002;190(3):175–182. doi:10.1097/00005053-200203000-00006
28. Dobie DJ, Kivlahan DR, Maynard C, Bush KR, Bradley KA. Screening for post-traumatic stress disorder in female veteran's affairs patients: validation of the PTSD checklist. *Gen Hosp Psychiatry.* 2002;24(6):367–374. doi:10.1016/S0163-8343(02)00207-4
29. Lang AJ, Laffaye C, Satz LE, Dresselhaus TR, Stein MB. Sensitivity and specificity of the PTSD checklist in detecting PTSD in female veterans in primary care. *J Trauma Stress.* 2003;16(3):257–264. doi:10.1023/A:1023796007788
30. Liu X, Yang Y, Ping Y, Xun Z, Xiong G. A study of the relationship between mental health and menstrual abnormalities in female middle school students from postearthquake Wenchuan. *Biosci Trends.* 2010;4(1):4–8.
31. Brewin CR. Systematic review of screening instruments for adults at risk of PTSD. *J Trauma Stress.* 2005;18(1):53. doi:10.1002/jts.20007
32. Tabachnick BG, Fidell LS. *Using Multivariate Statistics.* 5th ed. Boston (MA): Pearson Education; 2007.
33. Devellis R, Devellis R, Devellis RF. *Scale Development: Theory and Applications.* Sage publications; 1991.
34. Ribeiro F, Mussi FC, Pires C, Silva R, Macedo T, Santos C. Stress level among undergraduate nursing students related to the training phase and sociodemographic factors. *Rev Lat Am Enfermagem.* 2020;28. doi:10.1590/1518-8345.3036.3209
35. Stott J, Spector A, Orrell M, Scior K, Sweeney J, Charlesworth G. Limited validity of the hospital anxiety and depression scale (Hads) in dementia: evidence from a confirmatory factor analysis. *Int J Geriatr Psychiatry.* 2017;32(7):805–813. doi:10.1002/gps.4530
36. Kline RB. *Principles and Practice of Structural Equation Modeling.* 2nd ed. Guilford publications; 2004.
37. Hair JFJ, Black WC, Babin BJ, Anderson RE, Tatham RL. *Multivariate Data Analysis.* Pearson Education; 2006.
38. Schumacker RE, Lomax RG, Beginner's A. Guide to structural equation modeling. *Technometrics.* 2004;47(4):522.
39. Encantado J, Marques MM, Palmeira AL, Sebire SJ, Gouveia MJ. Development and cross-cultural validation of the goal content for weight maintenance scale (GCWMS). *Eat Weight Disord.* 2021;26:1–12.
40. Hair JF, Hult G, Ringle CM, Sarstedt M. *A Primer on Partial Least Squares Structural Equation Modeling (PLS-SEM).* 2nd ed. Sage publications; 2016.
41. Henseler JR, Ringle CM, Sarstedt M. A new criterion for assessing discriminant validity in variance-based structural equation modeling. *J Acad Mark Sci.* 2015;43(1):115–135. doi:10.1007/s11747-014-0403-8
42. McLaughlin KA, Koenen KC, Friedman MJ, et al. Subthreshold posttraumatic stress disorder in the world health organization world mental health surveys. *Biol Psychiatry.* 2015;77(4):375–384. doi:10.1016/j.biopsych.2014.03.028
43. Rosendal S, Mortensen EL, Andersen HS, Heir T. Use of health care services before and after a natural disaster among survivors with and without PTSD. *Psychiatr Serv.* 2014;65(1):91–97. doi:10.1176/appi.ps.201200535
44. Orsillo SM. *Measures for Acute Stress Disorder and Posttraumatic Stress Disorder.* US: Springer; 2002.
45. Horowitz M, Wilner N, Alvarez W. Impact of Event Scale: a measure of subjective stress. *Psychosom Med.* 1979;41(3):209–218. doi:10.1097/00006842-197905000-00004
46. Mcalonan G, Lee A, Cheung V, et al. Immediate and sustained psychological impact of an emerging infectious disease outbreak on health care workers. *Can J Psychiatry.* 2007;52(4):241–247. doi:10.1177/070674370705200406
47. Tam CWC, Pang EPF, Lam LCW, Chiu HFK. Severe acute respiratory syndrome (SARS) in Hong Kong in 2003: stress and psychological impact among frontline healthcare workers. *Psychol Med.* 2004;34(7):1197–1204. doi:10.1017/S0033291704002247
48. Couarraze S, Delamarre L, Marhar F, et al. The major worldwide stress of healthcare professionals during the first wave of the COVID-19 pandemic - The international COVISTRESS survey. *PLoS One.* 2021;16(10):e0257840. doi:10.1371/journal.pone.0257840
49. Ma Y, Rosenheck R, He H. Psychological stress among health care professionals during the 2019 novel coronavirus disease outbreak: cases from online consulting customers. *Intensive Crit Care Nurs.* 2020;61:102905. doi:10.1016/j.iccn.2020.102905
50. Kang C, Tong J, Meng F, Feng Q, Yang J. The role of mental health services during the COVID-19 outbreak in China. *Asian J Psychiatr.* 2020;52:102176. doi:10.1016/j.ajp.2020.102176
51. Han B, Ma C, Liu Z, et al. Perceived psychological feelings make important contributions to the symptoms of common mental disorders of medical staff during the COVID-19. *Front Psychiatry.* 2021;12:738610. doi:10.3389/fpsy.2021.738610
52. Wang L, Zhang M, Liu G, et al. Psychological impact of coronavirus disease (2019) (COVID-19) epidemic on medical staff in different posts in China: a multicenter study. *J Psychiatr Res.* 2020;129:198–205. doi:10.1016/j.jpsychires.2020.07.008
53. Kheradmand A, Mahjani M, Pirsalehi A, Fatemizadeh S, Khoshgoui B. Mental health status among healthcare workers during COVID-19 pandemic. *Iran J Psychiatry.* 2021;16(3):250.
54. Larsson M, Hägerhed-Engman L, Moniruzzaman S, Janson S, Sundell J, Bornehag CG. Can we trust cross-sectional studies when studying the risk of moisture-related problems indoor for asthma in children? *Int J Environ Health Res.* 2011;21(4):237–247. doi:10.1080/09603123.2010.533368

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>