

# Effects of Opium Addiction and Cigarette Smoking on Hematological Parameters

Gholamabbas Shahabinejad MSc<sup>1</sup>, Majid Sirati-Sabet PhD<sup>2</sup>, Mohammad Kazemi-Arababadi PhD<sup>3</sup>, Saeideh Nabati MSc<sup>4</sup>, Gholamreza Asadikaram PhD<sup>5</sup>

## Original Article

### Abstract

**Background:** The aim of the present study was to investigate the effects of opium addiction and cigarette smoking on the complete blood count (CBC).

**Methods:** Eighty-six male subjects, including 31 opium-addicted cigarette smokers (OACS), 19 opium-addicted non-cigarette smokers (OANCS), 17 non-opium-addicted cigarette smokers (NOACS), and 19 non-opium-addicted non-cigarette smokers (NOANCS) participated in this study. The CBC test was measured in all individuals.

**Findings:** The OACS had significantly higher white blood cell (WBC), lymphocyte, and red blood cell (RBC) count but lower in mean corpuscular volume (MCV) compared to NOANCS. The OANCS had significantly higher lymphocyte in comparison with NOACS. Our results demonstrated that the number of WBC, lymphocytes, and RBC were significantly higher, while, MCV was lower in OANCS subjects when compared to NOACS. The OACS had significantly higher level of lymphocyte in comparison with NOACS. The mean number of lymphocyte in OANCS was found significantly higher than NOACS. The smokers were shown to have significantly higher levels of WBC compared to NOANCS.

**Conclusion:** Our results showed that opium-addiction, especially when associated with cigarette smoking, has intensive effects on hematological factors and these alteration might leads to greater risk for developing atherosclerosis, cardiovascular diseases, and imbalance in immune system.

**Keywords:** Cigarette smoking; Addiction; Opium; Complete blood count

**Citation:** Shahabinejad G, Sirati-Sabet M, Kazemi-Arababadi M, Nabati S, Asadikaram G. **Effects of Opium Addiction and Cigarette Smoking on Hematological Parameters.** *Addict Health* 2016; 8(3): 179-85.

**Received:** 20.02.2016

**Accepted:** 28.04.2016

1- Neuroscience Research Center, Institute of Neuropharmacology AND Department of Biochemistry, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

2- Associate Professor, Department of Biochemistry, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3- Assistant Professor, Department of Laboratory Sciences, School of Paramedicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

4- PhD Student, Department of Biochemistry, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

5- Professor, Endocrinology and Metabolism Research Center, Institute of Basic and Clinical Physiology Sciences AND Department of Biochemistry, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Correspondence to: Gholamreza Asadikaram PhD, Email: gh\_asadi@kmu.ac.ir

## Introduction

Cigarette smoking is considered as a major leading causes of health problem in word over.<sup>1</sup> There are more than 4000 chemicals found in cigarette smoke, and a cigarette smoker is exposed to a number of harmful substances including nicotine, free radicals, carbon monoxide and other gaseous products.<sup>1,2</sup> In a number of studies, it has been found that smokers have higher white blood cell (WBC) counts than nonsmokers.<sup>2</sup> Tarazi et al. have reported increased red blood cell (RBC), several RBC-related parameters, and hemoglobin (Hb) concentration in cigarette smokers.<sup>3</sup> It is suggested that increase in hemoglobin level in blood of smokers may be a compensatory mechanism. However, all smokers does not show increase in hemoglobin level and this relates to smoker situation.<sup>4</sup> The effects of opium addiction on the number of RBC, the levels of some RBC-related parameters, and total and differential counts of peripheral WBC have shown elsewhere.<sup>5</sup> Opium can be used to synthesize some medications such as morphine, codeine, noscapine, and papaverine.<sup>6</sup> There are many reports regarding the effects of opium derivatives on biological systems. The presence of specific morphine receptors on lymphocytes has been reported. It is suggested that opioid receptor acts in both autocrine and paracrine fashion.<sup>7</sup> Although it is proposed that morphine affects some immune cells indirectly, but morphine can also directly affect macrophages and polymorphonuclear (PMN) leukocytes functions, and regulate expression of some type of T-cell surface markers.<sup>8</sup> Some researcher reported that the endogenous opioid peptides, including  $\beta$ -endorphin and the dynorphin peptides, and exogenous alkaloids such as morphine plays an important role in the lymphocytes and other immune cells function.<sup>9</sup> Although heroin induced increase in circulating leukocytes, but it decrease the number of splenic leukocytes by induction of leukocytes apoptosis.<sup>10</sup> Ke et al.,<sup>11</sup> have reported that noscapine plays key antitumor activity without the interference of immune responses. Moreover noscapine down-regulates proliferation of leukemia cell lines<sup>12</sup> and it induces chromosomal loss, hyperdiploidy and hypodiploid in human lymphocytes in in vivo

conditions.<sup>13</sup> Obviously these effects can directly influence the lymphocytes function, and thereby affecting cytokine networks and therefore, impact immune cells differentiation and proliferation. Papaverine, other opium alkaloids, induced apoptosis in different cell lines.<sup>14</sup> Patently, these events can influx metabolism and function of lymphocytes. It is also reported that papaverine, by stimulation of production of transforming growth factor- $\beta$  (TGF- $\beta$ ) which has potent immunosuppression effects, inserts a highly fatal syndrome as constrictive bronchiolitis.<sup>15</sup> However, opium contains more than 20 alkaloids, and more than 70 ingredients.<sup>16</sup> Thus, its effect is clearly different from morphine, noscapine, codeine and papaverine. Our previous studies on the effects of opium on apoptosis,<sup>17</sup> immune functions,<sup>18</sup> hematological parameters<sup>5</sup> and also the influence of cigarette smoking on cell functions, led us to evaluate the biological effects of opium and cigarette on peripheral blood cells of opium-addicted, cigarette smokers opium-addicted, and cigarette smokers non-opium-addicted individuals.

## Methods

### *Subject selection*

Subjects participated in this study consisted of 86 males, including 31 opium-addicted cigarette smokers (OACS), 19 opium-addicted non-cigarette smokers (OANCS), 17 non-opium-addicted cigarette smokers (NOACS), and 19 non-opium-addicted non-cigarette smokers (NOANCS) individuals. All subjects were aged from 19 to 56 years. Exclusion criteria included any kind of disease (either symptoms or diagnosis), consumption of any medication that might interfere with assessed parameters, consumption of other component of opium (heroin, morphine etc.) and withdrawal drugs such as methadone or other withdrawal regimes. Opium addicted subjects had been using opium at least 500 mg per day for more than 1 year prior to sample preparation. The method that was used by all subjects was smoking opium using opium pipe (Bafur). A cigarette smoker was defined as one who smoked at least 5 cigarettes per day for more than 1 year but did not use any other kind of tobacco. Each individual in the control group were selected with similar attributes regarding age, residency and BMI with no addiction to

opioids or cigarette in case group. This study was approved by the ethical committee of the Rafsanjan University of Medical Sciences, Iran, and written informed consent was filled out by the participants, prior to sample collection.

### Blood cells counting

Peripheral blood samples were collected from the subjects at 8-9 A.M., before consuming opium or cigarette and CBC was tested immediately. The total cell count was made using a hematology analyzer (Sysmex KX-21N) and Gimsa's stain and a light microscope were used for differential WBC counts (lymphocytes, monocytes, neutrophils, basophils and eosinophils per 100 WBC), which were based on morphological characteristics of the cells.

Results are presented in mean  $\pm$  standard errors (SE). To compare the mean values between groups, the one-way analysis of variance (ANOVA) was used. Statistical analysis was performed by SPSS software (version 18, SPSS Inc., Chicago, IL, USA) for Windows.  $P < 0.05$  was considered statistically significant.

## Results

Mean number of total WBC were significantly higher in OACS, OANCS and NOACS than NOANCS ( $P < 0.001$ ).

The mean percent of lymphocytes in OACS and OANCS was significantly higher when compared with NOACS and NOANCS ( $P < 0.001$ ). The numbers of RBC were also

significantly increased in OACS and OANCS when compared with NOANCS ( $P = 0.018$ ).

MCV was also significantly different among the enrolled groups ( $P = 0.001$ ). MCV was significantly decreased in OACS and OANCS when compared with NOANCS.

The mean of neutrophils count ( $P = 0.168$ ), Hct ( $P = 0.925$ ), Hb ( $P = 0.913$ ), and platelets count ( $P = 0.084$ ) was not different among the studied groups. The results are shown in details in table 1.

## Discussion

Our result revealed the profound effects of opium and cigarette on the total and differential counts of peripheral WBC, lymphocytes, RBC, and MCV (as a RBC-related parameter).

We have showed that the mean number of total WBC in opium and cigarette consumers was significantly higher than NOANCS groups. The precise mechanisms by which opium and cigarette could strongly elevate WBC counts and also can alter the differential WBC counts have yet to be identified. It may be hypothesized that opium and cigarette lead to liberation of WBC from bone marrow in indirect format by induction of pro-inflammatory cytokines which have been reported by our previous investigations.<sup>19</sup> Additionally, the number of peripheral WBCs was influenced by the modification of the expression of adhesion molecules on the endothelial cells and the differentiation of leukocytes from bone marrow stem cells.<sup>20</sup>

**Table 1.** Comparison of hematological indices between participants groups with and without opium addiction or smoking

Variable	OACS	OANCS	NOACS	NOANCS	P
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	
WBC (Number/dl)*	7077 $\pm$ 261	7595 $\pm$ 443	7247 $\pm$ 551	5326 $\pm$ 152	< 0.001
Lymphocytes (Percent)**	45.70 $\pm$ 1.74	43.90 $\pm$ 1.68	35.90 $\pm$ 2.63	35.40 $\pm$ 1.09	< 0.001
Neutrophils (Percent)	49.00 $\pm$ 1.15	48.60 $\pm$ 2.09	56.60 $\pm$ 5.37	52.00 $\pm$ 1.01	0.168
RBC (Million/dl) <sup>†</sup>	5.70 $\pm$ 0.12	5.70 $\pm$ 0.16	5.40 $\pm$ 0.12	5.20 $\pm$ 0.12	0.018
MCV <sup>††</sup>	80.30 $\pm$ 1.41	78.70 $\pm$ 1.47	83.20 $\pm$ 1.67	87.00 $\pm$ 0.93	0.001
Hb	14.90 $\pm$ 0.24	14.80 $\pm$ 0.40	14.60 $\pm$ 0.31	14.80 $\pm$ 0.24	0.913
Hct	45.50 $\pm$ 0.76	44.90 $\pm$ 1.17	45.00 $\pm$ 0.74	44.90 $\pm$ 0.71	0.925
Plt	267.00 $\pm$ 17.90	240.00 $\pm$ 16.90	235.00 $\pm$ 12.30	211.00 $\pm$ 9.39	0.084

\*The post hoc (Tukey) test showed that the number of total WBC and lymphocytes were significantly increased in OACS, and NOACS in comparison to NOANCS, \*\*The post hoc (Tukey) test showed that the percent of lymphocytes in OACS and OANCS were significantly increased when compared with NOACS and NOANCS, <sup>†</sup>The post hoc (Tukey) test revealed that the number of RBC were significantly increased in OACS and OANCS when compared with NOANCS, <sup>††</sup>The post hoc (Tukey) test demonstrated that MCV were significantly decreased in OACS and OANCS when compared with NOANCS.

WBC: White blood cell; RBC: Red blood cell; MCV: Mean corpuscular volume; Hb: Hemoglobin; Hct: Hematocrit; Plt: Platelet; OACS: Opium-addicted cigarette smokers; OANCS: Opium-addicted non-cigarette smokers; NOACS: Non-opium-addicted cigarette smokers; NOANCS: Non-opium-addicted non-cigarette smokers; SE: Standard error

Moreover, some opium side effects on the number of WBCs may be mediated through infection.<sup>5</sup> The release of catecholamines which are important factor to elevate the leukocyte count are induced by morphine.<sup>21</sup> Opium components change some multi-functional cytokines, such as Interleukin-6 (IL-6), which induces inflammatory process that leads to elevations in the peripheral blood leukocyte count.<sup>22</sup> Consistent with our results, Asakura et al.<sup>23</sup> showed that the risk of acute infection is higher in heroin addicted patients. As we know clearly today, the hematopoiesis is affected by a complex network of cytokines.<sup>24</sup> The serious effects of opium on the secretion of some cytokines such as TGF- $\beta$ , IL-4, IL-5, IL-10 and IFN- $\gamma$  has been shown.<sup>18</sup> Hence, the opium or some of its components interfere with cytokine network and consequently influence the production of WBC and lymphocytes. These findings indicated that the opium addiction have deep adverse effects on WBC and lymphocytes counts. Moreover, the opium or some of its derivatives may affect lymphopoiesis via interference with cytokines which are the responsible factors for differentiation of lymphocytes from bone marrow stem cells.<sup>25</sup> On the other hand cigarette smoking suppresses the immune response which may contribute to an increased risks of periodontal disease in smokers.<sup>26</sup> Our results showed significant differences in hematological parameters of subjects, as the OACS and OANCS participants had significantly higher lymphocyte compared to NOACSs. Our results demonstrated that the number of lymphocytes and WBC were significantly higher in OANCS subjects when compared to NOANCS. The cigarette smokers had significantly higher levels of WBC in comparison with NOANCS. We found that regular smokers had significantly higher WBC count compared to non-smokers. The high WBC count in male smokers in this study is consistent with other published reports.<sup>27</sup> Vulser et al.<sup>28</sup> observed that median total leukocyte count was 36% higher in current smokers as compared to non-smokers. Although, it is reported that inflammatory stimulation of the bronchial tract induces an increase in inflammatory markers in the blood, but the mechanism for smoking-induced increase in WBC count is not clear yet.<sup>29</sup> In addition, inducing effect of nicotine

on blood lymphocyte counts has also been suggested. While leukocytosis may simply be a marker of smoking-induced tissue damage, the high count can promote cardiovascular diseases through multiple pathologic mechanisms that mediate inflammation, plug the microvasculature, induce hypercoagulability, and promote infarct expansion.<sup>30</sup> Seinen et al. found that the greater the number of cigarettes smoked, the lower the hypoxanthine-guanine phosphoribosyltransferase (HGPRT) activity, and that HGPRT activity was higher in smokers who had started smoking later. On the other hand, it was demonstrated that HGPRT activity was inversely related to the WBC.<sup>31</sup> In fact, several studies have shown that WBC count is an independent predictor of cardiovascular diseases and its related complications.<sup>32</sup> The high WBC count in our male smoking subjects may also suggest that they might be at greater risk for developing atherosclerosis and cardiovascular diseases. We observed that RBC values were significantly higher in OACS in comparison to NOANCS, that is consistent with other investigations.<sup>33</sup> It is reported that high level of RBC, WBC and hematocrit are associated with blood viscosity and clotting in the individuals.<sup>34</sup> Interestingly, our results also demonstrated that the numbers of RBCs were significantly increased in the opium and cigarette addicted individuals.

It has been established that fibrinogen levels are higher in smokers than in non-smokers, and it has been estimated that the increasing risk of cardiac disease in smokers may be associated with higher fibrinogen levels through arterial wall infiltration and its effects on blood viscosity, platelet aggregation, and fibrin formation. This elevation may lead to congenital heart disease, pulmonary fibrosis and elevated erythropoietin level and, therefore, polycythemia.<sup>4</sup> Cigarette smoking developed a combined polycythemia to chronic hypoxia, leading to increased RBC production, with concomitant plasma volume reduction.<sup>35</sup> During cigarette smoking, carbon monoxide (CO) is produced by the imperfect ignition of its material. CO displaces oxygen from hemoglobin in RBCs, which decreases the release of oxygen to tissues.<sup>36</sup> Thus, the oxygen content of blood in the presence of carbon monoxide is much lower than normal. Obviously, this event leads to polycythemia. MCV, MCH and MCHC are

important RBC indices that represent average size and hemoglobin composition of the RBCs.<sup>37</sup> We found that MCV level in OACS and OANCS groups was significantly decreased compared with NOANCS. MCV indicates the size of a RBCs and presence of red cells smaller or larger than normal size means the person has anemia. A lowered level of MCV in our study indicates that subjects might suffer from macrocytic anemia that usually caused by iron and folic acid deficiencies. Furthermore, we demonstrated that in opium-addicted group the RBC counts was significantly increased when compared with control group. The reasons for these differences are remained to be cleared. According to the results of the present study the RBC indices are markedly decreased. It may be due to the either direct or indirect impact of opium derivatives on both bone marrow and RBC progenitor cells and these are in parallel with the results of Chan et al.<sup>38</sup> Further studies should be conducted in order to obtain a better understanding of the mechanisms that induce

these alterations in peripheral blood cells of opium-addicted subjects.

### Conclusion

Our findings showed that opium-addiction, especially when associated with cigarette smoking, induced serious adverse effects on hematological factors and these alterations might lead to imbalance in immune system and acute infection. On the whole, based on the findings of the present study and the results from similar research, opium may have a serious influx on the bone marrow including stem cells and progenitor cells of RBC.

### Conflict of Interests

The Authors have no conflict of interest.

### Acknowledgements

We would like to thank all the participants in our study.

### References

1. Condoluci A, Mazzara C, Zoccoli A, Pezzuto A, Tonini G. Impact of smoking on lung cancer treatment effectiveness: a review. *Future Oncol* 2016; 12(18): 2149-61.
2. Van Tiel E, Peeters PH, Smit HA, Nagelkerke NJ, Van Loon AJ, Grobbee DE, et al. Quitting smoking may restore hematological characteristics within five years. *Ann Epidemiol* 2002; 12(6): 378-88.
3. Tarazi IS, Sirdah MM, El Jeady H, Al Haddad RM. Does cigarette smoking affect the diagnostic reliability of hemoglobin alpha 2 delta 2 (HbA2)? *J Clin Lab Anal* 2008; 22(2): 119-22.
4. Fernandez JA, Prats JM, Artero JV, Mora AC, Farinas AV, Espinal A, et al. Systemic inflammation in 222,841 healthy employed smokers and nonsmokers: white blood cell count and relationship to spirometry. *Tob Induc Dis* 2012; 10(1): 7.
5. Asadikaram G, Sirati-Sabet M, Asiabanha M, Shahrokhi N, Jafarzadeh A, Khaksari M. Hematological changes in opium addicted diabetic rats. *Int J High Risk Behav Addict* 2013; 1(4): 141-8.
6. Schiff PL Jr. Opium and its alkaloids. *Am J Pharm Educ* 2002; 66(2): 186-94.
7. Liang X, Liu R, Chen C, Ji F, Li T. Opioid system modulates the immune function: a review. *Transl Perioper Pain Med* 2016; 1(1): 5-13.
8. Franchi S, Castelli M, Moretti S, Panerai A, Sacerdote P. Evaluation of murine macrophage cytokine production after in vivo morphine treatment. *Methods Mol Biol* 2015; 1230: 253-61.
9. Suzuki S, Chuang LF, Doi RH, Chuang RY. Morphine suppresses lymphocyte apoptosis by blocking p53-mediated death signaling. *Biochem Biophys Res Commun* 2003; 308(4): 802-8.
10. Brown JN, Ortiz GM, Angel TE, Jacobs JM, Gritsenko M, Chan EY, et al. Morphine produces immunosuppressive effects in nonhuman primates at the proteomic and cellular levels. *Mol Cell Proteomics* 2012; 11(9): 605-18.
11. Ke Y, Ye K, Grossniklaus HE, Archer DR, Joshi HC, Kapp JA. Noscapine inhibits tumor growth with little toxicity to normal tissues or inhibition of immune responses. *Cancer Immunol Immunother* 2000; 49(4-5): 217-25.
12. Hiser L, Herrington B, Lobert S. Effect of noscapine and vincristine combination on demyelination and cell proliferation in vitro. *Leuk Lymphoma* 2008; 49(8): 1603-9.
13. Schuler M, Muehlbauer P, Guzzie P, Eastmond DA. Noscapine hydrochloride-induced numerical aberrations in cultured human lymphocytes: a comparison of FISH detection methods and multiple end-points. *Mutagenesis* 2003; 18(3): 235-42.
14. Sajadian S, Vatankhah M, Majdzadeh M, Kouhsari SM, Ghahremani MH, Ostad SN. Cell cycle arrest and apoptogenic properties of opium alkaloids

- noscipine and papaverine on breast cancer stem cells. *Toxicol Mech Methods* 2015; 25(5): 388-95.
15. Svetlecic J, Molteni A, Herndon B. Bronchiolitis obliterans induced by intratracheal papaverine: a novel animal model. *Lung* 2004; 182(2): 119-34.
  16. Troy DB, Remington JP, Beringer P. Remington: the science and practice of pharmacy. Philadelphia, PA: Lippincott Williams and Wilkin; 2006.
  17. Karam GA, Rashidinejad HR, Aghaee MM, Ahmadi J, Rahmani MR, Mahmoodi M, et al. Opium can differently alter blood glucose, sodium and potassium in male and female rats. *Pak J Pharm Sci* 2008; 21(2): 180-4.
  18. Asadikaram G, Asiabanha M, Sayadi A, Jafarzadeh A, Hassanshahi G. Impact of opium on the serum levels of TGF-beta in diabetic, addicted and addicted-diabetic rats. *Iran J Immunol* 2010; 7(3): 186-92.
  19. Nabati S, Asadikaram G, Arababadi MK, Shahabinejad G, Rezaeian M, Mahmoodi M, et al. The plasma levels of the cytokines in opium-addicts and the effects of opium on the cytokines secretion by their lymphocytes. *Immunol Lett* 2013; 152(1): 42-6.
  20. Eisenstein TK, Hilburger ME. Opioid modulation of immune responses: effects on phagocyte and lymphoid cell populations. *J Neuroimmunol* 1998; 83(1-2): 36-44.
  21. Carvalho TT, Borghi SM, Pinho-Ribeiro FA, Mizokami SS, Cunha TM, Ferreira SH, et al. Granulocyte-colony stimulating factor (G-CSF)-induced mechanical hyperalgesia in mice: Role for peripheral TNFalpha, IL-1beta and IL-10. *Eur J Pharmacol* 2015; 749: 62-72.
  22. McCarty MF. Interleukin-6 as a central mediator of cardiovascular risk associated with chronic inflammation, smoking, diabetes, and visceral obesity: down-regulation with essential fatty acids, ethanol and pentoxifylline. *Med Hypotheses* 1999; 52(5): 465-77.
  23. Asakura H, Kawamoto K, Igimi S, Yamamoto S, Makino S. Enhancement of mice susceptibility to infection with *Listeria monocytogenes* by the treatment of morphine. *Microbiol Immunol* 2006; 50(7): 543-7.
  24. Sioud M, Floisand Y. NOD2/CARD15 on bone marrow CD34+ hematopoietic cells mediates induction of cytokines and cell differentiation. *J Leukoc Biol* 2009; 85(6): 939-46.
  25. Vallejo R, de Leon-Casasola O, Benyamin R. Opioid therapy and immunosuppression: a review. *Am J Ther* 2004; 11(5): 354-65.
  26. Stampfli MR, Anderson GP. How cigarette smoke skews immune responses to promote infection, lung disease and cancer. *Nat Rev Immunol* 2009; 9(5): 377-84.
  27. Lavi S, Prasad A, Yang EH, Mathew V, Simari RD, Rihal CS, et al. Smoking is associated with epicardial coronary endothelial dysfunction and elevated white blood cell count in patients with chest pain and early coronary artery disease. *Circulation* 2007; 115(20): 2621-7.
  28. Vulser H, Wiernik E, Tartour E, Thomas F, Pannier B, Czernichow S, et al. Smoking and the Association Between Depressive Symptoms and Absolute Neutrophil Count in the Investigations Preventives et Cliniques Cohort Study. *Psychosom Med* 2015; 77(9): 1039-49.
  29. Smith MR, Kinmonth AL, Luben RN, Bingham S, Day NE, Wareham NJ, et al. Smoking status and differential white cell count in men and women in the EPIC-Norfolk population. *Atherosclerosis* 2003; 169(2): 331-7.
  30. Zhang B, Gao C, Hou Q, Yin J, Xie L, Pu S, et al. Different independent susceptibility markers for first-ever cerebral infarction and myocardial infarction in young patients. *J Neurol* 2012; 259(7): 1420-5.
  31. Seinen ML, van Asseldonk DP, de Boer NK, Losekoot N, Smid K, Mulder CJ, et al. The effect of allopurinol and low-dose thiopurine combination therapy on the activity of three pivotal thiopurine metabolizing enzymes: results from a prospective pharmacological study. *J Crohns Colitis* 2013; 7(10): 812-9.
  32. Madjid M, Awan I, Willerson JT, Casscells SW. Leukocyte count and coronary heart disease: implications for risk assessment. *J Am Coll Cardiol* 2004; 44(10): 1945-56.
  33. Hashemi Domeneh B, Tavakoli N, Jafari N. Blood lead level in opium dependents and its association with anemia: A cross-sectional study from the capital of Iran. *J Res Med Sci* 2014; 19(10): 939-43.
  34. Rezende SM, Lijfering WM, Rosendaal FR, Cannegieter SC. Hematologic variables and venous thrombosis: red cell distribution width and blood monocyte count are associated with an increased risk. *Haematologica* 2014; 99(1): 194-200.
  35. Raval M, Paul A. Cerebral venous thrombosis and venous infarction: case report of a rare initial presentation of smoker's polycythemia. *Case Rep Neurol* 2010; 2(3): 150-6.
  36. White JL, Farrier EG. Method for preparing carbon fuel for smoking articles and product produced thereby (Publication Number: CA 1284025 C). [Patent]. 1991.
  37. Gomperts ED, Forman HJ. Solution of carbon monoxide for treatment of disease, including sickle cell disease (Publication Number: EP 2663193 A1). [Patent]. 2013.
  38. Chan YY, Yang SN, Lin JC, Chang JL, Lin JG, Lo WY. Inflammatory response in heroin addicts undergoing methadone maintenance treatment. *Psychiatry Res* 2015; 226(1): 230-4.

## اثرات اعتیاد به تریاک و مصرف سیگار بر شاخص‌های خونی

غلام‌عباس شهابی‌نژاد<sup>۱</sup>، دکتر مجید سیرتی ثابت<sup>۲</sup>، دکتر محمد کاظمی عرب‌آبادی<sup>۳</sup>، سعیده نباتی<sup>۴</sup>،  
دکتر غلامرضا اسدی کرم<sup>۵</sup>

### مقاله پژوهشی

### چکیده

**مقدمه:** هدف از انجام مطالعه حاضر، ارزیابی اثرات اعتیاد به تریاک و سیگار بر شمارش کامل سلول‌های خونی (Complete blood count) یا CBC) بود.

**روش‌ها:** ۸۶ مرد شامل ۳۱ فرد معتاد سیگاری، ۱۹ فرد معتاد غیر سیگاری، ۱۷ فرد سیگاری غیر معتاد و ۱۹ فرد بدون اعتیاد به تریاک و سیگار در مطالعه حاضر شرکت کردند و آزمایش CBC بر روی خون تمام افراد انجام گرفت.

**یافته‌ها:** میزان WBC (White blood cell)، لنفوسیت‌ها و RBC (Red blood cell) در افراد معتاد سیگاری بالاتر، اما میزان MCV (Mean corpuscular volume) در مقایسه با گروه شاهد (غیر معتاد و غیر سیگاری) کمتر بود. میزان لنفوسیت‌ها در افراد معتاد سیگاری در مقایسه با افراد سیگاری غیر معتاد، بیشتر به دست آمد. تعداد WBC، لنفوسیت‌ها و RBC در افراد معتاد سیگاری و افراد سیگاری غیر معتاد به طور معنی‌داری بالاتر، اما میزان MCV در مقایسه با افراد سیگاری غیر معتاد پایین‌تر مشاهده شد. افراد معتاد سیگاری میزان بالاتری از لنفوسیت‌ها در مقایسه با افراد سیگاری غیر معتاد داشتند. میانگین تعداد لنفوسیت‌ها در افراد معتاد سیگاری نسبت به افراد سیگاری غیر معتاد بیشتر بود. میزان WCB در افراد سیگاری نسبت به گروه شاهد (غیر معتاد و غیر سیگاری) افزایش بیشتری را نشان داد.

**نتیجه‌گیری:** نتایج مطالعه حاضر نشان داد که اعتیاد به تریاک به ویژه وقتی با مصرف سیگار همراه باشد، اثرات مخربی بر شاخص‌های خونی اعمال می‌کند. این تغییرات می‌تواند منجر به افزایش خطر ابتلا به آترواسکلروز، بیماری‌های قلبی-عروقی و عدم تعادل سیستم ایمنی گردد.

**واژگان کلیدی:** مصرف سیگار، اعتیاد، تریاک، CBC

**ارجاع:** شهابی‌نژاد غلام‌عباس، سیرتی ثابت مجید، کاظمی عرب‌آبادی محمد، نباتی سعیده، اسدی کرم غلامرضا. اثرات اعتیاد به تریاک و مصرف سیگار بر شاخص‌های خونی. مجله اعتیاد و سلامت ۱۳۹۵؛ ۸ (۳): ۱۸۵-۱۷۹.

تاریخ پذیرش: ۹۵/۲/۹

تاریخ دریافت: ۹۴/۱۲/۱

۱- مرکز تحقیقات علوم اعصاب، پژوهشکده نوروفارماکولوژی و گروه بیوشیمی، دانشکده پزشکی، دانشگاه علوم پزشکی کرمان، کرمان، ایران

۲- دانشیار، گروه بیوشیمی، دانشکده پزشکی، دانشگاه علوم پزشکی شهید بهشتی، تهران، ایران

۳- استادیار، گروه علوم آزمایشگاهی، دانشکده پیراپزشکی، دانشگاه علوم پزشکی رفسنجان، رفسنجان، ایران

۴- دانشجوی دکتری، گروه بیوشیمی، دانشکده پزشکی، دانشگاه علوم پزشکی شهید بهشتی، تهران، ایران

۵- استاد، مرکز تحقیقات غدد و متابولیسم، پژوهشکده علوم پایه و بالینی فیزیولوژی و گروه بیوشیمی، دانشکده پزشکی، دانشگاه علوم پزشکی کرمان، کرمان، ایران

Email: gh\_asadi@kmu.ac.ir

نویسنده مسئول: دکتر غلامرضا اسدی کرم