



# Establishment of a Bama miniature pig burn model with different burn depths

Xingxia Hao<sup>1,2,3#</sup>, Yunfei Chi<sup>1,2#</sup>, Hailiang Bai<sup>1,2</sup>, Shaozeng Li<sup>4</sup>, Shaofang Han<sup>2</sup>, Hongjie Duan<sup>2</sup>, Yang Chang<sup>2</sup>, Fangchao Hu<sup>1,2</sup>, Bin Wei<sup>1,2</sup>, Jinguang Zheng<sup>1,2</sup>, Jiake Chai<sup>2</sup>, Lingying Liu<sup>3,5</sup>

<sup>1</sup>Chinese PLA Medical School (Chinese PLA General Hospital), Beijing, China; <sup>2</sup>Senior Department of Burns and Plastic Surgery, The Fourth Medical Center of PLA General Hospital, Beijing, China; <sup>3</sup>The Inner Mongolia Medical University, Hohhot, China; <sup>4</sup>Department of Clinical Laboratory, The Fourth Medical Center of PLA General Hospital, Beijing, China; <sup>5</sup>Department of Nutrition, The Fourth Medical Center of PLA General Hospital, Beijing, China

*Contributions:* (I) Conception and design: X Hao, L Liu, Y Chi, J Chai; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: X Hao, L Liu, Y Chi; (V) Data analysis and interpretation: X Hao, Y Chi, H Bai, S Li, S Han, H Duan, Y Chang, F Hu, B Wei, J Zheng, L Liu; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

#These authors contributed equally to this work.

*Correspondence to:* Jiake Chai. Senior Department of Burns and Plastic Surgery, The Fourth Medical Center of PLA General Hospital, No. 51 Fucheng Road, Haidian District, Beijing 100037, China. Email: cjk304@126.com; Lingying Liu. Department of Nutrition, The Fourth Medical Center of PLA General Hospital, No. 51 Fucheng Road, Haidian District, Beijing 100037, China. Email: Liulingyingyisheng@163.com.

**Background:** The skin morphological characteristics of the Bama miniature pig are very similar to those of humans; thus, the Bama miniature pig is an ideal choice for establishing a skin burn model.

**Methods:** In this study, 6 ordinary, male, Bama miniature pigs (weight: 23–28 kg and length: 71–75 cm) were used to establish burn models. A mixture of 1 mg of Ketamine and Sumianxin II was used for Bama miniature pigs anesthetizing, and 1 mg of Pentobarbital sodium was added as necessary. The different burn depths were made using a continuous pressure of 1 kg and contact times of 0 s, 10 s, 15 s, 20 s, 25 s, 30 s, 35 s, 40 s, and 45 s by the newly invented electronic burn instrument. The burned tissues were collected and examined with hematoxylin and eosin (H&E) and Masson staining.

**Results:** Burning for 10–15 s caused a first-degree burn; the blood vessels in the superficial dermis were dilated and congested, and necrosis occurred above the basal layer of the epidermis. Burning for 20–25 s caused a superficial partial-thickness burn; the whole epidermal layer was necrotic, and the collagen fibers were slightly deformed. Burning for 30–35 s caused a deep partial-thickness burn; the whole epidermal layer and dermal layers were necrotic with leukocyte infiltration zones, and the collagen fibers were disordered, degenerated, and necrotized. Burning for 40–45 s caused a third-degree burn; the skin layers and adipose tissues were necrotic, and the thick blood vessels in the skin adipose tissues were full of disintegrated and agglutinated red blood cells.

**Conclusions:** Stable burn depth models of Bama miniature pigs were constructed using a new and innovative electronic burn instrument. Our findings provide a basis for further research on the burn mechanism and evaluations of therapeutic drugs.

**Keywords:** Burn model; burn pathophysiology; Bama miniature pig; skin morphological

Submitted Jul 28, 2022. Accepted for publication Sep 30, 2022.

doi: 10.21037/gs-22-476

View this article at: <https://dx.doi.org/10.21037/gs-22-476>

## Introduction

The incidence of burn injury, for which the elderly and children comprise high-risk groups, is increasing year by year (1). In hospitalized children with accidental injuries, burn injury ranks 2nd among the causes of injury (2). The annual incidence rate of burn injury in China ranges from about 1.5–2%; that is, 20 million people suffer from different degrees of burns, of whom about 5% require hospitalization (3). In addition to having a considerable incidence rate, severe burns also cause disfigurement or even accidental death. Burns can be divided into the following 3 degrees and 4 levels according to damage severity: first-, second- and third-degree burns, of which second-degree burns can be further divided into superficial second-degree burns and deep second-degree burn. First-degree, superficial second-degree, and deep second-degree burns generally take about 5–7 days, 2–3 weeks, and 3–4 weeks to heal, respectively. However, some of the more serious deep second-degree burn and third-degree burns require a longer time to repair, and some cannot even heal naturally, and require surgical skin grafting (4).

There have been significant advances in burn treatments; however, a new, effective treatment mode and drug still need to be developed. A large and ideal mammalian model with different degrees of burns needs to be established to observe skin repair, study the burn mechanism, and evaluate the potential of novel therapeutics and drugs. An animal model should be selected that is similar to the human structure, function, metabolism style, and disease characteristics. To date, mice (5), rats (6), rabbits (7), pigs (8), and dogs (9) have been commonly used to establish burn models; however, each has various problems.

The skin structure of the Bama miniature pig is very similar to that of the human in morphology, physiology, and pharmacology, including in relation to hair density, epidermal thickness, and humoral and metabolic changes in burned skin (10,11). Importantly, the burned skin of both pigs and humans repair through epithelization, so it is an ideal animal model for researching burn mechanism and evaluating novel therapeutics, and the data might provide theoretical foundation for further clinical transformation application. But the model also has a number of disadvantages, including high costs, and difficulties associated with the large size of the animals, and the complexity of the operation.

Depth-burn models with large animals, such as pigs and dogs, are prepared with combustible liquids, such

as absolute ethanol, combustible solids, such as jellied gasoline, thermal burns, such as hot-water burns, steam-induced burns, copper column boiled with hot-water burns, electrothermal burns, flame burns, radiation burns, chemical burns, and inhalation burns (12,13). However, these methods have different limitations, including unsteady temperatures, uncontrollable pressure on the contact surface, and inconsistent burn depth. In this study, a Bama miniature pig burn model with different degrees of burns was prepared using a new innovative electronic burn instrument (ZL201920043476.1). The new electronic burn instrument is characterized by accurate temperature control, uniform depth, and a controllable range of burn injury, and a more stable and reliable burn animal model was established using this burn instrument compared with other method. Our findings provide a basis for further research on the burn mechanism and evaluations of therapeutic drugs. We present the following article in accordance with the ARRIVE reporting checklist (available at <https://gs.amegroups.com/article/view/10.21037/gS-22-476/rc>).


## Methods

### *Animal care*

All the experiments adhered to procedures consistent with the International Guiding Principles for Biomedical Research Involving Animals issued by the Council for the International Organizations of Medical Sciences (CIOMS) and were approved by the Institutional Animal Care and Use Committee at The Fourth Medical Center of PLA General Hospital (the ethics approval ID: 2021KY033-KS001). A protocol was prepared before the study without registration. A total of 6 ordinary male Bama miniature pigs were purchased from the Beijing Strong Century Minipigs Breeding Base. The minipigs were housed at the Laboratory Animal Center of The Fourth Medical Center of PLA General Hospital (Beijing, China), and kept on a 12:12-h light–dark cycle, at room temperature maintained between 20 and 22 with a humidity range of 50–60%, for 1 week before the experiment for adaptation.

### *Establishment of a Bama miniature pig burn model with different degrees*

When the 6 male Bama miniature pigs reached a weight of 23–28 kg and had a mean length of 71–75 cm, they were used to make burn models. Before the experiment, the hair



**Figure 1** Electronic scalding instrument. (A) Schematic diagram. Number 10: burned unit [including number 11 (copper plate with a diameter of 5 cm), number 12 (outer protective layer of steel pipe), and number 13 (handle)]. Number 20: temperature control unit [including number 21 (button for setting temperature), number 22 (temperature display screen), and number 23 (on-off button)]. Number 30: supply electricity unit. (B) Actual structure diagram. (C) The removable copper plate of the scalding instrument.

of each pig was shaved and depilated on the back. Each pig was then anesthetized with an intramuscular injection of 1 mg of Ketamine and Sumianxin II. The anesthesia was maintained with 3% Pentobarbital sodium (0.2 mL/kg) as needed. To obtain consistent and controllable burn wounds, a new electronic burn instrument (ZL201920043476.1) was invented by our team (see *Figure 1*). In this study, 8 round burn wounds with a diameter of 5 cm and different burn degrees were prepared separately on both sides of each pig's back by applying a continuous pressure of 1 kg perpendicular to the spine using the electronic burn instrument at 100 °C. The different burn degrees were determined by the contact time of the electronic burn instrument, which were as follows: 0 s (control), 10 s, 15 s, 20 s, 25 s, 30 s, 35 s, 40 s, and 45 s. The physiological state and burn wounds of the Bama miniature pigs were then closely observed.

#### ***Specimen collection and preparation of formalin-fixed paraffin-embedded (FFPE) tissue slides***

The skin burn tissue on both sides of each pig's back was cut at 24 h after the burn to the muscular layer. The cut site was in the middle of the burn wound. The skin samples were divided into small pieces, rinsed with phosphate buffered solution and fixed in 4% paraformaldehyde for 7 d at room temperature. Subsequently, the skin burn tissue was immersed in gradient ethanol, ethanol-xylene mixture, pure xylene, and pre-melted paraffin wax 1 by 1 to embed the skin burn tissue into paraffin. The paraffin specimens were sectioned in a plane perpendicular to the incision, while the section thickness was 5  $\mu$ m.


#### ***Hematoxylin and eosin (H&E) staining***

The sections of burn tissue for 0 s (control), 10 s, 15 s, 20 s, 25 s, 30 s, 35 s, 40 s, and 45 s were successively immersed 1 by 1 in dimethylbenzene, 2 times for 10 min each time. The deparaffinized sections were immersed in gradient ethanol and then rehydrated by flowing distilled water, 3 times for 1 min each time, stained with hematoxylin dye solution for 5–10 min, and stained with eosin solution for 5 min, dehydrated in gradient ethanol, and hyalinized in pure xylene 3 times for 2 min each time, and finally sealed by the appropriate amount of neutral gum. After drying, the skin morphology was observed under an optical microscope (Olympus, USA).

#### ***Masson staining***

The pre-treatment of the sections of burn tissue was consistent with that of H&E staining. The pre-treated sections were immersed in potassium dichromate overnight at room temperature and then stained in hematoxylin solution for 3–5 minutes. After being washed with flowing distilled water for 1–3 min, the blue was reversed by immersing them in ethanol-hydrochloric acid solution. Subsequently, the sections were transferred into a dye solution composed of ponceau and acid fuchsin for 10–15 min, phosphomolybdic acid aqueous solution for 3–5 min, aniline blue staining solution for 6 min, glacial acetic acid (1% v/v) for 2 min, and 95% ethanol and xylene for 5 s. The sections were dried in a ventilated place, and then covered with a cover slip with neutral balsam. After drying, the sections were observed under an optical microscope





**Figure 2** The establishment of pig models with different burn depths. (A) Diameter of copper plate: 5 cm. (B) The earmark of the Bama miniature pig. (C) The burn model establishment: scalding conditions as follows: 100 °C; a continuous pressure of 1 kg perpendicular to the spine, and a contact time of 0 s, 10 s, 15 s, 20 s, 25 s, 30 s, 35 s, 40 s, and 45 s, respectively.

(Olympus, USA); blue indicated normal collagen fiber structure, and red indicated collagen fiber degeneration and necrosis.

## Results

### *A new electronic burn instrument*

Previous studies have shown that the methods for making burn models have the disadvantages of an uncontrollable and unstable temperature, contact pressure, and burn area, which result in inconsistent burn degrees and poor experimental repeatability. Thus, a portable electronic burn instrument was invented according to the specific experimental requirements, for which a patent has been applied for in China (No. ZL 201920043476.1). The mechanical structure of this instrument is shown in *Figure 1A,1B*. The copper plate of the electronic burn instrument that comes in direct contact with the pig skin is removable, and it uses double electron tubes for heating, and thus has a rapid temperature rise of <1 min. The temperature of the burn components is constant, and the change is very low after contact with the pig skin and can be controlled within  $\pm 0.1$  °C. The temperature control element is removable and has a variety of burn components, of which the shape can be round or square, and the size of burned component ranges from 1–10 cm (see *Figure 1C*). The contact pressure of the electronic burn instrument on the pig skin can also be adjusted to establish pig models with different burn degrees.

Thus, the new electronic burn instrument solved the


problems related to an accurate temperature, pressure control and a consistent burn degree, and thus ensured the good repeatability and consistency of the experiment.

### *Preparation of the Bama miniature pig burn model*

In total, 6 Bama miniature pigs were used to establish the structure burn model using the new electronic burn instrument. The injury conditions were as follows: 100 °C, continuous pressure of 1 kg perpendicular to the spine, a wound surface diameter of 5 cm (see *Figure 2A*), contact for 0 s (control), 10 s, 15 s, 20 s, 25 s, 30 s, 35 s, 40 s, and 45 s, respectively. Finally, we made 8 wounds with different burn degrees on both sides of the back of each pig with its own earmark (see *Figure 2B*). As *Figure 2C* shows, the wound sizes prepared at the different contact times were nearly the same and the wound edges were neat without ulceration and infection. No obvious blisters were observed to form in any of the burn wound tissues due to the thick cuticle of the pig skin, but the burn marks of pig skin gradually deepened with the extension of the burn time.

### *H&E staining*

To further determine the depth of the burn, we observed the histopathological changes of the burned skin tissues by H&E staining detection. As *Figure 3A* shows, for normal skin (i.e., skin with a burn time of 0 s), the skin structure maintained its integrity and the collagen fibers were



**Figure 3** H&E staining of the Bama miniature pigs with different burn depths at the following different contact times: (A-I) 0 s, 10 s, 15 s, 20 s, 25 s, 30 s, 35 s, 40 s, and 45 s, respectively (optical microscope,  $\times 50$  and  $\times 400$ ). H&E, hematoxylin and eosin.

arranged densely and orderly. As *Figure 3B,3C* show, for the first-degree burns (i.e., skin treated at 100 °C for 10 and 15 s), the blood vessels in the superficial dermis were dilated and congested, and necrosis occurred above the basal layer of the epidermis. As *Figure 3D,3E* show, for the superficial partial-thickness burns (i.e., skin treated at 100 °C for 20 and 25 s), the whole epidermal layer was necrotic, and the dermal papilla layer was also severely damaged. For the deep partial-thickness burns, (i.e., skin treated at 100 °C for 30 and 35 s), the epidermis and dermis were necrotic with leukocyte infiltration zones, but the skin appendages in the deep dermis remained alive (see *Figure 3F,3G*). For the third-degree burns, (i.e., skin treated at 100 °C for 40 and 45 s), the skin layer and adipose tissue were all necrotic, thick blood vessels could be observed in the skin adipose tissue, and were filled with disintegrating and agglutinating red blood cells (see *Figure 3H,3I*). Thus, treatment at 100 °C for 30 s or more resulted in deeper skin burn injuries.

### Masson staining


Masson staining was performed to observe the collagen fiber structure changes of the burned skin. Consistent with the H&E staining results, the Masson staining results showed that the epidermis and dermis structure of the normal skin was also complete, and the skin injury became

more and more serious with the extension of the burn time. The collagen fibers were arranged neatly in normal skin (i.e., skin treated at 100 °C for 0 s; see *Figure 4A*), and there was almost no damage to the collagen fibers in the first-degree burns. Consistent with the H&E results, the Masson results also clearly showed that the blood vessels in the superficial dermis were dilated and congested, and necrosis occurred above the basal layer of epidermis in the first-degree burns (i.e., skin treated at 100 °C for 10 and 15 s; see *Figure 4B,4C*). In the superficial partial-thickness burns (i.e., skin treated at 100 °C for 20 and 25 s), the collagen fibers were slightly disordered and injured (see *Figure 4D,4E*). In the deep partial-thickness burns (i.e., skin treated at 100 °C for 30 and 35 s), the structure of collagen fibers was partially degenerated and necrotized (see *Figure 4F,4G*). In the third-degree burns (i.e., skin treated at 100 °C for 40 and 45 s), the collagen fibers were severely degenerated, or even completely necrotized (see *Figure 4H,4I*). The Masson staining results indicated that the collagen fibers were severely damaged with treatment at 100 °C for 30 s or more.

### Discussion

Burns are a global health problem and cause serious physical and psychological harm to patients, especially children and the elderly. At present, about 265,000 people





**Figure 4** Masson staining of the Bama miniature pigs with different burn depths at the following different contact time: (A-I) 0 s, 10 s, 15 s, 20 s, 25 s, 30 s, 35 s, 40 s, and 45 s, respectively (optical microscope,  $\times 50$  and  $\times 400$ ).

worldwide have died of burns (14). It has been reported that the incidence of burns was 0.19% in the United States, 2.43% in Southeast Asia, and 1.87% in the Mediterranean (15-18). In China, >20 million patients suffer from different degrees of burns (19). With the continuous development and improvement of burn treatments, most burn patients can enter the rehabilitation period smoothly. However, the dysfunction and disfigurements caused by scar contracture still occur frequently in the process of rehabilitation (20). Thus, the development of effective burn treatment methods and drugs is a research hotspot. A burn animal model provides the necessary basis for burn research. The selection of an appropriate burn animal model largely depends on practical problems, such as the experimental purpose, experimental condition, and experimental cost.

Bama miniature pigs are ideal to establish experimental animal models to study skin burns, as they are similar to humans in terms of their body size, their physiological condition, and disease development. Most of the skin of Bama miniature pig is covered with white hair, which is convenient for experimental observation and operation (11). According to Liu, the skin morphological characteristics of the Bama miniature pig are similar to those of the human, including in relation to the thickness of the skin layer, the development of the surface vascular system, the structure of the dermal epidermal interface, and the extracellular

matrix (10). However, the skin of Bama miniature pigs lacks many antigenic determinants corresponding to those of the skin of humans, and there is no cross reaction with human antiserum, which is helpful to further study the immunological characteristics of human skin (10). Additionally, the drug metabolism process of Bama miniature pig skin is similar to that of human skin, so the applicability, dose safety, and toxicological characteristics of burn drugs can be evaluated well (21). Thus, Bama miniature pig skin has unique value in skin research.

In the establishment of an animal burn model, it is difficult to determine the precise area and depth of injury due to differences in each animal body shape and skin structure/thickness. Thus, it is necessary to individualize the injury and clarify the burn depth in combination with wound conditions and pathological examinations. Previously, there was no standardized, high-precision and controllable effective method for preparing a Bama miniature pig burn model. Common methods for preparing burn models of pigs, dogs, and other large animals include thermal burns (hot-water burns, steam burns, and hot-metal burns), electric-light source burns, flame burns (burns with combustible liquids, such as absolute ethanol, and combustible solids such as solidified gasoline powder), electric burns, radiation burns, chemical burns, and inhalation burn. However, these methods have various

limitations, such as unstable temperatures, uncontrollable contact pressures, and inconsistent burn depths (12,13). For example, when a copper column heated by boiling water comes in contact with animal skin, the copper column cools, and thus the temperature drops. Similarly, a combustible liquid may extend to an unexpected, undesignated skin area, making it difficult to control the burn area. Additionally, the previous methods for preparing Bama miniature pig burn models were performed mechanically according to the preparation conditions of other animal models, which have not been explored or improved.

To overcome these defects, a new electronic burn instrument was invented. The heated copper plate of the new electronic burn instrument comes in direct contact with the pig skin, so the area and location of burns are strictly controlled. The copper plate is removable, the size ranges from 1–10 cm, and can be round or square in shape; thus, the size of the burn area can be flexibly adjusted, preventing an insufficient burn depth caused by too small a burn area, and a systemic reaction caused by a too large burn area. The copper plate of the new electronic burn instrument that comes in direct contact with the pig skin is heated by double electron tubes. The temperature of the whole copper plate is uniform, constant, and changes little after coming in contact with the pig skin, and can be controlled within  $\pm 0.1$  °C. Additionally, the contact pressure of the copper plate on the pig skin can be adjusted according to the experimental purpose and physiological state of the Bama miniature pigs. The precise copper plate temperature and contact pressure on the pig skin ensure the consistency and controllability of the burn depth.

The new electronic burn instrument is suitable for preparing burn models of Bama miniature pigs, and other large animals. Additionally, the preparation conditions of an animal burn model, including the copper plate temperature, contact pressure on the skin, and burn area, need to be explored to prepare the different burn depths to meet the experimental designs of different studies. Further, the histopathological changes after injury at different times should be described to stabilize the animal model for further mechanism research and new treatment evaluations (22). The deep partial-thickness and third-degree burns belong to deep burns in clinic, and burned wound healing is relatively slow, leaves severe scars, and may require skin grafting operation. Burn severity generally directly correlates to the total body surface area (TBSA) and burned deep of the injured tissue. Above 30% TBSA

of deep burns in adult and 10% TBSA of deep burns in children, they are considered as severe burn and can induce multiple organ injury and excessive systemic inflammation. Severe burn is a complex and severe traumatic injury with high mortality in the acute phase, and prompt and appropriate treatment of severe burns is necessary, especially in the phase. In this study, the results showed that the preparation condition of 100 °C, a continuous pressure of 1 kg perpendicular to the spine, and a contact time with the pig skin of 30 s or more was optimum for establishing deeper burns in the Bama miniature pigs. Notably, pre-experiments were necessary to adjust the burn conditions according to the physiological state and age of the Bama miniature pigs and the experimental purpose.

In conclusion, this study provided an accurate and controllable stabilization method for the preparation of Bama miniature pig models with different burn depths using a newly invented electronic burn instrument. Further, these burn models provide a basis for further research on the burn mechanism and evaluations of therapeutic drugs.

## Acknowledgments

*Funding:* This work was supported by the Major Project of Military Logistical Support Department (Nos. ALB19J001, AWS15J003), the Natural Science Foundation of Beijing (No. 7222179), and the National Natural Science Foundation of China (No. 81772067), Health and Technology of Inner Mongolia (No. 202202146).

## Footnote

*Reporting Checklist:* The authors have completed the ARRIVE reporting checklist. Available at <https://gs.amegroups.com/article/view/10.21037/gS-22-476/rc>

*Data Sharing Statement:* Available at <https://gs.amegroups.com/article/view/10.21037/gS-22-476/dss>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://gs.amegroups.com/article/view/10.21037/gS-22-476/coif>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are

appropriately investigated and resolved. All the experiments adhered to procedures consistent with the International Guiding Principles for Biomedical Research Involving Animals issued by the Council for the International Organizations of Medical Sciences (CIOMS) and were approved by the Institutional Animal Care and Use Committee at The Fourth Medical Center of PLA General Hospital (the ethics approval ID: 2021KY033-KS001).

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Riedlinger DI, Jennings PA, Edgar DW, et al. Scald burns in children aged 14 and younger in Australia and New Zealand—an analysis based on the Burn Registry of Australia and New Zealand (BRANZ). *Burns* 2015;41:462-8.
- Jin Y, Duan L, Wang Y, et al. 844 Trend and current characteristics of burn from chinese national injury surveillance system from 2006 to 2013. *Inj Prev* 2016;22:4.
- Sheng Z, Guo Z. Editors. Critical burn treatment and rehabilitation. First edition. Beijing: Science Press; 2000:136-7.
- Itakussu EY, Morita AA, Kakitsuka EE, et al. Instruments to assess function or functionality in adults after a burn injury: A systematic review. *Burns* 2021;47:999-1011.
- Zhang T, Zhang R, Xu B, et al. Spinal endomorphins attenuate burn-injury pain in male mice by inhibiting p38 MAPK signaling pathway through the mu-opioid receptor. *Eur J Pharmacol* 2021;903:174139.
- Shen C. 650 Exploring the Mechanism of Wound Healing in Rats of Different Ages Using a Burn Model. *Journal of Burn Care & Research* 2020;Supplement\_1:S171.
- Djerrou Z, Maameri Z, Hamdi-Pacha Y, et al. Effect of virgin fatty oil of Pistacia lentiscus on experimental burn wound's healing in rabbits. *Afr J Tradit Complement Altern Med* 2010;7:258-63.
- Sun Y, Cao Y, Zhao R, et al. The Role of Autologous PRP on Deep Partial-Thickness Burn Wound Healing in Bama Pigs. *J Burn Care Res* 2020;41:657-62.
- Atiba A, Ghazy A. The Effects of Topical Dimethyle Sulfoxide on Second-Degree Burn Wound Healing in Dogs. *Alex J Vet Sci* 2015;45:6-12.
- Liu Y, Chen JY, Shang HT, et al. Light microscopic, electron microscopic, and immunohistochemical comparison of Bama minipig (*Sus scrofa domestica*) and human skin. *Comp Med* 2010;60:142-8.
- Chen JY, Hu J, Wei H. Comparative Study on Skin between BAMA Miniature Pig and Human. *Chinese Journal of Comparative Medicine* 2006;16:288-91.
- Borman H, Bilezikci B, Maral T. The use of superficial and deep partial-thickness burned tissue for flap fabrication. *Burns* 2007;1:S53-4.
- Shukla SK, Sharma AK, Shaw P, et al. Creation of rapid and reproducible burn in animal model with a newly developed burn device. *Burns* 2020;46:1142-9.
- Forbinake NA, Ohandza CS, Fai KN, et al. Mortality analysis of burns in a developing country: a CAMEROONIAN experience. *BMC Public Health* 2020;20:1269.
- Sorenson TJ, Mohr WJ, Mahajan AY. Pediatric Hand Burns Requiring Emergency Care in the United States. *J Burn Care Res* 2021. [Epub ahead of print]. doi: 10.1093/jbcr/irab144.
- Van Niekerk A, Rode H, Laflamme L. Incidence and patterns of childhood burn injuries in the Western Cape, South Africa. *Burns* 2004;30:341-7.
- Othman N, Kendrick D. Epidemiology of burn injuries in the East Mediterranean Region: a systematic review. *BMC Public Health* 2010;10:83.
- Van Yperen DT, Van Lieshout EMM, Verhofstad MHJ, et al. Epidemiology of burn patients admitted in the Netherlands: a nationwide registry study investigating incidence rates and hospital admission from 2014 to 2018. *Eur J Trauma Emerg Surg* 2022;48:2029-38.
- Zhu L, Zhang H, Shi F, et al. Epidemiology and outcome analysis of scalds in children caused by "guo lian kang": an 11-year review in a burn center in China. *Burns* 2015;41:289-96.
- Hawkins H, Jay J, Finnerty CC. Pathophysiology of the Burn Scar. *Total Burn Care (Fifth Edition)* 2018:466-75.e3.
- Liu Y, Zeng BH, Shang HT, et al. Bama miniature pigs (*Sus scrofa domestica*) as a model for drug evaluation for



- humans: comparison of in vitro metabolism and in vivo pharmacokinetics of lovastatin. *Comp Med* 2008;58:580-7.
22. Davenport L, Dobson G, Letson H. A new model for standardising and treating thermal injury in the rat.

*MethodsX* 2019;6:2021-7.

(English Language Editor: L. Huleatt)

**Cite this article as:** Hao X, Chi Y, Bai H, Li S, Han S, Duan H, Chang Y, Hu F, Wei B, Zheng J, Chai J, Liu L. Establishment of a Bama miniature pig burn model with different burn depths. *Gland Surg* 2022;11(10):1647-1655. doi: 10.21037/gs-22-476