

Introduction: The various restrictions and physical distancing introduced in Russia in early months of COVID-19 pandemic could have a particular impact on people who use alcohol and create new needs for prevention and treatment of associated disorders.

Objectives: The study was aimed at assessing changes in alcohol consumption among users of online social networks in Russia.

Methods: During June-September 2020, 1518 adult users (18+) of the most popular social networks in Russia (Odnoklassniki, VKontakte, Facebook, Twitter), completed an anonymous online survey regarding alcohol use changes in the first months of COVID-19 pandemic. Binary logistic regression was used to estimate associations of increased alcohol consumption with socio-demographic and pandemic-related factors.

Results: 35.4% of men and 25.6% of women increased their frequency of drinking; 24.9% of men and 17.7% of women increased their typical one-time volume of alcohol consumption, and 28.5% of men and 27.9% of women increased frequency of heavy episodic drinking in the early months of pandemic. After mutual adjustment of independent variables, age 18-29 years (OR=1.710; 95% CI 1.002-2.917), very strong restrictions of everyday life (3.127; 1.011-9.675) and severe negative consequences in relation to professional or financial situation due to spread of SARS-CoV-2 (2.247; 1.131-4.465) were positively associated with increase in frequency of drinking. The odds of increase in frequency of heavy episodic drinking were more than twice higher (2.329; 1.001-5.428) among those who experienced severe negative consequences to their professional/financial situation. Higher typical frequency, larger one-time volume of alcohol use, and higher frequency of heavy episodic drinking before pandemic were positively associated with increase of alcohol consumption in early months of pandemic.

Conclusions: Timely monitoring of changes in alcohol consumption during pandemic can allow prevention of alcohol-related disorders, including mental disorders, among users of online social networks.

Disclosure: No significant relationships.

Keywords: social networks; Russia; alcohol; Covid-19

EPP0580

COVID-19: first and second wave impact on outpatient service users of FACT or autism teams in the Netherlands

S. Castelein^{1,2*}, I.D.C. Van Balkom³ and J. Bruins¹

¹Lentis Psychiatric Institute, Lentis Research, Groningen, Netherlands;

²University of Groningen, Clinical Psychology, Groningen,

Netherlands and ³Lentis Psychiatric Institute, Autism Team Northern-netherlands, Jonx Department Of (youth) Mental Health And Autism, Zuidlaren, Netherlands

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.789

Introduction: Most research on COVID-19 effects has focused on the general population. Here we measure its impact on Dutch FACT and autism outpatient service users during both waves.

Objectives: This study aimed to: 1) investigate participants' mental health, 2) assess experiences with outpatient services, and 3) assess respondents' experiences with governmental measures in the Netherlands during the first and second wave of COVID-19.

Methods: Respondents (wave 1: n=100; wave 2: n=150) reported on mental health, experiences with outpatient care, government measures and information services in an online survey.

Results: Findings demonstrate happiness was rated an average of 6 out of 10, 70% of respondents scored below average on resilience, positive consequences for mental health (ordered world, reflection time) during both waves were similar, and prominent negative consequences included decreased social interactions and increased or new problems regarding mental health and daily functioning from wave 1-2. Lifestyle changed in 50% in both waves, although only slightly attributed to the pandemic. Substance use during both waves hardly changed. Mental healthcare continuation was highly appreciated in both waves (75-80% scored ≥ 7 on 10-point scale). (Video)calling was the most frequently mentioned positive care experience; missing face-to-face contact with care providers considered most negative. COVID-19 measures were less doable in the second wave. Vaccination willingness approximated 70%.

Conclusions: Results show a nuanced, but clear picture of experiences during both waves. Continuation of services through telehealth was well-received. Monitoring of long-term impact is needed.

Disclosure: No significant relationships.

Keywords: Covid-19; TeleHealth; Psychosis; government measures

EPP0581

Assessment of Relapse in Patients with Schizophrenia during COVID-19 pandemic

E. Abdelfattah^{1*}, M. Bassiony^{1,2}, M. Sehlo¹, A. Zayed¹ and S. Atwa¹

¹Faculty of medicine, Zagazig University, Psychiatry, Zagazig, Egypt

and ²Faculty of medicine Zagazig university, Psychiatry, Zagazig, Egypt

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.790

Introduction: People with schizophrenia could be more substantially influenced by the emotional stress brought on by the COVID-19 pandemic, resulting in relapses or worsening of an already existing mental health condition because of the high susceptibility to stress.

Objectives: This study aimed to assess the presence of relapse and its risk factors among patients with schizophrenia during the era of COVID 19 pandemic.

Methods: This study included 90 adults who met DSM-5 criteria for schizophrenia and were diagnosed by (SCID-I) Arabic version and who are following up at the outpatient psychiatric clinic, Zagazig University Hospital, Egypt, Positive and Negative Syndrome Scale (PANSS) measuring severity of symptoms, Compliance Rating Scale (CRS) and World Health Organization Disability Assessment Schedule 2.0 (WHODAS 2.0). scales were conducted on those patients before COVID-19 pandemic from January to April 2019 and repeated on September to November 2020 during COVID 19 pandemic to compare clinical parameters between those two periods as to detect any deterioration in their clinical state.

Results: The mean score of compliance rating scale (CRS) was decreased after COVID-19 pandemic compared to before COVID 19 ($P < 0.001$). The mean score of PANSS scale positive subscale (P) had increased after COVID-19 pandemic compared to before COVID 19 ($P < 0.001$).

Conclusions: There was a deterioration of the clinical state of schizophrenic patients during COVID-19 pandemic especially the positive symptoms and following up the news about COVID-