

Interleukin-17 activates JAK2/STAT3, PI3K/Akt and nuclear factor- κ B signaling pathway to promote the tumorigenesis of cervical cancer

YANFEI BAI^{1*}, HAITAO LI^{2*} and RUI LV³

¹Department of Obstetrics and Gynecology, Lanzhou Second People's Hospital, Lanzhou, Gansu 730000;

²Department of Gynecology, Affiliated Hospital of Gansu Medical College, Lanzhou, Gansu 744000;

³Department of Gynecologic Oncology, Gansu Provincial Cancer Hospital, Lanzhou, Gansu 730030, P.R. China

Received August 25, 2020; Accepted April 27, 2021

DOI: 10.3892/etm.2021.10726

Abstract. Interleukin (IL)-17 has been regarded as a significant factor in inflammation. In addition, IL-17 is known to be involved in the progression of cancers; however, the function of IL-17 in cervical cancer remains unclear. In the present study, cell viability was detected by Cell Counting Kit-8 assay. Quantitative PCR and western blotting were performed to detect gene and protein expression levels, respectively, in cancer cells or tissues. Ki-67 staining was used to evaluate cell proliferation. Wound-healing assay was used to detect cell migration. Moreover, Transwell assay was performed to investigate the invasion of cervical cancer cells. The results revealed that IL-17 significantly promoted the proliferation of cervical cancer cells. Additionally, IL-17 notably enhanced the migration and invasion of cervical cancer cells *in vitro*. IL-17 promoted the progression of cervical cancer via the activation of JAK2/STAT3 and PI3K/Akt/NF- κ B signaling. In conclusion, IL-17 was a key regulator during the progression of cervical cancer through the JAK2/STAT3 and PI3K/Akt/nuclear factor- κ B signaling pathway, which may serve as a novel target for the treatment of cervical cancer.

Introduction

Cervical cancer ranks third in terms of incidence of malignant tumors worldwide, and is the most frequent type of gynecological cancer in developing countries (1-3). The increasing trend in cervical cancer in developing countries is attributed to the

early beginning of sexual activities, certain sexual behaviors such as high number of partners, early age at first intercourse, infrequent use of condoms, multiple pregnancies with *Chlamydia* association, and immunosuppression with human immunodeficiency virus, which is associated with higher risk of infection by human papillomavirus (HPV) (4,5). In USA, the HPV16 and 18 types are detected in 70% of high-grade squamous intraepithelial lesions, as well as in invasive cervical cancer cases (6,7). To date, chemotherapy and surgery are the two major strategies for treating cervical cancer (8). However, the prognosis of cervical cancer remains poor. Thus, it is urgent to identify novel methods for the treatment of cervical cancer.

Recent studies have shown that interleukin (IL)-17 plays an important role in cervical cancer (9,10). IL-17A is a member of the IL-17 family, which has been regarded as a pro-inflammatory cytokine (11). In addition, IL-17 is secreted by various cells, including T helper cells, CD8⁺ T cells, $\gamma\delta$ T cells and natural killer cells in the tumor microenvironment (12). IL-17 and its receptor are expressed in a variety of cell types, including fibroblasts and tumor cells, leading to the secretion of pro-inflammatory cytokines, such as IL-6, various chemokines and metalloproteinases (13). Previous studies have shown that an inflammatory environment may lead to tumor growth by generating tumor-promoting cytokines, decreasing cytotoxic T cells and developing myelogenous inhibitory cells, and further production can promote tumor growth (14-16). Besides, previous findings have revealed that the risk of cervical cancer is associated with IL-17 polymorphism in both Chinese and Western populations (17,18). However, the mechanism by which IL-17 modulates the development of cervical cancer remains unclear.

Thus, the present study aimed to explore the potential molecular mechanism of IL-17 in cervical cancer. The results may provide experimental basis for the possibility of using IL-17 as a key marker to predict prognosis of cervical cancer.

Materials and methods

Sample collection. In total, 30 pairs of cervical cancer samples and adjacent normal tissues were collected from Gansu Provincial Cancer Hospital between June 2018 and June 2019.

Correspondence to: Dr Rui Lv, Department of Gynecologic Oncology, Gansu Provincial Cancer Hospital, 2 Xiaoxihu East Street, Lanzhou, Gansu 730030, P.R. China
E-mail: ruilv811@163.com

*Contributed equally

Key words: interleukin-17, cervical cancer, JAK2/STAT3, PI3K/Akt, nuclear factor- κ B

The clinical and pathological data of these patients (n=30) were collected with their written informed consent. The patient exclusion criteria were as follows: i) Patients (women) who suffered from other diseases and were currently under treatment; ii) pregnant and lactating women; iii) patients allergic to probiotics or have used/are using antibiotics recently; and iv) alcoholics (people who drink ≥ 5 bottles of beer at a time, or the alcohol content in the blood reaches ≥ 0.08). The patient inclusion criteria were as follow: Women (≥ 18 years old) who have been diagnosed with cervical cancer and have undergone surgery. Each tissue sample was stored at -80°C until RNA extraction. In addition, the serum samples were collected from the patients. The present study was approved by the Ethics Committee of Gansu Provincial Cancer Hospital. The distribution of age and sex (30 females; mean age, 52 years; age range, 32-68 years) among the patients with cervical cancer was presented in Table I.

Meanwhile, serum was also collected from healthy donors (n=50; age, 21-58 years; sex, 28 males and 22 females). The informed consent was also obtained from healthy individuals for blood donation in the present study. The inclusion criteria of individuals without cervical cancer as control blood donors were as follows: i) Aged from 20-60 years old; and ii) no history of cancer.

Cell culture. The HeLa cell line was obtained from the Shanghai Cell Bank of Chinese Academy of Sciences. Cells were cultured in DMEM (Thermo Fisher Scientific, Inc.) with 10% fetal bovine serum (FBS; Thermo Fisher Scientific, Inc.), 1% penicillin (Thermo Fisher Scientific, Inc.) and streptomycin (Thermo Fisher Scientific, Inc.) at 37°C in the presence of 5% CO_2 .

Cell transfection. HeLa cells were seeded at a density of 3×10^5 cells/well in a 6-well plate and cultured until 70% confluence. Then, the cells were transfected with small interfering RNA (si)STAT3 (10 nM), siJAK2 (10 nM) or negative control (empty vector, siNC, 10 nM) using Lipofectamine[®] 2000 reagent (Thermo Fisher Scientific, Inc.). For siRNA knockdown, the sequence of siRNA targeting JAK2 (siJAK2) or STAT3 (siSTAT3) was designed and synthesized from Shanghai GenePharma Co., Ltd. The efficiency of transfection was detected by reverse transcription-quantitative PCR (RT-qPCR). The sequences of siRNAs were as follows: siNC, 5'-ACGUGACACGUUCGGAGAAUU-3'; siJAK2, 5'-ATCATGUUUGAGACCUUAAA-3'; siSTAT3, 5'-CUUUGAGGUCAGCCGACUCU-3'. After 24 h of transfection, transfected cells were used in subsequent experiments.

RT-qPCR. Total RNAs were extracted from tissues or cell lines with TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.). RT-qPCR was conducted with PrimeScript RT Reagent kit (Takara Bio, Inc.) and SYBR Premix Ex Taq II kit (Takara Bio, Inc.). The temperature and duration of RT were as follows: 37°C for 60 min and 85°C for 5 min. The thermocycling conditions were as follows: Initial denaturation for 10 min at 95°C ; 40 cycles of 95°C for 15 sec and 60°C for 30 sec; and final extension for 1 min at 60°C . The primers were purchased from Nanjing Jinsirui Biotechnology Co., Ltd. β -actin was used as the internal control. The primers were

as follows: STAT3, 5'-CATCCTGAAGCTGACCCAGG-3'; STAT3 reverse, 5'-TCCTCACATGGGGGAGGTAG-3'; JAK2 forward, 5'-GAGACAACCTGTGACGGGCTT-3'; JAK2 reverse, 5'-GCTCAGCTCCCACTCACATC-3'; IL-17 forward, 5'-CCTTGGAAATCTCCACCGCAA-3'; IL-17 reverse, 5'-GAGCTCTTAGGCCACATGGT-3'; IL-17A forward, 5'-CTACAACCGATCCACCTCACC-3'; IL-17A reverse, 5'-AGCCCACGGACACCAGTATC-3'; IL-17F forward, 5'-CTGTGCCAGGAGGTAGTATGA-3'; IL-17F reverse, 5'-TTGATGCAGCCCAAGTTCCTA-3'; β -actin forward, 5'-GTCCACCGCAAATGCTTCTA-3'; and β -actin reverse, 5'-TGCTGTCACCTTCACCGTTC-3'. The $2^{-\Delta\Delta\text{Cq}}$ method (19) was used to measure relative expression.

Wound-healing assay. HeLa cells (5×10^3 per well) were plated into a 24-well Cell Culture Cluster. Once cells reached 80-90% confluence, the layer of cells was scratched perpendicular with a small pipette head. After washing with PBS for 3 times, serum-free medium was used for further culture, and the scratch widths at 0 and 24 h were recorded under an optical light microscope (magnification, x200). The experiment was repeated 3 times.

Enzyme-linked immunosorbent assay (ELISA). The levels of IL-17 in tissues of patients were detected using an ELISA kit (Hangzhou Multisciences Biotech Co., Ltd., cat. no. 70-EK117/2-96), according to the manufacturer's instructions.

Transwell assay. For cell invasion analysis, Transwell assay was performed. The upper chamber was pre-treated with $100 \mu\text{l}$ Matrigel. HeLa cells were seeded into the upper chamber in medium with 1% FBS, and the density was adjusted to $\sim 1.0 \times 10^6$ cells per chamber. RPMI-1640 medium with 10% FBS was added to the lower chamber. After 24 h of incubation at 37°C , the Transwell chamber was rinsed twice with PBS (5 min each time), fixed with 5% glutaraldehyde at 4°C , stained with 0.1% crystal violet at room temperature for 30 min, washed twice with PBS and observed under an optical light microscope (magnification, x200). The number of cells invading the Matrigel was regarded to represent the invasion ability.

Cell Counting Kit-8 (CCK-8) assay. HeLa cells were seeded in 96-well plates (5×10^3 per well) overnight. Then, cells were treated with 0, 5, 10, 50 or 50 ng/ml IL-17 for 72 h. Next, $10 \mu\text{l}$ CCK-8 reagent was added to each well and further incubated for 2 h at 37°C . Finally, the absorbance was measured at 450 nm using a microplate reader (Thermo Fisher Scientific, Inc.).

Western blotting. Total protein was isolated from tissue or cell lysates using RIPA buffer (Shanghai GenePharma Co., Ltd.), and quantified using a BCA protein assay kit (Beyotime Institute of Biotechnology). Proteins ($30 \mu\text{g/lane}$) were resolved on 10% SDS-PAGE gel, and then transferred to PVDF membranes (Bio-Rad Laboratories, Inc.). After blocking with 5% skimmed milk for 1 h at room temperature, the membranes were incubated with primary antibodies at 4°C overnight, and then incubated with an HRP-conjugated secondary anti-rabbit antibody (1:5,000; cat. no. ab7090; Abcam) at

Table I. Distribution of sex and age among patients with cervical cancer.

Sex	Age, years	Stage
Female	53	IIA2
Female	52	IIA2
Female	61	IIA1
Female	55	IIA1
Female	57	IIA1
Female	48	IIA2
Female	55	IIA1
Female	61	IB2
Female	49	IB1
Female	61	IIA1
Female	62	IIA2
Female	52	IB1
Female	46	IB2
Female	61	IIA1
Female	32	IIA2
Female	42	IIA1
Female	56	IB1
Female	51	IB2
Female	56	IIA1
Female	45	IIA1
Female	50	IIA1
Female	34	IB2
Female	55	IIA1
Female	55	IB1
Female	53	IIA1
Female	44	IB1
Female	68	IIA1
Female	41	IB1
Female	59	IIA1
Female	50	IIA1

IA1-IB1, early cervical cancer; IB2-IIA2, advanced cervical cancer.

room temperature for 1 h. Membranes were scanned using an Odyssey Imaging System and analyzed with Odyssey v2.0 software (LI-COR Biosciences). The visualization was performed using an ECL chemiluminescent kit (Beyotime Institute of Biotechnology) according to the manufacturer's instructions. The primary antibodies used in the present study were as follows: Anti-JAK2 (1:1,000; cat. no. ab108596; Abcam), anti-IL-17A (1:1,000; cat. no. ab79056; Abcam), anti-IL-17F (1:1,000; cat. no. ab168194; Abcam), anti-STAT3 (1:1,000; cat. no. ab68153; Abcam), anti-vascular endothelial growth factor (VEGF; 1:1,000; cat. no. ab32152; Abcam), anti-Akt (1:1,000; cat. no. ab8805; Abcam), anti-p65 (1:1,000; cat. no. ab32536; Abcam), anti-p-STAT3 (1:1,000; cat. no. ab267373; Abcam), p-p65 (1:1,000; cat. no. ab76302; Abcam), anti-p-JAK2 (1:1,000; cat. no. ab32101; Abcam), anti-p-Akt (1:1,000; cat. no. ab76302; Abcam) and anti-GAPDH (1:1,000; cat. no. ab8245; Abcam). GAPDH was used as an internal control.

Immunofluorescence. Cervical cancer cells (1×10^4 per well) were seeded in 24-well plates overnight and treated as following: Control, IL-17, IL-17 plus siRNA-NC, IL-17 plus siRNA-STAT3 or IL-17 plus siRNA-JAK2 for 72 h. After that, the cells were prefixed in 4% paraformaldehyde at 4°C for 10 min, and fixed in pre-cold methanol at 4°C for another 10 min. Next, cells were incubated with primary antibodies overnight at 4°C: Anti-Ki67 (Abcam; cat. no. ab15580; 1:1,000). The nuclei were stained with DAPI (Beyotime Institute of Biotechnology). Goat anti-rabbit IgG antibody (Abcam; cat. no. ab150077; 1:5,000) was used as the secondary antibody. The samples were visualized by fluorescence microscope (magnification, x200; Olympus CX23; Olympus Corporation) immediately.

Statistical analysis. In total, 3 independent experiments were performed, and all data are expressed as the mean \pm standard deviation. GraphPad Prism 7 (GraphPad Software, Inc.) was used for all statistical analyses. The comparison between two groups was analyzed using paired Student's t-test (Figs. 1A and 2) or unpaired Student's t-test (Fig. 1B). One-way ANOVA followed by Tukey's test was used for comparisons between multiple groups. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

IL-17 mRNA expression level is upregulated in cervical tumor tissues compared with that in normal tissues. In order to investigate the role of IL-17 in the progression of cervical cancer, RT-qPCR was employed. As indicated in Fig. 1A, the expression level of IL-17 was significantly upregulated in tumor tissues compared with that in normal tissues; to verify this result, ELISA was performed. The results demonstrated that the level of IL-17 in serum of patients with cervical cancer was significantly increased (Fig. 1B). Taken together, the results showed that IL-17 was upregulated during the tumorigenesis of cervical cancer.

JAK2/STAT3, NF- κ B, VEGF and PI3K signaling are involved in the tumorigenesis of cervical cancer. For the purpose of exploring the role of JAK2/STAT3, NF- κ B, VEGF and PI3K signaling in development of cervical cancer, western blotting was used. As shown in Fig. 2A-C, the expression levels of JAK2 and STAT3 in tumor tissues was notably increased compared with that in normal tissues. These data suggested that JAK2/STAT3 was involved in the pathogenesis of cervical cancer. Similarly, NF- κ B, VEGF and PI3K/Akt signaling were also upregulated in tumor tissues (Fig. 2A and D-F). These results indicated that JAK2/STAT3, NF- κ B, VEGF and PI3K/Akt were activated in the tumorigenesis of cervical cancer.

IL-17 promotes the growth of HeLa cells. To verify the function of IL-17 in the progression of cervical cancer, CCK-8 assay was performed. As shown in Fig. 3A, IL-17 notably increased the proliferation of HeLa cells. Moreover, a concentration of 50 ng/ml exhibited the most proliferative effect. Therefore, 50 ng/ml was used in the following experiments. Next, RT-qPCR and western blotting were used to detect the




Figure 1. IL-17 is upregulated in cervical cancer tissues. (A) Reverse transcription-quantitative PCR was used to detect the expression level of IL-17 in cervical cancer tissue or adjacent normal tissues. ** $P < 0.01$ via paired Student's t-test. (B) Enzyme-linked immunosorbent assay was performed to detect the level of IL-17 in the serum of patients with cervical cancer or healthy individuals. ** $P < 0.01$ via unpaired Student's t-test. IL, interleukin.




Figure 2. JAK2/STAT3, NF- κ B, VEGF and PI3K are involved in tumorigenesis of cervical cancer. (A) Protein expression levels of JAK2, p-JAK2, STAT3, p-STAT3, p-p65, Akt, p-Akt and VEGF in tissues detected by western blotting. Relative expression levels of (B) p-JAK2 (normalized to JAK2), (C) p-STAT3 (normalized to STAT3), (E) p-p65 (normalized to p65) and (F) p-Akt (normalized to Akt) were quantified. Relative expression level of (D) VEGF was normalized to GAPDH. ** $P < 0.01$ vs. control. The data were analyzed using paired Student's t-test. NF- κ B, nuclear factor- κ B; VEGF, vascular endothelial growth factor; p-, phosphorylated.

transfection efficiency. The data demonstrated that the expression of JAK2 in HeLa cells was significantly decreased by knockdown of JAK2 (Fig. 3B-D). Similarly, the expression of STAT3 in cervical cancer cells was notably inhibited after STAT3 silencing (Fig. 3B-D). These results suggested that

JAK2 and STAT3 siRNA were stably transfected into HeLa cells. Then, the results of Ki-67 staining demonstrated that IL-17 significantly promoted the proliferation of cervical cancer cells. However, knockdown of JAK2 or STAT3 partially rescued the proliferative effect of IL-17 (Fig. 3E). Overall,




Figure 3. IL-17 promotes the proliferation of cervical cancer cells. Cells were treated with different concentrations of IL-17 for 24–96 h. Then, (A) cell viability was detected by Cell Counting Kit-8 assay. (B–D) HeLa cells were transfected with JAK2 siRNA or STAT siRNA for 24 h. Then, reverse transcription-quantitative PCR and western blotting were performed to detect the transfection efficiency. (E) HeLa cells were untreated or with IL-17, IL-17 plus siNC or IL-17 plus siSTAT3 for 24 h. Then, cell proliferation was detected by Ki-67 staining. Red fluorescence indicates Ki-67. Blue fluorescence indicates DAPI. Scale bar, 100 μ m. ** P <0.01 vs. control; ## P <0.01 vs. IL-17. IL, interleukin; siRNA, small interfering RNA; NC, negative control.

these data suggested that IL-17 could promote the growth of cervical cancer cells via activation of JAK2/STAT3 signaling.

IL-17 significantly promotes the migration and invasion of cervical cancer cells. To further investigate the effect of IL-17 on the migration and invasion of cervical cancer cells, wound-healing and Transwell assays were performed. As shown in Fig. 4A and B, the migration and invasion of cervical cancer cells were notably inhibited in the presence of IL-17, which were partially reversed by downregulation of JAK2 or STAT3. These data confirmed that IL-17 could promote the migration and invasion of cervical cancer cells via the JAK2/STAT3 signaling pathway.

IL-17 promotes the tumorigenesis of cervical cancer via upregulation of IL-17A and IL-17F. IL-17A and IL-17F are two major isoforms of IL-17 that can regulate the cancer tumorigenesis (20,21). Thus, these two isoforms were selected for investigations. As indicated in Figs. 5A and B and S1A–D, the levels of IL-17A and IL-17F in cervical cancer cells were significantly upregulated by IL-17, while JAK2 or STAT3

knockdown reversed this phenomenon. Taken together, IL-17 promotes the tumorigenesis of cervical cancer via the upregulation of IL-17A and IL-17F.

IL-17 promotes the progression of cervical cancer through JAK2/STAT3, PI3K/Akt and NF- κ B signaling. To further verify the mechanism by which IL-17 modulates the progression of cervical cancer, western blotting was used. The results indicated that the expression levels of VEGF, phosphorylated (p)-JAK2, p-STAT3, p-Akt and p-p65 were significantly upregulated by IL-17, which was notably rescued by silencing of JAK2 or STAT3 (Figs. 6A and B, 7A and B). Taken together, the results confirmed that IL-17 promoted the progression of cervical cancer through the upregulation of JAK2/STAT3 signaling.

Discussion

The IL-17 family was identified in 1993 by gene screening of mouse T cells. It was originally named CTLA-8, and was regarded as a cytokine called IL-17 in 1995 (22). The IL-17




Figure 4. IL-17 significantly promotes the migration and invasion of cervical cancer cells. (A) Migration of HeLa cells was detected by wound-healing assay. Scale bar, 100 μ m. (B) Invasion of HeLa cells was tested by Transwell assay. Scale bar, 100 μ m. ** $P < 0.01$. IL, interleukin; siRNA, small interfering RNA; NC, negative control.

family contains 6 members (23). IL-17 is an important inflammatory regulator that may activate tissue responses and guide immune defense (24). Previous studies have reported that IL-17 could be involved in tumorigenesis due to the fact that inflammatory factors in the tumor microenvironment can promote tumors to produce cytokines and decrease cytotoxic T cells, thereby promoting tumor growth (12,25). However, the mechanism by which IL-17 regulates the development of cervical cancer remains unclear. The present study is the first to identify that IL-17 could promote the progression of cervical cancer via the activation of JAK2/STAT3. Moreover, Zhang *et al* (26) found that IL-17 could be closely associated

with the progression of thyroid cancer. Besides, a previous study indicated that IL-17 significantly promoted the occurrence of biliary tract cancer via self-producing cytokines (27). The present data further supplemented these previous results, indicating that IL-17 could act as a JAK2/STAT3 promoter during the occurrence of multiple diseases. According to Song *et al* (20), IL-17 could act as an oncogene in laryngeal cancer via the activation of PI3K/AKT/FAS/FASL signaling. Consistently, the present finding suggested that PI3K/Akt signaling could be activated by IL-17 in cervical cancer. It has been reported that IL-17 could activate PI3K/AKT in malignant tumors (28,29). Thereby, the function of IL-17 might




Figure 5. IL-17 promotes the tumorigenesis of cervical cancer via upregulation of IL-17A and IL-17F. (A and B) HeLa cells were transfected with JAK2 siRNA or STAT3 siRNA. The expression levels of IL-17A and IL-17F were detected in HeLa cells by western blotting. The relative expression levels were quantified by normalization to GAPDH. ** $P < 0.01$. IL, interleukin; siRNA, small interfering RNA; NC, negative control.

contribute to the consistence. On the other hand, IL-12, IL-23 and IL-17 are known to be pro-inflammatory cytokines in immunology (30), and TGF- β is an immunosuppressive cytokine (31). A previous report indicated that IL-12, IL-23 and IL-17 could be upregulated in acute lymphoblastic leukemia, while TGF- β was downregulated (32). Consistently, the present data indicated that IL-17 was upregulated in cervical cancer. Meanwhile, IL-12, IL-23 and TGF- β will be investigated in the future.

According to the literature, the JAK2/STAT3 signaling pathway is commonly associated with the metastasis of malignant tumors (33-35). Moreover, the JAK2/STAT3 signaling pathway has been also found to regulate the process of cancer metastasis (36,37). Therefore, it was hypothesized that the knockdown of IL-17 could mediate the JAK2/STAT3 signaling pathway and suppress the growth of cervical cancer cells. As expected, IL-17 could activate JAK2/STAT3 signaling. The present findings are consistent with those from previous studies (38,39), indicating that JAK2/STAT3 signaling could play a key role during tumorigenesis.

The present study also revealed that IL-17 could activate NF- κ B and PI3K/Akt signaling. Various evidence suggest that numerous signaling pathways are involved in the regulation of tumorigenesis, and the JAK2/STAT3, PI3K/Akt and NF- κ B signaling axes have been shown to play important roles in this process (40,41). It has been previously confirmed that the STAT3 transcription factor could be constitutively activated through phosphorylation by upstream JAK kinases in various

cancer types, including gastric cancer, glioma and esophageal cancer, in response to various stimuli such as cytokines and growth factors (42,43). STAT3 was activated to translocate to the nucleus, where it promotes cell proliferation and cell cycle progression, and inhibits apoptosis by activating the transcription of downstream oncogenes, such as Bax and Bcl-2 (44,45). The PI3K/Akt signaling pathway is also involved in the regulation of multiple cellular functions. Once activated, AKT phosphorylates a variety of substrates, resulting in cell cycle progression and inhibition of apoptosis (46,47). NF- κ B is a major transcription factor that is involved in the inflammatory regulation of cells by responding to pro-inflammatory stimuli (48). In the present study, IL-17 increased the expression of p-Akt and p-p65. However, the knockdown of JAK2 or STAT3 significantly reversed the effect of IL-17 on these two signaling pathways. Increasing reports have indicated that the STAT3, NF- κ B and PI3K/Akt signaling pathways could exert their function interactively or independently in different cellular contexts (49-51). These data were similar to the results of the present study, which indicated that IL-17 was the upstream factor of the aforementioned three signaling pathways in the tumorigenesis of cervical cancer. The present findings also indicated that knockdown of JAK2 or STAT3 could reverse the effect of IL-17 on the expression levels of PI3K- and NF- κ B-associated proteins. Therefore, it is urgent to determine whether there is an association between STAT3, Akt and NF- κ B signaling pathways in IL-17-treated cervical cancer cells.




Figure 6. IL-17 promotes the progression of cervical cancer through JAK2/STAT3, PI3K/Akt and NF- κ B signaling. (A and B) HeLa cells were transfected with JAK2 siRNA. Then, the protein expression of p65, p-p65 (normalized to p65), STAT3, p-STAT3 (normalized to STAT3), Akt, p-Akt (normalized to Akt), JAK2, p-JAK2 (normalized to JAK2) and VEGF in HeLa cells were detected by western blotting. The relative expression of VEGF was quantified normalized to GAPDH. ** $P < 0.01$. IL, interleukin; NF- κ B, nuclear factor NF- κ B; p-, phosphorylated-; siRNA, small interfering RNA; VEGF, vascular endothelial growth factor; NC, negative control.

Epigenetic modifications often play important roles in cancer tumorigenesis (52). In the present study, STAT3 was demonstrated to play a key role in IL-17-mediated cervical

cancer progression. According to Zhang *et al.* (53), STAT3 could induce the transcription of the DNA methyltransferase 1 gene (DNMT1) in malignant T lymphocytes. Based




Figure 7. STAT3 siRNA reversed the effect of IL-17 on JAK2/STAT3, PI3K/Akt and NF- κ B signaling. (A and B) HeLa cells were transfected with STAT3 siRNA. Then, the protein expression of p65, p-p65 (normalized to p65), STAT3, p-STAT3 (normalized to STAT3), Akt, p-Akt (normalized to Akt), JAK2, p-JAK2 (normalized to JAK2) and VEGF in HeLa cells were detected by western blotting. The relative expression of VEGF was quantified normalized to GAPDH. ** $P < 0.01$. IL, interleukin; NF- κ B, nuclear factor NF- κ B; p-, phosphorylated-; siRNA, small interfering RNA; VEGF, vascular endothelial growth factor; NC, negative control.

on the aforementioned study (53), STAT3 activation might promote the transcription of DNMT1 in cervical cancer.

The present study is the first to explore the function of IL-17 in cervical cancer, and the first to identify that IL-17 could promote the progression of cervical cancer via the activation of JAK2/STAT3. In addition, IL-17 was demonstrated

to activate JAK2/STAT3, PI3K/Akt and NF- κ B in cervical cancer. However, the present study has the following limitations: i) More IL-17 isoforms need to be detected; ii) some rescue experiments are needed to further verify the association between IL-17 and NF- κ B signaling. Thereby, more investigations are required in the future.

In conclusion, IL-17 significantly promoted the progression of cervical cancer, which may serve as a potential novel target for the treatment of cervical cancer.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

Study design, literature research, experimental study was performed by YB, HL and RL. Data acquisition, data analysis and statistical analysis were performed by YB, RL and YB confirmed the authenticity of all the raw data. All authors were responsible for guarantor of integrity of entire study, manuscript preparation and manuscript editing, and all authors read and approved the final manuscript.

Ethics approval and consent to participate

The study was carried out in accordance with the World Medical Association Declaration of Helsinki approved by Gansu Provincial Cancer Hospital (approval no. GPCH20190220). The clinical and pathological data of patients were collected with their written informed consent.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Tsikouras P, Zervoudis S, Manav B, Tomara E, Iatrakis G, Romanidis C, Bothou A and Galazios G: Cervical cancer: Screening, diagnosis and staging. *J BUON* 21: 320-325, 2016.
2. Benard VB, Thomas CC, King J, Massetti GM, Doria-Rose VP and Saraiya M; Centers for Disease C and Prevention (CDC): Vital signs: Cervical cancer incidence, mortality, and screening-United States, 2007-2012. *MMWR Morb Mortal Wkly Rep* 63: 1004-1009, 2014.
3. Zeng SY, Liang MR, Li LY, Li L, Jiang W and Zhong ML: Application of transvaginal external fascia trachelectomy in the treatment of CIN and micro-invasive cervical cancer. *Zhonghua Zhong Liu Za Zhi* 35: 543-546, 2013 (In Chinese).
4. Gilham C, Sargent A, Kitchener HC and Peto J: HPV testing compared with routine cytology in cervical screening: Long-term follow-up of ARTISTIC RCT. *Health Technol Assess* 23: 1-44, 2019.
5. Hong DK, Kim SA, Lim KT, Lee KH, Kim TJ and So KA: Clinical outcome of high-grade cervical intraepithelial neoplasia during pregnancy: A 10-year experience. *Eur J Obstet Gynecol Reprod Biol* 236: 173-176, 2019.
6. Cheng X, Feng Y, Wang X, Wan X, Xie X and Lu W: The effectiveness of conization treatment for post-menopausal women with high-grade cervical intraepithelial neoplasia. *Exp Ther Med* 5: 185-188, 2013.
7. Ostojic DV, Vrdoljak-Mozetic D, Stemberger-Papic S, Finderle A and Eminovic S: Cervical cytology and HPV test in follow-up after conisation or LLETZ. *Coll Antropol* 34: 219-224, 2010.
8. Apgar BS, Kittendorf AL, Bettcher CM, Wong J and Kaufman AJ: Update on ASCCP consensus guidelines for abnormal cervical screening tests and cervical histology. *Am Fam Physician* 80: 147-155, 2009.
9. Guo N, Shen G, Zhang Y, Moustafa AA, Ge D and You Z: Interleukin-17 promotes migration and invasion of human cancer cells through upregulation of MTA1 expression. *Front Oncol* 9: 546, 2019.
10. Alves JJ, De Medeiros Fernandes TAA, De Araujo JM, Cobucci RN, Lanza DC, Bezerra FL, Andrade VS and Fernandes JV: Th17 response in patients with cervical cancer. *Oncol Lett* 16: 6215-6227, 2018.
11. Karabulut M, Usul Afsar C, Serimez M and Karabulut S: Serum IL-17 levels can be diagnostic for gastric cancer. *J BUON* 24: 1601-1609, 2019.
12. Iwakura Y, Ishigame H, Saijo S and Nakae S: Functional specialization of interleukin-17 family members. *Immunity* 34: 149-162, 2011.
13. Lotti F, Jarrar AM, Pai RK, Hitomi M, Lathia J, Mace A, Gantt GA Jr, Sukhdeo K, DeVecchio J, Vasanji A, *et al*: Chemotherapy activates cancer-associated fibroblasts to maintain colorectal cancer-initiating cells by IL-17A. *J Exp Med* 210: 2851-2872, 2013.
14. Flies EJ, Mavoa S, Zosky GR, Mantzioris E, Williams C, Eri R, Brook BW and Buettel JC: Urban-associated diseases: Candidate diseases, environmental risk factors, and a path forward. *Environ Int* 133: 105187, 2019.
15. Alshammari TK, Alghamdi H, Green TA, Niazy A, Alkahdar L, Alrasheed N, Alhosaini K, Alswayed M, Elango R, Laezza F, *et al*: Assessing the role of toll-like receptor in isolated, standard and enriched housing conditions. *PLoS One* 14: e0222818, 2019.
16. Nagarkoti S, Dubey M, Sadaf S, Awasthi D, Chandra T, Jagavelu K, Kumar S and Dikshit M: Catalase S-Glutathionylation by NOX2 and mitochondrial-derived ROS adversely affects mice and human neutrophil survival. *Inflammation* 42: 2286-2296, 2019.
17. Cong J, Liu R, Wang X, Sheng L, Jiang H, Wang W, Zhang Y, Yang S and Li C: Association between interleukin-17 gene polymorphisms and the risk of cervical cancer in a Chinese population. *Int J Clin Exp Pathol* 8: 9567-9573, 2015.
18. Miranda LN, Reginaldo FP, Souza DM, Soares CP, Silva TG, Rocha KB, Jatoba CA, Donadi EA, Andrade JM, Goncalves AK and Crispim JC: Greater expression of the human leukocyte antigen-G (HLA-G) and interleukin-17 (IL-17) in cervical intraepithelial neoplasia: Analytical cross-sectional study. *Sao Paulo Med J* 133: 336-342, 2015.
19. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
20. Song Y, Yang M, Zhang H, Sun Y, Tao Y, Li H, Zhang J, Li Y and Yang J: IL-17 affects the progression, metastasis, and recurrence of laryngeal cancer via the inhibition of apoptosis through activation of the PI3K/AKT/FAS/FASL pathways. *J Immunol Res* 2020: 2953191, 2020.
21. Lu W, He F, Lin Z, Liu S, Tang L, Huang Y and Hu Z: Dysbiosis of the endometrial microbiota and its association with inflammatory cytokines in endometrial cancer. *Int J Cancer* 148: 1708-1716, 2021.
22. Liao C, Zhang C, Jin L and Yang Y: IL-17 alters the mesenchymal stem cell niche towards osteogenesis in cooperation with osteocytes. *J Cell Physiol* 235: 4466-4480, 2020.
23. Guo JQ, Liu J and Lu B: Expression of gamma-delta T cells in immune microenvironment in children with Henoch-Schonlein purpura. *Zhongguo Dang Dai Er Ke Za Zhi* 21: 960-965, 2019 (In Chinese).
24. Siefker DT, Vu L, You D, McBride A, Taylor R, Jones TL, DeVincenzo J and Cormier SA: Respiratory syncytial virus disease severity is associated with distinct CD8⁺ T-cell profiles. *Am J Respir Crit Care Med* 201: 325-334, 2020.
25. Wang J, Lu L, Luo Z, Li W, Lu Y, Tang Q and Pu J: MiR-383 inhibits cell growth and promotes cell apoptosis in hepatocellular carcinoma by targeting IL-17 via STAT3 signaling pathway. *Biomed Pharmacother* 120: 109551, 2019.

26. Zhang N, Wang Q, Tian Y, Xiong S, Li G and Xu L: Expressions of IL-17 and TNF- α in patients with Hashimoto's disease combined with thyroid cancer before and after surgery and their relationship with prognosis. *Clin Transl Oncol* 22: 1280-1287, 2020.
27. Kinoshita M, Kobayashi S, Gotoh K, Kubo M, Hayashi K, Iwagami Y, Yamada D, Akita H, Noda T, Asaoka T, *et al*: Heterogeneity of Treg/Th17 according to cancer progression and modification in biliary tract cancers via self-producing cytokines. *Dig Dis Sci* 65: 2937-2948, 2020.
28. Amara S, Majors C, Roy B, Hill S, Rose KL, Myles EL and Tiriveedhi V: Critical role of SIK3 in mediating high salt and IL-17 synergy leading to breast cancer cell proliferation. *PLoS One* 12: e0180097, 2017.
29. Varikuti S, Oghumu S, Elbaz M, Volpedo G, Ahirwar DK, Alarcon PC, Sperling RH, Moretti E, Pioso MS, Kimble J, *et al*: STAT1 gene deficient mice develop accelerated breast cancer growth and metastasis which is reduced by IL-17 blockade. *Oncoimmunology* 6: e1361088, 2017.
30. Zou Y, Dai SX, Chi HG, Li T, He ZW, Wang J, Ye CG, Huang GL, Zhao B, Li WY, *et al*: Baicalin attenuates TNBS-induced colitis in rats by modulating the Th17/Treg paradigm. *Arch Pharm Res* 38: 1873-1887, 2015.
31. Song KH, Jung SY, Kang SM, Kim MH, Ahn J, Hwang SG, Lee JH, Lim DS, Nam SY and Song JY: Induction of immunogenic cell death by radiation-upregulated karyopherin alpha 2 in vitro. *Eur J Cell Biol* 95: 219-227, 2016.
32. Perez-Figueroa E, Sanchez-Cuaxospa M, Martinez-Soto KA, Sanchez-Zauco N, Medina-Sanson A, Jimenez-Hernandez E, Torres-Nava JR, Felix-Castro JM, Gomez A, Ortega E, *et al*: Strong inflammatory response and Th1-polarization profile in children with acute lymphoblastic leukemia without apparent infection. *Oncol Rep* 35: 2699-2706, 2016.
33. Souza CM, do Amaral CL, Souza SC, de Souza ACP, de Cassia Alves Martins I, Contieri LS, Milanski M, Torsoni AS, Ignacio-Souza LM and Torsoni MA: JAK2/STAT3 pathway is required for alpha7nAChR-dependent expression of POMC and AGRP neuropeptides in male mice. *Cell Physiol Biochem* 53: 701-712, 2019.
34. Fogg KC, Olson WR, Miller JN, Khan A, Renner C, Hale I, Weisman PS and Kreeger PK: Alternatively activated macrophage-derived secretome stimulates ovarian cancer spheroid spreading through a JAK2/STAT3 pathway. *Cancer Lett* 458: 92-101, 2019.
35. Wei L, Chen Y, Zhang C, Liu M and Xiong H: Leptin induces IL-6 and IL-8 expression through leptin receptor Ob-Rb in human dental pulp fibroblasts. *Acta Odontol Scand* 77: 205-212, 2019.
36. Song Q, Liu B, Li X, Zhang Q, Cao L, Xu M, Meng Z, Wu X and Xu K: MiR-26a-5p potentiates metastasis of human lung cancer cells by regulating ITG β 8-JAK2/STAT3 axis. *Biochem Biophys Res Commun* 501: 494-500, 2018.
37. Zhang L, Lu P, Guo X, Liu T, Luo X and Zhu YT: Inhibition of JAK2/STAT3 signaling pathway protects mice from the DDP-induced acute kidney injury in lung cancer. *Inflamm Res* 68: 751-760, 2019.
38. Wu D, Dong W, Fang K and Wang M: As₄S₄ exhibits good killing effect on multiple myeloma cells via repressing SOCS1 methylation-mediated JAK2/STAT3 signaling pathway. *Technol Cancer Res Treat* 18: 1533033819896806, 2019.
39. Jing W, Guo X, Wang G, Bi Y, Han L, Zhu Q, Qiu C, Tanaka M and Zhao Y: Breast cancer cells promote CD169⁺ macrophage-associated immunosuppression through JAK2-mediated PD-L1 upregulation on macrophages. *Int Immunopharmacol* 78: 106012, 2020.
40. Wang X, Yin H, Zhang L, Zheng D, Yang Y, Zhang J, Jiang H, Ling X, Xin Y, Liang H, *et al*: The construction and analysis of the aberrant lncRNA-miRNA-mRNA network in non-small cell lung cancer. *J Thorac Dis* 11: 1772-1778, 2019.
41. Li X, Mak VCY, Zhou Y, Wang C, Wong ESY, Sharma R, Lu Y, Cheung ANY, Mills GB and Cheung LWT: Deregulated Gab2 phosphorylation mediates aberrant AKT and STAT3 signaling upon PIK3R1 loss in ovarian cancer. *Nat Commun* 10: 716, 2019.
42. Deng F, Wang S, Zhang L, Xie X, Cai S, Li H, Xie GL, Miao HL, Yang C, Liu X and Xia Z: Propofol through upregulating caveolin-3 attenuates post-hypoxic mitochondrial damage and cell death in H9C2 cardiomyocytes during hyperglycemia. *Cell Physiol Biochem* 44: 279-292, 2017.
43. Li YL, Gao L, Zucker IH and Schultz HD: NADPH oxidase-derived superoxide anion mediates angiotensin II-enhanced carotid body chemoreceptor sensitivity in heart failure rabbits. *Cardiovasc Res* 75: 546-554, 2007.
44. Wei Z, Jiang X, Qiao H, Zhai B, Zhang L, Zhang Q, Wu Y, Jiang H and Sun X: STAT3 interacts with Skp2/p27/p21 pathway to regulate the motility and invasion of gastric cancer cells. *Cell Signal* 25: 931-938, 2013.
45. Vona R, Gambardella L, Cittadini C, Straface E and Pietraforte D: Biomarkers of oxidative stress in metabolic syndrome and associated diseases. *Oxid Med Cell Longev* 2019: 8267234, 2019.
46. Li K: Iron pathophysiology in friedreich's ataxia. *Adv Exp Med Biol* 1173: 125-143, 2019.
47. Cho TM, Kim JY, Kim YJ, Sung D, Oh E, Jang S, Farrand L, Hoang VH, Nguyen CT, Ann J, *et al*: C-terminal HSP90 inhibitor L80 elicits anti-metastatic effects in triple-negative breast cancer via STAT3 inhibition. *Cancer Lett* 447: 141-153, 2019.
48. Ucci M, Di Tomo P, Tritschler F, Cordone VGP, Lanuti P, Bologna G, Di Silvestre S, Di Pietro N, Pipino C, Mandatori D, *et al*: Anti-inflammatory role of carotenoids in endothelial cells derived from umbilical cord of women affected by gestational diabetes mellitus. *Oxid Med Cell Longev* 2019: 8184656, 2019.
49. Li Y, Cui N, Zheng PS and Yang WT: BMX/Etk promotes cell proliferation and tumorigenicity of cervical cancer cells through PI3K/AKT/mTOR and STAT3 pathways. *Oncotarget* 8: 49238-49252, 2017.
50. Steelman LS, Abrams SL, Whelan J, Bertrand FE, Ludwig DE, Basecke J, Libra M, Stivala F, Milella M, Tafuri A, *et al*: Contributions of the Raf/MEK/ERK, PI3K/PTEN/Akt/mTOR and Jak/STAT pathways to leukemia. *Leukemia* 22: 686-707, 2008.
51. Yang X, Yan H, Jiang N, Yu Z, Yuan J, Ni Z and Fang W: IL-6 Trans-signaling drives a STAT3-dependent pathway that leads to structural alterations of peritoneal membrane. *Am J Physiol Renal Physiol* 318: F338-F353, 2020.
52. Buttura JR, Provisor Santos MN, Valieris R, Drummond RD, Defelicibus A, Lima JP, Calsavara VF, Freitas HC, Cordeiro de Lima VC, Fernanda Bartelli T, *et al*: Mutational signatures driven by epigenetic determinants enable the stratification of patients with gastric cancer for therapeutic intervention. *Cancers (Basel)* 13: 490, 2021.
53. Zhang Q, Wang HY, Woetmann A, Raghunath PN, Odum N and Wasik MA: STAT3 induces transcription of the DNA methyltransferase 1 gene (DNMT1) in malignant T lymphocytes. *Blood* 108: 1058-1064, 2006.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.