

Cross-sectional study of smoking exposure: no differential effect on OCT metrics in a cohort of MS patients

Mattia Rosso, Dorlan J Kimbrough, Cindy T Gonzalez, Bonnie I Glanz, Brian C Healy, Maria Assunta Rocca , Massimo Filippi , Howard Weiner and Tanuja Chitnis 

Multiple Sclerosis Journal—
Experimental, Translational
and Clinical

January–March 2019, 1–9

DOI: 10.1177/
2055217319828400

© The Author(s), 2019.
Article reuse guidelines:
sagepub.com/journals-
permissions

Abstract

Background: Optical coherence tomography (OCT) provides quantitative measures of retinal layer thickness. Cigarette smoking is a risk factor for multiple sclerosis (MS) onset and disease severity: its effects on OCT metrics have not been assessed.

Objective: The objective of this study was to assess the effect of smoking history on retinal nerve fiber layer (RNFL) and ganglion cell-inner plexiform (GCIP) of MS patients by OCT.

Methods: 112 MS patients were recruited from the Brigham and Women's Hospital. Spectralis OCT scans were acquired to measure GCIP, peripapillary RNFL, and total macular volume. Multivariable linear mixed effects regression model assessed RNFL and GCIP change with fixed effects for smoking history while adjusting for optic neuritis eye status, age, disease duration, sex, baseline EDSS, and disease modifying therapies (DMTs).

Results: Smoking histories were available for 102 patients: 46 (45.10%) had a history of smoking cigarettes and 56 (54.90%) never smoked. No statistically significant differences were found between ever-smokers and never-smokers with respect to GCIP, RNFL, and macular volume.

Conclusion: Our study shows no significant difference in retinal thickness between ever-smokers and never-smokers. If confirmed, this result suggests mechanistic differences between the retina and other central nervous system (CNS) compartments in response to smoking and should be noted when considering OCT as a surrogate measure of CNS activity.

Keywords: Imaging, multiple sclerosis, optical coherence tomography, cigarette smoking, optic neuritis, retinal nerve fiber layer, ganglion cell-inner plexiform layer

Date received: 25 September 2018; accepted: 31 December 2018

Introduction

Among the recognized environmental risk factors for multiple sclerosis (MS), the most evidence has been acquired for low vitamin D levels, low of exposure to ultraviolet light, Epstein–Barr virus, and, importantly, cigarette smoking.^{1,2} As the incidence of MS rose consistently during the 20th century, cigarette smoking became a widespread habit in the general population in Europe and North America. Since the 1960s, epidemiological studies identified cigarette smoking as a risk factor for MS. Smoking has been associated with a dose dependent increased

susceptibility to MS and, to a lesser extent, with the risk of conversion from clinically isolated syndrome (CIS) to clinically definite MS, and with an increased susceptibility to secondary progressive MS conversion.^{3–8} MS patients who smoke also seem to be predisposed to faster rates of disability accrual, as shown by higher Expanded Disability Status Scale (EDSS) scores.^{3,5,9–11}

The impact of smoking in MS patients has also been assessed through imaging studies: on magnetic resonance imaging, smokers show greater lesion

Correspondence to:
Tanuja Chitnis,
Partners Multiple Sclerosis
Center, 60 Fenwood Road,
Office 9002K, Boston, MA
02115, USA.
tchitnis@partners.org
Mattia Rosso,
Dorlan J Kimbrough,
Cindy T Gonzalez,
Bonnie I Glanz,
Brian C Healy,
Ann Romney Center for
Neurologic Disease, Harvard
Medical School, Boston,
MA, USA



Maria Assunta Rocca,
Massimo Filippi,
Institute of Experimental
Neurology, Vita-Salute San
Raffaele University,
Milan, Italy

Howard Weiner,
Tanuja Chitnis
Ann Romney Center for
Neurologic Disease, Harvard
Medical School, Boston,
MA, USA
Department of Neurology,
Brigham and Women's
Hospital, Boston, MA, USA

volumes, higher brain T2 lesion loads, and more severe brain atrophy than non-smokers.^{4,5} Spectral-domain optical coherence tomography (OCT) measurement of the retina yields high resolution imaging and quantification of the peripapillary retinal nerve fiber layer (RNFL) and segmented macular layers. RNFL and ganglion cell-inner plexiform layer (GCIP) have been used as outcome measures in a wide variety of studies: both appear to correlate with visual function and global disability (EDSS). In particular, GCIP appears to correlate with brain T2 lesion load, brain atrophy and EDSS-measured disability progression, rendering it a more robust measure than RNFL as it is less prone to artifacts and in the acute setting is less affected by post-optic neuritis associated swelling. The correlation between brain and retinal atrophy seems to result from a pathology of anterior visual pathway: while retrograde neurodegeneration results in GCIP thinning, anterograde neurodegeneration results in brain atrophy.^{12–16} Pathology studies have shown that anterior visual pathway damage is common among MS patients and therefore, it is a good hallmark of overall MS-related neurodegeneration.¹⁷

We utilized OCT in an attempt to examine the impact of cigarette smoking on retinal measures in MS patients. To better characterize our population, we stratified out patients according to their smoking history (ever-smokers vs. never-smokers).

Materials and methods

Subjects

Subjects enrolled in the Comprehensive Longitudinal Investigation of Multiple Sclerosis at the Brigham and Women's Hospital (CLIMB) who also met the further inclusion criteria for this cross-sectional analysis: (1) enrolled in the longitudinal OCT subgroup, (2) were over the age of 18, and (3) met the 2010 McDonald diagnostic criteria for MS or CIS. As part of the demographic survey, patients provided both basic personal data such as gender, date of birth, optic neuritis (ON) history, and smoking history categorized as being never-smokers or ever-smokers. When providing ON history, each patient was asked to include the dates of past ON event and the laterality of each ON episode (bilateral, left, right). Healthy control (HC) subjects were also included and provided basic demographic data. All the data was then stored on the internal relational database of the CLIMB study. Exclusion criteria for MS and HC subjects included previous history of ophthalmological disorders, e.g., clinically-significant cataracts,

glaucoma, diabetes and/or diabetic retinopathy, macular degeneration, and retinal detachment; refractive errors exceeding ± 6 diopters; history of immunological and/or neurological disorders such as neuromyelitis optica, systemic lupus erythematosus, asthma, autoimmune thyroiditis, and rheumatoid arthritis. Additionally, we excluded from the study all MS patients with an acute use of steroids or acute ON within three months of OCT scans. Ten MS patients were excluded from the primary analysis due to their unknown status for smoking history.

All MS patients underwent standardized clinical exams performed by a neurologist, and OCT scanning. HCs also underwent visual acuity testing, and OCT scanning. Visual acuity was assessed for one eye at a time while the non-tested fellow eye was covered by the patient. The score of each chart was based on the number of letters correctly identified, which ranged from 0 to 60. Each patient was examined under reproducible conditions: the distance from the chart, the lighting of the room, and the test instructions were standardized. Also, all study participants provided written informed consent for their participation and the Partners Human Research Committee provided Institutional Review Board (IRB) approval.

Optical coherence tomography

Retinal imaging was performed by trained examiners using Spectralis OCT (software version 6.0.9.0, Heidelberg Engineering, Heidelberg, Germany). Peripapillary RNFL and macular volume scans were obtained from both eyes of each participant (without dilation), with automated segmentation providing measures of the GCIP. Scans were obtained using high resolution and automatic real-time tracking for B-scan averaging to enhance quality and were subsequently manually inspected to corroborate accurate fixation and centering.

Statistical analysis

We compared the demographic and clinical characteristics between ever-smoker and never-smoker MS patients at the baseline OCT exam. The continuous variables were described by mean and standard deviation (SD), while the categorical variables were described by counts and percentages. Means were compared with either a two-sample t-test or a Mann–Whitney U test for non-normal data, while proportions were compared with either a chi-square test or Fisher's exact test for contingency tables.

We then compared RNFL and GCIP measurements between MS patients and HC. Since both eyes of

each subject were included in this analysis, linear mixed models with unstructured correlation structures were used to adjust for the within-subject correlation. Multivariable linear mixed effects regression model assessed differences in RNFL and GCIP measurements with fixed effects for MS versus HC while adjusting for ON eye status, age, and sex.

We assessed the relationship of RNFL and GCIP measurements between ever-smokers and never-smokers using mixed linear regression models. Multivariable linear mixed effects regression model with unstructured correlation structures assessed differences in RNFL and GCIP measurements with fixed effects for smoking history while adjusting for ON eye status, age, disease duration, sex, baseline EDSS, and disease modifying therapies (DMTs). All statistical analyses were performed using SAS version 9.4 (Statistical Analytical System, SAS Institute).

Results

Demographics

Out of 112 MS patients, 102 had information available on cigarette smoking. 46 (45.10%) were ever-smokers and 56 (54.90%) were never-smokers. Table 1 displays demographic and baseline clinical characteristics of MS patients grouped according to their smoking status. Ever-smoker patients had longer disease duration at visit (17 vs. 12 years, $p=0.0038$) and higher EDSS at visit (3 vs. 2, $p=0.0017$) than never-smokers. There was no statistically significant difference in terms of their age at the baseline visit (48 vs. 45, $p=0.1540$) or in their history of ON (54% vs. 43%, $p=0.2478$).

MS and healthy controls

57 HC and 112 MS patients were included in our preliminary analysis which focused on comparing the two groups in RNFL and GCIP measures. Supplementary table 1 shows their baseline demographic and clinical characteristics. HC were younger (38 y.o. vs. 46 y.o., $p<0.0001$) and more racially diverse ($p<0.0001$) compared to MS patients.

As expected, MS patients had lower retinal thickness compared to HC. MS patients had lower total macular volume compared to HC (HC: 1.94 mm^3 vs. MS: 1.77 mm^3 , $\beta=0.11$, $p=0.0106$), lower GCIP inner ring thickness (HC: $93.36\text{ }\mu\text{m}$ vs. MS: $.26\text{ }\mu\text{m}$, $\beta=11.81$, $p<0.0001$), and thinner GCIP outcomes across all individual quadrants (Table 2). HC had greater mean global RNFL ($99.62\text{ }\mu\text{m}$ vs. $85.99\text{ }\mu\text{m}$, $\beta=12.14$, $p<0.0001$) and greater RNFL

papillomacular bundle (PMB) thickness ($58.16\text{ }\mu\text{m}$ vs. $48.19\text{ }\mu\text{m}$, $\beta=8.40$, $p=0.0015$) compared to MS patients (Table 3).

Relationship of smoking history to GCIP and RNFL

We tested the association of GCIP and RNFL measurements between ever-smokers versus never-smokers MS patients, including nine RNFL and fourteen GCIP sectors (Tables 4 and 5). We failed to find any significant effect on GCIP metrics or RNFL measurements between ever-smokers versus never-smokers MS patients after adjusting for ON eye status, age, disease duration, sex, baseline EDSS and DMTs. The distribution in GCIP total macular volume, GCIP inner ring thickness, RNFL mean global, and RNFL PMB between ever-smokers and never-smokers are shown in Figure 1.

Discussion

In this cross-sectional study, our aim was to determine whether or not a history of cigarette smoking was associated with retinal damage quantified by RNFL, GCIP, and macular volume measures. We hypothesized that patients who had an ever-smoker history would have significantly thinned retina layers compared to patients who were never-smokers. Our results showed no significant differences in the thickness of retinal layers between ever-smoker and never-smoker MS patients.

While cigarette smoking has been consistently associated with MS, the exact mechanism of cigarette smoking toxicity has not been fully understood. The composition of cigarette smoke is complex, comprising more than 7000 chemicals, including nicotine and its derivatives along with a large number of compounds that form the cigarette smoke condensate.¹⁸ According to both experimental and epidemiological studies, it is the smoke condensate rather than nicotine that is responsible for the major toxic effects on MS patients.^{19,20} Some of these toxic compounds can directly damage axons (e.g. nitric oxide) and compounds that trigger demyelination (e.g. cyanates).²¹ Also, cigarette smoke has been implicated in the activation of lung macrophages, which may incite the proinflammatory cascade of MS.^{20,22,23} Further, microglial activation in the central nervous system (CNS) has also been suggested as an important hub in the process of demyelination.¹⁹

We expected smoking to be detrimental to MS patients via either direct retinal toxicity or through proinflammatory activation of the immune system

Table 1. Demographics characteristics between ever-smokers versus never-smokers MS patients.

Variables	Ever-smokers	Never-smokers	<i>p</i> -value
<i>N</i>	46	56	
Optic neuritis, <i>n</i> (%)			0.2478 ^b
Yes	25 (54.35%)	24 (42.86%)	
No	21 (45.65%)	32 (57.14%)	
Sex, <i>n</i> (%)			0.5042 ^b
Female	34 (73.91%)	38 (67.86%)	
Male	12 (26.09%)	18 (32.14%)	
Age at OCT exam date, years (mean ± SD)	47.92 ± 9.08	45.39 ± 8.64	0.1540 ^a
Race, <i>n</i> (%)			0.5985 ^c
Native American or Alaska Native	0 (0.00%)	1 (1.79%)	
Black or African American	1 (2.17%)	2 (3.57%)	
More than one race	0 (0.00%)	1 (1.79%)	
White	45 (97.83%)	52 (92.86%)	
Ethnicity, <i>n</i> (%)			0.6249 ^c
Hispanic or Latino	1 (2.17%)	3 (5.36%)	
Not Hispanic or Latino	45 (97.83%)	52 (92.86%)	
Unknown or not reported	0 (0.00%)	1 (1.79%)	
Age at first symptom, years (mean ± SD)	30.89 ± 11.09	33.45 ± 8.61	0.1916 ^a
Disease duration at visit, years (mean ± SD)	17.02 ± 10.23	11.93 ± 7.06	0.0038 ^a
EDSS at visit, (mean ± SD)	3.36 ± 2.26	2.04 ± 1.51	0.0017 ^d
Disease category, <i>n</i> (%)			0.1492 ^c
CIS	3 (6.52%)	2 (3.57%)	
Primary progressive MS	2 (4.35%)	1 (1.79%)	
Secondary progressive MS	10 (21.74%)	5 (8.93%)	
Relapsing–remitting MS	31 (67.39%)	48 (85.71%)	
DMT			0.6740 ^c
Aubagio	1 (2.17%)	0 (0.00%)	
Avonex	3 (6.52%)	4 (7.14%)	
Betaseron	1 (2.17%)	2 (3.57%)	
Cellcept	5 (10.87%)	2 (3.57%)	
Copaxone	5 (10.87%)	13 (23.21%)	
Gilenya	8 (17.39%)	10 (17.86%)	
Imuran	8 (17.39%)	8 (14.29%)	
None	2 (4.35%)	1 (1.79%)	
Rebif	1 (2.17%)	0 (0.00%)	
Rituxan	6 (13.04%)	8 (14.29%)	
Tecfidera (BG-12)	6 (13.04%)	8 (14.29%)	
Tysabri	1 (2.17%)	0 (0.00%)	

MS = multiple sclerosis; OCT = optic coherence tomography; EDSS = Expanded Disability Status Scale; CIS = clinically isolated syndrome; SD = standard deviation.

^aMeans were compared with a two-sample *t*-test.

^bProportions were compared via a chi-square test for contingency tables.

^cProportions were compared via a Fisher's exact test for contingency tables.

^dRank scores were compared via the Mann–Whitney *U* test.

leading to a more active disease phenotype. In fact, smoking has been shown to have deleterious effects on both the retina and the optic nerve in the context of acute macular degeneration, tobacco–alcohol

amblyopia, and Cuban epidemic optic neuropathy.^{24,25} Furthermore, retinal thinning in MS patients can be due to a direct damage of the optic nerve or to a more widespread process of brain atrophy resulting

Table 2. Association of GCIP measurements between MS patients versus healthy controls.

GCIP	MS		HC		Adjusted estimate ^a	Adjusted <i>p</i> -value ^a
	<i>N</i>	Mean ± SD	<i>N</i>	Mean ± SD		
Total macular volume (mm ³)	220	1.77 ± 0.26	114	1.94 ± 0.28	0.11	0.0106
Mean central GCIP thickness (μm)	220	45.07 ± 13.95	111	38.44 ± 8.96	-7.85	0.0003
Mean GCIP N1 thickness (μm)	220	83.07 ± 16.43	111	94.15 ± 9.84	6.61	0.0044
Mean GCIP N2 thickness (μm)	220	59.54 ± 9.10	110	68.88 ± 6.43	8.34	<.0001
Mean GCIP S1 thickness (μm)	219	80.64 ± 14.32	111	94.92 ± 7.16	11.00	<.0001
Mean GCIP S2 thickness (μm)	219	54.21 ± 8.05	111	62.48 ± 5.31	7.07	<.0001
Mean GCIP T1 thickness (μm)	220	79.48 ± 15.39	111	90.79 ± 8.30	7.45	0.0007
Mean GCIP T2 thickness (μm)	218	61.41 ± 8.37	111	67.26 ± 5.38	8.34	<.0001
Mean GCIP I1 thickness (μm)	220	77.40 ± 16.62	111	93.58 ± 8.84	11.81	<.0001
Mean GCIP I2 thickness (μm)	219	58.79 ± 8.14	111	59.50 ± 6.15	0.06	0.9644
Center thickness (μm)	220	30.56 ± 19.24	111	15.53 ± 5.10	-14.37	<.0001
Central minimal thickness (μm)	220	13.05 ± 3.22	111	12.05 ± 3.09	-0.70	0.1769
Central maximal thickness (μm)	220	82.88 ± 19.92	111	78.61 ± 14.84	-8.53	0.0045
Inner ring thickness (μm)	219	80.26 ± 14.94	111	93.36 ± 7.95	11.81	<.0001

MS = multiple sclerosis; HC = healthy control; GCIP = ganglion cell-inner plexiform; SD = standard deviation.

^aMultivariable linear mixed effects model adjusted for age and sex.

Table 3. Association of RNFL measurements between MS patients versus healthy control.

RNFL	MS		HC		Adjusted estimate ^a	Adjusted <i>p</i> -value ^a
	<i>N</i>	Mean ± SD	<i>N</i>	Mean ± SD		
Mean global RNFL thickness (μm)	197	85.99 ± 14.16	108	99.62 ± 10.38	12.14	<.0001
Mean RNFL thickness, PMB (μm)	207	48.19 ± 15.45	111	58.16 ± 14.64	8.36	0.0015
N/T RNFL thickness ratio	196	1.15 ± 0.38	107	1.11 ± 0.51	0.02	0.7458
Mean RNFL thickness, NS (μm)	200	92.06 ± 22.80	110	104.4 ± 25.65	11.89	0.0018
Mean RNFL thickness, N (μm)	197	66.58 ± 18.16	107	75.90 ± 21.94	9.38	0.0038
Mean RNFL thickness, NI (μm)	202	103.3 ± 25.71	110	115.5 ± 26.73	11.62	0.0078
Mean RNFL thickness, TI (μm)	204	122.2 ± 26.18	109	139.9 ± 22.30	11.35	0.0046
Mean RNFL thickness, T (μm)	207	61.37 ± 17.71	111	74.92 ± 19.01	10.80	0.0003
Mean RNFL thickness, TS (μm)	205	116.2 ± 22.68	109	131.2 ± 21.89	12.75	0.0006

MS = multiple sclerosis; HC = healthy control; RNFL = retinal nerve fiber layer; PMB = papillomacular bundle thickness; SD = standard deviation.

^aMultivariable linear mixed effects model adjusted for age and sex.

in trans-synaptic retinal thinning: both relapses and brain atrophy can be facilitated by cigarette smoking.^{4,15,26-30}

To the best of our knowledge via review of available literature in PubMed, this study appears to be the first to address the association of cigarette ever-smokers versus never-smokers MS patients using OCT imaging. In previous studies of healthy individuals, the effects of cigarette smoking on retinal parameters have been inconsistent. While a few

studies showed a decrease in RNFL and choroidal thickness in smokers, other studies have failed to replicate those findings.³¹⁻³³ Cigarette smoking has been shown to play a role in increasing RNFL thinning in migraine patients.³⁴ In contrast, our study showed no effect of smoking on RNFL thinning in MS patients.

The strengths of our study were a large patient population, the presence of smokers and non-smokers, and controlling for history of ON. The main




Figure 1. Comparing RNFL and GCIP distributions between the ever-smokers versus never-smokers showing the median, mean (unfilled diamond shape), and interquartile range. (a) Comparing GCIP total macular volume between ever-smokers ($N=45$) vs. never-smokers ($N=60$). (b) Comparing GCIP inner ring thickness between ever-smokers ($N=44$) vs. never-smokers ($N=60$). (c) Comparing RNFL mean global thickness between ever-smokers ($N=40$) vs. never-smokers ($N=49$). (d) Comparing RNFL mean thickness PMB between ever-smokers ($N=41$) vs. never-smokers ($N=53$).

Table 4. Association of GCIP measurements between ever-smokers versus never-smokers MS patients.

GCIP	Ever-smokers		Never-smokers		Adjusted estimate ^a	Adjusted p -value ^a
	N	Mean \pm SD	N	Mean \pm SD		
Total macular volume (mm^3)	89	1.76 \pm 0.27	111	1.79 \pm 0.24	0.02	0.6912
Mean central GCIP thickness (μm)	89	45.66 \pm 13.62	111	45.03 \pm 13.62	-0.44	0.8718
Mean GCIP N1 thickness (μm)	89	82.39 \pm 16.22	111	83.69 \pm 15.36	2.91	0.3078
Mean GCIP N2 thickness (μm)	89	58.84 \pm 9.45	111	60.04 \pm 8.60	0.46	0.7963
Mean GCIP S1 thickness (μm)	88	79.93 \pm 15.03	111	81.47 \pm 12.94	2.46	0.3569
Mean GCIP S2 thickness (μm)	88	53.36 \pm 8.85	111	55.04 \pm 7.36	0.47	0.7814
Mean GCIP T1 thickness (μm)	89	79.02 \pm 15.30	111	79.95 \pm 14.95	2.91	0.2941
Mean GCIP T2 thickness (μm)	89	60.84 \pm 9.37	109	62.01 \pm 7.28	0.11	0.9479
Mean GCIP I1 thickness (μm)	89	75.99 \pm 17.82	111	78.36 \pm 14.80	2.41	0.4380
Mean GCIP I2 thickness (μm)	89	58.66 \pm 7.91	110	59.16 \pm 7.92	-0.81	0.6195
Center thickness (μm)	89	32.03 \pm 20.03	111	29.97 \pm 18.48	-1.81	0.6436
Central minimal thickness (μm)	89	12.88 \pm 3.83	111	13.36 \pm 2.66	-1.01	0.1157

(continued)

Table 4. Continued

GCIP	Ever-smokers		Never-smokers		Adjusted estimate ^a	Adjusted <i>p</i> -value ^a
	<i>N</i>	Mean ± SD	<i>N</i>	Mean ± SD		
Central maximal thickness (μm)	89	84.53 ± 18.46	111	82.65 ± 19.27	2.96	0.4205
Inner ring thickness (μm)	88	79.60 ± 15.23	111	80.87 ± 13.82	3.11	0.2504

MS = multiple sclerosis; GCIP = ganglion cell-inner plexiform; SD = standard deviation.
^aMultivariable linear mixed effects model adjusted for ON eye status, age, disease duration, sex, baseline EDSS, and DMT.

Table 5. Association of RNFL measurements between ever-smokers versus never-smokers MS patients.

RNFL	Ever-smokers		Never-smokers		Adjusted estimate ^a	Adjusted <i>p</i> -value ^a
	<i>N</i>	Mean ± SD	<i>N</i>	Mean ± SD		
Mean global RNFL thickness (μm)	80	86.25 ± 14.19	97	85.56 ± 14.07	1.97	0.4937
Mean RNFL thickness, PMB (μm)	82	47.01 ± 12.59	105	49.12 ± 17.63	-2.53	0.4620
N/T RNFL thickness ratio	80	1.18 ± 0.36	96	1.11 ± 0.38	0.05	0.4698
Mean RNFL thickness, NS (μm)	83	92.52 ± 21.93	97	91.30 ± 23.75	4.08	0.3536
Mean RNFL thickness, N (μm)	81	69.02 ± 20.87	96	64.47 ± 15.35	6.82	0.0530
Mean RNFL thickness, NI (μm)	80	103.4 ± 22.25	103	102.5 ± 28.77	1.80	0.7393
Mean RNFL thickness, TI (μm)	81	122.2 ± 27.08	103	122.4 ± 24.99	0.81	0.8661
Mean RNFL thickness, T (μm)	82	60.55 ± 17.21	105	62.03 ± 18.19	-1.05	0.7709
Mean RNFL thickness, TS (μm)	83	116.8 ± 23.08	102	114.2 ± 21.70	6.41	0.1393

MS = multiple sclerosis; RNFL = retinal nerve fiber layer; PMB = papillomacular bundle thickness; SD = standard deviation.
^aMultivariable linear mixed effects model adjusted for ON eye status, age, disease duration, sex, baseline EDSS, and DMT.

limitations of our study were the cross-sectional design, the lack of more quantitative data on smoking history (e.g. pack-year history), the absence of a control population of smokers and non-smokers, and the division between ever-smokers and never-smokers (i.e. not considering the distinction between past and active smokers). In regard to quantitative smoking data, previous studies showed that cigarette smoking becomes a relevant factor past a given dose threshold. By the same token, quitting can reverse at least some of the negative effects of smoking: the date of quitting is likely to be informative.^{11,35} Therefore, a precise smoking history and pack-year values may be integral to an optimal analysis of the impact of cigarette smoking.

In conclusion, this study suggests that cigarette smoking history may not have an impact on retinal deterioration in MS. The question is likely more complex than currently appreciated. Therefore, longitudinal studies will be crucial to identify how retinal structures and other sites in the CNS are selectively affected by cigarette smoking.

Our study suggests that compartments within the CNS respond differentially to neurodegenerative stressors, and that this factor should be accounted for when considering interventional studies using surrogate markers such as retinal measures for neurodegeneration in MS.

Conflict of Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The CLIMB study is supported in part by grants from the Nancy Davis Foundation without walls and Merck-Serono.

ORCID iD

Maria Assunta Rocca  <http://orcid.org/0000-0003-2358-4320>

Massimo Filippi  <http://orcid.org/0000-0002-5485-0479>
Tanuja Chitnis  <http://orcid.org/0000-0002-9897-4422>

Supplemental Material

Supplemental material for this article is available online.

References

1. Ascherio A, Munger KL, White R, et al. Vitamin D as an early predictor of multiple sclerosis activity and progression. *JAMA Neurol* 2014; 71: 306–314.
2. Belbasis L, Bellou V, Evangelou E, et al. Environmental risk factors and multiple sclerosis: an umbrella review of systematic reviews and meta-analyses. *Lancet Neurol* 2015; 14: 263–273.
3. Hernan MA, Jick SS, Logroscino G, et al. Cigarette smoking and the progression of multiple sclerosis. *Brain* 2005; 128: 1461–1465.
4. Zivadinov R, Weinstock-Guttman B, Hashmi K, et al. Smoking is associated with increased lesion volumes and brain atrophy in multiple sclerosis. *Neurology* 2009; 73: 504–510.
5. Healy BC, Ali EN, Guttmann CR, et al. Smoking and disease progression in multiple sclerosis. *Arch Neurol* 2009; 66: 858–864.
6. Degelman ML and Herman KM. Smoking and multiple sclerosis: a systematic review and meta-analysis using the Bradford Hill criteria for causation. *Mult Scler Relat Disord* 2017; 17: 207–216.
7. Di Pauli F, Reindl M, Ehling R, et al. Smoking is a risk factor for early conversion to clinically definite multiple sclerosis. *Mult Scler* 2008; 14: 1026–1030.
8. Poorolajal J, Bahrami M, Karami M, et al. Effect of smoking on multiple sclerosis: a meta-analysis. *J Public Health* 2017; 39: 312–320.
9. Wingerchuk DM. Smoking: effects on multiple sclerosis susceptibility and disease progression. *Ther Adv Neurol Disord* 2012; 5: 13–22.
10. Ramanujam R, Hedstrom AK, Manouchehrinia A, et al. Effect of smoking cessation on multiple sclerosis prognosis. *JAMA Neurol* 2015; 72: 1117–1123.
11. D’Hooghe MB, Haentjens P, Nagels G, et al. Alcohol, coffee, fish, smoking and disease progression in multiple sclerosis. *Eur J Neurol* 2012; 19: 616–624.
12. Gabilondo I, Martinez-Lapiscina EH, Martinez-Heras E, et al. Trans-synaptic axonal degeneration in the visual pathway in multiple sclerosis. *Ann Neurol* 2014; 75: 98–107.
13. Pfueller CF, Brandt AU, Schubert F, et al. Metabolic changes in the visual cortex are linked to retinal nerve fiber layer thinning in multiple sclerosis. *PLoS One* 2011; 6: e18019.
14. Sinnecker T, Oberwahrenbrock T, Metz I, et al. Optic radiation damage in multiple sclerosis is associated with visual dysfunction and retinal thinning—an ultrahigh-field MR pilot study. *Eur Radiol* 2015; 25: 122–131.
15. Reich DS, Smith SA, Gordon-Lipkin EM, et al. Damage to the optic radiation in multiple sclerosis is associated with retinal injury and visual disability. *Arch Neurol* 2009; 66: 998–1006.
16. Klistorner A, Garrick R, Barnett MH, et al. Axonal loss in non-optic neuritis eyes of patients with multiple sclerosis linked to delayed visual evoked potential. *Neurology* 2013; 80: 242–245.
17. Green AJ, McQuaid S, Hauser SL, et al. Ocular pathology in multiple sclerosis: retinal atrophy and inflammation irrespective of disease duration. *Brain* 2010; 133: 1591–1601.
18. Wang AL, Lukas TJ, Yuan M, et al. Changes in retinal pigment epithelium related to cigarette smoke: possible relevance to smoking as a risk factor for age-related macular degeneration. *PLoS One* 2009; 4: e5304.
19. Gao Z, Nissen JC, Ji K, et al. The experimental autoimmune encephalomyelitis disease course is modulated by nicotine and other cigarette smoke components. *PLoS One* 2014; 9: e107979.
20. Hedstrom AK, Baarnhielm M, Olsson T, et al. Tobacco smoking, but not Swedish snuff use, increases the risk of multiple sclerosis. *Neurology* 2009; 73: 696–701.
21. Smith KJ, Kapoor R, Hall SM, et al. Electrically active axons degenerate when exposed to nitric oxide. *Ann Neurol* 2001; 49: 470–476.
22. Hoser G, Domagala-Kulawik J, Droszcz P, et al. Lymphocyte subsets differences in smokers and non-smokers with primary lung cancer: a flow cytometry analysis of bronchoalveolar lavage fluid cells. *Med Sci Monit* 2003; 9: 310–315.
23. Odoardi F, Sie C, Streyll K, et al. T cells become licensed in the lung to enter the central nervous system. *Nature* 2012; 488: 675–679.
24. Velilla S, Garcia-Medina JJ, Garcia-Layana A, et al. Smoking and age-related macular degeneration: review and update. *J Ophthalmol* 2013; 2013: 895147.
25. Grzybowski A and Holder GE. Tobacco optic neuropathy (TON) – the historical and present concept of the disease. *Acta Ophthalmol* 2011; 89: 495–499.
26. Syc SB, Saidha S, Newsome SD, et al. Optical coherence tomography segmentation reveals ganglion cell layer pathology after optic neuritis. *Brain* 2012; 135: 521–533.
27. Gordon-Lipkin E, Chodkowski B, Reich DS, et al. Retinal nerve fiber layer is associated with brain atrophy in multiple sclerosis. *Neurology* 2007; 69: 1603–1609.
28. Petzold A, de Boer JF, Schippling S, et al. Optical coherence tomography in multiple sclerosis: a systematic review and meta-analysis. *Lancet Neurol* 2010; 9: 921–932.
29. Ratchford JN, Saidha S, Sotirchos ES, et al. Active MS is associated with accelerated retinal ganglion cell/inner plexiform layer thinning. *Neurology* 2013; 80: 47–54.

30. Saidha S, Al-Louzi O, Ratchford JN, et al. Optical coherence tomography reflects brain atrophy in multiple sclerosis: a four-year study. *Ann Neurol* 2015; 78: 801–813.
31. Dervisogullari MS, Totan Y, Tenlik A, et al. Effects of cigarette smoking on choroidal and retinal thickness and ocular pulse amplitude. *Cutan Ocul Toxicol* 2015; 34: 217–221.
32. Sigler EJ, Randolph JC, Calzada JI, et al. Smoking and choroidal thickness in patients over 65 with early-atrophic age-related macular degeneration and normals. *Eye* 2014; 28: 838–846.
33. Ulas F, Celik F, Dogan U, et al. Effect of smoking on choroidal thickness in healthy smokers. *Curr Eye Res* 2014; 39: 504–511.
34. Demirci S, Gunes A, Demirci S, et al. The effect of cigarette smoking on retinal nerve fiber layer thickness in patients with migraine. *Cutan Ocul Toxicol* 2016; 35: 21–25.
35. Manouchehrinia A, Tench CR, Maxted J, et al. Tobacco smoking and disability progression in multiple sclerosis: United Kingdom cohort study. *Brain* 2013; 136: 2298–2304.