

Research article

Open Access

Small noncoding RNA GcvB is a novel regulator of acid resistance in *Escherichia coli*

Ye Jin¹, Rory M Watt², Antoine Danchin³ and Jian-dong Huang*¹

Address: ¹Department of Biochemistry, Li Ka Shing Faculty of Medicine, University of Hong Kong, Pok Fu Lam, Hong Kong SAR, PR China, ²Oral Biosciences, Faculty of Dentistry, University of Hong Kong, Pok Fu Lam, Hong Kong SAR, PR China and ³Genetics of Bacterial Genomes, CNRS URA2171, Institut Pasteur, 28, rue du Docteur Roux, Paris Cedex 15, 75724, France

Email: Ye Jin - yejin@hkucc.hku.hk; Rory M Watt - rmwatt@hku.hk; Antoine Danchin - antoine.danchin@normalesup.org; Jian-dong Huang* - jdhuang@hkucc.hku.hk

* Corresponding author

Published: 20 April 2009

Received: 26 January 2009

BMC Genomics 2009, 10:165 doi:10.1186/1471-2164-10-165

Accepted: 20 April 2009

This article is available from: <http://www.biomedcentral.com/1471-2164/10/165>

© 2009 Jin et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The low pH environment of the human stomach is lethal for most microorganisms; but not *Escherichia coli*, which can tolerate extreme acid stress. Acid resistance in *E. coli* is hierarchically controlled by numerous regulators among which are small noncoding RNAs (sncRNA).

Results: In this study, we individually deleted seventy-nine sncRNA genes from the *E. coli* K12-MG1655 chromosome, and established a single-sncRNA gene knockout library. By systematically screening the sncRNA mutant library, we show that the sncRNA GcvB is a novel regulator of acid resistance in *E. coli*. We demonstrate that GcvB enhances the ability of *E. coli* to survive low pH by upregulating the levels of the alternate sigma factor RpoS.

Conclusion: GcvB positively regulates acid resistance by affecting RpoS expression. These data advance our understanding of the sncRNA regulatory network involved in modulating acid resistance in *E. coli*.

Background

The ability to survive the extremely low pH environment present within the stomach (pH values < 2.5) is essential for colonization of the intestine by commensals and for the pathogenesis of enteric bacteria. Three distinct acid resistance (AR) systems are responsible for protecting bacterial cells from acid shock [1-3]. AR system 1 is a glucose-repressed system and does not require any external amino acids to function at low pH [1]. Unidentified components of this system are controlled by the alternate sigma factor RpoS, with the underlying mechanism remaining unclear [1]. In contrast, AR systems 2 and 3 provide AR by consuming intracellular protons via the glutamate (AR2) and

arginine (AR3) decarboxylation reactions [2,3]. AR2 is the most efficient AR system. It involves two isozymes of glutamate decarboxylase, encoded by genes *gadA* and *gadB*, and a membrane-associated antiporter (*GadC*) that exchanges external glutamate for the intracellular decarboxylation product γ -aminobutyric acid (GABA) [2].

AR2 is hierarchically regulated by multiple regulators, which include small noncoding RNAs (sncRNA) DsrA and possibly GadY. DsrA activates the expression of *rpoS* [4] and several AR genes including *hdeAB*, *gadAX*, and *gadBC* [5]. GadY, whose effects on AR have yet to be determined, controls the synthesis of GadX and GadW, which are reg-

ulators of the GadA and GadB glutamate decarboxylases [6,7]. In recent years, an increasing number of sncRNAs have been identified, but the functions of the vast majority of these are unknown. This prompted us to investigate whether sncRNAs in addition to DsrA and GadY, were involved in AR; and if so, to establish how they exerted their regulatory actions.

In the present study, seventy-nine snRNAs identified prior to 2006 [8-10] were selected for the construction of a single-gene knockout library within *E. coli* K12-MG1655. Comparison of each mutant with the wild type strain enabled us to identify the sncRNA GcvB as a previously unknown regulator of AR in *E. coli*. We demonstrate that GcvB positively regulates AR predominantly by upregulating the levels of the RpoS transcription factor (Sigma S). Our findings provide further insight into the regulatory roles played by sncRNAs in AR, which contributes to virulence of *E. coli*.

Results

Construction of single-sncRNA gene knockout mutant collection in *E. coli* K-12 MG1655

To systematically characterize sncRNAs, we established a single-sncRNA gene knockout library within *E. coli* K12-MG1655. Recombineering techniques were used to replace the target gene with a selectable chloramphenicol-resistance cassette, which was generated by polymerase chain reaction (PCR) using primers containing homology (45 nt) to the gene-flanking regions. This homologous recombination-based approach enabled precise gene deletion with minimal interference to adjacent genes (polar effects) [11,12]. In total, seventy-nine genes encoding sncRNAs identified prior to 2006 [8-10] were deleted individually from the MG1655 chromosome (see additional file 1).

The sncRNA GcvB is a novel regulator of AR in *E. coli*

We investigated the involvement of sncRNAs in the AR of *E. coli* by challenging the sncRNA mutants as well as the wild type strain by incubation in LB medium at pH 2.0 (for 15 or 30 minutes), and comparing their percent survival and/or rates of recovery after pH neutralization (termed acid recovery). Unless otherwise stated, bacterial cells were grown for 5 hrs to the early stationary phase before the acid challenge. As expected, cells lacking DsrA, which is a known AR regulator, were sensitive to the acid challenge, confirming previous findings (Fig. 1A) [4,5]. However, deleting *gadY* did not affect acid survival (Fig. 1A), suggesting that GadY is not required for AR in spite of its positive control of GadX, a transcriptional activator of the GadA and GadB glutamate decarboxylases [6,7]. Interestingly, the absence of the sncRNA GcvB resulted in a significant reduction in acid survival (Fig. 1A, B). The same AR phenotypes in the sncRNA mutants were also




Figure 1

Regulatory role of the small noncoding RNA GcvB in acid resistance in *Escherichia coli*. (A) MG1655-K12 wild type strain (wt) and various deletion mutants were grown in Luria Broth broth (pH 7.0) for 5 hr at 37°C. Cells were acid shocked at pH 2.0 for 30 min, and then neutralized by adding alkalinized LB (pH 9.3). Cells were allowed to grow for an additional 3 hrs before measurement of optical density at 600 nm (OD600). Two-tailed paired t tests were employed to compare each mutant with the wt in terms of acid resistance. (B) Because the *gcvB* mutant displayed a lower growth rate than the wt (data not shown), percent acid survival was determined to confirm the data of acid recovery measured as OD600. Cell viability was determined by plate counts and percent acid survival was calculated as the number of cfu remaining after the 30 min-acid treatment divided by the initial cfu at time zero (quoted as a percentage). (C) A plasmid expressing GcvB under the control of its native promoter was transformed into a *gcvB* mutant of *E. coli* MG1655. The mutant carrying an empty vector was used as a control. Cells were incubated for 5 hrs before being acid challenged for 15 min. Error bars represent standard deviation (* P -value < 0.05).

observed during the exponential growth phase (data not shown), suggesting that GcvB exerts constant AR regulatory functions throughout the various growth phases. We also individually deleted *gcvA* and *ygdl* which are the two genes immediately adjacent to *gcvB*. Both mutants displayed comparable AR as the wild type (Fig. 1A). This shows that the observed acid vulnerability in the *gcvB* mutant is not due to the disruption of the activities of its adjacent genes.

Next, we cloned the gene encoding GcvB (together with its native promoter) into plasmid pET32a (+) and overexpressed the sncRNA in the *E. coli* MG1655 *gcvB* null mutant. The *gcvB* mutant carrying an empty vector served as a negative control. We found that overproducing GcvB restored the ability of the *gcvB* mutant to survive at low pH (Fig 1C). These observations together with the knockout data revealed that GcvB acts as a positive regulator of AR.

The known GcvB target *OppA* plays a minor, if any, role in GcvB-mediated AR

GcvB has been demonstrated to repress the translation of *oppA* and *dppA* in *E. coli* by masking the ribosome binding sites of their mRNAs, which encode periplasmic substrate-binding proteins of ABC uptake systems for amino acids and peptides [13-15]. To determine whether these two proteins played a role in GcvB-mediated AR, we individually deleted their encoding genes, and examined the ability of the resultant mutants to survive at low pH. We found that deleting *oppA* slightly increased AR, as revealed by their acid recovery (OD600) and percent acid survival (cfu) levels, while the absence of *dppA* had little effect on AR (Fig. 2A, B). Thus *DppA* is not involved in GcvB-mediated AR, and *OppA* plays a minor, if any, role in this process. We reasoned that the GcvB sncRNA may be exerting its effects on AR via a previously unidentified molecular target.

GcvB mediates AR by upregulating *RpoS* expression

RpoS is the central regulator of glutamate-dependent AR2 genes such as *gadA*, *gadBC* [1], *gadX* [16], *gadE* [17], and *gadY* [6,7]. It is also required for the AR1 system [1,18]. We therefore set out to determine whether GcvB regulated AR by modulating *RpoS* expression levels. We found that deleting *gcvB* did not lead to a further reduction in acid survival levels in an *rpoS* mutant background (Fig. 3A). More importantly, the absence of *rpoS* completely abolished AR elicited by the plasmid-encoded GcvB (Fig. 3B), suggesting that GcvB-mediated AR is dependent on *RpoS*. Next, an *rpoS-lacZ* translational fusion was constructed to examine whether the presence or absence of the GcvB sncRNA significantly affected *rpoS* expression levels. Deleting *gcvB* reduced the expression levels of the *rpoS-lacZ* fusion by 50% (Fig. 4A), whereas *gcvB* overexpression increased it 3-fold (Fig. 4B). Furthermore, deleting *oppA*,




Figure 2
Effects of individually deleting *oppA* and *dppA* on acid resistance in *Escherichia coli*. The acid-stress survival abilities of two individual deletion mutant strains lacking previously identified targets of GcvB were compared with those of the MG1655 wild type (wt) strain. The abilities of strains to survive a 30 minute acid shock were quantified by two methods: (A) acid recovery measured as optical density at 600 nm (OD600). (B) Percent survival determined by colony forming units (cfu). Error bars represent standard deviation (* P-value < 0.05).

whose absence had previously been shown to slightly increase AR levels, had no effect on *rpoS-lacZ* expression (Fig. 4C). This indicates that the GcvB-mediated regulation of *RpoS* expression does not involve *OppA*.

We then investigated whether GcvB indirectly controls *RpoS* production via an upstream regulator. In the literature, there are numerous reports outlining mechanisms by which the synthesis of *RpoS* is regulated at the transcriptional [19-22], translational and posttranslational levels [23-25]. Of these regulators of *RpoS* expression, Crp [20], H-NS [23], GadX [22], and a hybrid sensor kinase called BarA [21] have previously been associated with AR. Although BarA acts as a positive regulator of *rpoS* transcription, its effect is restricted to the exponential phase. It




Figure 3
Effects of deleting *rpoS* on GcvB-mediated acid resistance. (A) Acid resistance in an *rpoS* mutant and an *rpoS gcvB* double mutant. (B) Effects of GcvB overproduction on acid resistance in an *rpoS* mutant. Cell cultures of *E. coli* MG1655 mutant strains were grown in LB broth (pH 7.0) at 37°C for 5 hrs and then acid challenged (pH 2.0) for 15 min. After neutralization by adding alkalinized LB (pH 9.3), cells were cultured for an additional 3 hrs before measurement of optical density at 600 nm (OD600 nm). Error bars represent standard deviation (* *P*-value < 0.05).

is therefore unlikely that BarA is a potential target for GcvB. Consequently, we examined whether GcvB exerts its positive control of RpoS through Crp, H-NS or GadX. We found that deleting *gcvB* reduced acid survival levels in an *hns* mutant background. This was also the case for *crp*- or *gadX*-deficient strains (Fig. 5). Consistent with the knockout data, *gcvB* overexpression increased AR levels in the *hns*-, *crp*- or *gadX*-deficient strains (Fig. 6). These results suggest that Hns, Crp and GadX are not involved in the GcvB-mediated regulation of *rpoS* expression.

rpoS transcription is driven by a major promoter *rpoSp* and two weak promoters *nlpDp1* and 2 [26]. To examine whether GcvB affected *rpoS* expression at the transcrip-




Figure 4
Effects of GcvB sncRNA and OppA protein on the expression of RpoS in a chromosomal *rpoS-lacZ* reporter strain. (A) Effects of deleting *gcvB* on the expression of an *rpoS-lacZ* reporter gene fusion in *E. coli* MG1655. (B) Effect of GcvB overproduction on the expression of the *rpoS-lacZ* fusion. GcvB production was driven by its native promoter on a pET32a plasmid construct. (C) Effect of deleting *oppA* on the expression of the *rpoS-lacZ* reporter. Beta-galactosidase activities due to translation of chromosomal *lacZ* fusions were normalized to those of MG1655 carrying an *rpoS-lacZ* fusion with or without an empty vector (controls). Error bars represent standard deviation (* *P*-value < 0.05).




Figure 5
Effects of deleting *gcvB* on acid resistance in mutants lacking *hns*, *crp* or *gadX*. Cell cultures of *E. coli* MG1655 mutant strains containing single deletions of the *hns*, *crp* or *gadX* genes were grown in LB broth (pH 7.0) at 37°C for 5 hrs and then acid challenged (pH 2.0) for 15 min (*gadX* mutant) or 30 min (*hns* and *crp* mutants). After neutralization by adding alkalized LB (pH 9.3), cells were cultured for an additional 3 hrs before measurement of optical density at 600 nm (OD600 nm). Error bars represent standard deviation (* *P*-value < 0.05).

tional level, we deleted the region spanning the *rpoSp*, *nlpDp1* and 2 promoter sites (Fig. 4). As removing these promoters will eliminate *rpoS* expression, a constitutive promoter encoded in the knockout cassette (described in Materials and Methods) was used to drive *rpoS* transcription in the absence of *rpoSp*, *nlpDp1* and 2. The resulting mutant was referred to as *d*₅₆₇₋₁₃₄₂. As shown in Fig. 6, replacing the native promoters of *rpoS* with this heterologous promoter had little effect on GcvB-induced AR. These results indicate that GcvB does not regulate *rpoS* expression at the transcriptional level.

Discussion

In this study, we constructed a library comprising single-gene knockout mutants of seventy-nine sncRNA genes in *E. coli*. We used this mutant collection to identify sncRNAs involved in acid resistance (AR) within this bacterium. Consequently, we were able to identify the sncRNA GcvB as a novel regulator of AR.

DppA and OppA, which were previously identified as targets of GcvB, were shown not to be responsible for GcvB-mediated AR. While the *dppA* deletion did not affect AR, the absence of OppA slightly improved the ability of *E. coli* to survive under extreme acid conditions. Since GcvB-mediated inhibition of *oppA* and *dppA* only weakly increased AR, we reasoned that the GcvB sncRNA may be regulating AR via some other mechanistic pathway.




Figure 6
Effects of GcvB overexpression on acid resistance in mutants lacking *hns*, *crp*, *gadX* or native promoters of *rpoS*. The mutant lacking the native promoters of *rpoS* was named *d*₅₆₇₋₁₃₄₂. Cell cultures of *E. coli* MG1655 strains containing single deletions were grown in LB broth (pH 7.0) at 37°C for 5 hrs and then acid challenged (pH 2.0) for 15 min (*gadX* mutant) or 30 min (*hns* and *crp* mutants and *d*₅₆₇₋₁₃₄₂). After neutralization by alkalized LB (pH 9.3), cells were cultured for an additional 3 hrs before measurement of optical density at 600 nm (OD600 nm). Error bars represent standard deviation (* *P*-value < 0.05).

Upon *rpoS* deletion, GcvB had no influence on AR. Furthermore, we found that GcvB positively regulated *rpoS* expression. Deleting *oppA* had no effect on the levels of RpoS, providing evidence that the GcvB-mediated control of RpoS does not involve the OppA protein. Also, GcvB does not regulate RpoS via H-NS, Crp, or GadX; as GcvB-mediated AR was unaffected when these genes were singularly deleted. The region spanning from -567 to -1342 nt harbors all three of the of the *rpoS* promoters (*rpoSp*, *nlpDp1* and 2). Replacing these native promoters with the constitutive promoter from a Cm knockout cassette had no affect on GcvB-mediated AR. This suggests that GcvB does not regulate *rpoS* expression at the transcriptional level.

Like GcvB, the sncRNAs DsrA and RprA also act as positive regulators of AR in *E. coli*. DsrA and RprA bind to the same region of the 5' leader of *rpoS* mRNA. A sequence within this region was previously predicted to form a self-inhibitory stem loop structure that probably occludes the Shine-Dalgarno ribosome binding site and prevents translation [24,27]. DsrA [24,27] and RprA [28,29] bind to this stem-loop structure, free the ribosome binding site, and thereby mediate *rpoS* translation. This region also contains RNase III cleavage sites that result in rapid decay of the *rpoS* transcript [30]. The base-pairing of DsrA within this region stabilizes *rpoS* mRNA by creating an alternative RNase III cleavage site [30]. As for GcvB, it regulates *rpoS* expression

by an unknown mechanism. Computational sequence analyses using the RNAhybrid server [31] have indicated that GcvB does not contain any extensive regions of sequence complementary to the 5' leader of *rpoS* mRNA (data not shown). As such, the mechanism by which GcvB regulates *rpoS* expression remains a subject of further investigation.

Conclusion

Here we reveal that the sncRNA GcvB positively regulates the ability of *E. coli* to survive low pH conditions by upregulating the expression of RpoS. Our findings provide insight into the control of AR by GcvB. However, further detailed studies will be required to investigate the precise nature of the GcvB/*rpoS* interaction at the molecular level.

Methods

Bacterial strains, plasmids, and growth conditions

E. coli K12-MG1655 was used for all gene deletion and overexpression experiments. The *E. coli* expression vector pET32a (+) (Novagen, Madison, WI) was used for the overexpression of sncRNAs. All bacterial strains were grown at 37°C, with shaking at 230 rpm, in Luria-Bertani (LB) medium supplemented with antibiotics when required. Bacterial growth was monitored by measuring optical density at 600 nm (OD600). The antibiotics ampicillin (50 µg/ml), kanamycin (50 µg/ml), and chloramphenicol (12.5 µg/ml) were used for selection.

Gene deletions

Gene deletion mutations were constructed using the λ -Red recombination system. *E. coli* MG1655 was transformed with plasmid pSim6 (a gift from Dr. Donald Court) [32] on which the expression of the λ recombination proteins is induced at 42°C. PCR fragments encompassing a loxP-cm-loxP ('floxed') chloramphenicol resistance cassette with homology (45 nt) to the regions immediately flanking each deletion locus were transformed into MG1655 harboring pSim6. After induction of λ red, recombinants were selected for chloramphenicol (cm) resistance, and were further verified by colony PCR.

Gene cloning

The selectable loxP-cm-loxP cassette was first inserted immediately after the stop codon of the gene of interest on the chromosome (as described above). We then used recombineering to 'clone' this gene of interest preceded by its native promoter and 'floxed' cm cassette downstream of the T7 promoter in a pET32a expression vector. Specifically, we PCR amplified the gene of interest (including its native promoter) and adjacent 'floxed' cm cassette using primers that contained homology (45 nt) to the plasmid insertion site. The PCR product and expression vector were co-transformed into DY330[12] after induction of λ red. Recombinants were selected for chloramphenicol

resistance and verified by PCR and sequencing. The native promoter drove the gene overexpression in MG1655 without induction.

Construction of chromosomal lacZ translational fusions and beta-galactosidase assays

First, the loxP-cm-loxP selectable cassette was inserted immediately after the stop codon of the *lacZ* gene on the MG1655 chromosome using λ red recombination (as described above). Next, the *lacZ*-loxP-cm-loxP cassette was PCR amplified and inserted (in frame) immediately prior to the stop codon of the target gene on the chromosome. The inserted *lacZ* fragment started from the 8th codon of *lacZ* gene and co-transcribed and -translated with the fused genes. The expression of gene-*lacZ* fusions was quantified using a beta-galactosidase assay kit from Thermo Fisher Scientific (Rockford, Illinois, USA).

Acid resistance assay

Overnight cultures were diluted 1:50 and incubated at 37°C for 5 hrs in LB medium. 3 volumes of acidified LB medium (pH 1.9) was then added to 1 volume of the cell culture so that the final pH value was approximately 2.0. The acid challenge was maintained for 15 min for cells that were acid sensitive, and 30 min for acid-resistant cells. The acid challenge was stopped by adding 3 volumes of alkalized LB medium (pH 9.3), making the final pH value approximately 7.0. For samples that displayed similar growth rates, the neutralized cell cultures were allowed to grow for 3 hrs following the acid challenge, and acid survival levels were determined by measurement of optical density at 600 nm (OD600, termed acid recovery). When samples with different growth rates were examined, cell viability was determined by plate counts and percentage acid survival was calculated as the number of number of colony forming units (cfu) remaining after the acid treatment divided by the initial cfu at time zero.

Statistical analysis

Two-tailed paired t-tests were used for the comparison of means obtained from the beta-galactosidase and acid survival assays. *P* values of < 0.05 were considered statistically significant.

Abbreviations

AR: acid resistance; sncRNA: small noncoding RNA; UTR: untranslated region.

Authors' contributions

YJ planned and designed the experiments, performed the experiments and the data analysis, wrote the main draft of the paper, and generated the figures. RW conceived of the study and helped to draft the manuscript. AD aided in the experimental design and in editing the manuscript. JH provided guidance over the entire project, and aided in

data analysis and editing of the manuscript. All authors read and approved the final manuscript.

Additional material

Additional file 1

Small noncoding RNAs (sncRNA) that were selected for construction of a single-sncRNA gene knockout library within E. coli K12-MG1655. This table summarizes positions and functions of the small noncoding RNAs that were included in a single-sncRNA gene knockout library within E. coli K12-MG1655.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-10-165-S1.doc>]

Acknowledgements

We thank D.L. Court for providing plasmid pSim6 for the recombineering experiments. This work was supported by HKU CRCG Seed Funding Programme for Basic Research and by a grant (HKU 7485/06M) from the Research Grant Council of Hong Kong to JDH. AD acknowledges support from the PROACTYS programme, grant CT-2006-029104 in an effort to define genes essential for the construction of a synthetic cell.

References

- Castanie-Cornet MP, Penfound TA, Smith D, Elliott JF, Foster JW: **Control of acid resistance in Escherichia coli.** *J Bacteriol* 1999, **181(11)**:3525-3535.
- Hersh BM, Farooq FT, Barstad DN, Blankenhorn DL, Slonczewski JL: **A glutamate-dependent acid resistance gene in Escherichia coli.** *J Bacteriol* 1996, **178(13)**:3978-3981.
- Lin J, Smith MP, Chapin KC, Baik HS, Bennett GN, Foster JW: **Mechanisms of acid resistance in enterohemorrhagic Escherichia coli.** *Appl Environ Microbiol* 1996, **62(9)**:3094-3100.
- Sledjeski DD, Gupta A, Gottesman S: **The small RNA, DsrA, is essential for the low temperature expression of RpoS during exponential growth in Escherichia coli.** *EMBO J* 1996, **15(15)**:3993-4000.
- Lease RA, Smith D, McDonough K, Belfort M: **The small noncoding DsrA RNA is an acid resistance regulator in Escherichia coli.** *J Bacteriol* 2004, **186(18)**:6179-6185.
- Opdyke JA, Kang JG, Storz G: **GadY, a small-RNA regulator of acid response genes in Escherichia coli.** *J Bacteriol* 2004, **186(20)**:6698-6705.
- Tramonti A, De Canio M, De Biase D: **GadX/GadW-dependent regulation of the Escherichia coli acid fitness island: transcriptional control at the gadY-gadW divergent promoters and identification of four novel 42-bp GadX/GadW-specific binding sites.** *Mol Microbiol* 2008, **70(4)**:965-982.
- Griffiths-Jones S, Moxon S, Marshall M, Khanna A, Eddy SR, Bateman A: **Rfam: annotating non-coding RNAs in complete genomes.** *Nucleic Acids Res* 2005:D121-124.
- Vallenet D, Labarre L, Rouy Z, Barbe V, Bocs S, Cruveiller S, Lajus A, Pascal G, Scarpelli C, Medigue C: **MaGe: a microbial genome annotation system supported by synteny results.** *Nucleic Acids Res* 2006, **34(1)**:53-65.
- Vogel J, Bartels V, Tang TH, Churakov G, Slatger-Jager JG, Huttenhofer A, Wagner EG: **RNomics in Escherichia coli detects new sRNA species and indicates parallel transcriptional output in bacteria.** *Nucleic Acids Res* 2003, **31(22)**:6435-6443.
- Ellis HM, Yu D, DiTizio T, Court DL: **High efficiency mutagenesis, repair, and engineering of chromosomal DNA using single-stranded oligonucleotides.** *Proc Natl Acad Sci USA* 2001, **98(12)**:6742-6746.
- Yu D, Ellis HM, Lee EC, Jenkins NA, Copeland NG, Court DL: **An efficient recombination system for chromosome engineering in Escherichia coli.** *Proc Natl Acad Sci USA* 2000, **97(11)**:5978-5983.
- Urbanowski ML, Stauffer LT, Stauffer GV: **The gcvB gene encodes a small untranslated RNA involved in expression of the dipeptide and oligopeptide transport systems in Escherichia coli.** *Mol Microbiol* 2000, **37(4)**:856-868.
- Urban JH, Vogel J: **Translational control and target recognition by Escherichia coli small RNAs in vivo.** *Nucleic Acids Res* 2007, **35(3)**:1018-1037.
- Sharma CM, Darfeuille F, Plantinga TH, Vogel J: **A small RNA regulates multiple ABC transporter mRNAs by targeting C/A-rich elements inside and upstream of ribosome-binding sites.** *Genes Dev* 2007, **21(21)**:2804-2817.
- Tramonti A, Visca P, De Canio M, Falconi M, De Biase D: **Functional characterization and regulation of gadX, a gene encoding an AraC/XylS-like transcriptional activator of the Escherichia coli glutamic acid decarboxylase system.** *J Bacteriol* 2002, **184(10)**:2603-2613.
- Sayed AK, Odom C, Foster JW: **The Escherichia coli AraC-family regulators GadX and GadW activate gadE, the central activator of glutamate-dependent acid resistance.** *Microbiology* 2007, **153(Pt 8)**:2584-2592.
- Small P, Blankenhorn D, Welty D, Zinser E, Slonczewski JL: **Acid and base resistance in Escherichia coli and Shigella flexneri: role of rpoS and growth pH.** *J Bacteriol* 1994, **176(6)**:1729-1737.
- Mika F, Hengge R: **A two-component phosphotransfer network involving ArcB, ArcA, and RssB coordinates synthesis and proteolysis of sigmaS (RpoS) in E. coli.** *Genes Dev* 2005, **19(22)**:2770-2781.
- Lange R, Hengge-Aronis R: **The cellular concentration of the sigma S subunit of RNA polymerase in Escherichia coli is controlled at the levels of transcription, translation, and protein stability.** *Genes Dev* 1994, **8(13)**:1600-1612.
- Mukhopadhyay S, Audia JP, Roy RN, Schellhorn HE: **Transcriptional induction of the conserved alternative sigma factor RpoS in Escherichia coli is dependent on BarA, a probable two-component regulator.** *Mol Microbiol* 2000, **37(2)**:371-381.
- Hommais F, Krin E, Coppee JY, Lacroix C, Yeramian E, Danchin A, Bertin P: **GadE (YhiE): a novel activator involved in the response to acid environment in Escherichia coli.** *Microbiology* 2004, **150(Pt 1)**:61-72.
- Brescia CC, Kaw MK, Sledjeski DD: **The DNA binding protein H-Ns binds to and alters the stability of RNA in vitro and in vivo.** *J Mol Biol* 2004, **339(3)**:505-514.
- Lease RA, Cusick ME, Belfort M: **Riboregulation in Escherichia coli: DsrA RNA acts by RNA:RNA interactions at multiple loci.** *Proc Natl Acad Sci USA* 1998, **95(21)**:12456-12461.
- Zhang A, Altuvia S, Tiwari A, Argaman L, Hengge-Aronis R, Storz G: **The OxyS regulatory RNA represses rpoS translation and binds the Hfq (HF-I) protein.** *EMBO J* 1998, **17(20)**:6061-6068.
- Lange R, Fischer D, Hengge-Aronis R: **Identification of transcriptional start sites and the role of ppGpp in the expression of rpoS, the structural gene for the sigma S subunit of RNA polymerase in Escherichia coli.** *J Bacteriol* 1995, **177(16)**:4676-4680.
- Majdalani N, Cunnig C, Sledjeski D, Elliott T, Gottesman S: **DsrA RNA regulates translation of RpoS message by an anti-antisense mechanism, independent of its action as an antisense of transcription.** *Proc Natl Acad Sci USA* 1998, **95(21)**:12462-12467.
- Majdalani N, Hernandez D, Gottesman S: **Regulation and mode of action of the second small RNA activator of RpoS translation, RprA.** *Mol Microbiol* 2002, **46(3)**:813-826.
- Updegrove T, Wilf N, Sun X, Wartell RM: **Effect of Hfq on RprA-RpoS mRNA pairing: Hfq-RNA binding and the influence of the 5' rpoS mRNA leader region.** *Biochemistry* 2008, **47(43)**:11184-11195.
- Resch A, Afonyushkin T, Lombo TB, McDowall KJ, Blasi U, Kabardin VR: **Translational activation by the noncoding RNA DsrA involves alternative RNase III processing in the rpoS 5'-leader.** *RNA* 2008, **14(3)**:454-459.
- Kruger J, Rehmsmeier M: **RNAhybrid: microRNA target prediction easy, fast and flexible.** *Nucleic Acids Res* 2006:W451-454.
- Datta S, Costantino N, Court DL: **A set of recombineering plasmids for gram-negative bacteria.** *Gene* 2006, **379**:109-115.