


Circumstances of clinical fracture events with advancing age in older men

Kristine E. Ensrud^{1,2,*} , Sheena Patel³, Lisa Langsetmo^{2,4}, Peggy M. Cawthon³, Howard A. Fink^{2,4}, John T. Schousboe^{5,6}, Douglas C. Bauer⁷, Jane A. Cauley⁸, Eric S. Orwoll⁹, for the Osteoporotic Fractures in Men (MrOS) Study

¹Division of Epidemiology & Community Health, University of Minnesota, Minneapolis, MN 55454, USA

²Center for Care Delivery and Outcomes Research, VA Health Care System, Minneapolis, MN 55417, USA

³California Pacific Medical Center Research Institute, San Francisco, CA 94107, USA

⁴Department of Medicine, University of Minnesota, Minneapolis, MN 55455, USA

⁵HealthPartners Institute, Bloomington, MN 55425, USA

⁶Division of Health Policy & Management, University of Minnesota, Minneapolis, MN 55455, USA

⁷Department of Medicine, University of California, San Francisco, CA 94115, USA

⁸Department of Epidemiology, University of Pittsburgh, Pittsburgh, PA 15261, USA

⁹Department of Medicine, Oregon Health & Science University, Portland, OR 97239, USA

*Corresponding author: Kristine E. Ensrud, Division of Epidemiology & Community Health, University of Minnesota, Mailing Address: One Veterans Drive (111-0), Minneapolis, MN 55417, USA (ensru001@umn.edu)

Abstract

To characterize fracture circumstances by age at time of fracture among community-dwelling older men, we analyzed incident clinical fractures in the Osteoporotic Fractures in Men study. Participants were contacted every 4 mo to identify fractures confirmed by radiographic reports. Fractures were classified as fall- or non-fall-related and further categorized by degree of trauma: minimal (fall from \leq standing height), moderate (fall on stairs, steps or curb), or severe (fall from $>$ standing height) for fall-related fractures; and minimal (eg, coughing), moderate (eg, collisions with objects during normal activity without falling), or severe (eg, motor vehicle accident) for non-fall-related fractures. Of 2351 clinical fractures in 1424 men during an average follow-up of 9.9 yr, 12.7% occurred at age <75 yr, 15.7% at age 75–79 yr, 26.1% at age 80–84 yr, and 45.5% at age ≥ 85 yr. A total of 1891 fractures (80.4% of all fractures) were the result of a fall. The proportion of fall-related fractures steadily rose with increasing age at time of fracture, ranging from 65.8% in men <75 yr rising to 84.7% in men ≥ 85 yr ($p < .001$). Most fall-related fractures (76.4%) were due to minimal trauma; the proportion of all fractures due to a fall with minimal trauma steadily rose with increasing age ($p < .001$). In contrast, approximately half of non-fall-related fractures (53.5%) were due to severe trauma. The proportion of all fractures due to severe trauma (whether fall-related or not) declined with advancing age ($p < .001$). In conclusion, the most frequently occurring fracture circumstance in older men was a fall from \leq standing height. This circumstance was increasingly common with advancing age occurring in 7 of every 10 fractures in men ≥ 85 yr, while a fracture (fall-related or not) due to severe trauma was less common with advancing age. Findings have implications for development of fracture prevention strategies in older men.

Keywords: older men, fracture circumstances, falls, degree of trauma, epidemiology

Lay Summary

This study characterized fracture circumstances by age at time of fracture in 5994 community-dwelling older men. Of 2351 fractures occurring during 10 yr, 80% were due to a fall. Three-quarters of fall-related fractures were due to minimal trauma, while half of non-fall-related fractures were due to severe trauma. The proportion of fractures that were fall-related with minimal trauma rose with increasing age, while the proportion of fractures with severe trauma (whether fall-related or not) declined with advancing age.

Introduction

Fall prevalence in older adults is modestly higher among women compared with men, but fall rates rise with advancing age in both sexes.^{1–3} Among a nationally representative sample of 149 319 non-institutionalized US adults ≥ 65 yr,⁴ 30.3% of women and 26.5% of men reported falling at least once in the preceding 12 mo; fall prevalence rose in a graded manner with older age from 26.7% among those age 65–74 yr, to 29.8% among those age 75–84 yr, to 36.5% among those age ≥ 85 yr. Similarly, among a representative sample of 4365 community-dwelling Canadian adults age ≥ 65 yr,⁵

fall prevalence steadily rose from 15.8% in men and 17.8% in women age 65–69 up to 23.2% in men and 30.5% in women age ≥ 85 yr. Most falls result in no injury, but 30% of falls result in an injury requiring medical attention or activity restriction and 10%–15% of falls result in fractures.^{1,6} While previous studies have reported higher rates of injurious falls in both sexes with advancing age,^{4,7} it is uncertain whether the proportion of fall-related fractures rises with increasing age, especially among older men who have a lower prevalence of fractures.⁸ Of additional importance, while the percentage of fractures due to severe trauma in older adults is higher in

Received: October 18, 2024. Revised: January 2, 2025. Accepted: January 7, 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of The American Society for Bone and Mineral Research.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

men versus women,^{9,10} it is unknown if degree of trauma in fractures in men varies with increasing age.

Thus, to better characterize fracture circumstances among community-dwelling older men according to age at time of fracture, we analyzed incident clinical fractures experienced by participants enrolled in the Osteoporotic Fractures in Men (MrOS) study.

Materials and methods

Study population

A total of 5994 community-dwelling men ≥ 65 yr old were enrolled in MrOS between 2000 and 2002.¹¹ Participants were recruited from population-based listings in 6 US regions.¹² The study was approved by institutional review boards.

Clinical fractures

Participants were contacted tri-annually after the baseline examination (over 98% of follow-up contacts completed in active survivors) to ask about fractures. Self-reported fractures were confirmed by radiographic reports.¹³ For self-reported spine fractures, the community spinal imaging study was also obtained. Incident clinical vertebral fractures were confirmed by the study radiologist using the Genant method¹⁴ to establish that the community study showed a new deformity of a higher grade than was present in the same vertebra on study spine films obtained prior to the self-reported event.¹⁵

Fracture circumstances

When a fracture was reported, clinic staff interviewed the participant and completed a fracture report form. The fracture was classified as fall- or non-fall-related and coded according to degree of trauma. For fall-related fractures, degree of trauma was classified: minimal (fall from \leq standing height), moderate (fall on stairs, steps or curb), or severe (fall from $>$ standing height). For non-fall-related fractures, degree of trauma was considered: minimal (eg, coughing), moderate (eg, collisions with objects during normal activity without falling), or severe (eg, motor vehicle accident).

Analytical sample

A total of 2351 confirmed fractures experienced by 1424 men during a mean (SD) follow-up of 9.9 (5.7) yr comprised the analytical sample.

Statistical analyses

Fracture (and not participants with versus without incident fracture) was the unit of analysis in our study. Using chi-square tests, the proportion of fall-related (and non-fall-related) fractures and the proportion of all fractures that were fall-related (and non-fall-related) due to a specific degree of trauma were compared across age groups (<75 yr, 75-79 yr, 80-84 yr, ≥ 85 yr) at the time of the fracture.

Results

Of the 2351 fractures, 298 (12.7%) occurred in 221 men age 65-74 yr at the time of fracture; 370 (15.7%) occurred in 299 men age 75-79 yr at the time of fracture; 613 (26.1%) occurred in 454 men age 80-84 yr at the time of fracture; and 1070 (45.5%) occurred in 686 men age ≥ 85 yr at the

time of fracture (Table 1). A total of 1891 fractures (80.4% of all fractures) were due to a fall. The proportion of fall-related fractures rose in a graded manner with increasing age at time of fracture, ranging from 65.8% in men <75 yr rising to 84.7% in men ≥ 85 yr ($p < .001$). The proportion of non-fall-related fractures steadily declined with older age, ranging from 34.2% in men <75 yr declining to 15.3% in men ≥ 85 yr ($p < .001$).

Of the 1891 fall-related fractures, 76.4% were due to minimal trauma, 14.7% were due to moderate trauma, and 8.9% were due to severe trauma. In contrast, of the 460 non-fall-related fractures, 21.3% were due to minimal trauma, 25.2% were due to moderate trauma, and 53.5% were due to severe trauma.

The proportion of all fractures related to a fall and due to minimal trauma rose with increasing age (34.9% of fractures in men <75 yr rising to 70.9% of fractures in men ≥ 85 yr [$p < .001$]), whereas the proportion of all fractures related to fall and due to severe trauma declined with advancing age (18.5% of fractures in men <75 yr declining to 3.8% of fractures in men ≥ 85 yr [$p < .001$]) (Figure 1).

The proportion of all fractures not related to a fall and due to severe trauma declined with increasing age (19.5% of fractures in men <75 yr declining to 8.4% of fractures in men ≥ 85 yr [$p < .001$]). Similarly, the proportion of all fractures not related to a fall and due to moderate trauma declined with advancing age (10.7% of fractures in men <75 yr declining to 2.6% of fractures in men ≥ 85 yr [$p < .001$]).

Discussion


Among this cohort of community-dwelling older men, the proportion of fall-related fractures steadily rose (and the proportion of non-fall-related fractures steadily declined) with increasing age at time of fracture. Fall-related fractures due to minimal trauma were increasingly common with advancing age and comprised an increasing proportion of all fractures. In contrast, the proportion of all fractures due to severe trauma, whether fall-related or not, steadily declined with advancing age.

Fracture circumstances with advancing age have not been previously investigated in men, but our findings are in general agreement with those from a longitudinal global database of 60 393 postmenopausal women that captured incident fractures with annual questionnaires during 3 yr of follow-up.¹⁶ Among 4122 women reporting first incident fractures, the proportion of hip and non-hip non-vertebral fractures with fall as the cause increased with advancing age and the proportion of these fracture types with severe trauma as the cause declined with increasing age.

Study strengths include prospective long-term follow-up, radiographic confirmation of incident fractures and standardized assessment of circumstances, and degree of trauma. Limitations include the lack of diversity in race/ethnicity in the predominantly non-Hispanic White cohort.¹¹ In addition, our study was descriptive in design and the unit of analysis was fracture. We did not evaluate characteristics of men with versus without incident fracture or determine whether any association of a candidate risk factor with incident fracture remained after accounting for potential confounders. Previous MrOS publications have identified independent risk factors for clinical fracture outcomes in community-dwelling older

Table 1. Fracture category according to age group at time of fracture.

Fracture category	Overall (<i>n</i> = 2351)	Age group at time of fracture			
		65-74 yr (<i>n</i> = 298)	75-79 yr (<i>n</i> = 370)	80-84 yr (<i>n</i> = 613)	≥85 yr (<i>n</i> = 1070)
Fall-related fracture, <i>n</i> (%)	1891 (80.4)	196 (65.8)	285 (77.0)	504 (82.2)	906 (84.7)
Non-fall-related fracture, <i>n</i> (%)	460 (19.6)	102 (34.2)	85 (23.0)	109 (17.8)	164 (15.3)
Degree of trauma, <i>n</i> (%)					
Minimal trauma fall (fall from ≤ standing height)	1445 (61.5)	104 (34.9)	212 (57.3)	370 (60.4)	759 (70.9)
Moderate trauma fall (fall on stairs, steps, or curb)	278 (11.8)	37 (12.4)	36 (9.7)	99 (16.2)	106 (9.9)
Severe trauma fall (fall from > standing height)	168 (7.2)	55 (18.5)	37 (10.0)	35 (5.7)	41 (3.8)
Minimal trauma other than a fall	98 (4.2)	12 (4.0)	16 (4.3)	24 (3.9)	46 (4.3)
Moderate trauma other than a fall	116 (4.9)	32 (10.7)	22 (6.0)	34 (5.6)	28 (2.6)
Severe trauma other than a fall	246 (10.5)	58 (19.5)	47 (12.7)	51 (8.3)	90 (8.4)

**Figure 1.** Proportion of clinical fractures in each age group according to fall relationship and degree of trauma. A total of 298 fractures occurred in 221 men age 65-74 yr at time of fracture, 370 fractures occurred in 299 men age 75-79 yr at time of fracture, 613 fractures occurred in 454 men age 80-84 yr at time of fracture, and 1070 fractures occurred in 686 men age ≥85 yr at time of fracture.

men including any non-vertebral,¹³ hip¹⁷ and vertebral¹⁸ fractures.

Our results suggest that fracture prevention strategies in younger old men should focus on minimizing risk-taking behaviors, while those in oldest old men should focus on fall prevention. To better guide development of these strategies, future studies should characterize location (eg, outdoors vs indoors) and specific activity at time of fracture occurrence.

Acknowledgments

None.

Author contributions

Kristine E. Ensrud (Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Writing—original draft),

Sheena Patel (Formal analysis, Validation, Writing—review & editing), Lisa Langsetmo (Conceptualization, Writing—review & editing), Peggy M. Cawthon (Conceptualization, Funding acquisition, Investigation, Writing—review & editing), Howard A. Fink (Writing—review & editing), John T. Schousboe (Writing—review & editing), Douglas C. Bauer (Investigation, Project administration, Writing—review & editing), Jane A. Cauley (Funding acquisition, Investigation, Writing—review & editing), and Eric S. Orwoll (Conceptualization, Funding acquisition, Investigation, Project administration, Writing—review & editing).

Funding

The Osteoporotic Fractures in Men (MrOS) Study is supported by National Institutes of Health funding. The following institutes provide support under the following grant numbers: the National Institute on Aging (NIA: U01 AG027810, U01 AG042124, U01 AG042139, U01 AG042140, U01 AG042143, U01 AG042145, U01 AG042168,

and R01 AG066671), the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS: U01 AR066160), and the National Center for Advancing Translational Sciences (NCATS: UL1 TR002369).

Conflicts of interest

The authors have nothing to disclose.

Data availability

Data from MrOS are available via <https://mrosonline.ucsf.edu/>.

References

- Ensrud KE. Epidemiology of fracture risk with advancing age. *J Gerontol A Biol Sci Med Sci*. 2013;68(10):1236–1242. <https://doi.org/10.1093/gerona/glt092>
- Gale CR, Cooper C, Aihie SA. Prevalence and risk factors for falls in older men and women: the English longitudinal study of ageing. *Age Ageing*. 2016;45(6):789–794. <https://doi.org/10.1093/ageing/afw129>
- Li Y, Hou L, Zhao H, Xie R, Yi Y, Ding X. Risk factors for falls among community-dwelling older adults: a systematic review and meta-analysis. *Front Med (Lausanne)*. 2022;9:1019094. <https://doi.org/10.3389/fmed.2022.1019094>
- Bergen G, Stevens MR, Burns ER. Falls and fall injuries among adults aged ≥ 65 years—United States, 2014. *MMWR Morb Mortal Wkly Rep*. 2016;65(37):993–998. <https://doi.org/10.15585/mmwr.mm6537a2>
- Chang VC, Do MT. Risk factors for falls among seniors: implications of gender. *Am J Epidemiol*. 2015;181(7):521–531. <https://doi.org/10.1093/aje/kwu268>
- Berry SD, Miller RR. Falls: epidemiology, pathophysiology, and relationship to fracture. *Curr Osteoporos Rep*. 2008;6(4):149–154. <https://doi.org/10.1007/s11914-008-0026-4>
- Johnson S, Kelly S, Rasali D. Differences in fall injury hospitalization and related survival rates among older adults across age, sex, and areas of residence in Canada. *Inj Epidemiol*. 2015;2(1):24. <https://doi.org/10.1186/s40621-015-0056-1>
- Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A. Incidence and economic burden of osteoporosis-related fractures in the United States, 2005–2025. *J Bone Miner Res*. 2007;22(3):465–475. <https://doi.org/10.1359/jbmr.061113>
- Ensrud KE, Blackwell TL, Cawthon PM, et al. Degree of trauma differs for major osteoporotic fracture events in older men versus older women. *J Bone Miner Res*. 2016;31(1):204–207. <https://doi.org/10.1002/jbmr.2589>
- Prior JC, Langsetmo L, Lentle BC, et al. Ten-year incident osteoporosis-related fractures in the population-based Canadian multicentre osteoporosis study—comparing site and age-specific risks in women and men. *Bone*. 2015;71:237–243. <https://doi.org/10.1016/j.bone.2014.10.026>
- Orwoll E, Blank JB, Barrett-Connor E, et al. Design and baseline characteristics of the osteoporotic fractures in men (MrOS) study—a large observational study of the determinants of fracture in older men. *Contemp Clin Trials*. 2005;26(5):569–585. <https://doi.org/10.1016/j.cct.2005.05.006>
- Blank JB, Cawthon PM, Carrion-Petersen ML, et al. Overview of recruitment for the osteoporotic fractures in men study (MrOS). *Contemp Clin Trials*. 2005;26(5):557–568. <https://doi.org/10.1016/j.cct.2005.05.005>
- Lewis CE, Ewing SK, Taylor BC, et al. Predictors of non-spine fracture in elderly men: the MrOS study. *J Bone Miner Res*. 2007;22(2):211–219. <https://doi.org/10.1359/jbmr.061017>
- Genant HK, Wu CY, van Kuijk C, Nevitt MC. Vertebral fracture assessment using a semiquantitative technique. *J Bone Miner Res*. 1993;8(9):1137–1148. <https://doi.org/10.1002/jbmr.5650080915>
- Ensrud KE, Harrison SL, Cauley JA, et al. Impact of competing risk of mortality on association of weight loss with risk of central body fractures in older men: a prospective cohort study. *J Bone Miner Res*. 2017;32(3):624–632. <https://doi.org/10.1002/jbmr.3020>
- Costa AG, Wyman A, Siris ES, et al. When, where and how osteoporosis-associated fractures occur: an analysis from the global longitudinal study of osteoporosis in women (GLOW). *PLoS One*. 2013;8(12):e83306. <https://doi.org/10.1371/journal.pone.0083306>
- Cauley JA, Cawthon PM, Peters KE, et al. Risk factors for hip fracture in older men: the osteoporotic fractures in men study (MrOS). *J Bone Miner Res*. 2016;31(10):1810–1819. <https://doi.org/10.1002/jbmr.2836>
- Freitas SS, Barrett-Connor E, Ensrud KE, et al. Rate and circumstances of clinical vertebral fractures in older men. *Osteoporos Int*. 2008;19(5):615–623. <https://doi.org/10.1007/s00198-007-0510-1>