




CORRECTION

Open Access



Correction to: Dose–response relationship of pulmonary disorders by inhalation exposure to cross-linked water-soluble acrylic acid polymers in F344 rats

Tomoki Takeda^{1*†} , Shotaro Yamano^{1*†} , Yuko Goto¹, Shigeyuki Hirai¹, Yusuke Furukawa¹, Yoshinori Kikuchi¹, Kyohei Misumi¹, Masaaki Suzuki¹, Kenji Takanobu¹, Hideki Senoh¹, Misae Saito¹, Hitomi Kondo¹, George Daghlian², Young-Kwon Hong², Yasuhiro Yoshimatsu³, Masanori Hirashima³, Yoichiro Kobashi⁴, Kenzo Okamoto⁵, Takumi Kishimoto⁶ and Yumi Umeda¹ 

Correction to: *Particle and Fibre Toxicology* (2022) 19:27
<https://doi.org/10.1186/s12989-022-00468-9>

Following publication of the original article [1], the authors reported that the Graphical Abstract was missing.

The Graphical Abstract has been provided in this Correction.

The original article [1] has been updated.

The original article can be found online at <https://doi.org/10.1186/s12989-022-00468-9>.

*Correspondence: tomoki-takeda@jbrcc.johas.go.jp; shotaro-yamano@pulmpath.com


†Tomoki Takeda and Shotaro Yamano have contributed equally to this work

¹ Japan Bioassay Research Center, Japan Organization of Occupational Health and Safety, Hadano, Kanagawa 257-0015, Japan
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



Graphic abstract



Characteristics of CWAAP-induced rat toxicity

	Male	Female
Organ-targeting	Lung (alveoli predominant) and MLN	
NOAEC	0.3 mg/m ³	0.3 mg/m ³
LOAEC	1 mg/m ³	1 mg/m ³
Reversibility	1 mg/m ³	1 mg/m ³
Toxicologically irreversible	3 and 10 mg/m ³	3 and 10 mg/m ³
Gender differences	Females recovering less than males	

Representative images of lung after 13w exposure



Author details

¹Japan Bioassay Research Center, Japan Organization of Occupational Health and Safety, Hadano, Kanagawa 257-0015, Japan. ²Department of Surgery, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA. ³Division of Pharmacology, Niigata University Graduate School of Medical and Dental Sciences, Niigata 951-8510, Japan. ⁴Department of Pathology, Tenri Hospital, Tenri, Nara 632-8552, Japan. ⁵Department of Pathology, Hokkaido Chuo Rosai Hospital, Japan Organization of Occupational Health and Safety, Iwamizawa, Hokkaido 068-0004, Japan. ⁶Director of Research and Training Center for Asbestos-Related Diseases, Okayama, Okayama 702-8055, Japan.

Published online: 13 May 2022

Reference

1. Takeda T, Yamano S, Goto Y, et al. Dose–response relationship of pulmonary disorders by inhalation exposure to cross-linked water-soluble acrylic acid polymers in F344 rats. *Part Fibre Toxicol.* 2022;19:27. <https://doi.org/10.1186/s12989-022-00468-9>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.