



## Correction to: Cancer survivors who fully participate in the PROFILES registry have better health-related quality of life than those who drop out

Imogen Ramsey<sup>1,2</sup> · Belle H. de Rooij<sup>2,3</sup> · Floortje Mols<sup>2,3</sup> · Nadia Corsini<sup>1</sup> · Nicole J. E. Horevoorts<sup>2,3</sup> · Marion Eckert<sup>1</sup> · Lonneke V. van de Poll-Franse<sup>2,3,4</sup>

Published online: 21 November 2019  
© The Author(s) 2019

**Correction to: Journal of Cancer Survivorship**  
<https://doi.org/10.1007/s11764-019-00793-7>

The original version of the article contains a mistake in figure 2 labeling.

The part labels of figure 2i and 2j is incorrectly labelled as 2a and 2b. The correct figure 2 is given below.

The original article has been corrected.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s11764-019-00793-7>


---

✉ Imogen Ramsey  
[imogen.ramsey@unisa.edu.au](mailto:imogen.ramsey@unisa.edu.au)

<sup>1</sup> Rosemary Bryant AO Research Centre, School of Nursing and Midwifery and UniSA Cancer Research Institute, University of South Australia, Adelaide, Australia

<sup>2</sup> The Netherlands Comprehensive Cancer Organization, Utrecht, the Netherlands

<sup>3</sup> Department of Medical and Clinical Psychology, CoRPS—Center of Research on Psychology in Somatic Diseases, Tilburg University, Tilburg, the Netherlands



**Fig. 2** Unadjusted mean functioning scores on the EORTCQLQ-C30 (a–h) and anxiety and depressive symptoms on the HADS (i, j) according to time of dropout (range: 0–100 and 0–21, respectively). Note: EORTC QLQ-C30 scales range from 0 to 100; higher scores reflect better perceived HRQOL. HADS scales range from 0 to 21; higher scores re-

fect higher prevalence of anxiety and depressive symptoms. *p* values indicate significant group differences between slopes and baseline scores compared with full responders in multilevel mixed models adjusted for time, age, sex, socioeconomic status, education, marital status, comorbidity, disease stage, and treatment received