

POSTER PRESENTATION

Open Access

Repeated methylene blue administration produces analgesia in experimental pain

A Dondas¹, A Luca¹, T Alexa^{1*}, VA Grigoras², OC Mungiu¹, CR Bohotin¹

From The European Headache and Migraine Trust International Congress
London, UK. 20-23 September 2012

Introduction

Methylene blue (MB), a widely used inhibitor of NO activity/production, is also a reduction-oxidation agent that can act both as a powerful antioxidant and as an enhancer of the electron transport chain. Furthermore, it prevents formation of mitochondrial oxygen free radicals and promotes oxygen consumption (Atamna et al. 2010,,Rojas et al. 2012).

Purpose

The aim of the study was to investigate the effects of chronic (14 days) MB administration on experimental pain in mice.

Methods

Sixteen Swiss male mice were divided into 2 groups: control group (n=8) and MB group (n=8); both groups received daily injections, for 14 days, either with saline 20 μ l i.p. (control group) or with MB 5 mg/kg b.w. i.p (MB group). Nociceptive tests (tail flick and hot plate) as well as mechanical and thermal withdrawal thresholds were measured every two days before MB/ saline administration. Before and two hours after the last dose was administered (day 14), each group was evaluated for the nociceptive tests and heat/mechanical hyperalgesia; results were compared with paired Student's t test. After nociceptive tests, the mice received 20 μ l of 5% formalin into the upper right lip and the intensity of the orofacial pain was assessed. The results were compared with those from saline group using unpaired Student's t test.

Results

Chronic administration of MB increased tail flick and hot plate latencies (p=0.03, p=0.02). We also noted an increase in the reaction time for thermal hyperalgesia assessed by Hargreaves method (p=0.04). As for the formalin-induced orofacial pain, MB produced a significant analgesic effect on both phases (p=0.03).

Conclusion

Our study demonstrates that chronic administration of MB has analgesic effects on acute nociception as well as on the orofacial inflammatory pain; further studies must be conducted in order to elucidate the mechanism by which the methylene blue exerts its antinociceptive effect.

Acknowledgements

Research supported by Executive Agency for Higher Education and Research Funding (UEFISCSU) Romania project PN-II-ID-PCE-2011-3-0875.

Author details

¹Centre for the Study and Therapy of Pain, University of Medicine and Pharmacy, Grigore. T. Popa, Romania. ²Department of Neurology, Rehabilitation Hospital, University of Medicine and Pharmacy, Grigore. T. Popa, Romania.

Published: 21 February 2013

References

1. Atamna H, Kumar R: Protective role of methylene blue in Alzheimer's disease via mitochondria and cytochrome c oxidase. *J Alzheimers Dis* 2010, **20**:S439-52.
2. Rojas JC, Bruchey AK, Gonzalez-Lima F: Neurometabolic mechanisms for memory enhancement and neuroprotection of methylene blue. *Prog Neurobiol* 2012, **96**:32-45.

doi:10.1186/1129-2377-14-S1-P86

Cite this article as: Dondas et al.: Repeated methylene blue administration produces analgesia in experimental pain. *The Journal of Headache and Pain* 2013 **14**(Suppl 1):P86.

¹Centre for the Study and Therapy of Pain, University of Medicine and Pharmacy, Grigore. T. Popa, Romania
Full list of author information is available at the end of the article