



Differential gene expression analysis of human cumulus cells

Sirin Bakti Demiray¹, Ege Nazan Tavmergen Goker², Erol Tavmergen², Ozlem Yilmaz³, Nilufer Calimlioglu², Huseyin Okan Soykam⁴, Gulperi Oktem³, Ugur Sezerman⁵

¹Assisted Reproduction Unit, Tepecik Education and Research Hospital, Izmir; Departments of ²Obstetrics and Gynecology and ³Histology and Embryology, Ege University Faculty of Medicine, Izmir; ⁴Epigenetiks Genetics Bioinformatics Software Inc., Istanbul; ⁵Department of Biostatistics and Bioinformatics, Acibadem Mehmet Ali Aydinlar University, Institute of Health Sciences, Istanbul, Turkey

Objective: This study was performed to explore the possibility that each oocyte and its surrounding cumulus cells might have different genetic expression patterns that could affect human reproduction.

Methods: Differential gene expression analysis was performed for 10 clusters of cumulus cells obtained from 10 cumulus-oocyte complexes from 10 patients. Same procedures related to oocyte maturation, microinjection, and microarray analyses were performed for each group of cumulus cells. Two differential gene expression analyses were performed: one for the outcome of clinical pregnancy and one for the outcome of live birth.

Results: Significant genes resulting from these analyses were selected and the top 20 affected pathways in each group were analyzed. Circadian entrainment is determined to be the most affected pathway for clinical pregnancy, and proteoglycans in cancer pathway is the most affected pathway for live birth. Circadian entrainment is also amongst the 12 pathways that are found to be in top 20 affected pathways for both outcomes, and has both lowest *p*-value and highest number of times found count.

Conclusion: Although further confirmatory studies are necessary, findings of this study suggest that these pathways, especially circadian entrainment in cumulus cells, may be essential for embryo development and pregnancy.

Keywords: Assisted reproductive techniques; Circadian clocks; Cumulus cells; Granulosa cells; Infertility; Live birth; Microarray analysis; Ovarian follicle; Pregnancy; Proteoglycans

Introduction

The prevalence of infertility is increasing, and new treatment options have been developed and collectively termed assisted reproductive techniques (ARTs) [1,2]. One of the most frequently used ART

method is intracytoplasmic sperm injection (ICSI). A very important step in human ARTs is noninvasive oocyte selection, followed by transfer of appropriate embryos because conducting this step properly reduces the risk of multiple gestations and increases the probability of pregnancy and live birth [3]. However, the morphological parameters used for embryo selection are not always efficient because embryos that have good morphology may not always have normal chromosomes [4].

Cumulus cells (CCs) are the follicle cells surrounding the oocyte in cumulus-oocyte complexes [5], which play an active role in the development process of oocytes [1]. Therefore, studies of cumulus-oocyte complexes may yield valuable insights into pregnancy and live birth mechanisms. Moreover, as CCs are considered to be waste tissue, use of these cells is not restricted by ethical considerations. For this reason, they are an ideal material for noninvasive methods to de-

Received: Nov 16, 2018 · Revised: Jan 9, 2019 · Accepted: Feb 2, 2019

Corresponding author: **Sirin Bakti Demiray**

Assisted Reproduction Unit, Tepecik Education and Research Hospital, Izmir 35180, Turkey

Tel: +90-2324443560-2013 Fax: +90-5062455604

E-mail: sirinbaktidemiray@gmail.com

*This work was supported by the University Scientific Research Project Fund (Ege-BAPTIP-00014, 2011).

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

termine markers of oocyte and embryo quality in ICSI and to predict potential success in terms of clinical pregnancy and live birth in humans. Ample evidence exists that successful follicle maturation in the mammalian ovary involves efficient paracrine communication between oocytes and granulosa cells (GCs), and good paracrine communication requires appropriate genetic control. CCs have the potential to predict oocyte competence. Thus, the expression profiles of CCs may be used as a parameter to facilitate optimal embryo selection [4], and may therefore be related to pregnancy and live birth outcomes. Recent reports have provided a plethora of information about GC morphology and/or their genetic expression profiles related to ARTs. Gene expression microarray analyses are used for mRNA profiling [6], and this technique has been applied to predict differences in genetic expression between CCs grouped according to various parameters of oocyte and embryo competence. However, a disadvantage of analyzing grouped CCs is that pooled genetic information is analyzed using microarray sequencing, rather than CCs belonging to a specific oocyte.

Enrichment of the transcriptome analysis results of CCs could provide information about critical cellular mechanisms involved in clinical pregnancy and live birth, potentially facilitating the development of novel ARTs and therapies. This study was performed to explore the possibility that specific oocytes and their surrounding CCs might have different genetic expression patterns that could affect human reproduction. To this end, this study sought to identify differentially expressed genes in cases of clinical pregnancy and live birth in comparison to non-pregnancy and stillbirth, respectively. These results were further enriched by pathway analysis using the PANOGA tool, providing valuable insights into the cellular mechanisms involved in clinical pregnancy and live birth, which may be possible to exploit in order to predict the outcomes of clinical pregnancy and live birth after embryo transfer.

Methods

This prospective case series included 10 women (age, 23–35 years) who were referred to Ege University Medical Faculty Hospital for idiopathic infertility and received ART therapy. The total sample size was 10 oocyte-cumulus complexes obtained over a period of 14 months between October 2011 and December 2012. CC gene expression in 10 clusters of CCs obtained from 10 cumulus-oocyte complexes from 10 patients was individually evaluated by microarrays using a NimbleGen human gene expression $12 \times 135K$ microarray kit (Roche, Penzberg, Germany). Although very few CCs were associated with each oocyte, each oocyte from which a transferred embryo developed was analyzed individually, along with its surrounding CCs, through microarray analysis. The results were reported using clinical

pregnancy and live birth as outcome variables. This study included only patients with unexplained infertility, and excluded cases of infertility based on male factors and other female factors in the standard patient population. All participating patients provided oral and written consent. Approval for this study was granted by the Ethics Committee for Human Research of Ege University (11-2, 9; 28.03.2011). All procedures were applied in accordance with the ethical standards described in the 1964 Declaration of Helsinki and its later amendments.

1. Ovarian stimulation protocol and oocyte pick-up

Ovarian stimulation therapy was applied to women prior to ICSI. Recombinant follicle-stimulating hormone (FSH, 75–225 IU; Gonal F, Serono International, Geneva, Switzerland) was administered according to the patient's ovarian response. Then, 0.1 mg of a gonadotropin-releasing hormone (GnRH) analogue (Decapeptyl; Ferring, Kiel, Germany) was started on the 21st day of the previous menstrual cycle. Transvaginal ultrasonography was performed at regular intervals during induction, and blood estradiol and progesterone levels were measured in order to assess follicular development. Human chorionic gonadotropin (hCG, 5,000–10,000 IU; Ovitrelle, Merck-Serono, Rome, Italy) was administered when the follicles achieved a sufficient diameter. After 34 to 36 hours, follicles were aspirated transvaginally into sterile tubes under anesthesia [7].

2. Evaluation of oocyte quality and collection of CCs

Cumulus-oocyte complexes were incubated in an incubator at 37°C in an atmosphere containing 5% CO₂. CCs were dissected from cumulus-oocyte complexes using a sterile pipette and immediately transferred into a sterile tube individually. The CCs surrounding the same oocyte were taken as individual groups, and were not pooled together. The CCs were then stored at –80°C for further analyses for about 3 months [8]. The remaining CCs adjacent to the oocytes were removed from the oocytes with enzymatic and mechanical methods using a sterile pipette and hyaluronidase (Hyase, Vitrolife, Frolunda, Sweden). Oocytes were evaluated morphologically and according to their nuclear maturation.

3. ICSI and fertilization

Approximately 30 to 60 minutes after denudation, mature oocytes were subjected to microinjection procedures. After microinjection, all oocytes were incubated in a fertilization medium using an incubator at 37°C with an atmosphere of 5% CO₂ (individually, not pooled). After 18 hours, the morphology of the pronuclei was evaluated to determine whether fertilization of the oocytes had occurred. Normal fertilization was defined as the presence of two pronuclei, with the first and second polar bodies and cytoplasmic localization of the polar bodies.

4. Embryo transfer and monitoring of pregnancy

About 20 hours after fertilization of an egg, the first mitotic division began and two-cell embryos were formed. Starting then, the embryos were evaluated according to the shape, size, and number of blastomeres, rate of fragmentation, rate of multinucleation, and degree of compaction [9,10]. On day 4 after fertilization, developing embryos were evaluated for compaction, morulation, and blastocoel formation. The inner cell mass and trophectoderm cell mass were evaluated for the compaction and cavitation processes of blastomeres. On day 4 after fertilization, laser-assisted hatching was applied to the embryos, and embryo transfer was performed. Beta-hCG levels in the blood of the patients were measured to determine pregnancy 12 to 14 days after embryo transfer. The patients were monitored to determine whether there was a gestational sac and to measure the fetal heart rate with ultrasound.

5. Microarray protocol

Gene expression in CCs was individually evaluated by microarrays in 10 clusters of CCs obtained from 10 cumulus-oocyte complexes from 10 different patients using a NimbleGen human gene expression 12 × 135K microarray kit according to the manufacturer’s instructions. Twelve-sample hybridization was performed on a single microarray slide, with three probes per gene. Total RNA was isolated using an RNeasy kit (Qiagen, Hilden, Germany) according to the manufacturer’s instructions. cDNA synthesis from total RNA was carried out using a Superscript Double Chain cDNA synthesis kit (Invitrogen Life Technologies, Carlsbad, CA, USA). cDNA was marked using a single color DNA labeling kit (NimbleGen). cDNA was hybridized using the NimbleGen Hybridization System. Washing and drying of the array were carried out with a NimbleGen MS 2000 microarray scanner, and raw probe expression data were collected using DEVA Software (Roche Nimblegen, Meylan, France).

6. Microarray data analysis

Microarray data resulting from the sequencing of 10 CCs, each surrounding an oocyte from one of 10 different patients, were analyzed to identify differentially expressed genes associated with the dichotomous outcomes of clinical pregnancy (pregnant vs. nonpregnant) and live birth (livebirth vs. stillbirth). Two differential gene expression analyses were performed: one for the outcome of clinical pregnancy and one for the outcome of live birth. Data were preprocessed using the oligo package [11] in R (www.R-project.org, 2016) and statistical analyses for differential gene expression analyses were performed using the limma package [12] in R. Several probes can be used to detect the transcriptome level of a single gene. In such cases, we used the probe with the most significantly differentiated value of significance for that gene. Significant genes resulting from these analyses (with $p < 0.05$) were selected and input into the PANOGA tool [13] for pathway analysis.

In order to determine the functionally enriched pathways of genes, PANOGA tool first aims to determine active subnetworks associated with those genes and their interactions via a protein-to-protein network. The BioGrid 3.4.141 protein-protein interaction (PPI) network [14] is loaded into Cytoscape [15] and the list of affected genes and their p -values are mapped onto the PPI network. To determine the active subnetworks of interacting proteins that are possibly associated with a disease, greedy search implementation of JActive Modules [16] is used. For functional enrichment of subnetworks, PANOGA tool computes the proportion of genes in a subnetwork that are also found in a human Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway, compared to the overall proportion of genes described for that pathway. For this purpose, ClueGO [17] plugin of Cytoscape is utilized. ClueGO works by facilitating a biological interpretation through visualization of functionally grouped terms in form of networks and charts. In order to determine statistical significance of an enrichment of the identified subnetwork, two-sided (enrichment/depletion) test based on hypergeometric distribution is applied, and the resulting p -values are corrected using the Bonferroni method for multiple testing.

Since PANOGA involves optimization procedures and stochastic processes, it was run 20 times for each case, and resulting KEGG [18] pathways were summarized across all runs for the outcomes of clinical pregnancy and live birth. In our analysis, we reported KEGG pathway ID, KEGG pathway name, significance of each pathway, pathway-relevant genes that are found in the original PANOGA input, and number of times each pathway found. The more frequently a pathway is found in an analysis, the less likely it is to be a random occurrence.

Results

Clinical pregnancy rates and live birth rates were used to assess the

Table 1. The laboratory and clinical results of 10 patients with unexplained infertility who underwent assisted reproductive technologies and whose cumulus cells underwent microarray analysis

Patient no.	Diagnosis	Age (yr)	Clinical pregnancy (+/-)	Live birth (+/-)
1	Idiopathic infertility	33	-	-
2	Idiopathic infertility	29	-	-
3	Idiopathic infertility	29	-	-
4	Idiopathic infertility	31	+	+
5	Idiopathic infertility	35	+	+
6	Idiopathic infertility	25	+	-
7	Idiopathic infertility	27	-	-
8	Idiopathic infertility	34	-	-
9	Idiopathic infertility	27	-	-
10	Idiopathic infertility	28	-	-

success of ICSI. Clinical pregnancy was observed in three of the 10 patients, and live birth occurred in two patients (Table 1). A total of 4,371 significant genes for clinical pregnancy and 1,975 significant

genes for live birth were enriched to 152 and 151 KEGG pathways, respectively, through 20 PANOGA runs. The top 20 affected pathways for clinical pregnancy and live births are summarized in Tables 2 and

Table 2. The top 20 statistically significant pathways for clinical pregnancy according to p-values

KEGG ID	KEGG term	p-value	Times found
04713	Circadian entrainment	6.04E ⁻³⁰	2,142
04610	Complement and coagulation cascades	7.22E ⁻²⁷	326
03050	Proteasome	8.73E ⁻²⁶	159
04015	Rap1 signaling pathway	1.14E ⁻²⁵	1,087
05032	Morphine addiction	1.48E ⁻²⁵	580
05200	Pathways in cancer	3.22E ⁻²³	182
04724	Glutamatergic synapse	1.06E ⁻²²	1,073
04080	Neuroactive ligand-receptor interaction	1.17E ⁻²¹	33
04020	Calcium signaling pathway	4.89E ⁻²¹	656
04014	Ras signaling pathway	9.22E ⁻²¹	413
04510	Focal adhesion	9.83E ⁻²¹	233
04725	Cholinergic synapse	2.27E ⁻²⁰	1,121
04728	Dopaminergic synapse	3.76E ⁻²⁰	1,568
04012	ErbB signaling pathway	4.36E ⁻²⁰	1,750
04062	Chemokine signaling pathway	7.46E ⁻²⁰	241
04727	GABAergic synapse	7.60E ⁻²⁰	547
04723	Retrograde endocannabinoid signaling	9.58E ⁻²⁰	653
04210	Apoptosis	3.84E ⁻¹⁹	1,690
04720	Long-term potentiation	7.52E ⁻¹⁹	3,695
04144	Endocytosis	1.51E ⁻¹⁸	58

KEGG, Kyoto Encyclopedia of Genes and Genomes.

Table 3. The top 20 statistically significant pathways for live birth according to p-values

KEGG ID	KEGG term	p-value	Times found
05205	Proteoglycans in cancer	1.29E ⁻²⁴	532
05200	Pathways in cancer	5.51E ⁻²³	168
04210	Apoptosis	1.97E ⁻²⁰	1119
04014	Ras signaling pathway	1.13E ⁻¹⁹	213
04010	MAPK signaling pathway	1.22E ⁻¹⁹	283
05032	Morphine addiction	2.78E ⁻¹⁹	366
04510	Focal adhesion	2.32E ⁻¹⁸	162
04062	Chemokine signaling pathway	5.91E ⁻¹⁸	322
04722	Neurotrophin signaling pathway	1.81E ⁻¹⁷	1,087
04723	Retrograde endocannabinoid signaling	5.18E ⁻¹⁷	260
04064	NF-kappa B signaling pathway	8.13E ⁻¹⁷	190
04012	ErbB signaling pathway	1.17E ⁻¹⁶	879
04610	Complement and coagulation cascades	1.93E ⁻¹⁶	105
04015	Rap1 signaling pathway	2.99E ⁻¹⁶	307
04350	TGF-beta signaling pathway	3.12E ⁻¹⁶	321
04668	TNF signaling pathway	4.70E ⁻¹⁶	588
05202	Transcriptional misregulation in cancer	8.38E ⁻¹⁶	341
04713	Circadian entrainment	1.21E ⁻¹⁵	286
04725	Cholinergic synapse	1.85E ⁻¹⁵	344
05223	Non-small cell lung cancer	2.06E ⁻¹⁵	932

KEGG, Kyoto Encyclopedia of Genes and Genomes; MARK, mitogen-activated protein kinases; NF, nuclear factor; TGF, transforming growth factor; TNF, tumor necrosis factor.

3, respectively. The two outcomes had 12 pathways in common, with different levels of significance, showing the involvement of certain cellular mechanisms in both clinical pregnancy and live birth. The 12 pathways that were significantly affected for both outcomes were: circadian entrainment, complement and coagulation cascades, Rap1 signaling pathway, morphine addiction, pathways in cancer, Ras signaling pathway, focal adhesion, cholinergic synapse, ErbB signaling pathway, chemokine signaling pathway, retrograde endocannabinoid signaling, and apoptosis. Circadian entrainment had a lower *p*-value (6.04×10^{-30}) than other pathways. The pathways that were different between live birth and clinical pregnancy were primarily several signaling pathways involved in neurological signal trafficking. Unique clinically-relevant pathways include proteasome and endocytosis.

We also conducted analyses of genes associated with both outcomes, and the results for the top 20 affected genes are summarized in Tables 4, 5 for clinical pregnancy and live birth, respectively. None of the top 50 significant genes for either outcome is in common. The most distinct significantly differentially expressed genes from both analyses that were also directly involved in the most affected pathways are reported below.

In clinical pregnancy, *PRKACB* ranks 16th among the most significantly altered genes, and is found in 13 of the top 20 significantly affected pathways. *COL4A* ranks 7th among the most significantly altered genes, and takes part in two of the top 20 significantly affected

pathways. *FGG* and *C4BPA* rank 21st and 24th, respectively, among the most significantly altered genes, and are also found in the complement and coagulation pathway. *SLC8A2* ranks 27th among the most significantly altered genes, and plays a role in the calcium signaling pathway.

In the live birth transcriptome analysis, *BDNF* ranks 10th among the most significantly altered genes, and is found in two of the top 20 significantly affected pathways. *NR4A3* ranks 2nd among the most significantly altered genes, and is found in the transcriptional misregulation in cancer pathway. *ENDR* ranks 22nd among the most significantly altered genes, and plays a role in pathways in cancer. *MAGI1* ranks 26th among the most significantly altered genes, and is also found in the RAP1 signaling pathway.

Since our method maps the significant genes to a PPI network and finds active subnetworks involving many of the affected genes, it also takes into account the indirect effects of genes through cascades of interactions with certain cellular processes. Therefore, some of the genes might not be directly involved in the affected pathway, but instead may affect some of the genes in that pathway through interactions of proteins.

With this in mind, we investigated distances between genes within the PPI network for the top 20 significant genes resulting from differential expression analysis and for genes associated with the most significant pathways of both outcomes; circadian entrainment for clinical pregnancy and proteoglycans in cancer for live birth. Distance

Table 4. The top 20 statistically significant genes for clinical pregnancy according to *p*-values

Gene	logFC	<i>p</i> -value
<i>PPHLN1</i>	-1.747187984	2.10E ⁻⁰⁶
<i>ZFP57</i>	-2.993479513	1.73E ⁻⁰⁵
<i>CD7</i>	1.634646302	3.73E ⁻⁰⁵
<i>ARHGEF4</i>	-2.972436989	4.99E ⁻⁰⁵
<i>SOX8</i>	-2.706781964	5.05E ⁻⁰⁵
<i>TBX4</i>	1.296222769	5.19E ⁻⁰⁵
<i>COL4A6</i>	1.564709693	5.35E ⁻⁰⁵
<i>DKFZp686L1814</i>	-1.307955019	5.80E ⁻⁰⁵
<i>MAWBP</i>	1.57256687	5.95E ⁻⁰⁵
<i>MUC17</i>	1.477177013	6.87E ⁻⁰⁵
<i>C15orf33</i>	-1.600279744	6.92E ⁻⁰⁵
<i>WASPIP</i>	1.440695515	8.23E ⁻⁰⁵
<i>FCRLM2</i>	1.184684422	8.41E ⁻⁰⁵
<i>SPANXE</i>	1.349434304	8.50E ⁻⁰⁵
<i>FLJ39502</i>	3.287634413	8.57E ⁻⁰⁵
<i>PRKACB</i>	-1.650870764	8.68E ⁻⁰⁵
<i>ELA2</i>	1.178323805	9.02E ⁻⁰⁵
<i>CNIH3</i>	-1.976904531	9.03E ⁻⁰⁵
<i>SSPN</i>	-1.749017451	9.14E ⁻⁰⁵
<i>DDAH1</i>	-1.723440567	9.28E ⁻⁰⁵

Table 5. The top 20 statistically significant genes for live birth according to *p*-values

Gene	logFC	<i>p</i> -value
<i>MYOC</i>	3.737516546	7.04E ⁻⁰⁷
<i>NR4A3</i>	4.324984553	3.15E ⁻⁰⁶
<i>FLJ20701</i>	-3.669639519	3.27E ⁻⁰⁶
<i>BRF1</i>	1.90998993	5.11E ⁻⁰⁶
<i>STXBP5L</i>	2.772004737	7.36E ⁻⁰⁶
<i>CR1L</i>	3.329420892	9.84E ⁻⁰⁶
<i>BTBD11</i>	-3.294941796	1.91E ⁻⁰⁵
<i>IGSF4</i>	-2.403987308	3.59E ⁻⁰⁵
<i>LDHAL6B</i>	3.312196874	3.62E ⁻⁰⁵
<i>BDNF</i>	-2.058433847	3.71E ⁻⁰⁵
<i>BTG2</i>	2.49767883	4.33E ⁻⁰⁵
<i>EYA2</i>	2.034444882	4.99E ⁻⁰⁵
<i>DKFZP686A01247</i>	1.748650995	5.13E ⁻⁰⁵
<i>RPS6KA1</i>	-1.64992345	6.60E ⁻⁰⁵
<i>EPHA1</i>	1.863062351	6.77E ⁻⁰⁵
<i>KIAA1345</i>	-1.718641218	7.74E ⁻⁰⁵
<i>ACACB</i>	2.346361813	9.70E ⁻⁰⁵
<i>LOC440508</i>	2.954390096	0.000103611
<i>DDX12</i>	2.218784482	0.000107346
<i>KIAA1841</i>	1.78790507	0.000115263

reported herein is the number of proteins in the PPI network in between two interested gene.

In circadian entrainment pathway, cumulative average distance over all genes is found to be 3.27. *CALML3* is found to be the gene with lowest average distance of 2.6 and *MTNR1B* is found to be the gene with highest average distance of 4.

In proteoglycans in cancer pathway, cumulative average distance over all genes is found to be 2,91. *TP53* is to be the gene with lowest average distance of 2.07 and *IL12B* is found to be the gene with highest average distance of 4.46. In pathway analysis, we focused only on the top 20 pathways ranked by statistical significance. The relevance of these pathways to clinical pregnancy and live birth was established via a thorough literature search, which is summarized in the Discussion section.

Discussion

1. Clinical pregnancy

Our analysis determined that circadian entrainment is the most sig-

nificantly affected pathway for clinical pregnancy. The affected genes and their interactions in circadian entrainment pathway are shown in Figure 1. Circadian entrainment is stabilized by the maintenance of regular light/dark and sleep/wake cycle and plays a critical role in steroid biosynthesis, including luteinizing hormone (LH) and GnRH secretions in pregnancy. It acts as a regulator of many bioprocesses, with the circadian clock regulating the timing of ovulation and progesterone secretion in the GCs and luteal cells of ovarian follicles. Disruption of the female circadian clock cycle has a negative effect on women becoming pregnant [19]. Melatonin also has an impact on reproduction, especially during pregnancy, by protecting against molecular damage and cellular dysfunction [20].

The complement and coagulation cascades are other pathways that showed a significant difference between CCs depending on whether clinical pregnancy occurred. In a previous study that compared gene expression in CCs and GCs surrounding oocytes, it has been reported that there were differentially expressed genes related to the complement and coagulation cascades between these groups of cells [21]. Proteasomal proteolysis has multiple functions in the




Figure 1. Significantly altered genes in the circadian entrainment pathway and their interactions for clinical pregnancy, denoted by red pathway nodes. Data from Kanehisa Laboratories. It is permitted to be used for academic purposes. Full definitions and descriptions for the pathway nodes can be found at https://www.genome.jp/kegg-bin/show_pathway?map=hsa04713&show_description=show.

process of oocyte meiosis resumption and in other steps of nuclear maturation [22,23]. The results of current study also indicate that proteasome pathway might impact clinical pregnancy outcomes. Previous studies have shown that RAP1 signaling pathway, RAS signaling pathway, and ErbB signaling pathway have an impact on reproduction [24,25]. The RAS-mediated ERK1/2 pathway in pituitary cells, which is affected by elevated GnRH, is crucial for stimulating the LH surge and ovulation [26]. In current study, it is also demonstrated that RAP1 signaling pathway, RAS signaling pathway, and ErbB signaling pathway may play active roles in clinical pregnancy. Furthermore, genes related to morphine addiction, cancer, calcium signaling, and endocytosis pathways demonstrated significant differences in expression in CCs depending on whether clinical pregnancy was achieved. The presence of opioid receptor subtypes has been demonstrated in preimplantation embryos and in oviduct, and these have active roles in embryo development [27]. The hippo tumor suppressor pathway has functions in follicle cells related to oocyte polarity and development of the body axes in *Drosophila* [28]. Gonadotropin paracrine factors induce Ca²⁺ mobilization in CCs, which affects gamete development and function, and therefore fertilization and reproduction [29]. Furthermore, rat GCs may utilize FSH for biological processes through endocytosis [30].

Genes involved in neurological signaling pathways, such as glutamatergic synapse, cholinergic synapse, dopaminergic synapse, gamma-aminobutyric acid (GABA)-ergic synapse, long-term potentiation, and neuroactive ligand-receptor interaction pathways show significant differences in CCs depending on whether clinical pregnancy occurred. These pathways and related genes may play functional role in pregnancy, as they affect gonadotropin hormone secretion by regulating hormonal signaling pathways. Genes responsive to neurotransmitters, such as dopamine, glycine, and GABA may play important roles in folliculogenesis and oocyte maturation [31]. GABA is important for regulation of reproduction via GnRH neurons [32]. Genes that encode receptors for GABA and glutamate have been determined to be related with oocyte maturation [33]. In a recent study, significant differences have been reported in dopaminergic synapse, mitogen-activated protein kinases (MAPK) signaling, steroid hormone biosynthesis, steroid biosynthesis, GnRH signaling, oocyte meiosis, progesterone-mediated oocyte maturation, calcium signaling, and G-protein coupled receptor signaling pathways in goose ovarian tissue across different reproductive periods [34]. With the induction or inactivation of the LH choriogonadotropin receptor, the neuroactive ligand-receptor interaction pathway led to changes in the menstrual cycle, LH levels, ovarian size, and response to LH and hCG [33].

2. Live birth

In this microarray analysis of human CCs, significant differences are observed in genes related to several hormonal, metabolic, apoptotic, and neuronal pathways. These pathways may be associated with live birth, as shown by their differential expression according to whether live birth occurred.

In this study, genes related to proteoglycans in the cancer pathway, apoptosis pathways, tumor necrosis factor (TNF) and MAPK pathways, retrograde endocannabinoid signaling, and the transcription factor nuclear factor- κ B shows a significant association with success of live birth. CCs and GCs synthesize proteoglycan by gonadotropin stimulation to produce an elastic matrix with mucus, and the synthesis of proteoglycan changes during follicular development and atresia [35]. Proapoptotic and antiapoptotic pathways in GCs are very important in ovarian follicles and affect reproductive outcomes of various mammals, including humans [36,37]. During mammalian ovulation, TNF maintains apoptosis at the ovarian surface epithelium to promote stigma development and ovulation [38]. In a recent microarray analysis of gene expression in human GCs and CCs, it has been shown that TNF has been highly expressed in GCs [21]. Jun N-terminal kinase, p38 MAPK, and ERK signaling pathways coordinate proapoptotic and antiapoptotic factors in GCs to regulate follicular atresia [39]. The maturation of the mouse cumulus-oocyte complex depends on MAPK pathway in CCs [40]. Cannabinoid receptors are expressed in the human ovarian medulla and cortex [41]. The endocannabinoid system mediates pregnancy, implantation, sperm motility, sperm-oocyte interactions, the capacitation process, and embryo development [42]. The focal adhesion pathway and chemokine signaling pathway displays significant differences according to live birth outcomes of the patients. There is a gap junction-dependent interaction between GCs and oocytes [43]. The chemokine signaling pathway plays an active role in the regulation of the cumulus extracellular matrix to promote successful fertilization by stimulating integrin attachment to extracellular matrix proteins [44]. According to our results, neurotrophic factor pathways and transforming growth factor-beta (TGF- β) pathways seem to be correlated with patients' live birth outcome. A family of neurotrophic factors is expressed in ovarian follicles as antiapoptotic factors that promote the developmental competence of oocytes [45]. The TGF- β superfamily is a diverse group of proteins with critical roles in various physiological processes, and the members of this superfamily include several subfamilies, such as the bone morphogenetic protein (BMP) subfamily and anti-Müllerian hormone (AMH; also termed as Müllerian-inhibiting substance). These are expressed in the ovary and regulate folliculogenesis [46-48]. The results of our recent study also demonstrate that there is a significant difference in the expression of BMP2 in the CCs of good-quality oocytes, and subsequently good embryos [49]. CCs associat-




Figure 2. Significantly altered genes in the circadian entrainment pathway and their interactions for live birth, denoted by red pathway nodes. Data from Kanehisa Laboratories. It is permitted to be used for academic purposes. Full definitions and descriptions for the pathway nodes can be found at https://www.genome.jp/kegg-bin/show_pathway?map=hsa04713&show_description=show.

ed either with developmentally incompetent or competent antral oocytes have been compared using microarrays with the whole transcriptome, and increased AMH gene expression has been shown to be correlated with oocyte competence [47]. Using gene expression profiles with microarray techniques, it is thought that CCs could have the potential to be used as a biomarker to predict oocyte quality and the pregnancy outcome of an embryo [1,50].

Although it was not as significant as in the clinical pregnancy analysis, circadian entrainment is also found to be significantly altered in the live birth analysis. Circadian entrainment ranks 18th among the most significantly altered pathways in the live birth analysis, whereas it ranks first in the clinical pregnancy analysis. This striking difference between the role of circadian entrainment in clinical pregnancy and in live birth may have resulted from the fact that different genes and interactions are significantly altered in the circadian entrainment pathway in the live birth analysis in comparison to the clinical pregnancy analysis. These altered genes and interactions can be seen in Figure 2.

Circadian entrainment has lower *p*-value (6.04×10^{-30}) than the

proteoglycans in cancer pathway, which has a *p*-value of 1.29×10^{-24} . This discrepancy may have occurred because 30 genes are associated with circadian entrainment pathway, whereas 17 genes are found in the proteoglycans in cancer pathway. Furthermore, the genes in the circadian entrainment pathway are generally connected to differentially expressed genes with higher significance than is the case for the genes in proteoglycans in cancer pathway. This shows that affected genes in the circadian entrainment pathway is part of many active subnetworks supporting the significance of that pathway. Overall, both pathways have relatively low cumulative average distances, supporting the fact that on average, in PPI, genes associated with significantly affected pathways are found in close proximity to the significantly altered genes found in the differential expression analysis.

In this study, significant differences are observed in the genes related to several pathways depending on the outcomes of clinical pregnancy and/or live birth. These results shed light on cellular factors that may play crucial role in clinical pregnancy and live birth. However, these results must be supported by further studies, and results

may differ between patients and cycles. Nonetheless, the results of this study suggest that these pathways, especially the circadian entrainment pathway in CCs, may play essential roles in clinical pregnancy and live birth.

In conclusion, the results of this study provide new insights into human CCs and new information on the crosstalk between different signaling pathways during pregnancy. Due to their ethical advantages, current methods and those of similar noninvasive studies using CCs are of great importance. These results may further enhance the reliability of oocyte and/or embryo selection criteria and could promote the maintenance of excellent clinical outcomes in *in vitro* fertilization (IVF)/ICSI programs. These findings will be of value for further studies conducted to improve the success of IVF/ICSI programs.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

Acknowledgments

We would like to thank Duygu Unuvar Purcu (Biologist) for technical help in some experiments.

ORCID

Sirin Bakti Demiray <https://orcid.org/0000-0002-6261-1267>
 Huseyin Okan Soykam <https://orcid.org/0000-0002-8955-2412>
 Ugur Sezerman <https://orcid.org/0000-0003-0905-6783>

Author contributions

Conceptualization: SBD, GO. Data curation: SBD, ENTG, ET, NC. Formal analysis: HOS, US. Funding acquisition: SBD, GO. Methodology: SBD, OY. Project administration: SBD, GO, OY. Visualization: SBD, HOS. Writing – original draft: SBD, US, HOS. Writing - review & editing: SBD, US, HOS.

References

1. Assidi M, Montag M, Van der Ven K, Sirard MA. Biomarkers of human oocyte developmental competence expressed in cumulus cells before ICSI: a preliminary study. *J Assist Reprod Genet* 2011;28:173-88.
2. Balasch J. Investigation of the infertile couple: investigation of the infertile couple in the era of assisted reproductive technology: a time for reappraisal. *Hum Reprod* 2000;15:2251-7.
3. Cillo F, Brevini TA, Antonini S, Paffoni A, Ragni G, Gandolfi F. Asso-

- ciation between human oocyte developmental competence and expression levels of some cumulus genes. *Reproduction* 2007;134:645-50.
4. Bettegowda A, Patel OV, Lee KB, Park KE, Salem M, Yao J, et al. Identification of novel bovine cumulus cell molecular markers predictive of oocyte competence: functional and diagnostic implications. *Biol Reprod* 2008;79:301-9.
5. Matzuk MM, Burns KH, Viveiros MM, Eppig JJ. Intercellular communication in the mammalian ovary: oocytes carry the conversation. *Science* 2002;296:2178-80.
6. True L, Feng Z. Immunohistochemical validation of expression microarray results. *J Mol Diagn* 2005;7:149-51.
7. Fenwick J, Platteau P, Murdoch AP, Herbert M. Time from insemination to first cleavage predicts developmental competence of human preimplantation embryos *in vitro*. *Hum Reprod* 2002;17:407-12.
8. Feuerstein P, Cadoret V, Dalbies-Tran R, Guerif F, Bidault R, Royere D. Gene expression in human cumulus cells: one approach to oocyte competence. *Hum Reprod* 2007;22:3069-77.
9. della Ragione T, Verheyen G, Papanikolaou EG, Van Landuyt L, Devroey P, Van Steirteghem A. Developmental stage on day-5 and fragmentation rate on day-3 can influence the implantation potential of top-quality blastocysts in IVF cycles with single embryo transfer. *Reprod Biol Endocrinol* 2007;5:2.
10. van Montfoort AP, Geraedts JP, Dumoulin JC, Stassen AP, Evers JL, Ayoubi TA. Differential gene expression in cumulus cells as a prognostic indicator of embryo viability: a microarray analysis. *Mol Hum Reprod* 2008;14:157-68.
11. Carvalho BS, Irizarry RA. A framework for oligonucleotide microarray preprocessing. *Bioinformatics* 2010;26:2363-7.
12. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015;43:e47.
13. Bakir-Gungor B, Egemen E, Sezerman OU. PANOGA: a web server for identification of SNP-targeted pathways from genome-wide association study data. *Bioinformatics* 2014;30:1287-9.
14. Stark C, Breitkreutz BJ, Reguly T, Boucher L, Breitkreutz A, Tyers M. BioGRID: a general repository for interaction datasets. *Nucleic Acids Res* 2006;34:D535-9.
15. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 2003;13:2498-504.
16. Ideker T, Ozier O, Schwikowski B, Siegel AF. Discovering regulatory and signalling circuits in molecular interaction networks. *Bioinformatics* 2002;18 Suppl 1:S233-40.
17. Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M,

- Kirilovsky A, et al. ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* 2009;25:1091-3.
18. Kanehisa M, Goto S. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 2000;28:27-30.
 19. Sellix MT, Menaker M. Circadian clocks in mammalian reproductive physiology: effects of the "other" biological clock on fertility. *Discov Med* 2011;11:273-81.
 20. Reiter RJ, Tan DX, Korkmaz A, Rosales-Corral SA. Melatonin and stable circadian rhythms optimize maternal, placental and fetal physiology. *Hum Reprod Update* 2014;20:293-307.
 21. Burnik Papler T, Vrtacnik Bokal E, Maver A, Kopitar AN, Lovrecic L. Transcriptomic analysis and meta-analysis of human granulosa and cumulus cells. *PLoS One* 2015;10:e0136473.
 22. Huo LJ, Fan HY, Zhong ZS, Chen DY, Schatten H, Sun QY. Ubiquitin-proteasome pathway modulates mouse oocyte meiotic maturation and fertilization via regulation of MAPK cascade and cyclin B1 degradation. *Mech Dev* 2004;121:1275-87.
 23. Yi YJ, Nagyova E, Manandhar G, Prochazka R, Sutovsky M, Park CS, et al. Proteolytic activity of the 26S proteasome is required for the meiotic resumption, germinal vesicle breakdown, and cumulus expansion of porcine cumulus-oocyte complexes matured in vitro. *Biol Reprod* 2008;78:115-26.
 24. Wayne CM, Fan HY, Cheng X, Richards JS. Follicle-stimulating hormone induces multiple signaling cascades: evidence that activation of Rous sarcoma oncogene, RAS, and the epidermal growth factor receptor are critical for granulosa cell differentiation. *Mol Endocrinol* 2007;21:1940-57.
 25. Gloaguen P, Crepieux P, Heitzler D, Poupon A, Reiter E. Mapping the follicle-stimulating hormone-induced signaling networks. *Front Endocrinol (Lausanne)* 2011;2:45.
 26. Fan HY, Liu Z, Mullany LK, Richards JS. Consequences of RAS and MAPK activation in the ovary: the good, the bad and the ugly. *Mol Cell Endocrinol* 2012;356:74-9.
 27. Chen Y, Kong S, Tang X, Fu Y, Wang B, Zhang S, et al. Preimplantation mouse embryo is a target for opioid ligand-receptor signaling. *Biol Reprod* 2014;91:4.
 28. Yu J, Poulton J, Huang YC, Deng WM. The hippo pathway promotes Notch signaling in regulation of cell differentiation, proliferation, and oocyte polarity. *PLoS One* 2008;3:e1761.
 29. Chen H, Kui C, Chan HC. Ca(2+) mobilization in cumulus cells: role in oocyte maturation and acrosome reaction. *Cell Calcium* 2013;53:68-75.
 30. Sanford JC, Batten BE. Endocytosis of follicle-stimulating hormone by ovarian granulosa cells: analysis of hormone processing and receptor dynamics. *J Cell Physiol* 1989;138:154-64.
 31. Devjak R, Fon Tacer K, Juvan P, Virant Klun I, Rozman D, Vrtacnik Bokal E. Cumulus cells gene expression profiling in terms of oocyte maturity in controlled ovarian hyperstimulation using GnRH agonist or GnRH antagonist. *PLoS One* 2012;7:e47106.
 32. Watanabe M, Fukuda A, Nabekura J. The role of GABA in the regulation of GnRH neurons. *Front Neurosci* 2014;8:387.
 33. Huang X, Hao C, Shen X, Liu X, Shan Y, Zhang Y, et al. Differences in the transcriptional profiles of human cumulus cells isolated from MI and MII oocytes of patients with polycystic ovary syndrome. *Reproduction* 2013;145:597-608.
 34. Luan X, Liu D, Cao Z, Luo L, Liu M, Gao M, et al. Transcriptome profiling identifies differentially expressed genes in Huoyan goose ovaries between the laying period and ceased period. *PLoS One* 2014;9:e113211.
 35. Salustri A, Camaioni A, Di Giacomo M, Fulop C, Hascall VC. Hyaluronan and proteoglycans in ovarian follicles. *Hum Reprod Update* 1999;5:293-301.
 36. Tiwari M, Prasad S, Tripathi A, Pandey AN, Ali I, Singh AK, et al. Apoptosis in mammalian oocytes: a review. *Apoptosis* 2015;20:1019-25.
 37. Hennebold JD. Preventing granulosa cell apoptosis through the action of a single microRNA. *Biol Reprod* 2010;83:165-7.
 38. Murdoch WJ, Lund SA. Prostaglandin-independent anovulatory mechanism of indomethacin action: inhibition of tumor necrosis factor alpha-induced sheep ovarian cell apoptosis. *Biol Reprod* 1999;61:1655-9.
 39. Peter AT, Dhanasekaran N. Apoptosis of granulosa cells: a review on the role of MAPK-signalling modules. *Reprod Domest Anim* 2003;38:209-13.
 40. Su YQ, Denegre JM, Wigglesworth K, Pendola FL, O'Brien MJ, Eppig JJ. Oocyte-dependent activation of mitogen-activated protein kinase (ERK1/2) in cumulus cells is required for the maturation of the mouse oocyte-cumulus cell complex. *Dev Biol* 2003;263:126-38.
 41. El-Talatini MR, Taylor AH, Elson JC, Brown L, Davidson AC, Konje JC. Localisation and function of the endocannabinoid system in the human ovary. *PLoS One* 2009;4:e4579.
 42. Battista N, Meccariello R, Cobellis G, Fasano S, Di Tommaso M, Pirazzi V, et al. The role of endocannabinoids in gonadal function and fertility along the evolutionary axis. *Mol Cell Endocrinol* 2012;355:1-14.
 43. McGinnis LK, Kinsey WH. Role of focal adhesion kinase in oocyte-follicle communication. *Mol Reprod Dev* 2015;82:90-102.
 44. Yodoi R, Tamba S, Morimoto K, Segi-Nishida E, Nishihara M, Ichikawa A, et al. RhoA/Rho kinase signaling in the cumulus mediates extracellular matrix assembly. *Endocrinology* 2009;150:3345-52.
 45. Ikeda S, Yamada M. Midkine and cytoplasmic maturation of

- mammalian oocytes in the context of ovarian follicle physiology. *Br J Pharmacol* 2014;171:827-36.
46. Erickson GF, Shimasaki S. The spatiotemporal expression pattern of the bone morphogenetic protein family in rat ovary cell types during the estrous cycle. *Reprod Biol Endocrinol* 2003;1:9.
 47. Vigone G, Merico V, Prigione A, Mulas F, Sacchi L, Gabetta M, et al. Transcriptome based identification of mouse cumulus cell markers that predict the developmental competence of their enclosed antral oocytes. *BMC Genomics* 2013;14:380.
 48. Knight PG, Glister C. TGF-beta superfamily members and ovarian follicle development. *Reproduction* 2006;132:191-206.
 49. Demiray SB, Yilmaz O, Goker EN, Tavmergen E, Calimlioglu N, Sezerman U, et al. Expression of the bone morphogenetic protein-2 (BMP2) in the human cumulus cells as a biomarker of oocytes and embryo quality. *J Hum Reprod Sci* 2017;10:194-200.
 50. Hamel M, Dufort I, Robert C, Leveille MC, Leader A, Sirard MA. Identification of follicular marker genes as pregnancy predictors for human IVF: new evidence for the involvement of luteinization process. *Mol Hum Reprod* 2010;16:548-56.