

Detection of Estrogen-Independent Growth-Stimulating Activity in Breast Cancer Tissues: Implication for Tumor Aggressiveness

Yuri Yamaguchi · Yuko Seino · Hiroyuki Takei ·
Masafumi Kurosumi · Shin-ichi Hayashi

Received: 17 July 2013 / Accepted: 24 October 2013 / Published online: 8 November 2013
© The Author(s) 2013. This article is published with open access at Springerlink.com

Abstract Estrogen and various growth factors affecting tumor behavior are present in the breast cancer microenvironment, but their comprehensive effects and signal crosstalks are different in each case. However, there is no system to evaluate the factors, detected in individual breast cancer cases, that regulate ER activity and tumor progression. In this study, we analyzed the effects of individual breast cancer extracts by our original system using an estrogen-signal reporter cell line, MCF-7-E10, which we previously established. MCF-7-E10 cell line is stably transfected by an estrogen response element (ERE)-green fluorescent protein (GFP) gene; it expresses GFP when estrogen receptors (ERs) are activated by estrogen or growth factor signal-mediated ER phosphorylation. Using this cell line, we analyzed the comprehensive effects of factors derived from breast cancer tissues on ER activity and growth of MCF-7-E10 cells for each case. We also analyzed

relationships between these activities and clinicopathologic characteristics of patients who provided cancer specimens. The breast cancer extracts, which reflect the combined activities of growth factors present in individual cases, stimulated MCF-7-E10 cell growth in an estrogen-independent manner, and specifically stimulated growth of other breast cancer cell lines, regardless of ER expression. High growth-promoting activities were seen in tumor regions of specimens with tumors > 10 mm in size, HER2 intrinsic subtype, and scirrhous and solid-tubular carcinoma histological subtypes. Anti-human hepatocyte growth factor (HGF) antibody and an inhibitor for insulin-like growth factor-1 (IGF-1) receptor inhibited MCF-7-E10 cell growth by the breast cancer extracts, indicating that signal pathways via HGF or IGF-1 receptor significantly affect breast cancer. These data suggest that growth factors other than estrogen in the tumor extract significantly affect breast cancer aggressiveness in an estrogen-independent manner, and could be useful therapeutic targets.

Y. Yamaguchi (✉) · Y. Seino
Research Institute for Clinical Oncology, Saitama Cancer Center,
818 Komuro, Ina-machi, Saitama-ken 362-0806, Japan
e-mail: yamaguchi@cancer-c.pref.saitama.jp

Y. Seino
e-mail: y-seino@cancer-c.pref.saitama.jp

H. Takei
Division of Breast Surgery, Saitama Cancer Center Hospital,
Saitama, Japan
e-mail: h-takei@cancer-c.pref.saitama.jp

M. Kurosumi
Department of Pathology,
Saitama Cancer Center Hospital, Saitama, Japan
e-mail: mkurosumi@cancer-c.pref.saitama.jp

Y. Seino · S.-i. Hayashi
Department of Molecular and Functional Dynamics, Graduate
School of Medicine, Tohoku University, Sendai, Japan

S.-i. Hayashi
e-mail: shin@med.tohoku.ac.jp

Keywords Tumor microenvironment ·
Estrogen-independent · HGF · IGF1-R · Breast cancer

Background

The tumor microenvironment is enriched in factors such as growth factors, cytokines and chemokines, and critically affects initiation and progression of various tumor types [1–5]. For postmenopausal women with low levels of plasma estrogen, breast cancer growth and progression are mainly caused by estrogen produced locally in the tumor microenvironment [6–8]. Intratumoral production of estrogen is induced by aromatase, a key enzyme in estrogen biosynthesis, which is expressed by carcinoma-associated stromal fibroblasts [7–10]. Aromatase is a target of endocrine therapy for breast

cancers; aromatase inhibitors attenuate estrogen biosynthesis in treating hormone-responsive breast cancer [11, 12]. Estrogen stimulates breast cancer growth via expression of a diverse set of growth-related genes in tumor cells, and through activation of estrogen receptor (ER), a transcription factor [13, 14]. ER α is a primary predictive marker for hormonal therapy in breast cancer, but approximately one-third of ER+ patients do not respond to this therapy, suggesting that ER α is not a perfect predictor for hormonal therapy. To shed light on these issues, and to study the molecular basis for breast cancer, we first focused on analysis of estrogen signals by development of a custom-made cDNA microarray, and provided novel diagnostic and prognostic estrogen-induced genes [15–17].

In addition to the genomic pathway, estrogen induces non-genomic pathways by interacting with signal cascades for growth factors [4, 13], such as epidermal growth factor (EGF) and insulin-like growth factor-1 (IGF-1), which activate ER α in an estrogen-independent manner by phosphorylating several ER α sites using their downstream signal kinases, including MAPK and PI3K [4, 5, 18–20]. Growth factors are produced by malignant cells themselves, adjacent tumor stromal fibroblasts and inflammatory cells in the microenvironment.

To analyze the carcinoma-associated fibroblasts-induced ER activation in individual breast cancers, we established an estrogen response element (ERE)-green fluorescent protein (GFP) assay system. It allows us to detect estrogen- and phosphorylation-dependent ER-activating ability of stromal fibroblasts adjacent to tumor cells under coculture with MCF-7-E10 cells, a clone of MCF-7 stably transfected with the *ERE-GFP* gene [20]. Using this system, we examined relationships between ER-activating ability of stromal fibroblasts and clinicopathological characteristics. We found that, although ER-activating abilities of stromal fibroblasts vary among breast cancers, they are higher in breast cancers from postmenopausal patients than in those from premenopausal patients [20]. This is in accordance with the fact that intratumoral estrogen production causes progression of postmenopausal breast cancers [6–8]. ER-activating abilities of fibroblasts in grade 3 breast cancers are lower than in grade 1 breast cancers, suggesting that the grade 3 microenvironment stimulates proliferation of breast cancer cells via an estrogen-independent pathway [20].

In the breast cancer microenvironment, various growth factors and cytokines reportedly interact to control tumor growth, as described above. However, their significance in tumor growth in vivo and in response to hormonal therapy are unclear. To investigate the role of breast cancer-derived factors on breast cancer growth, we studied effects of the supernatants of minced breast cancer tissues on MCF-7-E10 cell growth. Tissue supernatant, unlike tissue extract or conditioned

medium of tissue prepared after culture for several days in vitro, reflects the comprehensive effects of factors detected in the tumor in vivo. In addition to estrogen-related signals, we found that the breast cancer-derived factors effectively stimulate MCF-7-E10 cell growth via an estrogen-independent pathway.

Methods

Cells & Cell Culture

Cell lines used in this study were cultured in RPMI1640 medium (GIBCO) supplemented with 10 % FCS (Tissue Culture Biologicals) at 37 °C in a humidified atmosphere of 5 % CO₂. We previously established the estrogen-signal reporter cell line MCF-7-E10 derived from MCF-7 cells by stable transfection with an *ERE-GFP* reporter plasmid [20]. To analyze the effect of breast cancer tissue supernatant (BCTS) on ER activity in MCF-7-E10 cells, cells were precultured in estrogen-deprived medium (phenol red-free RPMI1640 medium supplemented with 10 % dextran-coated, charcoal-treated FCS) for 3 days.

Preparation of BCTS

Breast cancer specimens were processed within 1 h after surgical resection. After being weighed, specimens were transferred to tubes containing phenol red- and serum-free RPMI 1640 medium at 100 mg/ml, and minced to particles < ~1 mm³ in size. The suspension was centrifuged (600 × *g*, 10 min, 4 °C) and the supernatant was further centrifuged (12,000 × *g*, 10 min, 4 °C) to obtain BCTS. The protein concentration of each sample was determined using BCA Protein Assay Reagent (PIERCE).

Human breast cancer tissues were obtained by surgery at the Saitama Cancer Center Hospital (Saitama, Japan) after informed consent had been obtained from the patients. The Saitama Cancer Center Ethics Committee approved this study. In the clinicopathological classifications of the patients (Table 1), ER and progesterone receptor (PgR) status was determined using monoclonal anti-ER α antibody 1D5 (Dako, Glostrup, Denmark) and monoclonal anti-PgR antibody PgR636 (Dako), and evaluated on the basis of Allred scoring [21]. HER2 protein expression was scored as 0, 1+, 2+ or 3+ using the HerceptTest™ (Dako); *HER2* genome status was evaluated by fluorescent in situ hybridization (FISH) using PathVysion HER-2 DNA Probe Kit (Abbott Laboratories, Abbott Park, IL, USA). According to the ASCO/CAP guidelines [22], absolute *HER2* gene/chromosome 17 copy number ratios

Table 1 Clinical characteristics of patients

	No. of patients ^a
Age (y)	
< 51	26
≥ 50	65
Unknown	2
Menopausal status	
Pre	28
Post	64
Male	1
Tumor diameter (cm)	
≤ 1	15
> 1	75
Unknown	3
ER	
Positive	69
Negative	24
PgR	
Positive	52
Negative	41
HER2	
0	37
1	31
2	10
3	13
Unknown	2
Stage	
0	1
I	37
II	42
III	11
Unknown	2
Histology	
Scirrhous	41
Solidtubular	23
Papillotubular	13
Mucinous	7
Apocrine	2
DCIS	3
Unknown	4
Grade	
1	18
2	17
3	47
Unknown	11
Nodal status	
Negative	38
Positive	53
Unknown	2

^a Total number of patients was 93

greater than 2.2 and less than 1.8 indicated *HER2* amplification (positive) and *HER2* non-amplification (negative), respectively. Histologic grading was evaluated according to the Elston and Ellis grading scheme [23].

Cell Growth Assay

After 3 days of culture in estrogen-deprived medium, cells were seeded at $1 \times 10^3/150 \mu\text{l}$ in a 96-well multi-dish culture plate, or at $1 \times 10^4/1 \text{ ml}$ in a 24-well plate, with or without BCTS at indicated protein concentrations for 4 days. Viable cells were examined using a Cell Counting Kit-8 assay according to manufacturer's instructions (Dojindo Laboratories, Japan).

Evaluation of ER Activity

ER activities in MCF-7-E10 cells, which had been transfected with *ERE-GFP*, after incubation with BCTS or E₂, were monitored through GFP expression [20]. To quantify GFP expression, cells expressing GFP were counted under a fluorescence microscope after the cells were harvested by treatment with trypsin. Data are presented as percentage of cells expressing GFP.

Quantification of Growth Factors in BCTS by ELISA

Human EGF and IGF-1 levels in BCTS were quantified by ELISA using Quantikine (R&D Systems, MN, USA) specific for each growth factor.

Materials

Unless otherwise stated, all other materials were from Sigma-Aldrich Inc. (St. Louis, MO, USA). Inhibitors for EGF receptor and IGF receptor, and normal mouse IgG were from Calbiochem. Mouse anti-human HGF monoclonal antibody was from the Institute of Immunology (Tokyo, Japan). Mouse IgG1 antibody (Chemicon International, CA, USA) was used as an isotype control. IGF-1 receptor inhibitor, AG1024, and EGF receptor inhibitor, AG1478, were from Chemicon International.

Statistical Analysis

Statistical analyses were performed using the Stat Flex version 6.0 software program (Artech Co., Ltd., Osaka, Japan). In comparisons among groups, ANOVA and two-sample *t*-tests were used to assess the statistical significance of differences.

Data are expressed as means \pm S.D. $P < 0.05$ was considered statistically significant.

Results

BCTS Stimulates Breast Cancer Cell Growth in Both Estrogen-Dependent and -Independent Manners

In the tumor microenvironment, many growth factors, cytokines and chemokines directly and indirectly control growth. To study their comprehensive influence on breast cancer aggressiveness, we first analyzed effects of BCTS on MCF-7-E10 cell growth (Fig. 1), which allowed us to examine the total effect of breast cancer-derived factors secreted from tumor and stromal cells, as they exist in vivo, on growth and estrogen-related signals of breast cancer cells. BCTS dose-dependently stimulated MCF-7-E10 cell growth (Fig. 1a). Although activities varied among specimens, more than 60 % showed higher growth-stimulating activity than with estrogen (Fig. 1b).

To examine the specificity of target cells, we studied the effect of BCTS on growth of other tumor cell lines, including

a breast cancer cell line, T47D, a lung adenocarcinoma cell line, PC9, and a cervical cancer cell line, HeLa (Fig. 2a). The growth of T47D, another ER + human breast cancer cell line, was stimulated by BCTS while growth of PC9 was not increased. HeLa cell growth was rather inhibited by BCTS. The growth of MDA-MB-231 cells, an ER- human breast cancer cell line, was also stimulated by the extracts (data not shown). These results suggest that BCTS specifically stimulated breast cancer cell growth regardless of ER expression.

Next, to see whether growth-stimulating activity in the tissue supernatant affected only the tumoral region, we analyzed extracts of tumoral regions and non-tumoral regions 2 cm distal to the tumor. The tumoral regions had more growth-stimulating activity than the non-tumoral regions (Fig. 2b), suggesting that the tumoral regions have an abundance of growth-stimulating activities for breast cancer cells.

To see if ER activation was required for BCTS-induced growth stimulation, we analyzed GFP expression in MCF-7-E10 cells, and found growth stimulation was not necessarily accompanied by ER activation (Fig. 3a). We next examined effects of anti-estrogen agents such as tamoxifen and fulvestrant on BCTS-induced growth stimulation, and found that high growth-stimulating activities were resistant to fulvestrant (Fig. 3b) and tamoxifen (Fig. 3c). These results indicate that, in addition to an ER-dependent pathway, BCTS stimulates breast cancer growth via an ER-independent pathway.

Growth-Stimulating Activity Correlated with Clinicopathological Characteristics

We analyzed the relationships between ER-independent growth-stimulating activity detected in BCTS and clinicopathologic characteristics of the specimens' donors (Fig. 4). Although BCTS growth-stimulating activity did not correlate with expression of ER α or PgR, stage, menopausal status, grade or nodal status (data not shown), specimens from tumors larger than 10 mm showed higher growth-stimulating activity than those smaller than 10 mm (Fig. 4a). Breast cancers are categorized into four intrinsic subtypes according to gene-expression profile: luminal A (ER + and/or PgR+, HER2-), luminal B (ER + and/or PgR+, HER2+), HER2 (ER-, PgR-, HER2+) and basal-type (ER-, PgR-, HER2-) [24, 25]. BCST derived from HER2 subtype showed slightly or significantly higher growth-stimulating activity than that from luminal B or basal types, respectively (Fig. 4b), suggesting that the tumor extracts of HER2 subtype have an abundance of growth factors stimulating their own receptors, including those of the ERBB family.

We next analyzed relationships between HER2 expression and growth-stimulating activity in ER- breast cancers, and found that the cases with high growth-stimulating activity differed significantly from those with low activity in terms





Fig. 1 BCTS effectively stimulated growth of MCF-7-E10 cells. After 3 days of culture in estrogen-deprived medium, MCF-7-E10 cells were cultured with breast cancer tissue supernatant at the indicated protein concentrations (a) or at 25 µg (b) in total 150 µl medium per well in 96-well plate for 4 days. The viable cells were examined using a Cell Counting Kit-8 assay. Values relative to control are shown. Data are presented as mean \pm SD of triplicate determinations. *, $P < 0.05$; **, $P < 0.01$


Fig. 2 Specificity of cell growth-stimulating activity detected in BCTS. **a** BCTS at 25 μg protein concentration in 150 μl medium per well in 96-well plate effectively stimulated growth of breast cancer cell lines, MCF-7-E10 and T47D which were precultured in estrogen-deprived medium for 3 days. The viable cells were examined using a Cell Counting Kit-8 assay. **b** Specificity for BCTS-derived region. T, tissue supernatant derived from tumor region; NT, tissue supernatant derived from the region 2 cm distal to the tumor region. Data are presented as mean ± SD of triplicate experiments. *, *P*<0.05



of HER2 expression; more specimens with high HER2 scores (score 3) were seen among cases with high activity (Fig. 4c). This difference could not be observed for ER + breast cancers.

Breast cancers have histological types that reflect biological characteristics. Invasive ductal carcinoma can be classified into three subtypes—papillotubular, solid-tubular and

Fig. 3 BCTS stimulated growth of MCF7-E10 cells in an estrogen-independent manner. **a** MCF7-E10 cells were cultured with BCTS at 25 μg protein concentration in 150 μl medium per well in 96-well plate for 4 days. Cell growth was examined using a Cell Counting Kit-8 assay and ER activities are shown as the percentage of MCF-7-E10 cells expressing GFP. **b, c** MCF-7-E10 cells were cultured with BCTS at 25 μg protein concentration in 150 μl medium per well in 96-well plate or the indicated concentrations in the presence or absence of anti-estrogen agents, fulvestrant (1 μM) or 4-hydroxy tamoxifen (4OHT, 1 μM), for 4 days. 17β-Estradiol (E₂) was also tested at 1 nM or the indicated concentrations. Cell growth was examined using a Cell Counting Kit-8 assay. Data are presented as mean ± SD of triplicate experiments






Fig. 4 Correlations between growth-stimulating activity and clinicopathological characteristics, intrinsic subtypes and histological subtypes. MCF7-E10 cells were cultured with BCTS at 25 μ g protein concentration in 150 μ l medium per well in 96-well plate for 4 days. Cell growth was examined as described in **Materials and Methods** for triplicate experiments, and the growth-stimulating activities are shown as the ratios

calculated relative to the control. Data are presented as mean \pm SD of triplicate experiments. High growth-stimulating activity in specimens was associated with tumor size (**a**), intrinsic subtype (**b**), HER2 expression in ER-negative breast cancer (**c**) or histological classifications (**d**). Differences between groups were determined by two-sample *t*-test. $P < 0.05$ was considered statistically significant

scirrhous carcinoma—which are related to prognosis. We previously reported their relative overall survival rates as papillotubular carcinoma > solid-tubular carcinoma > scirrhous carcinoma [26]. The more aggressive scirrhous carcinoma and solid-tubular carcinoma show higher growth-stimulating activity than do papillotubular carcinoma and mucinous types (Fig. 4d), suggesting that growth-stimulating activity is related to aggressiveness in breast cancer.

Growth Factors in BCTS Promote MCF-7-E10 Cell Growth

Growth-stimulating activity was heat labile and detectable in the fraction with an MW greater than 5 kDa (data not shown), suggesting that it could be derived from proteinous factors. Among various factors in the tumor microenvironment, HGF derived from stromal fibroblasts has been reported to stimulate growth of mouse mammary tumor cells in primary culture [27]; EGF and IGF-1 are known to activate ER via phosphorylation [18, 19]. To analyze the participation of these growth factors in tumor growth-stimulating activities found in BCTS, we first examined the effect of anti-HGF antibody on them. As shown in Fig. 5a, anti-HGF antibody, but not control IgG, effectively inhibited extract-stimulated growth of MCF-7-E10 cells. MCF-7 cells reportedly express c-Met, a receptor for HGF.


We next analyzed the roles of EGF and IGF-1, using the inhibitors specific for their receptors. IGF-R inhibitor dose-dependently inhibited the growth of MCF-7-E10 cells while EGF-R inhibitor, in contrast, stimulated their growth (Fig. 5b, c).

Finally, we analyzed growth factors present in BCTS using the enzyme immunoassay. HGF was detected in more than 70 % of the tested samples, whereas EGF was detected only in 3 out of 25 samples (Fig. 5d). Although the analysis using IGF-1R inhibitor suggested involvement of IGF-1 in the growth-stimulating effect of BCTS as described above, IGF-1 could not be detected in the enzyme immunoassay. This might be because of the immunoassay's sensitivity, or because other ligands for IGF-1R (such as IGF-II, insulin or unknown factors) might have been present in the tumor extracts. These results suggest that signal pathways via HGF or IGF-1R play a significant role in promoting the growth of breast cancer cells.

Discussion

The tumor microenvironment is apparently associated with important aspects of epithelial solid tumor progression, including tumor growth, angiogenesis and metastasis. In the tumor microenvironment, growth factors such as EGF, IGF-1, transforming growth factor- α , transforming growth factor β

Fig. 5 Detection of growth factors involved in growth-promoting activity for MCF-7-E10 cells in BCTS. MCF-7-E10 cells were incubated with BCTS in the presence of anti-HGF antibody (**a**), AG1024, IGF-1R inhibitor (**b**), or AG1478, EGFR inhibitor (**c**), at the indicated concentrations. For anti-HGF antibody treatment, BCTS was pre-incubated with anti-HGF antibody for 30 min at room temperature and was then used for assay. Mouse IgG1 antibody was used as an isotype control. **d** The concentrations of HGF and EGF detected in BCTS were analyzed by immunoassay using Quantikine (R&D Systems, MN, USA). **, $P < 0.01$, ***, $P < 0.001$



and stromal-derived factor-I reportedly affect breast cancer growth, directly or indirectly [1, 3, 4]; however, the combined effects of these factors and their signal interactions in vivo are unclear. In this study, using the supernatant of breast cancer tissues, we analyzed the comprehensive effects of breast cancer-derived factors and found that BCTS effectively and specifically stimulated breast cancer cell growth. In addition to estrogen, which is locally produced in the microenvironment in breast cancers of postmenopausal patients [6, 8], our results suggest that the tumor extracts also stimulated breast cancer cell growth in an estrogen-independent manner, as anti-estrogen agents such as tamoxifen and fulvestrant did not inhibit the effect of BCTS. Furthermore, clinicopathological data and BCTS-associated growth-stimulation correlated with tumor size and HER2 expression, indicating the physiological significance of growth-stimulating activity in BCTS. Thus, BCTS offers an appropriate means to analyze the combined effect of the breast cancer-derived factors on tumor cell behavior.

Although many growth factors might be present in BCTS, we found HGF and IGF-1R-related signals to affect the growth-stimulating activity of BCTS, because it was suppressed by anti-HGF antibody and IGF-1R inhibitor. HGF was detected in tissue extracts of more than 70 % of breast cancer specimens whereas EGF was detected in only 12 % (Fig. 5d). The growth-stimulating activities did not always correlate with HGF concentrations (data not shown), but this is expected, as growth-stimulating activities in the supernatant

are derived from the signal cross-talks of several factors. HGF, which acts through its receptor MET, is a multifunctional cytokine that induces cell survival, growth, differentiation and motility in most solid human cancers including colorectal, renal and breast cancers [28]. In normal epithelial cells, HGF, in combination with other growth factors, promotes mammary ductal morphogenesis [29]. Overexpression of both HGF and MET have been frequently reported in breast cancers, and are associated with poor prognosis [30]. HGF reportedly stimulates breast cancer growth in a paracrine fashion, in that HGF is produced primarily by stromal fibroblasts and acts on epithelial cells through its receptor MET [27, 31]. Stromal fibroblasts from breast cancer tissue produce large amounts of HGF compared with normal fibroblasts [30]. A c-Met-targeted therapy, ARQ197—which selectively targets c-Met tyrosine kinase—is currently in a phase II clinical trial [32]; SGX523—a novel ATP-competitive inhibitor, that is exquisitely selective for inhibition of MET-mediated signaling—is also being developed [33].

We found that IGF-1R signaling mediated the growth-stimulating activity of BCTS, because IGF-1R-specific inhibitor decreased the growth-stimulating effect of BCTS. IGF-1R-related signals are widely shown to induce cell proliferation and survival in breast cancer [34–36]; IGF-R1 activation protects breast cancer cells from apoptosis induced by various anticancer drugs [37]. While BCTS stimulated growth of MCF-7-E10 cells in an estrogen-independent manner, functional interactions between estrogen and IGF-1R signaling

pathways, including Ras/MAPK and PI3K/Akt have been reported [38] Estrogen also up-regulates IGF-1R expression in breast cancer [36]. However, we could not detect IGF-1 and stromal cell-derived growth factor-1 α in BCTS (data not shown), possibly because of the limit of sensitivity by the immunoassay used in our study; or that other ligands may be present in the breast cancer microenvironment that activate IGF-1R—including IGF-II, insulin and unknown factors [39]. Indeed, overexpression of IGF-1R in MCF-7 cells has been shown to induce IGF-1R tyrosine kinase activation in the absence of exogenous IGF-1 [40].

These results suggest that signaling pathways via HGF/c-Met or IGF-1R significantly affect breast cancer cell growth. However, growth-stimulating activity found in BCTS might be derived from orchestrated signal crosstalks of several factors, because recombinant growth factors, including HGF and IGF-1, could not induce MCF-7-E10 cell growth when used alone. Further investigations of these activities and the identification of the cellular sources of the growth factors are needed to identify the mechanisms of the growth-stimulating effect of breast cancer tissue supernatant, which may help design more effective targeted therapies for breast cancer.

Conclusions

The breast cancer microenvironment provides estrogen and growth factors that affect tumor behavior, but the comprehensive effects of these factors, including signal crosstalk, on progression of breast cancer remain unclear. Using an estrogen-signal reporter cell line, MCF-7-E10, stably transfected with the ERE-GFP gene, we analyzed the effect of factors present in breast cancer tissues to reflect the *in vivo* status of individual cases. We found that they stimulated growth of MCF-7-E10 cells in an estrogen-independent manner, and that growth-promoting activity is related to aggressiveness in breast cancer. Moreover, signal pathways via HGF and IGF-1 receptor were involved in these activities. Our study strongly suggests that the evaluation of comprehensive tumor-promoting activity for individual breast cancers is important in determining appropriate therapy.

Acknowledgments This study was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan and the Advanced Research for Medical Products Mining Programme of the National Institute of Biomedical Innovation (NIBIO).

Competing interest The authors declare that they have no competing interest.

Authors' contributions YY and SH were involved in experimental design, performed all experiments, and drafted the manuscript. YS assisted in experiments and performed statistical analysis of the data. HT participated in acquisition and interpretation of the clinical data of

patients. MK participated in experimental design and histological evaluation. All authors contributed to the analysis of data and approved the final manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Allinen M, Beroukhim R, Cai L, Brennan C, Lahti-Domenici J, Huang H, Porter D, Hu M, Chin L, Richardson A et al (2004) Molecular characterization of the tumor microenvironment in breast cancer. *Cancer Cell* 6:17–32
- Mueller MM, Fusenig NE (2004) Friends or foes: bipolar effects of the tumour stroma in cancer. *Nat Rev Cancer* 4:839–849
- Orimo A, Gupta PB, Sgroi DC, Arenzana-Seisdedos F, Delaunay T, Naeem R, Carey VJ, Richardson AL, Weinberg RA (2005) Stromal fibroblasts present in invasive human breast carcinomas promote tumor growth and angiogenesis through elevated SDF-1/CXCL12 secretion. *Cell* 121:335–348
- Yamaguchi Y (2007) Microenvironmental regulation of estrogen signals in breast cancer. *Breast Cancer* 14:175–181
- Yamaguchi Y, Hayashi S (2009) Estrogen-related cancer microenvironment of breast carcinoma. *Endocr J* 56:1–7
- O'Neill JS, Miller WR (1987) Aromatase activity in breast adipose tissue from women with benign and malignant breast diseases. *Br J Cancer* 56:601–604
- Santen RJ, Santner SJ, Pauley RJ, Tait L, Kaseta J, Demers LM, Hamilton C, Yue W, Wang JP (1997) Estrogen production via the aromatase enzyme in breast carcinoma: which cell type is responsible? *J Steroid Biochem Mol Biol* 61:267–271
- Simpson ER, Davis SR (2001) Minireview: aromatase and the regulation of estrogen biosynthesis—some new perspectives. *Endocrinology* 142:4589–4594
- Zhao Y, Agarwal VR, Mendelson CR, Simpson ER (1997) Transcriptional regulation of CYP19 gene (aromatase) expression in adipose stromal cells in primary culture. *J Steroid Biochem Mol Biol* 61:203–210
- Smith IE, Dowsett M (2003) Aromatase inhibitors in breast cancer. *N Engl J Med* 348:2431–2442
- Dowsett M, Cuzick J, Ingle J, Coates A, Forbes J, Bliss J, Buyse M, Baum M, Buzdar A, Colleoni M et al (2010) Meta-analysis of breast cancer outcomes in adjuvant trials of aromatase inhibitors versus tamoxifen. *J Clin Oncol* 28:509–518
- Hayashi S, Yamaguchi Y (2005) Estrogen signaling and prediction of endocrine therapy. *Cancer Chemother Pharmacol* 56:27–31
- Hayashi S, Niwa T, Yamaguchi Y (2009) Estrogen signaling pathway and its imaging in human breast cancer. *Cancer Sci* 100:1773–1778
- Hayashi S (2004) Prediction of hormone sensitivity by DNA microarray. *Biomed Pharmacother* 58:1–9
- Inoue A, Yoshida N, Omoto Y, Oguchi S, Yamori T, Kiyama R, Hayashi S (2002) Development of cDNA microarray for expression profiling of estrogen-responsive genes. *J Mol Endocrinol* 29:175–192
- Inoue A, Omoto Y, Yamaguchi Y, Kiyama R, Hayashi S (2004) Transcription factor EGR3 is involved in the estrogen-signaling pathway in breast cancer cells. *J Mol Endocrinol* 32:649–666
- Yoshida N, Omoto Y, Inoue A, Eguchi H, Kobayashi Y, Kurosumi M, Saji S, Suemasu K, Okazaki T, Nakachi K et al (2004) Prediction of prognosis of estrogen receptor-

- positive breast cancer with combination of selected estrogen-regulated genes. *Cancer Sci* 95:496–502
18. Kato S, Endoh H, Masuhiro Y, Kitamoto T, Uchiyama S, Sasaki H, Masushige S, Gotoh Y, Nishida E, Kawashima H et al (1995) Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science* 270:1491–1494
 19. Osborne CK, Shou J, Massarweh S, Schiff R (2005) Crosstalk between estrogen receptor and growth factor receptor pathways as a cause for endocrine therapy resistance in breast cancer. *Clin Cancer Res* 11:865s–70s
 20. Yamaguchi Y, Takei H, Suemasu K, Kobayashi Y, Kurosumi M, Harada N, Hayashi S (2005) Tumor-stromal interaction through the estrogen-signaling pathway in human breast cancer. *Cancer Res* 65:4653–4662
 21. Allred DC, Harvey JM, Berardo M, Clark GM (1998) Prognostic and predictive factors in breast cancer by immunohistochemical analysis. *Mod Pathol* 11:155–168
 22. Wolff AC, Hammond MEH, Schwartz JN, Hagerty KL, Allred DC, Cote RJ, Dowsett M, Fitzgibbons PL, Hanna WM, Langer A et al (2007) American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol* 25:118–145
 23. Elston CW, Ellis IO (1991) Pathological prognostic factors in breast cancer. I. The value of histological grade in breast cancer: experience from a large study with long-term follow-up. *Histopathology* 19:403–410
 24. Sørlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB, van de Rijn M, Jeffrey SS et al (2001) Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A* 98:10869–10874
 25. Sørlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, Deng S, Johnsen H, Pesich R, Geisler S et al (2003) Repeated observation of breast tumor subtypes in independent gene expression data sets. *Proc Natl Acad Sci U S A* 100:8418–8423
 26. Kurosumi M, Tabei T, Inoue K, Takei H, Ninomiya J, Naganuma R, Suemasu K, Higashi Y, Tsuchiya E (2003) Prognostic significance of scoring system based on histological heterogeneity of invasive ductal carcinoma for node-negative breast cancer patients. *Oncol Rep* 10:833–837
 27. Sasaki M, Enami J (1999) Mammary fibroblast-derived hepatocyte growth factor and mammogenic hormones stimulate the growth of mouse mammary epithelial cells in primary culture. *Endocrine J* 44:359–366
 28. Christensen JG, Burrows J, Salgia R (2005) c-Met as a target for human cancer and characterization of inhibitors for therapeutic intervention. *Cancer Lett* 225:1–26
 29. Yang Y, Spitzer E, Meyer D, Sachs M, Niemann C, Hartmann G, Weidner KM, Birchmeier C, Birchmeier W (1995) Sequential requirement of hepatocyte growth factor and neuregulin in the morphogenesis and differentiation of the mammary gland. *J Cell Biol* 131:215–226
 30. Yamashita J, Ogawa M, Yamashita S, Nomura K, Kuramoto M, Saishoji T, Shin S (1994) Immunoreactive hepatocyte growth factor is a strong and independent predictor of recurrence and survival in human breast cancer. *Cancer Res* 54:1630–1633
 31. Tyan SW, Kuo WH, Huang CK, Pan CC, Shew JY, Chang KJ, Lee EY, Lee WH (2011) Breast cancer cells induce cancer-associated fibroblasts to secrete hepatocyte growth factor to enhance breast tumorigenesis. *PLoS ONE* 6:e15313
 32. Munshi N, Sébastien J, Li Y, Chang-Rung C, France DS, Ashwell MA, Hill J, Moussa MM, Leggett DS, Li CJ (2010) ARQ197, a novel and selective inhibitor of the human c-Met receptor tyrosine kinase and antitumor activity. *Mol Cancer Ther* 9:1544–1553
 33. Buchanan SG, Hendle J, Lee PS, Smith CR, Bounaud PY, Jessen KA, Tang CM, Huser NH, Felce JD, Froning KJ et al (2009) SGX523 is an exquisitely selective, ATP-competitive inhibitor of the MET receptor tyrosine kinase with antitumor activity in vivo. *Mol Cancer Ther* 8:3181–3190
 34. Resnik JL, Reichart DB, Huey K, Webster NJG, Seely BL (1998) Elevated insulin-like growth factor I receptor autophosphorylation and kinase activity in human breast cancer. *Cancer Res* 58:1159–1164
 35. Law JH, Habibi G, Hu K, Masoudi H, Wang MY, Stratford AL, Park E, Gee JM, Finlay P, Jones HE et al (2008) Phosphorylated insulin-like growth factor-I/insulin receptor is present in all breast cancer subtypes and is related to poor survival. *Cancer Res* 68:10238–10246
 36. Maor S, Mayer D, Yarden RI, Lee AV, Sarfstein R, Werner H, Papa MZ (2006) Estrogen receptor regulates insulin-like growth factor-I receptor gene expression in breast tumor cells: involvement of transcription factor Sp1. *J Endocrinol* 191:605–612
 37. Dunn SE, Hardman RA, Kari FW, Barrett JC (1997) Insulin-like growth factor 1 (IGF-1) alters drug sensitivity of HBL100 human breast cancer cells by inhibition of apoptosis induced by diverse anticancer drugs. *Cancer Res* 57:2687–2693
 38. Stoica GE, Franke TF, Moroni M, Moroni M, Mueller S, Morgan E, Iann MC, Winder AD, Reiter R, Wellstein A, Martin MB, Stoica A (2003) Effect of estradiol on estrogen receptor-alpha gene expression and activity can be modulated by the ErbB2/PI 3-K/Akt pathway. *Oncogene* 22:7998–8011
 39. Macaulay VM (1992) Insulin-like growth factors and cancer. *Br J Cancer* 65:311–320
 40. Surmacz E, Burgaud JL (1995) Overexpression of insulin receptor substrate 1 (IRS-1) in the human breast cancer cell line MCF-7 induces loss of estrogen requirements for growth and transformation. *Clin Cancer Res* 1:1429–1436