

Meet Our Editorial Board Member

Stefan Boehm

Department of Neurophysiology and Neuropharmacology
Center for Physiology and Pharmacology
Medical University of Vienna
Schwarzspanierstrasse 17/I, A-1090 Vienna
Austria

Dr. Boehm obtained MD from the University of Vienna, Austria, in 1987 and received his postdoctoral education in Vienna and Frankfurt/Main, Germany. He had been a professor of neuropharmacology at the medical university of Graz and was appointed as a professor of neuropharmacology at the medical university of Vienna in 2007. Currently, he is head of the department of neurophysiology and neuropharmacology at the center for physiology and pharmacology and the director of the doctoral school. His research focuses on mechanisms underlying pre- as well as postsynaptic modulation of synaptic transmission and mechanisms of action of analgesic as well as anticonvulsant drugs.



SELECTED PUBLICATIONS

- [1] Boehm, S., Betz, H. Somatostatin inhibits excitatory transmission at rat hippocampal synapses *via* presynaptic receptors. *J. Neurosci.*, **1997**, *17*(11), 4066-75.
- [2] O'Connor, V., El Far, O., Bofill-Cardona, E., Nanoff, C., Freissmuth, M., Karschin, A., Airas, J.M., Betz, H., Boehm, S. Calmodulin dependence of presynaptic metabotropic glutamate receptor signaling. *Science*, **1999**, *286*(5442), 1180-4.
- [3] Boehm, S. ATP stimulates sympathetic transmitter release *via* presynaptic P2X purinoceptors. *J. Neurosci.*, **1999**, *19*(2), 737-46.
- [4] Boehm, S., Kubista, H. Fine tuning of sympathetic transmitter release *via* ionotropic and metabotropic presynaptic receptors. *Pharmacol. Rev.*, **2002**, *54*(1), 43-99.
- [5] Kubista, H., Edelbauer, H., Boehm, S. Evidence for structural and functional diversity among SDS-resistant SNARE complexes in neuroendocrine cells. *J. Cell Sci.*, **2004**, *117*(Pt 6), 955-66.
- [6] Lechner, S.G., Hussl, S., Schicker, K.W., Drobny, H., Boehm, S. Presynaptic inhibition *via* a phospholipase C- and phosphatidylinositol bisphosphate-dependent regulation of neuronal Ca²⁺ channels. *Mol. Pharmacol.*, **2005**, *68*(5), 1387-96.
- [7] Schicker, K., Hussl, S., Chandaka, G.K., Kosenburger, K., Yang, J.W., Waldhoer, M., Sitte, H.H. Boehm, S. A membrane network of receptors and enzymes for adenine nucleotides and nucleosides. *Biochim. Biophys. Acta*, **2009**, *1793*(2), 325-34.
- [8] Schicker, K.W., Chandaka, G.K., Geier, P., Kubista, H., Boehm, S. P2Y1 receptors mediate an activation of neuronal calcium-dependent K⁺ channels. *J. Physiol.*, **2010**, *588*(Pt 19), 3713-25.
- [9] Yousuf, A., Klinger, F., Schicker, K., Boehm, S. Nucleotides control the excitability of sensory neurons *via* two P2Y receptors and a bifurcated signaling cascade. *Pain*, **2011**, *152*(8), 1899-908.
- [10] Buchmayer, F., Schicker, K., Steinkellner, T., Geier, P., Stübiger, G., Hamilton, P.J., Jurik, A., Stockner, T., Yang, J.W., Montgomery, T., Holy, M., Hofmaier, T., Kudlacek, O., Matthies, H.J., Ecker, G.F., Bochkov, V., Galli, A., Boehm, S., Sitte, H.H. Amphetamine actions at the serotonin transporter rely on the availability of phosphatidylinositol-4, 5-bisphosphate. *Proc. Natl. Acad. Sci. U. S. A.*, **2013**, *110*(28), 11642-7.
- [11] Klinger, F., Bajric, M., Salzer, I., Dorostkar, M.M., Khan, D., Pollak, D.D., Kubista, H., Boehm, S., Koenig, X. δ Subunit-containing GABAA receptors are preferred targets for the centrally acting analgesic flupirtine. *Br. J. Pharmacol.*, **2015**, *172*(20), 4946-58.
- [12] Salzer, I., Gantumur, E., Yousuf, A., Boehm, S. Control of sensory neuron excitability by serotonin involves 5HT_{2C} receptors and Ca⁽²⁺⁾-activated chloride channels. *Neuropharmacology*, **2016**, *110*(Pt A), 277-286.
- [13] Salzer, I., Erdem, F.A., Chen, W.Q., Heo, S., Koenig, X., Schicker, K.W., Kubista, H., Lubec, G., Boehm, S., Yang, J.W. Phosphorylation regulates the sensitivity of voltage-gated Kv7.2 channels towards phosphatidylinositol-4, 5-bisphosphate. *J. Physiol.*, **2017**, *595*(3), 759-776.
- [14] Mayer, F.P., Schmid, D., Owens, W.A., Gould, G.G., Apuschkin, M., Kudlacek, O., Salzer, I., Boehm, S., Chiba, P., Williams, P.H., Wu, H.H., Gether, U., Koek, W., Daws, L.C., Sitte, H.H. An unsuspected role for organic cation transporter 3 in the actions of amphetamine. *Neuropsychopharmacology*, **2018**, *43*(12), 2408-2417.
- [15] Ray, S., Salzer, I., Kronschläger, M.T., Boehm, S. The paracetamol metabolite NAPQI reduces excitability in first and second order neurons of the pain pathway through actions on KV7 channels. *Pain*, **2019**, *160*(4), 954-964.
- [16] Hotka, M., Cagalinec, M., Hilber, K., Hool, L., Boehm, S., Kubista, H. L-type Ca²⁺ channel-mediated Ca²⁺ influx adjusts neuronal mitochondrial function to physiological and pathophysiological conditions. *Sci. Signal.*, **2020**, *13*(618).