



Neighborhood Factors Associated with COVID-19 Cases in California

Debora L. Oh¹ · Dan Meltzer¹ · Katarina Wang¹ · Alison J. Canchola¹ · Mindy C. DeRouen¹ · Corinne McDaniels-Davidson^{2,6} · Joseph Gibbons³ · Luis Carvajal-Carmona⁴ · Jesse N. Nodora^{5,6} · Linda Hill⁵ · Scarlett Lin Gomez^{1,7} · Maria Elena Martinez⁶

Received: 31 January 2022 / Revised: 28 October 2022 / Accepted: 1 November 2022
© W. Montague Cobb-NMA Health Institute 2022

Abstract

Background There is a need to assess neighborhood-level factors driving COVID-19 disparities across racial and ethnic groups.

Objective To use census tract-level data to investigate neighborhood-level factors contributing to racial and ethnic group-specific COVID-19 case rates in California.

Design Quasi-Poisson generalized linear models were used to identify neighborhood-level factors associated with COVID-19 cases. In separate sequential models for Hispanic, Black, and Asian, we characterized the associations between neighborhood factors on neighborhood COVID-19 cases. Subanalyses were conducted on neighborhoods with majority Hispanic, Black, and Asian residents to identify factors that might be unique to these neighborhoods. Geographically weighted regression using a quasi-Poisson model was conducted to identify regional differences.

Main Measures All COVID-19 cases and tests reported through January 31, 2021, to the California Department of Public Health. Neighborhood-level data from census tracts were obtained from American Community Survey 5-year estimates (2015–2019), United States Census (2010), and United States Department of Housing and Urban Development.

Key Results The neighborhood factors associated with COVID-19 case rate were racial and ethnic composition, age, limited English proficiency (LEP), income, household size, and population density. LEP had the largest influence on the positive association between proportion of Hispanic residents and COVID-19 cases (– 2.1% change). This was also true for proportion of Asian residents (– 1.8% change), but not for the proportion of Black residents (– 0.1% change). The influence of LEP was strongest in areas of the Bay Area, Los Angeles, and San Diego.

Conclusion Neighborhood-level contextual drivers of COVID-19 burden differ across racial and ethnic groups.

Keywords COVID-19 · Neighborhood · Disparities · California · Race

Introduction

In January 2020, California had only three confirmed cases of 2019 novel coronavirus (COVID-19). Within weeks, community transmission was confirmed, and the first

COVID-related deaths were reported. By September 2021, California had reported over 7.1 million COVID-19 cases and over 77,000 COVID-related deaths [1]. For a state with a population of close to 40 million, that equates to

✉ Debora L. Oh
Debora.Oh@ucsf.edu

¹ Department of Epidemiology and Biostatistics, University of California San Francisco, 550 16th Street, 2nd Floor, San Francisco, CA 94158, USA

² School of Public Health, San Diego State University, San Diego, CA, USA

³ Department of Sociology, San Diego State University, San Diego, CA, USA

⁴ Department of Biochemistry and Molecular Medicine, University of California Davis, Davis, CA, USA

⁵ Herbert Wertheim School of Public Health and Human Longevity Science, University of California San Diego, La Jolla, CA, USA

⁶ Moores Cancer Center, University of California San Diego, La Jolla, CA, USA

⁷ Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, USA

approximately 1 in 6 people infected (among those tested) and 1 in 500 people who have died.

The neighborhood where a person lives exposes them to social, structural, and environmental factors that have been shown to have complex effects on health beyond individual-level factors [2, 3]. Thus, it is perhaps no surprise that while SARS-CoV-2 may spread easily, it does not spread equally across neighborhoods. Inequities in SARS-CoV-2 exposure and COVID-19 outcomes are often driven by structural racism, an interconnected system of policies and institutions that maintains White privilege [4–9] and impacts health through various factors, such as differences in neighborhood factors, employment, and health care access [10–12]. Geographic areas with higher poverty, higher average household size, lower education, and more foreign born residents have been linked with higher COVID-19 case and death rates [13–19]. Since structural racism, such as historical redlining, has driven individuals of minoritized racial groups into racially segregated neighborhoods, neighborhood disparities can contribute to racial and ethnic disparities in health outcomes [20–22].

Use of more granular geographic units to assess environmental factors driving COVID-19 disparities can better inform policy changes and outreach strategies to groups facing structural inequities [15]. Although much has been published on racial disparities in COVID-19 outcomes, there is less known about the specific neighborhood-level factors contributing to inequities in burden of COVID-19 among racial and ethnic groups. Thus, we use census tract-level data to investigate neighborhood, social, and built environment factors that may account for COVID-19 case rates among racially minoritized populations in California. We hypothesize that the neighborhood-level contributors to COVID-19 case burden vary across racial and ethnic groups.

Methods

Data

Census tract-level data on cumulative numbers of COVID-19 cases and tests up to January 31, 2021, were obtained from the California Department of Public Health (CDPH). At that time, less than 2% of Californians were fully vaccinated, and thus, vaccinations had not likely begun to affect the trajectory of the pandemic. Tests were defined as the cumulative total number of COVID tests reported to CDPH. Cases were defined as the cumulative total number of COVID positive cases reported to CDPH via California Reportable Disease Information Exchange (CalREDIE), Los Angeles, and San Diego communicable disease reporting systems.

We used census tract-level data to represent neighborhoods in California. We selected variables based on a

conceptual model of COVID-19 infection that includes neighborhood-level race/ethnicity, socioeconomic status, immigration factors, and housing factors. Census tract-level data on sociodemographics of residents (racial and ethnic composition, sex, age, limited English proficiency (LEP), foreign born, recent immigrant, uninsured, average household size, percentage with less than a high school education, and severe overcrowding) were obtained from the American Community Survey 5-year estimates (2015–2019) [23]. For the race and ethnicity variables, White was defined as non-Hispanic White, Black was defined as Hispanic or non-Hispanic Black, Hispanic included all who identified as Hispanic/Latino, and Asian included all Asian ethnicities but not Native Hawaiians or Pacific Islanders. For all races and ethnicities, residents were included if they reported a certain race and ethnicity alone or in combination with other races and ethnicities. Each race/ethnicity was used as a separate variable in the model, rather than as one categorical variable. LEP was defined as reporting speaking English less than “very well.” Recent immigrant was defined as being born outside the USA and entering the country after 2010. Severe overcrowding was defined as household with more than 1.51 occupants per room [24]. Population density was derived using United States Census (2010) population counts and defined as population counts per square kilometer [25]. Data on low-income households were from the United States Department of Housing and Urban Development (HUD) [26]. HUD’s Extremely Low-Income (ELI) measure defines poor households as those earning below 30% of the area median income. Each factor was expressed as proportion of residents in the census tract (except for average household size and population density).

The neighborhood data analyzed during the current study from American Community Survey, United States Census, and United States Department of Housing and Urban Development are publicly available and complied for download at healthatlas.ucsf.edu. The COVID-19 data analyzed during the current study are not publicly available due to privacy agreements with the California Department of Public Health.

Statistical Analyses

To examine the effect of neighborhood-level factors on the number of COVID-19 cases, quasi-Poisson generalized linear models with census tract-level observations were run, offset for the natural logarithm of population size and adjusted for the number of tests and county (to account for differential COVID-19 policies and practices). Because the case data were overdispersed, we used a generalized linear model with a quasi-likelihood approach (quasi-Poisson) that adjusts count data for overdispersion by including a dispersion parameter [27]. Pearson correlations were used to identify highly correlated variables (> 0.70) which were removed

from the fully adjusted model. The exponentiated regression coefficient, rate ratio (RR) is interpreted as the relative rate of increase in COVID-19 case count per unit increase in neighborhood factor. All variables reported as a proportion were divided by 10 so the estimated effects (i.e., estimated change in the number of cases) can be interpreted as per a 10% change in the factor; similarly, population density was divided by 10,000 so the effects can be interpreted as per a 10,000-unit change in population density.

We estimated the relative contribution of each neighborhood factor to racial and ethnic associations with the number of COVID-19 cases. The baseline model included the proportion of Hispanic, Black, and Asian residents as three separate variables and was adjusted for number of tests and county. Covariables were added individually to the baseline model to determine their influence on COVID-19 associations with proportion Hispanic, Black, and Asian, by calculating the percent change in the racial/ethnic rate ratio when the covariable was added to the baseline model. For Hispanic, Black, and Asian separately, covariables were ranked in order of their influence (from most negative change to most positive change). In separate sequential models for Hispanic, Black, and Asian, covariables were added to the baseline model in the order of their influence. Subanalyses were conducted limited to census tracts with a majority (> 50%) of Hispanic, Black, and Asian residents to identify factors that might be unique to these neighborhoods.

Geographically weighted regression (GWR) using a quasi-Poisson model was conducted to identify regional differences in the associations identified in the previous model [28]. National Historical Geographic Information System spatial data for California Census tracts [29] was joined with California COVID-19 data and sociodemographic data. A quasi-Poisson GWR model with a log link and offset by the natural log of the total population was based on all variables of interest (proportion Asian, proportion Black, proportion Hispanic, proportion ≥ 65 years, proportion with limited English proficiency, proportion uninsured, proportion with extremely low income, severe overcrowding, proportion male, average household size, and population density). Data were then exported for mapping.

Quasi-Poisson models, visualizations, data preparation, and GWR were conducted in R (version 4.0.3) using RStudio (version 1.4.1106) with the tidyverse, sf, ggplot2, and spgwr packages. Maps were made in ArcGIS Pro (version 2.7.0).

Results

There was a total of 3,034,403 COVID-19 cases reported through January 31, 2021, from 58 counties and 8009 census tracts (out of a total of 8057 CA census tracts), with

a maximum number of cases in one census tract of 3692 cases, and a median of 302. The mean number of cases within deciles of each census tract variable is reported in Fig. 1. Case numbers decreased with increasing proportions of White or Asian residents and residents ≥ 65 years. Case numbers increased in census tracts with increasing proportions of Hispanic or Black residents, LEP, uninsured, less than high school degree, severe overcrowding, and higher average household size.

Baseline and fully adjusted models assessing associations of each neighborhood factor and COVID-19 cases are presented in Table 1. High Pearson correlations were identified between proportion of White and Hispanic (0.77), LEP and foreign born (0.85), and uninsured and not high school graduate (0.72). Variables for proportion White, foreign born, and not high school graduate were thus omitted from the fully adjusted model. In fully adjusted models, neighborhood factors that were independently associated with case rate included racial and ethnic composition, age, limited English proficiency, income, household size, and population density.

We assessed the relative contribution of each neighborhood factor on the association between each racial and ethnic group and COVID-19 cases (Table 2). Results of our sequential modeling (Fig. 2) show that proportion with LEP in a census tract had the largest influence on the positive association between the proportion of Hispanic residents and COVID-19 cases in a census tract, meaning that LEP explained some of the association (-2.1% change of estimate for Hispanic with the addition of LEP into the model). This was also true for proportion of Asian residents (-1.8% change), but not for the proportion of Black residents (-0.1% change). None of the assessed variables had a $> 1\%$ influence on the association between the proportion of Black residents and COVID-19 cases.

In a subanalysis of the census tracts with a majority of Hispanic residents (2520 census tracts with $> 50\%$ Hispanic residents), the effect of LEP on COVID-19 cases was attenuated (Table 3) as compared to when looking at all California census tracts. A 10% increase in residents with LEP was associated with a 1.8% (1.018, 95% CI 1.005–1.031) increase in COVID-19 cases (compared to 4.4% (1.044, 95% CI 1.034–1.054) in all California census tracts), after adjusting for other neighborhood factors. In census tracts with a majority of Asian residents (449 census tracts with $> 50\%$ Asians), a 10% increase in residents with LEP was associated with an 11% increase in COVID-19 cases. In census tracts with a majority of Black residents (72 census tracts with $> 50\%$ Black residents), the factor most strongly associated with COVID-19 cases was proportion of Hispanic residents.

In the GWR, 26 census tracts with a population of zero were excluded from the analysis. Results of the generalized





Fig. 1 Mean cumulative COVID-19 cases per 100,000 population (up to January 31, 2021) within each decile of variable for all census tracts in California

GWR, using an optimized bandwidth of 0.006218417, show that the strength of rate ratio estimates vary across California (Fig. 3, Supplemental Figure, and Supplemental Table). For percent LEP, the estimate was highest in many urban and suburban areas of the Bay Area, Los Angeles, and San Diego, as well as some other outlying areas.

Discussion

Results of our analysis using census tract-level data confirm the disproportionate burden of the COVID-19 pandemic in Hispanic and Black neighborhoods in California. Using census tract-level data on all COVID-19 cases in California through January 31, 2021, we observed that a higher proportion of Hispanic and Black residents was significantly associated with a higher COVID-19 case rate. This association persisted even after controlling for age and other neighborhood contextual factors. In addition, in fully adjusted models, we observed that proportion of residents with LEP in a neighborhood has a strong influence on racial and ethnic associations with COVID-19 cases for Hispanic and Asian residents but not Black residents. Associations between LEP and case rate are relatively strong in neighborhoods that are majority Asian and less so in neighborhoods that are majority Hispanic.

Existing studies have established the connection between neighborhood influences and health, specifically how residential segregation and inequalities in resources reinforce each other and lead to inequities in neighborhood physical and social environments and ultimately result in increased risk for adverse health outcomes [30–33]. Recent studies have investigated the neighborhood-level factors contributing to COVID-19 disparities across racial and ethnic groups [13–15, 18, 34–39]. However, few of these studies have analyzed subcounty-level data. At the state and county level, researchers have reported worse COVID-19 outcomes for geographic areas with higher proportions of Black and Hispanic residents [13, 14, 18]. However, there is great heterogeneity in neighborhood environments within states and counties, and smaller geographic units allow for a better understanding of specific neighborhood social and built environment factors that may be driving neighborhood disparities in COVID-19 burden. A study by Reitsma et al. analyzed individual COVID-19 case rates and stratified by subcounty Public Use Microdata Area (PUMA) level and reported that Hispanic residents in California are 8.1 times more likely to live in high-exposure-risk households than White residents and are overrepresented in cumulative cases

[40]. Another study from San Diego County in California analyzed spatiotemporal spread of COVID-19 at the zip code level and found the strongest associations with zip code-level proportions of Spanish-speaking residents and residents with less than 9th grade education [41]. Our study is one of the few statewide analyses of COVID-19 at the census tract level, and the only, to our knowledge, to examine this level of geography in California [34, 35].

The novelty of our analytic approach relates to our assessment of the relative contribution of each of the neighborhood social and built environment factors to COVID-19 cases separately for each group defined by race and ethnicity. A particularly notable result was the association between LEP and COVID-19 case burden specific to neighborhoods with higher proportions of Hispanic and Asian residents. Future studies should examine the extent to which lack of English proficiency in the neighborhood environment contributes to exposure risk or impedes public health efforts to control case rates. Ultimately, our results show the importance of expanding culturally and linguistically tailored public health strategies. For neighborhoods with high proportions of Hispanic and Asian residents, provision of more tailored measures such as translated resources and community-based educators are needed to reduce spread of COVID-19. Seto et al. reported similar findings to ours by assessing census tract-level neighborhood factors and COVID-19 testing in King County, Washington. The authors found a positive trend between proportion of linguistic isolation and COVID-19 case number. However, because this characteristic was highly correlated with race and ethnicity, it was dropped from the modeling [34]. Results from a single health system in Seattle, Washington, showed that despite availability of interpreter services, COVID-19 testing was lower and infection was higher in non-English speakers than English-speaking patients [38], suggesting that solutions need to go beyond providing interpreter services at the point of care.

The COVID-19 pandemic story for Black Californians is different than that for Hispanic or Asian Californians. The COVID-19 burden is clearly greater for census tracts with higher proportions of Black residents. However, none of the neighborhood factors included in this analysis accounted for case burden among predominantly Black neighborhoods. Similar to our study, Benitez et al. analyzed zip code-level data across six diverse cities, including San Diego, to assess COVID-19-related racial and ethnic disparities, showing a clear high burden among neighborhoods with a greater proportion of Hispanic and Black residents. After adjustment for neighborhood-level factors, the association between the proportion of Hispanic residents and COVID-19 cases and deaths was attenuated while that for the proportion of Black residents largely persisted [39]. We do not conceptualize this (or any other) ethnoracial inequity in COVID infection as resulting from purported ethnoracial biology or ethnoracial

Table 1 Associations between cumulative COVID-19 cases (up to January 31, 2021) and neighborhood factors at the census tract level in California ($n = 8009$)

	Minimally adjusted* rate ratio (95% CI)	Fully adjusted† rate ratio (95% CI)
% Asian‡	1.01 (1.01–1.02)	0.99 (0.99–1.00)
% Black‡	1.08 (1.07–1.08)	1.08 (1.07–1.08)
% Hispanic‡	1.18 (1.17–1.18)	1.15 (1.14–1.15)
% White‡	0.96 (0.93–0.99)	–
% Male‡	0.91 (0.90–0.92)	0.96 (0.95–0.98)
% Age 65+‡	1.05 (1.03–1.06)	1.02 (1.01–1.03)
% Limited English proficiency‡	1.04 (1.04–1.05)	1.04 (1.03–1.05)
% Foreign born‡	1.05 (1.04–1.06)	–
% Recent immigrant‡	0.98 (0.97–0.98)	0.97 (0.97–0.98)
% Uninsured‡	1.01 (0.99–1.02)	0.99 (0.97–1.00)
% Extremely low income‡	1.03 (1.02–1.03)	1.03 (1.02–1.04)
% Not HS graduate‡	1.01 (1.00–1.02)	–
Average household size	1.01 (1.00–1.02)	1.03 (1.02–1.04)
% Severe overcrowding‡	0.99 (0.97–1.00)	0.94 (0.92–0.96)
Population density§	1.04 (1.02–1.06)	1.06 (1.03–1.09)

*Quasi-Poisson models with observations on the census tract level with an offset for natural log of total population and adjusted for % Asian, % Black, % Hispanic, number of tests, and county (baseline model). Results for % Asian, % Black, and % White are taken from a single baseline model, all other models include one factor in addition to the baseline model

†Fully adjusted model included % Hispanic, % Black, % Asian, number of tests, county, % male, % age 65+, % limited English proficiency, % recent immigrant, % uninsured, % extremely low income, average household size, % severe overcrowding, and population density. % White, % foreign born, and % not high school graduate were not included in the fully adjusted model as they were highly correlated with other variables

‡Association shown for a 10% increase

§Persons/square kilometer. Association shown for a 10,000-unit increase

Table 2 Percent change in association between race or ethnicity and cumulative COVID-19 cases (up to January 31, 2021) for each neighborhood factor

	% Hispanic	% Black	% Asian
% Male	0.1%	0.0%	–0.1%
% Age 65+	0.7%	0.6%	0.3%
% Limited English proficiency	–2.1%	–0.1%	–1.8%
% Recent immigrant	–0.2%	0.1%	0.2%
% Uninsured	–0.1%	0.4%	0.0%
% Extremely low income	–0.6%	–0.3%	0.0%
Average household size	–0.2%	0.5%	–0.1%
% Severe overcrowding	0.2%	0.5%	0.1%
Population density	–0.2%	–0.1%	–0.1%

Baseline model included % Hispanic, % Black, and % Asian, and was adjusted for tests and county. Neighborhood factors were added individually to the baseline model and the percent change in the racial or ethnic rate ratio was calculated to determine the influence of each factor on the association between COVID-19 cases and % Hispanic, % Black, and % Asian

stereotypes around disease vulnerability or health behaviors; previous scholarship has addressed the lack of evidence for these suggestions and situated racial and ethnic COVID

disparities within existing frameworks outlining fundamental causes of health inequities [42–45]. Other structural and institutional factors not available within our study thus may be driving COVID-19 case burdens among neighborhoods with higher proportions of Black residents. It is also possible that anti-Black racism is woven so tightly into American society that its effects are not adequately captured by common neighborhood-level mediators we assessed like insurance status, household size, education, and population density. Addressing the health inequities and societal impacts of anti-Black racism will require significant, sustained, multi-sector investments and action [46, 47].

This study is ecological and our interpretations of associations between neighborhood factors and neighborhood COVID-19 case rates are not to be extrapolated to explain individual circumstances. Use of census tract boundaries for analysis requires acknowledgment of the modifiable areal unit problem (MAUP) which states that the drawing of boundaries has direct effects on results and contributes to issues with ecologic analysis [48]. Our attempt to adjust for differences in local-level policy was to control for county; however, regional/city policy or other differences may also occur that may be obscured in our analysis. Additionally, because of the dynamic nature of the pandemic and shifting




Fig. 2 Sequential models for cumulative COVID-19 cases (up to January 31, 2021) for neighborhood factors at the census tract level by race and ethnicity. Baseline model included % Hispanic, % Black, and % Asian, and was adjusted for tests and county. Neighborhood factors were added individually to the baseline model to determine their influence on the association between COVID-19 cases and %

Hispanic, % Black, and % Asian. Neighborhood factors were ranked from those with the largest negative change to the racial/ethnic parameter estimate to those with the largest positive change when the factor was added to the baseline model. Sequential models (shown here) added factors one by one to the model in the order of influence

Table 3 Associations between neighborhood factors and cumulative COVID-19 cases up to January 31, 2021, for majority Hispanic, Black, or Asian census tracts

	Rate ratio for > 50% Hispanic census tracts (n = 2520)* (95% CI)	Rate ratio for > 50% Black census tracts (n = 74)* (95% CI)	Rate ratio for > 50% Asian census tracts (n = 449)* (95% CI)
% Hispanic [†]	1.09 (1.08–1.11)	1.42 (1.23–1.64)	1.28 (1.22–1.34)
% Black [†]	1.03 (1.02–1.05)	1.15 (1.02–1.31)	1.46 (1.35–1.57)
% Asian [†]	0.98 (0.97–1.00)	1.03 (0.78–1.35)	1.02 (0.97–1.07)
% Male [†]	0.98 (0.96–1.01)	1.08 (0.95–1.22)	1.07 (0.96–1.18)
% Age 65 + [†]	1.07 (1.04–1.10)	1.05 (0.91–1.21)	1.15 (1.07–1.22)
% Limited English proficiency [†]	1.02 (1.00–1.03)	0.88 (0.72–1.07)	1.11 (1.06–1.16)
% Recent immigrant [†]	0.99 (0.98–1.01)	0.96 (0.90–1.02)	0.99 (0.96–1.02)
% Uninsured [†]	1.00 (0.98–1.02)	1.18 (0.97–1.44)	1.15 (1.03–1.28)
% Extremely low income [†]	1.02 (1.01–1.04)	1.02 (0.97–1.08)	1.02 (0.978–1.06)
Average household size	1.05 (1.03–1.07)	1.02 (0.87–1.19)	1.17 (1.08–1.26)
% Severe overcrowding [†]	0.95 (0.93–0.97)	1.04 (0.80–1.34)	1.00 (0.91–1.10)
Population density [‡]	1.03 (1.00–1.07)	1.01 (0.78–1.29)	1.09 (0.98–1.20)

*Fully adjusted models adjusted for all factors in this table, tests, and county

[†]Association shown for a 10% increase

[‡]Persons/square kilometer. Association shown for a 10,000-unit increase

local policies over time, looking at data over an entire time period may miss temporal shifts of relevance to understanding the spread and dynamically changing risk of COVID-19. The COVID-19 case data in this analysis included cases up until January 31, 2021, which minimizes the impact of vaccination status and the Delta and Omicron variant on cases. In addition, neighborhood, racial, and ethnic COVID disparities extend beyond infection rates to COVID-19-related disease severity and mortality [13–20]. These post-infection outcomes are likely strongly impacted by corresponding

disparities in infection rates [49], but further exacerbated by structural determinants of overall health (comorbidity burden) and health care [50, 51]. While we appreciate the importance of disparities in post-infection clinical outcomes and the structural and social determinants of these clinical outcomes, they were outside the scope of the present study. For the results of GWR, local coefficients for some census tracts in the resulting maps may not be significant and should be interpreted with caution. Furthermore, this study did not directly assess access to health care, which could




Fig. 3 Geographically weighted regression model for cumulative COVID-19 cases (up to January 31, 2021) showing rate ratios for limited English proficiency (LEP) in California

be an uncontrolled confounding variable. Nursing homes were also not accounted for, which could have caused census tracts with many nursing homes to have increased COVID-19 cases. Moreover, this study did not have a measure for the proportion of residents employed in essential work in a census tract.

Our results show that neighborhood-level contextual factors appear to drive some racial and ethnic disparities in

COVID-19 cases in California. These data should inform ongoing and future strategies for outreach to specific groups facing structural inequities. Future research can help better understand factors underlying disparities in COVID-19 transmission in different neighborhoods. Gathering qualitative information from key stakeholders in affected communities may help point toward the root of the issues as well as potential solutions. In order to protect the health of

all, the health of these vulnerable populations needs to be prioritized, and the drivers of these disparities need to be addressed.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s40615-022-01443-y>.

Acknowledgements Contributors: The authors thank the California Department of Public Health and the University of California COVID Modeling Consortium for providing data for this analysis as well as the UCSF School of Medicine Dean's Office of Population Health and Health Equity and Stamen Design for the development of the UCSF Health Atlas.

Author Contribution All authors contributed to the study as follows: study conception and design: DLO, DM, AJC, MCD, JG, SLG, MEM; data collection: DLO, DM; analysis and interpretation of results: DLO, DM, KW, AJC, MCD, CMD, JG, LCC, SLG, MEM; and draft manuscript preparation: all authors.

Funding This work was funded by the Population Health Data Initiative within the University of California San Francisco School of Medicine Dean's Office and the University of California San Francisco Executive Vice Chancellor's Office and grants U54CA132379, U54CA132384, UH3CA233314-02S1, and 5P30CA023100-33. LCC receives funding from the Auburn Community Cancer Endowed Chair in Basic Science, the California Initiative to Advance Precision Medicine (OPR18111), and grants from the National Institutes of Health (NHLBI 21–312-0217571-66106L, NIEHS 3P30ES023513-06S1, NCI R01CA223978, NCI 3P30CA093373, NCATS UL1 TR001860). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Declarations

Competing Interests CMD was a paid consultant for Gilead Sciences and has a close family member that is employed by Quidel Corporation.

References

- California State Government. Tracking COVID-19 in California. <https://covid19.ca.gov/state-dashboard/>. Accessed 27 Sept 2021.
- Diez Roux AV. Neighborhoods and health: what do we know? What should we do? *Am J Public Health*. 2016;106(3):430–1. <https://doi.org/10.2105/AJPH.2016.303064>.
- Duncan DT, Kawachi I. *Neighborhoods and health*. Oxford University Press. 2018.
- Bailey ZD, Feldman JM, Bassett MT. How structural racism works—racist policies as a root cause of US racial health inequities. *N Engl J Med*. 2021;384(8):768–73. <https://doi.org/10.1056/NEJMms2025396>.
- Hardeman RR, Murphy KA, Karbeah J, Kozhimannil KB. Naming institutionalized racism in the public health literature: a systematic literature review. *Public Health Rep Wash DC* 1974. 2018;133(3):240–9. <https://doi.org/10.1177/0033354918760574>.
- Yearby R, Mohapatra S. Law, structural racism, and the COVID-19 pandemic. *J Law Biosci*. 2020;7(1):036. <https://doi.org/10.1093/jlb/lsaa036>.
- Egede LE, Walker RJ. Structural racism, social risk factors, and COVID-19 — a dangerous convergence for Black Americans. *N Engl J Med*. 2020;383(12): e77. <https://doi.org/10.1056/NEJMp2023616>.
- Tan SB, deSouza P, Raifman M. Structural racism and COVID-19 in the USA: a county-level empirical analysis. *J Racial Ethn Health Disparities*. 2022;9:236–46. <https://doi.org/10.1007/s40615-020-00948-8>.
- Dickinson KL, Roberts JD, Banacos N, et al. Structural racism and the COVID-19 experience in the United States. *Health Secur*. 2021;19(S1):S-14. <https://doi.org/10.1089/hs.2021.0031>
- Chen YH, Glymour M, Riley A, et al. Excess mortality associated with the COVID-19 pandemic among Californians 18–65 years of age, by occupational sector and occupation: March through November 2020. *PLoS ONE*. 2021;16(6):e0252454. <https://doi.org/10.1371/journal.pone.0252454>.
- Chin T, Kahn R, Li R, et al. US-county level variation in intersecting individual, household and community characteristics relevant to COVID-19 and planning an equitable response: a cross-sectional analysis. *BMJ Open*. 2020;10(9):e039886. <https://doi.org/10.1136/bmjopen-2020-039886>.
- Jay J, Bor J, Nsoesie E, et al. Neighbourhood income and physical distancing during the COVID-19 pandemic in the United States. *Nat Hum Behav*. 2021;4(12):1294–302.
- Figueroa JF, Wadhwa RK, Mehtsun WT, Riley K, Phelan J, Jha AK. Association of race, ethnicity, and community-level factors with COVID-19 cases and deaths across U.S. counties. *Healthc Amst Neth*. 2021;9(1):100495. <https://doi.org/10.1016/j.hjdsi.2020.100495>.
- Adhikari S, Pantaleo NP, Feldman JM, Ogedegbe O, Thorpe L, Troxel AB. Assessment of community-level disparities in coronavirus disease 2019 (COVID-19) infections and deaths in large US metropolitan areas. *JAMA Netw Open*. 2020;3(7):e2016938. <https://doi.org/10.1001/jamanetworkopen.2020.16938>
- Commodore-Mensah Y, Cooper LA. Reversing the tide of racial and ethnic disparities in excess deaths during the COVID-19 pandemic. *Ann Intern Med*. 2021;174(12):1755–1756. <https://doi.org/10.7326/M21-3841>
- Martínez ME, Nodora JN, Carvajal-Carmona LG. The dual pandemic of COVID-19 and systemic inequities in US Latino communities. *Cancer*. 2021;127(10):1548–50. <https://doi.org/10.1002/ncr.33401>.
- Ingraham NE, Purcell LN, Karam BS, et al. Racial and ethnic disparities in hospital admissions from COVID-19: determining the impact of neighborhood deprivation and primary language. *J Gen Intern Med*. 2021;36(11):3462–70. <https://doi.org/10.1007/s11606-021-06790-w>
- Gross CP, Essien UR, Pasha S, Gross JR, Wang S yi, Nunez-Smith M. Racial and ethnic disparities in population-level COVID-19 mortality. *J Gen Intern Med*. 2020; 35(10):3097–99. <https://doi.org/10.1007/s11606-020-06081-w>
- Quan D, Luna Wong L, Shallal A, et al. Impact of race and socioeconomic status on outcomes in patients hospitalized with COVID-19. *J Gen Intern Med*. 2021;36(5):1302–09. <https://doi.org/10.1007/s11606-020-06527-1>
- Lynch EE, Malcoe LH, Laurent SE, Richardson J, Mitchell BC, Meier HCS. The legacy of structural racism: associations between historic redlining, current mortgage lending, and health. *SSM - Popul Health*. 2021;14:100793. <https://doi.org/10.1016/j.ssmph.2021.100793>.
- Nardone AL, Casey JA, Rudolph KE, Karasek D, Mujahid M, Morello-Frosch R. Associations between historical redlining and birth outcomes from 2006 through 2015 in California. *PLoS ONE*. 2020;15(8):e0237241. <https://doi.org/10.1371/journal.pone.0237241>.
- Nardone A, Casey JA, Morello-Frosch R, Mujahid M, Balmes JR, Thakur N. Associations between historical residential redlining and current age-adjusted rates of emergency department visits due

- to asthma across eight cities in California: an ecological study. *Lancet Planet Health*. 2020;4(1):e24–31. [https://doi.org/10.1016/S2542-5196\(19\)30241-4](https://doi.org/10.1016/S2542-5196(19)30241-4).
23. American Community Survey 5-year estimates, 2015–2019. U.S. Census Bureau. 2020. <https://data.census.gov/>. Accessed 17 Nov 2021
 24. Historical Census of Housing Tables: crowding. U.S. Census Bureau. 2000. <https://www.census.gov/data/tables/time-series/dec/coh-crowding.html>. Accessed 17 Nov 2021
 25. 2010 U.S. Decennial Census. U.S. Census Bureau. 2020. <https://data.census.gov>. Accessed 17 Nov 2021.
 26. Extremely low income households: HUD Comprehensive Housing Affordability Strategy (CHAS) data based on American Community Survey 5-year estimates, 2012–2016. U.S. Department of Housing and Urban Development. https://www.huduser.gov/portal/datasets/cp.html#2006-2017_data. Accessed 17 Nov 2021.
 27. Jay M. Ver Hoef, Peter L. Boveng. Quasi-Poisson vs. negative binomial regression: how should we model overdispersed count data. *Ecology*. 2007;88(11):2766–2772. <https://doi.org/10.1890/07-0043.1>.
 28. Fotheringham AS, Brunsdon C, Charlton M. *Geographically weighted regression: the analysis of spatially varying relationships*. John Wiley & Sons. 2003.
 29. Manson S, Schroeder J, Van Riper D, Kugler T, Ruggles S. IPUMS national historical geographic information system: version 15.0 . IPUMS. 2020. <https://www.nhgis.org/>. Accessed 29 Sept 2021.
 30. Diez Roux AV, Mair C. Neighborhoods and health. *Ann N Y Acad Sci*. 2010;1186:125–45. <https://doi.org/10.1111/j.1749-6632.2009.05333.x>.
 31. Pickett KE, Pearl M. Multilevel analyses of neighbourhood socioeconomic context and health outcomes: a critical review. *J Epidemiol Community Health*. 2001;55(2):111–22. <https://doi.org/10.1136/jech.55.2.111>.
 32. Kirby JB, Kaneda T. Neighborhood socioeconomic disadvantage and access to health care. *J Health Soc Behav*. 2005;46(1):15–31. <https://doi.org/10.1177/002214650504600103>.
 33. Gaskin DJ, Roberts ET, Chan KS, McCleary R, Buttorff C, Delarmente BA. No man is an island: the impact of neighborhood disadvantage on mortality. *Int J Environ Res Public Health*. 2019;16(7):E1265. <https://doi.org/10.3390/ijerph16071265>.
 34. Seto E, Min E, Ingram C, Cummings BJ, Farquhar SA. Community-level factors associated with COVID-19 cases and testing equity in King County, Washington. *Int J Environ Res Public Health*. 2020;17(24):9516. <https://doi.org/10.3390/ijerph17249516>
 35. Madhav KC, Oral E, Straif-Bourgeois S, Rung AL, Peters ES. The effect of area deprivation on COVID-19 risk in Louisiana. *PloS One*. 2020;15(12):e0243028. <https://doi.org/10.1371/journal.pone.0243028>
 36. Arasteh K. Prevalence of comorbidities and risks associated with COVID-19 among Black and Hispanic populations in New York City: an examination of the 2018 New York City Community Health Survey. *J Racial Ethn Health Disparities*. 2021;8(4):863–69. <https://doi.org/10.1007/s40615-020-00844-1>
 37. Khanijahani A, Tomassoni L. Socioeconomic and racial segregation and COVID-19: concentrated disadvantage and Black concentration in association with COVID-19 deaths in the USA. *J Racial Ethn Health Disparities*. 2022;9(1):367–375. <https://doi.org/10.1007/s40615-021-00965-1>
 38. Kim HN, Lan KF, Nkyekyer E, et al. Assessment of disparities in COVID-19 testing and infection across language groups in Seattle, Washington. *JAMA Netw Open*. 2020;3(9):e2021213. <https://doi.org/10.1001/jamanetworkopen.2020.21213>
 39. Benitez J, Courtemanche C, Yelowitz A. Racial and ethnic disparities in COVID-19: evidence from six large cities. *J Econ Race Policy*. 2020;3(4):243–61. <https://doi.org/10.1007/s41996-020-00068-9>.
 40. Reitsma MB, Claypool AL, Vargo J, et al. Racial/ethnic disparities in COVID-19 exposure risk, testing, and cases at the subcounty level in California. *Health Aff*. 2021;40(6). <https://doi.org/10.1377/hlthaff.2021.00098>
 41. Tsou MH, Xu J, Lin CD, et al. Analyzing socioeconomic factors and health disparity of COVID-19 spatiotemporal spread patterns at neighborhood levels in San Diego county. *medRxiv*. 2021. <https://doi.org/10.1101/2021.02.22.21251757>
 42. Chowkwanyun M, Reed AL. Racial health disparities and COVID-19 — caution and context. *N Engl J Med*. 2020;383(3):201–3. <https://doi.org/10.1056/NEJMp2012910>.
 43. Tsai J. COVID-19 is not a story of race, but a record of racism—our scholarship should reflect that reality. *Am J Bioeth AJOB*. 2021;21(2):43–7. <https://doi.org/10.1080/15265161.2020.1861377>.
 44. Deyrup A, Graves JL. Racial biology and medical misconceptions. *N Engl J Med*. 2022;386(6):501–3. <https://doi.org/10.1056/NEJMp2116224>.
 45. Webb Hooper M, Nápoles AM, Pérez-Stable EJ. COVID-19 and racial/ethnic disparities. *JAMA*. 2020;323(24):2466–7. <https://doi.org/10.1001/jama.2020.8598>.
 46. Lee RT, Perez AD, Boykin CM, Mendoza-Denton R. On the prevalence of racial discrimination in the United States. *PLoS ONE*. 2019;14(1):e0210698. <https://doi.org/10.1371/journal.pone.0210698>.
 47. Lloyd C. The Black American experience: here is what we have learned. Gallup Blog. <https://news.gallup.com/opinion/gallup/352763/black-american-experience-what-we-have-learned.aspx>. Published July 27, 2021. Accessed 2 Sept 2021.
 48. Wang Y, Di Q. Modifiable areal unit problem and environmental factors of COVID-19 outbreak. *Sci Total Environ*. 2020;740:139984. <https://doi.org/10.1016/j.scitotenv.2020.139984>.
 49. Zelner J, Trangucci R, Narahariseti R, et al. Racial disparities in coronavirus disease 2019 (COVID-19) mortality are driven by unequal infection risks. *Clin Infect Dis Off Publ Infect Dis Soc Am*. 2021;72(5):e88–95. <https://doi.org/10.1093/cid/ciaa1723>.
 50. Dalsania AK, Fastiggi MJ, Kahlam A, et al. The relationship between social determinants of health and racial disparities in COVID-19 mortality. *J Racial Ethn Health Disparities*. 2022;9(1):288–95. <https://doi.org/10.1007/s40615-020-00952-y>.
 51. Lopez L, Hart LH, Katz MH. Racial and ethnic health disparities related to COVID-19. *JAMA*. 2021;325(8):719–20. <https://doi.org/10.1001/jama.2020.26443>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.