

RESEARCH

Open Access

# Is hypoglycemia fear independently associated with health-related quality of life?

Lizheng Shi<sup>1\*</sup>, Hui Shao<sup>1</sup>, Yingnan Zhao<sup>2</sup> and Nina A Thomas<sup>3</sup>

## Abstract

**Objective:** Patients may fear the symptoms and consequences associated with hypoglycemia. We tested whether fear of hypoglycemia is independently associated with poorer health-related quality of life (HRQOL).

**Research design and methods:** Data were collected using direct-mail survey and enrollment information from adult commercial health plan enrollees with type 2 diabetes during a 12-month period (12/01/2008 to 11/30/2009). HRQOL was evaluated by the EuroQol (EQ)-5D index and 12-item Short Form Health Survey Mental Component Summary (SF-12 MCS) and Physical Component Summary (SF-12 PCS). Fear of hypoglycemia was assessed using the Hypoglycemia Fear Survey (HFS). Two ordinary least-squares (OLS) models of HRQOL controlling for demographics and illness characteristics were specified, and OLS regression coefficients and statistical inferences were compared. Model 1 included 1 variable of hypoglycemia symptoms; Model 2 included both hypoglycemia symptoms and HFS score.

**Results:** Of 3999 patients contacted, 813 responded to the survey. Model 1: hypoglycemia symptoms alone were associated with worse HRQOL (SF-12 MCS and SF-12 PCS scores and EQ-5D utility score; all  $P < 0.05$ ). Model 2: hypoglycemia symptoms were significantly associated only with SF-12 MCS score. HFS total score was significantly associated with all 3 HRQOL scores. Hypoglycemia symptoms, Hispanic ethnicity, and longer diabetes duration were associated with greater hypoglycemia fear. Higher income, white race, and treatment without sulfonylurea or insulin were associated with less hypoglycemia fear (all  $P < 0.05$ ).

**Conclusions:** In addition to the effect of symptomatic hypoglycemia on HRQOL, fear of hypoglycemia was independently associated with lower overall health status and mental and physical health.

**Keywords:** Antidiabetic agents, Health status, Hypoglycemia, Outcome assessment, Quality of life, Type 2 diabetes

Hypoglycemia is associated with increased risk of a variety of adverse clinical outcomes in patients with type 2 diabetes, including microvascular events (eg, nephropathy, retinopathy, neuropathy), macrovascular events (eg, heart failure, peripheral vascular disease, myocardial events, stroke), and death [1-3]. High rates of hypoglycemia symptoms (38%–63%) [4-6] and hypoglycemia episodes (14%–23%; based on low blood sugar) [7,8] are observed in patients with type 2 diabetes in general clinical practice. Hypoglycemia is primarily associated with treatment with insulin and insulin secretagogues [9]. A nationwide cohort study demonstrated that patients who experience hypoglycemia are more likely to be using insulin and sulfonylureas than those who do not

experience hypoglycemia. Hypoglycemia, use of insulin, and use of sulfonylureas are associated with increased risk for major cardiovascular (CV) events, CV disease, and death [3]. Risk of hypoglycemia also increases with age and duration of diabetes [10].

Patients may fear the symptoms associated with a mild episode of hypoglycemia as well as the consequences of severe hypoglycemia, which may include seizure or loss of consciousness [11].

Fear of hypoglycemia is a considerable problem in patients with diabetes [5,12] and may itself represent a barrier to glycemic control. Hypoglycemia is associated with lower health-related quality of life (HRQOL) [5,7,8,13], depression [7], and lower treatment satisfaction [4,5], which could undermine treatment adherence [7]. Fear of hypoglycemia may lead to preventive coping behaviors, such as taking less insulin or increasing food intake [12,14], that negatively

\* Correspondence: lshi1@tulane.edu

<sup>1</sup>Department of Global Health Systems and Development, Tulane University School of Public Health and Tropical Medicine, New Orleans, LA, USA  
Full list of author information is available at the end of the article

affect glycemic control. Fear of hypoglycemia may also lead physicians to limit the aggressiveness of therapy [15].

In a cross-sectional survey study, we found that symptomatic hypoglycemia was associated with worse HRQOL as well as greater fear of hypoglycemia [16]. Compared with patients who did not report a history of symptomatic hypoglycemia, patients with a confirmed history of symptomatic hypoglycemia had higher scores on the Hypoglycemia Fear Survey (HFS) Behavior and Worry subscales (indicating more fear of and worry about hypoglycemia; each  $P < 0.001$ ), lower scores on the EuroQol (EQ)-5D index (indicating poorer health status,  $P < 0.001$ ), and lower scores on the 12-item Short Form Health Survey (SF-12) Mental Component Summary (MCS) score ( $P < 0.001$ ) and Physical Component Summary (PCS) score ( $P = 0.002$ ). The objective of this retrospective analysis was to examine the relationships among symptomatic hypoglycemia, patient self-reported fear of hypoglycemia, and HRQOL while controlling for patient baseline demographic and disease characteristics. We tested the hypothesis that, beyond the effect of symptomatic hypoglycemia itself on patient HRQOL, fear of hypoglycemia is independently associated with lower HRQOL.

## Research design and methods

### Study design

Detailed methodology for this cross-sectional observational research study was previously published [16]. The study combined patient-reported data collected through mail surveys and retrospective analysis of administrative claims data (medical data, pharmacy data, and enrollment information) from a large US health plan (the Life Sciences Research Database). The study protocol was approved by a central institutional review board (IRB) prior to study initiation. And due to the fact that the patients in our study were identified from administrative claims data, the study concept was also approved by the large US health plan prior to commencement of the study. Participants were adult enrollees in the health plan with a diagnosis of type 2 diabetes and  $\geq 2$  pharmacy claims for an oral antidiabetic medication (OAD) with or without insulin identified in the administrative claims data from December 1, 2008, to November 30, 2009.

### Patient survey

For the survey, patients ( $N = 3999$ ) were contacted directly by mail. The patient survey assessed demographic variables, age at time of type 2 diabetes diagnosis, family history of diabetes, and smoking status. Patients were asked about their current OAD, duration of treatment, and modifications to their treatment within the previous 6 months. Medication adherence was assessed using the

Morisky self-report medication adherence 4-item questionnaire [17].

Fear of hypoglycemia was evaluated using the HFS [18]. This validated assessment lists 15 behaviors that patients may engage in to avoid low blood sugar, with potential consequences, and 18 items related to low blood sugar about which patients with diabetes may be concerned. The recall period was 6 months. A total score, Behavior subscale score (sum of the 15 behavioral items), and Worry subscale score (sum of the 18 worry items) were generated. In addition, patients were asked whether they had changed their medication or were afraid to change medications because of hypoglycemia.

Health-related quality of life was assessed using 2 instruments. Overall health status was assessed using the EQ-5D, a standardized measure that rates 5 dimensions of health (mobility, self-care, usual activities, pain/discomfort, anxiety/depression) across 3 levels (no problems, some/moderate problems, and extreme problems) [19,20]. The index value (ie, health utility) between 0 (death) and 1 (perfect health) was calculated using the value sets (weights) from a US population sample. The other instrument was the SF-12 [21], which can produce an MCS and a PCS. Summary scores range from 0 to 100, with a higher score indicating a better quality of life in mental and physical components. Hypoglycemia was defined based on American Diabetes Association symptoms [22] and blood glucose readings at the time that symptoms occurred.

The survey packet we used in this study contained a pre-paid cash incentive of \$10 to the patients, along with the invitation letter printed on letterhead, the IRB-approved consent form, the survey instrument printed in booklet form, and a postage-paid business reply envelope. We sent reminders to the patients 2 and 4 weeks following the initial survey to increase our response rate.

### Statistical analyses

Survey variables (eg, demographics and clinical characteristics) were analyzed descriptively, and bivariate comparisons of demographic characteristics and outcome measures were performed using appropriate tests (eg, *t*-test or chi-square test) based on the distribution of the measure. Multivariate analyses of HFS, SF-12, and EQ-5D outcomes were conducted using ordinary least squares. Specific predictors (ie, explanatory variables) to be included were selected primarily based on clinical rationale. Associations among symptomatic hypoglycemia, fear of hypoglycemia, and HRQOL were modeled in 2 ordinary least-squares regressions, controlling for sociodemographics, illness characteristics, and treatment factors [16]. Model 1 of the HRQOL =  $f$  (symptomatic hypoglycemia) tested the associations between symptomatic hypoglycemia and HRQOL, and based on previously published results [16], these were expected to

be statistically significant. Model 2, which was the primary assessment of the study hypothesis, added 1 additional variable of hypoglycemia fear and tested the impact that fear of hypoglycemia had on health status independently of the impact of hypoglycemia symptoms. Following standard procedure, regression diagnostics were performed to assess goodness of fit and violations of model assumptions (eg, multicollinearity, heteroskedasticity) for each model. Any violations of the model were noted and appropriate corrections made to the data. The fitted and observed data were examined to uncover outliers, their effect on the analysis, and possible misspecification of the initial equation. SAS 9.2 software (SAS Institute, Inc., Cary, NC) was used for statistical analyses [23]. The significance level was set at 0.05 (2-tailed).

Because different classes of antidiabetic drugs may be more likely to cause hypoglycemia [24-27], the association between fear for hypoglycemia and quality of life might also be different across different patient subgroups of antidiabetic medication users. Due to this concern, the same statistical analysis procedures were conducted on two patient subgroups of antidiabetic medication users: insulin users and sulfonylurea users.

## Results

### Patient survey results

Of 3999 patients contacted, 813 responded with a complete survey (response rate, 20.3%). Of these, 578 (71.1%) reported  $\geq 1$  hypoglycemia symptom within the previous 3 months. Among the population characteristics assessed, only OAD group and sex showed a statistically significant difference between patients with and without symptomatic hypoglycemia (Table 1). Patients in the OAD plus insulin group were more likely to report  $\geq 1$  hypoglycemia symptom (83.3% [244/293]) compared with those in the sulfonylurea without insulin group (69.9% [172/246]) or the nonsulfonylurea OAD without insulin group (59.1% [162/274];  $P < 0.001$ ). Significantly more women (77.1% [263/341]) than men (66.7% [315/472]) reported  $\geq 1$  hypoglycemia symptom.

Symptomatic hypoglycemia was associated with more fear of hypoglycemia (higher total HFS score) and worse HRQOL (lower SF-12 MCS, SF-12 PCS, and EQ-5D scores; each,  $P < 0.05$ ; Figure 1). Table 2 presents the associations between the HFS and various factors. The presence of symptomatic hypoglycemia was associated with greater fear of hypoglycemia and had the strongest association (regression coefficient  $> 10$  points of HFS total score,  $P < 0.0001$ ). Other factors also showed statistically significant associations (Table 2). Higher income was associated with less fear of hypoglycemia ( $P < 0.0001$ ), as was white racial categorization compared with nonwhite racial categorizations ( $P < 0.0001$ ). Hispanic ethnicity was associated with greater fear of hypoglycemia ( $P = 0.03$ ). Longer

duration with a diagnosis of diabetes and better self-reported adherence were also associated with greater fear of hypoglycemia (each  $P < 0.0001$ ).

The ordinary least-squares model of HRQOL using symptomatic hypoglycemia only and controlling for demographics and illness characteristics (Model 1, Table 3) yielded regression coefficients for symptomatic hypoglycemia on HRQOL scores. The associations between symptomatic hypoglycemia and EQ-5D score and SF-12 MCS score were statistically significant (each  $P < 0.05$ ). In Model 2, which included the additional variable of HFS, the HFS score was significantly associated with scores on all 3 HRQOL outcomes (each  $P < 0.0001$ ), whereas symptomatic hypoglycemia showed a statistically significant association only for the SF-12 MCS ( $P = 0.022$ ). The fully specified regression models for Table 3 are provided in Additional file 1: Tables S1.1-S1.6.

Subgroup analyses were conducted for insulin users (Table 4) and sulfonylurea users (Table 5). For insulin users ( $n = 293$ ), Model 1 showed that the symptomatic hypoglycemia was not associated with 3 HRQOL scores (each  $P > 0.05$ ). Model 2 showed that the HFS score was still significantly associated with scores on all 3 HRQOL outcomes (each  $P < 0.05$ ), whereas symptomatic hypoglycemia showed no statistically significant association with any of the 3 HRQOL outcomes (each  $P > 0.05$ ). For sulfonylurea users ( $n = 246$ ), Model 1 showed that the symptomatic hypoglycemia was associated with EQ-5D scores and SF-12 MCS scores (each  $P < 0.05$ ). Model 2 showed that the HFS score was significantly associated with scores on all 3 HRQOL outcomes (each  $P < 0.05$ ), whereas symptomatic hypoglycemia was only significantly associated with EQ-5D scores and SF-12 MCS scores (each  $P < 0.05$ ). The fully specified regression models for Table 4 and Table 5 are provided in Additional file 1: Tables S2.0-S3.5.

## Discussion

In patients with type 2 diabetes treated with antihyperglycemic therapy, hypoglycemia imposes a significant physical and psychological burden [28]. Previously, we demonstrated that hypoglycemia is significantly associated with poorer HRQOL as well as increased fear of hypoglycemia [16]. Other studies have shown an association between the severity of hypoglycemia and increased fear of hypoglycemia [29-31]. This retrospective analysis was conducted to test the hypothesis that fear of hypoglycemia is itself independently associated with poorer HRQOL. Results support this hypothesis, demonstrating a significant independent association between fear of hypoglycemia and scores on the EQ-5D and SF-12 PCS and MCS. This relationship was demonstrated for both insulin and sulfonylurea users. These results are consistent with previous cross sectional studies that have

**Table 1 Population characteristics between patients who reported versus those who did not report symptoms of hypoglycemia**


Characteristic, n (%)	No hypoglycemia (n = 235)	Hypoglycemia (n = 578)	P value*
Age, y			
<65	203 (86.4)	525 (90.8)	0.0603
≥65	32 (13.6)	53 (9.2)	
Sex			
Female	78 (33.2)	263 (45.5)	0.0013
Male	157 (66.8)	315 (54.5)	
Family history of diabetes			
No	58 (26.9)	110 (20.4)	0.0528
Yes	158 (73.2)	430 (79.6)	
Treatment			
Any OAD with insulin	49 (20.9)	244 (42.2)	
SU without insulin	74 (31.5)	172 (29.8)	<.0001
Non-SU OAD without insulin	112 (47.7)	162 (28.0)	
Diabetes treatment period, y			
<1	45 (19.2)	96 (16.7)	0.4765
1–2	70 (29.9)	151 (26.3)	
3–5	63 (26.9)	167 (29.1)	
6–10	36 (15.4)	91 (15.9)	
>10	20 (8.6)	69 (12.0)	
Ethnicity			
White			
No	52 (22.5)	161 (28.3)	0.0961
Yes	179 (77.5)	409 (71.8)	
Black or African American			
No	196 (84.9)	469 (82.3)	0.3806
Yes	35 (15.2)	101 (17.7)	
Hispanic or Latino			
No	209 (89.7)	525 (92.3)	0.2360
Yes	24 (10.3)	44 (7.7)	
Marital status			
Single, never married	25 (10.7)	68 (11.8)	0.6079
Married	167 (71.4)	390 (67.7)	
Divorced	32 (13.7)	82 (14.2)	
Separated	2 (0.9)	14 (2.4)	
Widowed	8 (3.4)	22 (3.8)	
Annual household income, \$US			
<25,000	13 (5.9)	44 (8.0)	0.7025
25,000–49,999	68 (30.6)	169 (30.8)	
50,000–74,999	42 (18.9)	116 (21.1)	
75,000–99,999	36 (16.2)	78 (14.2)	
>100,000	63 (28.4)	142 (25.9)	

**Table 1 Population characteristics between patients who reported versus those who did not report symptoms of hypoglycemia (Continued)**

Education			
Less than secondary	9 (3.9)	14 (2.5)	0.8425
Some secondary	4 (1.7)	14 (2.5)	
Secondary or equivalent	53 (22.8)	131 (22.9)	
Tertiary but no degree	60 (25.8)	166 (29.0)	
2-year undergraduate degree	28 (12.0)	59 (10.3)	
>2-year undergraduate degree	51 (21.9)	199 (20.8)	
Graduate school	28 (12.0)	69 (12.1)	
US Region			
Northeast	13 (5.5)	33 (5.7)	0.2144
Midwest	29 (12.3)	54 (9.3)	
South	181 (77.0)	440 (76.1)	
West	12 (5.1)	51 (8.8)	
Morisky adherence			
Low	123 (52.8)	265 (46.3)	0.3058
Relatively low	71 (30.5)	186 (32.5)	
Middle	28 (12.0)	77 (13.4)	
Relatively high	10 (4.3)	36 (6.3)	
High	1 (0.4)	9 (1.6)	

OAD = oral antidiabetic medication; SU = sulfonylurea.

\*Chi-square test.



**Table 2 Influencing factors of hypoglycemia fear survey total scores**

Variable (reference)	Regression coefficient	Standard error	P value	95% CI	
				Lower	Upper
Hypoglycemia	10.47	1.28	<0.0001	7.95	12.99
Age (<65 y)					
≥65 y	-1.23	1.91	0.52	-4.98	2.51
Sex (male)					
Female	-1.76	1.23	0.15	-4.17	0.66
Body mass index	-0.10	0.08	0.22	-0.25	0.06
Family history of diabetes	-0.58	1.37	0.67	-3.27	2.10
Treatment (SU without insulin)					
Non-SU OAD without insulin	-2.25	1.44	0.12	-5.09	0.58
Any OAD with insulin	2.09	1.42	0.14	-0.71	4.89
Duration of current diabetes medication	-0.70	0.47	0.14	-1.63	0.23
Ethnicity					
White (not white)	-4.56	1.35	<0.0001	-7.22	-1.91
Hispanic or Latino (not Hispanic or Latino)	4.74	2.13	0.03	0.56	8.92
Married (currently not married)	-0.04	1.29	0.97	-2.59	2.50
Household income level	-1.94	0.47	<0.0001	-2.86	-1.03
Education (less than secondary)	0.75	3.56	0.83	-6.25	7.75
Region (Northeast)					
Midwest	1.24	2.82	0.66	-4.29	6.77
South	1.41	2.33	0.55	-3.17	5.99
West	0.21	3.14	0.95	-5.96	6.37
Diabetes duration (y)	0.36	0.08	<0.0001	0.20	0.52
Morisky adherence	1.72	0.60	<0.0001	0.54	2.91

OAD = oral antidiabetic medication; SU = sulfonylurea.

shown an association between fear of hypoglycemia and HRQOL [29,32].

A significant association was observed between symptomatic hypoglycemia and EQ-5D and SF-12 MCS scores in Model 1, whereas only the association with SF-12 MCS was significant after incorporating the HFS into the model. In contrast, fear of hypoglycemia demonstrated a significant association with all 3 HRQOL outcomes. This finding suggests that, statistically, fear of hypoglycemia may be a more important predictor than

hypoglycemia itself of patient well-being and health status.

Multiple factors associated with fear of hypoglycemia were identified. Consistent with our study [16] and other reports [29,31,33], symptomatic hypoglycemia had the strongest (positive) association with fear of hypoglycemia. Thus, it appears that in patients with type 2 diabetes, experiencing an event of hypoglycemia increases rather than diminishes fear of such an event. An important clinical practice implication of this finding is the need for

**Table 3 Role of hypoglycemia fear on health-related quality of life using alternative models**

Variable	EQ-5D (US)			SF-12 MCS			SF-12 PCS		
	β	P value	95% CI for β	β	P value	95% CI for β	β	P value	95% CI for β
Model 1									
Hypo	0.041	0.002	-0.067 to -0.015	-3.628	<0.0001	-5.259 to -1.998	-0.974	0.247	-2.623 to 0.675
Model 2									
Hypo	-0.014	0.303	-0.040 to 0.013	-1.937	0.022	-3.591 to -0.283	0.579	0.500	-1.104 to 2.262
HFS	-0.003	<0.0001	-0.003 to -0.002	-0.162	<0.0001	-0.210 to -0.115	-0.149	<0.0001	-0.197 to -0.101

β = regression coefficient; EQ = EuroQol; Hypo = symptomatic hypoglycemia; HFS = Hypoglycemia Fear Survey; MCS = Mental Component Summary; PCS = Physical Component Summary; SF-12 = 12-item Short Form Health Survey.

**Table 4 Role of hypoglycemia fear on health-related quality of life using alternative models (Insulin Subgroup)**

Variable	EQ-5D (US)			SF-12 MCS			SF-12 PCS		
	$\beta$	P value	95% CI for $\beta$	$\beta$	P value	95% CI for $\beta$	$\beta$	P value	95% CI for $\beta$
Model 1									
Hypo	-0.001	0.96	-0.058 to 0.055	-2.506	0.137	-5.813 to 0.801	-1.668	0.359	-5.244 to 1.908
Model 2									
Hypo	0.029	0.311	-0.027 to 0.086	-0.285	0.863	-3.534 to 2.965	-0.358	0.847	-4.011 to 3.296
HFS	-0.003	<0.001	-0.004 to -0.001	-0.188	<0.0001	-0.259 to -0.116	-0.111	<0.007	-0.191 to -0.031

$\beta$  = regression coefficient; EQ = EuroQol; Hypo = symptomatic hypoglycemia; HFS = Hypoglycemia Fear Survey; MCS = Mental Component Summary; PCS = Physical Component Summary; SF-12 = 12-item Short Form Health Survey.

healthcare professionals to be cognizant of the fear of hypoglycemia and take steps to address this fear and related behaviors.

In the current analysis, Hispanic/Latino ethnicity, duration of diabetes, and self-reported adherence also showed a positive association with fear of hypoglycemia, whereas household income and white race showed a negative association. These data suggest that demographic variables might be helpful in predicting those patients most likely to experience fear of hypoglycemia. The positive association between duration of diabetes and fear of hypoglycemia could reflect greater insulin use in patients with a longer duration of disease. Alternatively, an independent association between these variables would seem to further indicate that experiencing hypoglycemia increases fear of the event.

As no causality can be inferred in this cross-sectional analysis, the positive relationship between self-reported adherence and fear of hypoglycemia is difficult to interpret. In a previous study, patients reporting hypoglycemia symptoms were more likely to report barriers to adherence (eg, bothered by medication side effects, unable to follow plans) [4]. In addition, data suggest that patients who have experienced a hypoglycemic episode are likely to initiate preventive behaviors that may include modification of their dosing regimen [14,16]. Although these findings may seem at odds with one another, it is reasonable to expect that patients adherent to strict glycemic control regimens would have more reason to be fearful of hypoglycemia. Additional studies with longitudinal data

collection are needed to fully understand the variables that predict fear of hypoglycemia.

Predictors of hypoglycemia in patients with type 2 diabetes are better understood. In the current analysis, more women versus men and more patients treated with OAD plus insulin versus sulfonylurea without insulin or nonsulfonylurea OAD without insulin reported hypoglycemia. The difference in hypoglycemia with respect to diabetes therapy is to be expected, given that insulin and insulin secretagogues are the most common cause of hypoglycemia in patients with type 2 diabetes [15]. The observation that more women than men reported hypoglycemic symptoms was unexpected but has been observed in another study [29] and could possibly reflect sex-related differences in self-reporting of hypoglycemic symptoms.

This analysis has some limitations. Because the study was cross-sectional, no causal relationships can be inferred from the findings. Data on the frequency and severity of hypoglycemic episodes were not collected so their contribution to fear of hypoglycemia could not be assessed. Also, because the survey design used a convenience sample of patients that included only those who were members of specific health plans, with voluntary participation, survey participants may not be representative of all patients with type 2 diabetes. The study also relied on patient self-report, and the survey had a patient recall period of 6 months. Longitudinal cohort studies are warranted to further understand the impact of hypoglycemia and to test potential interventions to address the issue of hypoglycemia fear to improve health

**Table 5 Role of hypoglycemia fear on health-related quality of life using alternative models (Sulfonylurea Subgroup)**

Variable	EQ-5D (US)			SF-12 MCS			SF-12 PCS		
	$\beta$	P value	95% CI for $\beta$	$\beta$	P value	95% CI for $\beta$	$\beta$	P value	95% CI for $\beta$
Model 1									
Hypo	-0.077	0.001	-0.123 to -0.031	-6.531	<0.001	-9.606 to -3.456	-1.815	0.241	-4.857 to 1.226
Model 2									
Hypo	-0.060	0.014	-0.107 to -0.013	-5.300	0.001	-8.453 to -2.147	0.056	0.971	-2.973 to 3.084
HFS	-0.002	0.014	-0.004 to -0.000	-0.140	0.008	-0.243 to -0.038	-0.213	<0.001	-0.311 to -0.114

$\beta$  = regression coefficient; EQ = EuroQol; Hypo = symptomatic hypoglycemia; HFS = Hypoglycemia Fear Survey; MCS = Mental Component Summary; PCS = Physical Component Summary; SF-12 = 12-item Short Form Health Survey.

for patients with diabetes. In conclusion, in addition to symptomatic hypoglycemia itself, fear of hypoglycemia was independently associated with lower HRQOL (overall health status, mental health, and physical health) in patients with type 2 diabetes. There is an unmet need for patient education programs that address patient fear of hypoglycemia and use of medications with a lower risk of hypoglycemia.

## Additional file

**Additional file 1: Table S1.1** Fully Specified Hypoglycemia Model 1: EQ-5D (US). **Table S1.2** Fully Specified Hypoglycemia Model 1: SF-12 MCS. **Table S1.3** Fully Specified Hypoglycemia Model 1: SF-12 PCS. **Table S1.4** Fully Specified Hypoglycemia Model 2: EQ-5D (US). **Table S1.5** Fully Specified Hypoglycemia Model 2: SF-12 MCS. **Table S1.6** Fully Specified Hypoglycemia Model 2: SF-12 PCS. **Table S2.0** Hypoglycemia Model 1: Insulin Subgroup Measured by EQ-5D(US). **Table S2.1** Hypoglycemia Model 1: Insulin Subgroup Measured by SF-12 MCS. **Table S2.2** Hypoglycemia Model 1: Insulin Subgroup Measured by SF-12 PCS. **Table S2.3** Hypoglycemia Model 2: Insulin Subgroup Measured by EQ-5D (US). **Table S2.4** Hypoglycemia Model 2: Insulin Subgroup Measured by SF-12 MCS. **Table S2.5** Hypoglycemia Model 2: Insulin Subgroup Measured by SF-12 PCS. **Table S3.0** Hypoglycemia Model 1: Sulfonylurea Subgroup Measured by EQ-5D (US). **Table S3.1** Hypoglycemia Model 1: Sulfonylurea Subgroup Measured by SF-12 MCS. **Table S3.2** Hypoglycemia Model 1: Sulfonylurea Subgroup Measured by SF-12 PCS. **Table S3.3** Hypoglycemia Model 2: Sulfonylurea Subgroup Measured by EQ-5D (US). **Table S3.4** Hypoglycemia Model 2: Sulfonylurea Subgroup Measured by SF-12 MCS. **Table S3.5** Hypoglycemia Model 2: Sulfonylurea Subgroup Measured by SF-12 PCS.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

LS is the guarantor of this article. HS performed data analysis. YZ reviewed the results and improved model specifications. LS, HS, YZ, and NAT reviewed, edited, and approved the manuscript.

## Acknowledgments

The authors thank Vivian Fonseca, MD, for his helpful discussions during the development of this study. All authors contributed to the concept and design of the study; the acquisition, analysis, and interpretation of data; development of the statistical analysis plan; and review/editing of the manuscript for content. All authors had full access to the data included in the manuscript and approved the final version.

L.S. has received grants (to Tulane) from Novartis, Takeda, and Genentech and has received honoraria for consulting from AstraZeneca. N.A.T. is an employee of Bristol-Myers Squibb. Y.Z. and H.S. report no potential conflicts. The study was sponsored by AstraZeneca (Wilmington, DE). Medical writing support for the preparation of this manuscript was provided by Deborah M. Campoli-Richards, BSPHarm, RPH, Nicole Strangman, PhD, and Janet E. Matsuura, PhD, of Complete Healthcare Communications, Inc. (Chadds Ford, PA), with funding from Bristol-Myers Squibb and AstraZeneca. Data from this study were presented at the International Society for Pharmacoeconomics and Outcomes Research 18th Annual International Meeting, May 18–22, 2013, New Orleans, LA.

## Author details

<sup>1</sup>Department of Global Health Systems and Development, Tulane University School of Public Health and Tropical Medicine, New Orleans, LA, USA.

<sup>2</sup>College of Pharmacy, Xavier University of Louisiana, New Orleans, LA, USA.

<sup>3</sup>US Health Services Research, Bristol-Myers Squibb, Princeton, NJ, USA.

Received: 5 May 2014 Accepted: 6 November 2014

Published online: 30 November 2014

## References

1. Zhao Y, Campbell CR, Fonseca V, Shi L: Impact of hypoglycemia associated with antihyperglycemic medications on vascular risks in veterans with type 2 diabetes. *Diabetes Care* 2012, **35**:1126–1132.
2. Zoungas S, Patel A, Chalmers J, De Galan BE, Li Q, Billot L, Woodward M, Ninomiya T, Neal B, MacMahon S, Grobbee DE, Pascal Kengne A, Marre M, Heller S: Severe hypoglycemia and risks of vascular events and death. *N Engl J Med* 2010, **363**:1410–1418.
3. Hsu PF, Sung SH, Cheng HM, Yeh JS, Liu WL, Chan WL, Chen CH, Chou P, Chuang SY: Association of clinical symptomatic hypoglycemia with cardiovascular events and total mortality in type 2 diabetes: a nationwide population-based study. *Diabetes Care* 2013, **36**:894–900.
4. Alvarez Guisasaola F, Tofe Povedano S, Krishnarajah G, Lyu R, Mavros P, Yin D: Hypoglycaemic symptoms, treatment satisfaction, adherence and their associations with glycaemic goal in patients with type 2 diabetes mellitus: findings from the Real-Life Effectiveness and Care Patterns of Diabetes Management (RECAP-DM) study. *Diabetes Obes Metab* 2008, **10**:25–32.
5. Marrett E, Stargardt T, Mavros P, Alexander CM: Patient-reported outcomes in a survey of patients treated with oral antihyperglycaemic medications: associations with hypoglycaemia and weight gain. *Diabetes Obes Metab* 2009, **11**:1138–1144.
6. Pollack MF, Purayidathil FW, Bolge SC, Williams SA: Patient-reported tolerability issues with oral antidiabetic agents: associations with adherence; treatment satisfaction and health-related quality of life. *Diabetes Res Clin Pract* 2010, **87**:204–210.
7. Green AJ, Fox KM, Grandy S: Self-reported hypoglycemia and impact on quality of life and depression among adults with type 2 diabetes mellitus. *Diabetes Res Clin Pract* 2012, **96**:313–318.
8. Williams SA, Pollack MF, Di Bonaventura M: Effects of hypoglycemia on health-related quality of life, treatment satisfaction and healthcare resource utilization in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract* 2011, **91**:363–370.
9. Inzucchi SE, Bergenstal RM, Buse JB, Diamant M, Ferrannini E, Nauck M, Peters AL, Tsapas A, Wender R, Matthews DR: Management of hypoglycemia in type 2 diabetes: a patient-centered approach: position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2012, **35**:1364–1379.
10. Brunton SA: Hypoglycemic potential of current and emerging pharmacotherapies in type 2 diabetes mellitus. *Postgrad Med* 2012, **124**:74–83.
11. Cryer PE: Hypoglycemia: still the limiting factor in the glycemic management of diabetes. *Endocr Pract* 2008, **14**:750–756.
12. Wild D, von Maltzahn R, Brohan E, Christensen T, Clauson P, Gonder-Frederick L: A critical review of the literature on fear of hypoglycemia in diabetes: implications for diabetes management and patient education. *Patient Educ Couns* 2007, **68**:10–15.
13. Pettersson B, Rosenqvist U, Deleskog A, Journath G, Wandell P: Self-reported experience of hypoglycemia among adults with type 2 diabetes mellitus (Exhype). *Diabetes Res Clin Pract* 2011, **92**:19–25.
14. Leiter LA, Yale J, Chiasson J, Harris S, Kleinstiver P, Sauriol L: Assessment of the impact of fear of hypoglycemic episodes on glycemic and hypoglycemia management. *Can J Diabetes* 2005, **29**:186–192.
15. Amiel SA, Dixon T, Mann R, Jameson K: Hypoglycaemia in type 2 diabetes. *Diabet Med* 2008, **25**:245–254.
16. Williams SA, Shi L, Brenneman SK, Johnson JC, Wegner JC, Fonseca V: The burden of hypoglycemia on healthcare utilization, costs, and quality of life among type 2 diabetes mellitus patients. *J Diabetes Complications* 2012, **26**:399–406.
17. Morisky DE, Green LW, Levine DM: Concurrent and predictive validity of a self-reported measure of medication adherence. *Med Care* 1986, **24**:67–74.
18. Cox DJ, Irvine A, Gonder-Frederick L, Nowacek G, Butterfield J: Fear of hypoglycemia: quantification, validation, and utilization. *Diabetes Care* 1987, **10**:617–621.
19. Matza LS, Boye KS, Yurgin N: Validation of two generic patient-reported outcome measures in patients with type 2 diabetes. *Health Qual Life Outcomes* 2007, **5**:47.



20. What is EQ-5D? [<http://www.euroqol.org>].
21. Ware J Jr, Kosinski M, Keller SD: A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996, **34**:220–233.
22. Living with diabetes. [<http://www.diabetes.org/living-with-diabetes/treatment-and-care/blood-glucose-control/hypoglycemia-low-blood.html>]
23. SAS Institute Inc: *SAS/STAT® User's Guide. Version 9.2*. Cary, NC: SAS Institute Inc; 2008.
24. Wrobel MP, Wystrychowski G, Psurek A, Szymborska-Kajaneck A, Strojek K: Association between hypoglycemia and the type of insulin in diabetic patients treated with multiple injections: an observational study. *Pol Arch Med Wewn* 2014, **124**:173–179.
25. Pitsillides AN, Anderson SM, Kovatchev B: Hypoglycemia risk and glucose variability indices derived from routine self-monitoring of blood glucose are related to laboratory measures of insulin sensitivity and epinephrine counterregulation. *Diabetes Technol Ther* 2011, **13**:11–17.
26. Kalra S, Deepak MC, Narang P, Singh V, Maheshwari A: Correlation between measures of hypoglycemia and glycemic improvement in sulfonylurea treated patients with type 2 diabetes in India: results from the OBSTACLE hypoglycemia study. *J Postgrad Med* 2014, **60**:151–155.
27. Odawara M, Kadowaki T, Naito Y: Incidence and predictors of hypoglycemia in Japanese patients with type 2 diabetes treated by insulin glargine and oral antidiabetic drugs in real-life: ALOHA post-marketing surveillance study sub-analysis. *Diabetol Metab Syndr* 2014, **6**:20.
28. Seaquist ER, Anderson J, Childs B, Cryer P, Dagogo-Jack S, Fish L, Heller SR, Rodriguez H, Rosenzweig J, Vigersky R: Hypoglycemia and diabetes: a report of a workgroup of the American Diabetes Association and The Endocrine Society. *Diabetes Care* 2013, **36**:1384–1395.
29. Marrett E, Radican L, Davies MJ, Zhang Q: Assessment of severity and frequency of self-reported hypoglycemia on quality of life in patients with type 2 diabetes treated with oral antihyperglycemic agents: a survey study. *BMC Res Notes* 2011, **4**:251.
30. Sheu WH, Ji LN, Nitiyanant W, Baik SH, Yin D, Mavros P, Chan SP: Hypoglycemia is associated with increased worry and lower quality of life among patients with type 2 diabetes treated with oral antihyperglycemic agents in the Asia-Pacific region. *Diabetes Res Clin Pract* 2012, **96**:141–148.
31. Stargardt T, Gonder-Frederick L, Krobot KJ, Alexander CM: Fear of hypoglycaemia: defining a minimum clinically important difference in patients with type 2 diabetes. *Health Qual Life Outcomes* 2009, **7**:91.
32. Solli O, Stavem K, Kristiansen IS: Health-related quality of life in diabetes: the associations of complications with EQ-5D scores. *Health Qual Life Outcomes* 2010, **8**:18.
33. Mitchell BD, Vietri J, Zagar A, Curtis B, Reaney M: Hypoglycaemic events in patients with type 2 diabetes in the United Kingdom: associations with patient-reported outcomes and self-reported HbA1c. *BMC Endocr Disord* 2013, **13**:59.

doi:10.1186/s12955-014-0167-3

Cite this article as: Shi *et al.*: Is hypoglycemia fear independently associated with health-related quality of life? *Health and Quality of Life Outcomes* 2014 **12**:167.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

