

RESEARCH ARTICLE

Association of psychotropic drug use with falls among older adults in Germany. Results of the German Health Interview and Examination Survey for Adults 2008-2011 (DEGS1)

Yong Du*, Ingrid-Katharina Wolf, Hildtraud Knopf

Department of Epidemiology and Health Monitoring, Robert Koch Institute, Berlin Germany

* duy@rki.de



OPEN ACCESS

Citation: Du Y, Wolf I-K, Knopf H (2017) Association of psychotropic drug use with falls among older adults in Germany. Results of the German Health Interview and Examination Survey for Adults 2008-2011 (DEGS1). PLoS ONE 12(8): e0182432. <https://doi.org/10.1371/journal.pone.0182432>

Editor: Alessandra Marengoni, University of Brescia, ITALY

Received: February 27, 2017

Accepted: July 18, 2017

Published: August 8, 2017

Copyright: © 2017 Du et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The authors confirm that some access restrictions apply to the data underlying the findings. The data set cannot be made publicly available because informed consent from study participants did not cover public deposition of data. However, the minimal data set underlying the findings is archived in the 'Health Monitoring' Research Data Centre at the Robert Koch Institute (RKI) and can be accessed by all interested researchers. On-site access to the data set is possible at the Secure Data Center of the

Abstract

Purpose

To investigate the association of psychotropic drug use with falls among older adults in Germany based on data from the National Health Interview and Examination Survey for Adults 2008–2011 (DEGS1).

Methods

DEGS1 collected data on drug use in the past 7 days and on falls occurred in the last 12 months. Study participants were older adults aged 65–79 years with complete data on drug use and falls (N = 1,833). Odds ratio (OR) and 95% confidence intervals (95% CI) were derived from logistic regression models adjusting for potential confounders including socio-demographic characteristics, health-related behaviors (alcohol drinking), body mass index and health conditions (frailty, vision impairment, disability, polypharmacy, blood pressure) as well as use of potential falls-risk-increasing drugs. SPSS complex sample methods were used for statistical analysis.

Results

Compared to people without falls, people with falls (n = 370) had a higher psychotropic drug use (33.1% vs. 20.7%, $p < .001$). After adjusting for potential confounders, use of psychotropic drugs overall was associated with a higher risk of falls (OR 1.64, 95% CI 1.14–2.37). This was particularly true for the use of synthetic psychotropic drugs (1.57, 1.08–2.28), antidepressants overall (2.88, 1.63–5.09) or synthetic antidepressants (2.66, 1.50–4.73), specifically, selective serotonin reuptake inhibitors (SSRIs) (6.22, 2.28–17.0). Similar results were found for recurrent falls.

RKI's 'Health Monitoring' Research Data Centre. Requests should be submitted to the 'Health Monitoring' Research Data Centre, Robert Koch Institute, Berlin, Germany (e-mail: fdz@rki.de).

Funding: This work was supported by the Federal Ministry of Health (<https://www.bundesgesundheitsministerium.de/>) and the Robert Koch Institute (www.rki.de). The German Health Interview and Examination Survey for Adults 2008-2011 was funded by the Federal Ministry of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Conclusions

Use of psychotropic drugs overall, especially synthetic antidepressants like SSRIs, is associated with higher risks of falls and recurrent falls among community dwelling older adults aged 65–79 years in Germany.

Introduction

As one of the major health problems, falls occur commonly and frequently among older adults with one third of adults aged ≥ 65 years falling at least once in a given year [1, 2]. Major consequences of falls for individuals include physical injuries and fractures leading to functional decline, disability and decreased quality of life. To society, falls impose high financial burdens and healthcare costs due to fall-caused hospitalization and mortality [3, 4]. In the EU approximately 2.3 million and in the US 2.8 million fall-related injuries are admitted to emergencies while 36,000 adults in the EU and 27,000 older adults in the US are reported to have died from falls each year [2, 5]. The health care expenditure for treating fall-related injuries is estimated to be €25 billion in the EU [5] and \$31 billion in the US [6]. In addition, many older adults are afraid of falling, which may result in psychological consequences such as mental stress, depression or anxiety [7].

Falls among older adults are largely preventable by identifying and controlling particularly modifiable risk factors [4, 8, 9]. Use of psychotropic drugs has been identified as an independent risk factor for falls in various studies including systematic reviews and meta-analyses [10–14]. Yet, most of previous studies on psychotropic drug use and falls considered only some of the important health conditions associated with falls such as vision impairment [15, 16], frailty [17], polypharmacy [18], use of potential fall risk-increasing drugs [12, 19] and disability [20]. Results of these studies may be confounded by unmeasured factors. So far, few studies investigating the association between falls and psychotropic drug use have controlled for these factors.

Fall-related injuries among older adults increase along with an aging population [21, 22]. Germany is currently the second oldest population in the world, with 20.9% of the population aged 65 years or over ($n = 16.9$ million) [23]. About 40% of women and 30% of men aged 65–90 years in Germany report any falls in the past 12 months [24]. Every one in five German adults aged 60–79 years used at least one psychotropic drug in the last 7 days [25]. Since psychotropic drug use is potentially a modifiable factor, further exploring the association of the use of psychotropic drugs, particularly specific subgroups of interest, with falls may provide insight into the prevention strategies of falls among older adults.

Population-based epidemiological studies on the association between psychotropic drug use and falls are lacking in Germany. Based on data of the most recent German Health Interview and Examination Survey for Adults (DEGS1) conducted in 2008–2011, we investigate the use of overall psychotropic drugs, major subgroups of psychotropic drugs as well as specific drugs of interest in relation to any falls and repeated falls after controlling for important health conditions and other factors that are likely to be associated with falls.

Methods

Data source: German Health Interview and Examination Survey for Adults 2008–2011 (DEGS1)

The German Health Interview and Examination Survey for Adults, wave 1 (DEGS1) was carried out by the Robert Koch Institute from November 2008 to December 2011 with the aim to provide representative data on the health of adults aged 18–79 years living in Germany [26, 27]. Details of study design, sampling strategy and protocol have been published previously [26, 27]. In Brief, DEGS1 used a two-stage random sampling procedure. First, 180 representative communities (primary sample units, PSU) were randomly sampled from a complete list of German communities proportional to community size. Second, age and sex stratified random samples were drawn within PSUs from local population registries. DEGS1 had a complex design. On one side, DEGS1 invited all surviving participants of the German National Health Interview and Examination Survey 1998 (GNHIES98) in order to establish a survey panel component. On the other side, DEGS1 also recruited new participants in order to maintain a nationally representative sample at the population level [26]. The response rate was 64% for 28–79 year-old re-participants from GNHIES98 ($n = 3795$) and 42% for newly recruited 18–79 year-old participants ($n = 4,192$) [26, 27]. Of 3,795 re-participants, 2,923 persons, together with all newly recruited ($n = 4,192$), completed both the interview and examination survey parts, amounting to a nationally population-representative sample of adults aged 18–79 years ($N = 7,115$) [26, 27].

DEGS1 was approved by the Federal and State Commissioners for Data Protection and the Charité-Universitätsmedizin Berlin ethics committee (No. EA2/047/08). Survey participants provided written informed consent prior to interviews and examinations.

Data collection, study population and definition of study variables

Data collection in DEGS1 included self-administered questionnaires, standardized health professional administered computer-assisted personal interviews (CAPI), physiological measurements and laboratory tests [26].

Primary outcome: Falls and recurrent falls. In DEGS1, data of falls were collected only among participants 65 years and older. We asked: “*Within the past 12 months, have you fallen, stumbled or slipped, so that you became unbalanced and, as a result, rested on the floor or lower surface?*” Those who answered with ‘Yes’ were defined as people with falls. Among people with falls, we further asked how many times the falls occurred. Those who answered 2 times or more were defined as people with recurrent falls [24].

Primary independent variable: Psychotropic drug use. As part of CAPI, detailed information on drug use (such as daily doses, routes of application and duration of use) in the past 7 days prior to the medical interview was recorded by trained health professionals [28]. In the invitation letter, participants were asked to bring the original packages of all medications used during the past seven days—prescribed and Over-The-Counter (OTC) products—to the examination site for the purpose of documentation and verification. This permitted automated recording of unique product identifiers and drug coding according to the WHO “Anatomical Therapeutic Chemical” (ATC) classification system [28]. Definition of psychotropic drugs has been published previously [25, 29]. Briefly, we included drugs belonging to the nervous system class (ATC code N00) as well as opiates used as antitussives (ATC code R05DA) and aspirin-caffeine combination preparations (ATC code N02BA71) [25, 29] (Appendix Table 1a). We excluded other analgesics and antipyretics such as aspirin and paracetamol (ATC code N02B), local anesthetics (ATC code N01B), homeopathic drugs, and drugs with indistinct active

Table 1. Descriptive characteristics of study population. German national health interview and examination survey 2008–2011 (DEGS1).

		Total (N = 1833)			People with falls (n = 370)			People without falls (n = 1463)			p			
		%*	95% CI*		n	%*	95% CI*		n	%*		95% CI*		n
Sex	<i>Men</i>	46.0	43.5	48.4	911	34.2	28.6	40.3	142	49.0	46.1	52.0	769	.000
	<i>Women</i>	54.0	51.6	56.5	922	65.8	59.7	71.4	228	51.0	48.0	53.9	694	
Age group	<i>65–69</i>	34.8	32.3	37.5	736	31.4	25.9	37.4	139	35.7	32.9	38.7	597	.212
	<i>70–74</i>	42.8	39.9	45.9	744	42.4	35.8	49.2	146	43.0	39.7	46.3	598	
	<i>75–79</i>	22.3	19.8	25.1	353	26.2	21.0	32.3	85	21.3	18.6	24.3	268	
Living alone	<i>Yes</i>	23.0	20.4	25.9	395	32.0	25.5	39.3	98	20.7	18.1	23.5	297	.001
	<i>No</i>	77.0	74.1	79.6	1434	68.0	60.7	74.5	271	79.3	76.5	81.9	1163	
Community size	<i>Rural towns</i>	15.6	10.4	22.6	301	14.3	8.7	22.6	55	15.9	10.7	23.0	246	.404
	<i>Small cities</i>	26.4	19.6	34.5	430	24.4	17.1	33.6	73	26.9	19.9	35.3	357	
	<i>Middle cities</i>	27.8	21.2	35.5	546	27.0	19.6	36.0	113	28.0	21.2	36.0	433	
	<i>Large cities</i>	30.3	23.3	38.3	556	34.3	25.9	43.7	129	29.2	22.2	37.4	427	
Region of residence	<i>Eastern</i>	23.0	17.3	30.0	621	23.1	16.4	31.6	119	23.0	17.2	30.1	502	.138
	<i>Northern</i>	15.8	10.4	23.3	236	14.3	8.9	22.3	47	16.2	10.6	23.9	189	
	<i>Central</i>	34.5	27.1	42.8	567	39.7	30.8	49.4	123	33.2	25.8	41.5	444	
	<i>Southern</i>	26.7	19.9	34.7	409	22.8	15.8	31.8	81	27.7	20.5	36.2	328	
Social status	<i>Lower</i>	26.0	22.6	29.8	345	24.4	18.1	32.0	66	26.5	22.8	30.5	279	.767
	<i>Middle</i>	59.4	55.8	62.9	1105	61.4	54.2	68.1	221	58.9	54.7	62.9	884	
	<i>Upper</i>	14.6	12.5	16.9	376	14.2	10.7	18.7	80	14.7	12.4	17.3	296	
Body mass index, kg/m²	<i><25</i>	20.3	18.2	22.6	400	20.0	15.3	25.8	83	20.4	17.9	23.0	317	.215
	<i>25–30</i>	44.0	41.2	46.9	823	39.5	33.6	45.8	146	45.2	41.9	48.5	677	
	<i>> = 30</i>	35.7	32.7	38.8	592	40.5	34.2	47.1	139	34.4	31.0	38.0	453	
Recognized disability	<i>Yes</i>	28.3	25.2	31.6	463	40.6	33.9	47.7	127	25.0	21.8	28.6	336	.000
	<i>No</i>	71.7	68.4	74.8	1322	59.4	52.3	66.1	238	75.0	71.4	78.2	1084	
Polypharmacy	<i>Yes</i>	35.0	32.4	37.7	641	39.2	33.2	45.5	152	33.9	31.0	36.9	489	.115
	<i>No</i>	65.0	62.3	67.6	1192	60.8	54.5	66.8	218	66.1	63.1	69.0	974	
Daily drinking	<i>Yes</i>	18.1	15.7	20.8	348	14.9	10.6	20.4	63	19.0	16.3	22.0	285	.163
	<i>No</i>	81.9	79.2	84.3	1381	85.1	79.6	89.4	287	81.0	78.0	83.7	1094	
Vision impairment	<i>No</i>	77.9	75.2	80.5	1393	75.2	68.4	81.0	269	78.6	75.3	81.6	1124	.179
	<i>Slight</i>	18.2	15.8	20.9	290	18.5	13.5	24.8	61	18.1	15.4	21.2	229	
	<i>Severe</i>	3.9	2.8	5.3	66	6.3	3.4	11.2	19	3.3	2.2	4.7	47	
Frailty	<i>No</i>	58.6	55.7	61.5	1129	52.8	45.8	59.6	201	60.2	56.7	63.5	928	.019
	<i>Pre-Frailty</i>	38.9	36.0	41.9	649	42.5	35.7	49.6	149	38.0	34.7	41.4	500	
	<i>Frailty</i>	2.5	1.7	3.5	43	4.7	2.7	8.1	17	1.9	1.2	3.0	26	
Continuous variables		Mean*	95% CI*		n	Mean*	95% CI*		n	Mean*	95% CI*		n	
	<i>SBP, mm Hg</i>	130.3	129.2	131.5	1828	128.1	125.9	130.3	367	130.9	129.7	132.1	1461	.015
	<i>DBP, mm Hg</i>	72.8	72.2	73.5	1828	71.2	70.0	72.4	367	73.3	72.5	74.0	1461	.003

*Weighted and standardized to the population of 31.12.2010. P values: Comparison between people with falls and without falls.

Community size: Rural towns (<5,000 inhabitants), small cities (5,000-<20,000 inhabitants), middle-sized cities (20,000-<100,000 inhabitants) and large cities (100,000 inhabitants or more)

Region of residence: Eastern Germany (Berlin, Brandenburg, Mecklenburg-Vorpommern, Sachsen, Sachsen-Anhalt and Thüringen); Northern Germany (federal states: Bremen, Hamburg, Niedersachsen and Schleswig-Holstein); Central Germany (Hessen, Nordrhein-Westfalen, Rheinland-Pfalz and Saarland); Southern Germany (Baden-Württemberg and Bayern).

Missing values: Living alone (n = 4), social status (n = 7), recognized disability (n = 48), vision impairment (n = 84), daily drinking (n = 104), frailty (n = 12).

<https://doi.org/10.1371/journal.pone.0182432.t001>

ingredients under the ATC class N00 [25, 29]. Psychotropic drugs with herbal active ingredients were considered and coded separately under specific subgroups (S1 Table).

For the present analyses, individuals aged 65–79 years with complete data on falls and psychotropic drug use were included as the study population (N = 1,833) (Table 1).

Covariables. A number of covariables that are likely to be associated with falls were investigated in this study. This included socio-demographic characteristics (age, sex, community size, region of residence, individual information on household size—namely number of persons living in a household—, income, profession and educational attainment), health-related behaviors, and dietary habits such as alcohol consumption. Information on socio-demographic characteristics, health-related behavior and dietary habits were collected by self-administered questionnaires. Community size was classified as rural towns, small cities, middle-sized cities and large cities based on population density according to an established German classification system. Regions of residence were grouped into 4 commonly described geographical areas: Northern Germany (federal states: *Bremen, Hamburg, Niedersachsen and Schleswig-Holstein*); Central Germany (*Hessen, Nordrhein-Westfalen, Rheinland-Pfalz and Saarland*); Southern Germany (*Baden-Württemberg and Bayern*); Eastern Germany (*Berlin, Brandenburg, Mecklenburg-Vorpommern, Sachsen, Sachsen-Anhalt and Thüringen*). According to the number of persons living in a household, living alone was defined if only one single person was reported in a household. Socio-economic status (SES) was classified as ‘lower’, ‘middle’ and ‘upper’ using an established index including information on education, professional status and household income [30].

Daily alcohol drinking was assumed if survey participants reported drinking any alcohol-containing beverages at least once a day in the past 12 months [25]. Body mass index (BMI) was calculated as the ratio of body weight (kg) and body height (m) squared and categorized into $<25 \text{ kg/m}^2$, $25\text{--}29.9 \text{ kg/m}^2$ and $\geq 30 \text{ kg/m}^2$. Several health conditions were considered in the present analysis based on findings of previous studies: Frailty [17], vision impairment [15, 16], disability [20], polypharmacy [14, 18, 31] and systolic blood pressure (SBP) [32]. Frailty was defined as having 3 and more of the following criteria: Exhaustion, low weight, low physical activity, low walking speed and low grip strength while pre-frailty was defined as having 1–2 of these components [33]. Survey participants were asked if they had any difficulties either in reading printed newspaper or in identifying the face of a person 4 meters away, using glasses or any other reading aid if necessary. Possible answer choices for the question were: (1) can read without difficulty, (2) can read with some difficulty, (3) can read with great difficulty and (4) unable to read at all. Participants in categories ‘3’ or ‘4’ were defined as experiencing ‘severe vision impairment’ and ‘2’ as experiencing ‘some vision impairment’. People were also asked if they had an officially certified disability (yes/no). Further, polypharmacy was assumed if five or more prescription medicines were used in the past 7 days. Three standardized blood pressure (BP) measurements were taken at three minutes intervals in upright sitting with an oscillometric device (Datascope Accutorr Plus). The mean of the second and third measures for systolic BP (SBP) and diastolic BP (DBP) was adopted [34].

A number of drugs have been reported to have the potential to increase the risks of falls (potential fall risk-increasing drugs) [19, 35]. Based on drug use data collected in DEGS1, we considered the major groups of potential falls risk-increasing drugs, namely antihypertensive medications (ATC code C02 antihypertensive, C03 diuretics, C07 beta-adrenergic antagonists, C08 calcium channel blockers and C09 agents acting on the renin-angiotensin system) [36], anti-diabetes medications (ATC code: A10) [19], nonsteroidal anti-inflammatory drugs (NSAIDs) (M01A, M02AA and N02BA) such as aspirin, ibuprofen and naproxen etc. [12, 37], statins (C10AA & C10BA) [38] and thyroid therapy medications (H03) [19].

Statistical analyses

All statistical analyses were performed using SPSS statistical software (version 20.0, SPSS Inc. Chicago, IL). In order to adjust for sample clustering effects, the SPSS complex samples module was used. Sampling weights were used to correct deviations in the sample from the structure of the German general population of the 31st December 2010 [27].

Participants' characteristics were summarized by people with falls and without falls in the past 12 months. Differences between people with and without falls were examined using second-order Rao-Scott chi-square tests for categorical variables and using General Linear Models (CSGLM) for continuous variables.

We fitted several logistic models with falls or recurrent fall events in the past 12 months as dependent variables and use of psychotropic drugs as the primary independent variable adjusting risk factors of falls. Model 1 was adjusted for sex and age group. Model 2 was additionally adjusted for social status, community size, region of residence and living alone. Based on model 2, model 3 was further adjusted for health behaviors including daily alcohol drinking, BMI and health conditions including frailty, polypharmacy, disability, vision impairment and SBP. Based on model 3, model 4 was further adjusted for the use of potential falls-risk-increasing drugs including antihypertensives, antidiabetic medications, NSAIDs, statins and thyroid therapy medications. For each of these models, we first looked at the use of overall psychotropic drugs, all synthetic psychotropic drugs and phyto-medicines, and then the use of major subgroups of psychotropic drugs as well as specific drugs or drug classes of interest.

A total of 246 or 13.4% of study participants had missing observations in some variables with a range of 0.2% for living alone, 2.6% for recognized disability, 4.6% for vision impairment, to 5.7% for daily drinking. The numbers of persons with missing values were explicitly stated for each variable (Table 1). Persons with missing values were excluded from the analyses, with pairwise deletion for descriptive and listwise deletion for multivariable analyses (i.e. multivariable models were restricted to participants with valid data on all model covariables). Statistical significance was defined at $p < 0.05$ based on two-sided tests.

Results

Table 1 shows the descriptive characteristics of the study population stratified by people with and without falls in the past 12 months. Of 1,833 older adults, approximately 20% lived alone or drank daily; more than one third had a BMI ≥ 30 kg/m² or used polypharmacy; about 40% had frailty or pre-frailty and 22% had a severe or slight vision impairment. 370 older adults reported any falls with a weighted prevalence of 20.7% (95% CI 18.7–23.0%). Of 370 people with falls, 149 (40.3%) reported repeated falls (data not shown in Table 1). Compared to people without falls, people with falls were more likely to be female, to live alone, and to have a recognized disability and a lower average SBP/DBP (Table 1). In addition, they had a higher proportion of frailty and pre-frailty (Table 1). No difference was found between people with and without falls in regard to the distribution of age group, community size, region of residence, social status, BMI, using polypharmacy, daily drinking and vision impairment (Table 1).

Table 2 shows the prevalence for the use of psychotropic drugs as well as potential falls risk-increasing drugs among people with and without falls. People with falls had overall a higher psychotropic drug use compared to people without falls (33.1% vs. 20.7%, $p < .001$). The same is true for the use of any synthetics (25.1% vs. 16.2%) and phyto-medicines (12.1% vs. 6.3%) as well as the subgroups of anti-depressants, -both synthetic antidepressants (Non-Selective Monoamino Reuptake Inhibitors, NSMRIs and Selective Serotonin Reuptake Inhibitors, SSRIs) and St. John's wort-, and antimentia drugs (mainly Ginkgo biloba) (Table 2). No

Table 2. Prevalence of psychotropic drug use and potential falls-risk-increasing drug use among older adults with and without falls. German national health interview and examination survey 2008–2011 (DEGS1).

	People with falls (n = 370)			People without falls (n = 1463)					
	%	95%CI	n	%	95%CI	n	p		
Psychotropic drugs									
Any psychotropic drugs (Synthetics & phytomedicines**)	33.1	27.4	39.2	113	20.7	18.0	23.7	300	.000
Any synthetic psychotropic drugs	25.1	19.9	31.3	89	16.2	13.6	19.1	231	.001
Any phytomedicines	12.1	7.8	18.1	33	6.3	4.9	8.0	93	.016
Any anti-depressants (St. John's wort and synthetical antidepressants)	15.3	10.8	21.2	46	6.1	4.6	8.1	77	.000
<i>St. John's wort</i> (N05CP03/N06AP01/51)	2.3	.8	6.1	5	.5	.2	1.0	9	.010
Any synthetical antidepressants	13.0	8.8	18.6	41	5.8	4.3	7.7	71	.001
NSMRIs (N06AA)	7.0	3.8	12.6	21	3.1	2.1	4.8	36	.019
SSRIs (N06AB)	4.3	2.4	7.6	14	1.3	.8	2.2	19	.002
Any hypnotics & sedatives (synth., antihistamine and phytoceuticals)	4.3	2.4	7.5	19	3.7	2.7	5.1	62	.660
Any synthetics. (N05C) and antihistamine (N05CM)	2.4	1.1	5.1	11	1.7	1.1	2.6	33	.493
Any synthetics (N05C)	2.4	1.1	5.1	11	1.3	.8	2.1	25	.207
Valerian (N05CP01/51)	1.8	.7	4.4	7	1.6	1.0	2.6	27	.887
Any benzodiazepines (N05BA, N05CD, N03AE01)	4.0	1.8	8.6	12	2.9	1.9	4.4	37	.485
Benzodiazepine-related drugs (N05CF)	5.2	2.7	9.5	20	3.7	2.6	5.4	52	.366
Narcotic analgesics (N02A)	6.6	4.1	10.5	23	4.1	3.0	5.6	61	.087
Any anti-dementia drugs (N06DA and Ginkgo biloba)	8.3	4.7	14.2	24	4.2	3.1	5.7	59	.046
Ginkgo biloba (N06DP01)	8.0	4.4	14.0	21	3.9	2.9	5.3	55	.037
Anti-epileptics (N03)	2.6	1.3	5.0	13	1.9	1.0	3.3	30	.448
Antiparkinson drugs (N04)	2.5	1.1	5.5	9	1.4	.7	2.6	21	.203
Potential falls-risk-increasing drugs									
Any antihypertensives (ATC-Code C02, C03, C07, C08, C09)	65.5	59.1	71.4	246	68.9	65.8	71.8	984	.307
Anti-diabetes medications (A10)	17.0	12.6	22.5	58	13.6	11.3	16.3	196	.193
NSAIDs (M01A, M02AA and N02BA)	29.1	23.9	34.9	107	19.0	16.1	22.2	260	.000
Statins (C10AA & C10BA)	25.4	20.6	30.8	106	27.0	24.3	29.8	399	.596
Thyroid therapy (H02)	24.2	18.6	30.9	86	17.6	14.9	20.6	253	.031

* Weighted and standardized to the population of 31.12.2010.

P-values: Comparison between people with falls and people without falls

**Phytomedicines: *St. John's wort* (N06AP), Ginkgo biloba (N06DP01), *Valerian* (N05CP).

Figures in bold denote statistical significance (p < .050)

NSMRIs: Non-Selective Monoamino Reuptake Inhibitors

SSRIs: Selective Serotonin Reuptake Inhibitors

NSAIDs: Nonsteroidal Anti-Inflammatory Drugs (M01A, M02AA and N02BA)

<https://doi.org/10.1371/journal.pone.0182432.t002>

difference was found between people with and without falls concerning the use of antihypertensives, anti-diabetes medications and statins. Yet, people with falls had a higher use for NSAIDs and thyroid therapy medicines (Table 2).

In logistic regression models adjusting for age and sex, use of overall psychotropic drugs and of synthetic psychotropic drugs was associated with falls, but this did not apply to the use of phytomedicines (Table 3). Further adjusting for confounders included in model 2, model 3 and model 4, the association remains robust (Table 3). The risk of any falls was 64% higher among psychotropic drug users compared to nonusers. In terms of major subgroups of psychotropic drugs, use of antidepressants overall increased the risk of falls significantly after considering all potential confounders (OR 2.88, 95% CI 1.63–5.09). Use of synthetic

Table 3. Association of psychotropic drug use with risks of falls. German national health interview and examination survey 2008–2011 (DEGS1).

	Model 1			Model 2			Model 3			Model 4		
	OR	95%CI		OR	95%CI		OR	95%CI		OR	95%CI	
Any psychotropic drugs (Synthetics & phytomedicines)	1.69	1.24	2.31	1.68	1.23	2.29	1.64	1.14	2.35	1.64	1.14	2.37
Any synthetic psychotropic drugs	1.58	1.12	2.24	1.61	1.14	2.29	1.52	1.05	2.21	1.57	1.08	2.28
Any phytomedicines	1.76	0.98	3.19	1.62	0.91	2.89	1.66	0.84	3.29	1.62	0.82	3.19
Any anti-depressants (St. John's wort and synthetic antidepressants)	2.39	1.43	4.01	2.33	1.39	3.90	2.69	1.50	4.83	2.88	1.63	5.09
<i>St. John's wort</i> (N05CP03/N06AP01/51)	4.36	1.30	14.6	4.66	1.21	18.0	4.22	0.62	28.9	3.71	0.64	21.6
Any synthetic antidepressants	2.10	1.21	3.62	2.01	1.16	3.47	2.43	1.36	4.34	2.66	1.50	4.73
NSMRIs (N06AA)	1.95	0.89	4.24	1.72	0.80	3.68	1.75	0.77	4.01	1.84	0.83	4.10
SSRIs (N06AB)	2.97	1.31	6.74	3.18	1.34	7.56	5.46	1.96	15.2	6.22	2.28	17.0
Any hypnotics & sedatives (synth., antihistamine and phytoceuticals)	0.99	0.49	2.00	1.05	0.52	2.13	0.88	0.42	1.83	0.79	0.37	1.71
Any synth. (N05C) and antihistamine (N05CM)	1.27	0.53	3.03	1.30	0.53	3.18	0.67	0.28	1.61	0.58	0.22	1.49
Any benzodiazepines (N05BA, N05CD, N03AE01)	1.20	0.47	3.07	1.22	0.48	3.14	0.90	0.24	3.39	0.90	0.25	3.28
Benzodiazepines and benzodiazepine-related drugs (N05CF)	1.22	0.57	2.60	1.24	0.58	2.66	0.89	0.32	2.44	0.85	0.31	2.32
Narcotic analgesics (N02A)	1.52	0.84	2.72	1.58	0.87	2.88	1.39	0.68	2.84	1.48	0.70	3.12
Any anti-dementia drugs (N06DA+ Ginkgo biloba)	1.81	0.90	3.68	1.53	0.78	3.03	1.60	0.71	3.64	1.57	0.69	3.60
Ginkgo biloba (N06DP01)	1.87	0.91	3.85	1.56	0.77	3.14	1.64	0.71	3.77	1.61	0.69	3.74
Anti-epileptics (N03)	1.58	0.62	3.98	1.76	0.75	4.17	1.36	0.51	3.61	1.47	0.58	3.73
Antiparkinson drugs (N04)	1.76	0.70	4.40	1.80	0.70	4.65	1.12	0.31	4.08	1.08	0.26	4.55
Valerian (N05CP01/51)	0.73	0.26	2.05	0.79	0.28	2.21	0.96	0.33	2.77	0.90	0.31	2.61

Odds ratio (OR) and 95% confidence intervals (95% CI) were derived from logistic regression models.

Model 1: adjusted for sex and age group (65–69, 70–74, 75–79 years)

Model 2: model 1+social status (lower, middle, upper), community size (rural area, small, middle and large city), region (East, north, central, south), living alone (yes, no),

Model 3: Model 2+ frailty (frailty, pre-frailty, no frailty), polypharmacy (yes, no), disability (yes, no), vision impairment (severe, slight, no), daily alcohol drinking (yes, no), BMI <25, 25–30, > = 30), SBP (continuous variables)

Model 4: Model 3+antihypertensives, antidiabetics, nonsteroidal anti-inflammatory drugs (NSAIDs), statins, thyroid therapy (yes, no)

NSMRIs: non-selective monoamine reuptake inhibitors

SSRIs: Selective serotonin reuptake inhibitors

Figures in bold denote statistical significance (p < .050)

<https://doi.org/10.1371/journal.pone.0182432.t003>

antidepressants (2.66, 1.50–4.73), but not St. John's wort was associated with any falls. Specifically, use of SSRIs (6.22, 2.28–17.0), but not non-selective monoamine reuptake inhibitors (NSMRIs) increased the risk of any falls (Table 3). For the use of other psychotropic drugs, no significantly increased risks were found (Table 3).

As with falls, use of overall psychotropic drugs was also associated with recurrent falls (adjusted OR 1.84, 95% CI 1.02–3.31) (Table 4). The odds ratios for the relationship between recurrent falls and the use of overall antidepressants, synthetic antidepressants, such as SSRIs, were consistently significant (p<0.05) in all 4 models. The odds ratios for narcotic analgesics (N02A) and anti-Parkinson drugs (N04) were found to be significant in model 1 and model 2, but not in model 3 and model 4 (Table 4).

Sensitivity analyses involved limiting to those who had used any psychotropic drugs over 12 months and results were similar. In the fully adjusted Model 4, the OR for any psychotropic drug use over 12 months was 1.70 (95% CI 1.10–2.63) for any falls and 2.10 (95% CI 1.04–4.22) for recurrent falls; the OR for any synthetic drug use over 12 months was 1.89 (95% CI 1.21–2.94) for any falls and 1.92 (95% CI 1.11–3.34) for recurrent falls (data not shown in Tables 3 and 4).

Table 4. Association of psychotropic drug use and risk of recurrent falls. German national health interview and examination survey 2008–2011 (DEGS1).

	Model 1			Model 2			Model 3			Model 4		
	OR	95%CI		OR	95%CI		OR	95%CI		OR	95%CI	
Any psychotropic drugs (Synthetics & phytomedicines)	1.97	1.27	3.05	1.96	1.25	3.08	1.83	0.99	3.35	1.84	1.02	3.31
Any synthetics	1.65	1.04	2.64	1.70	1.05	2.75	1.30	0.77	2.20	1.38	0.83	2.29
Any phytomedicines	1.60	0.74	3.44	1.44	0.64	3.24	1.76	0.58	5.35	1.69	0.57	5.00
Any anti-depressants (St. John's wort and all synthetic antidepressants)	2.09	1.14	3.82	2.08	1.10	3.95	2.74	1.36	5.51	3.15	1.60	6.23
<i>St. John's wort</i> (N05CP03/N06AP01/51)	1.10	0.22	5.58	1.19	0.20	7.16	3.30	0.46	23.7	2.76	0.30	25.3
Any synthetic antidepressants	2.16	1.15	4.07	2.12	1.08	4.15	2.53	1.22	5.27	3.01	1.48	6.09
NSMRIs (N06AA)	1.28	0.51	3.23	1.11	0.40	3.06	0.95	0.35	2.57	1.11	0.42	2.92
SSRIs (N06AB)	3.10	1.25	7.69	3.55	1.42	8.85	6.74	2.47	18.4	7.02	2.38	20.7
Any hypnotics & sedatives (synth., antihistamine and phytoceuticals)	0.74	0.27	2.08	0.80	0.29	2.22	0.89	0.28	2.88	0.72	0.22	2.40
Any synth. (N05C) and antihistamine (N05CM)	0.73	0.20	2.65	0.81	0.21	3.08	0.56	0.13	2.51	0.45	0.10	2.08
Benzodiazepines (N05BA, N05CD, N03AE01)	1.23	0.40	3.78	1.28	0.42	3.92	0.93	0.16	5.22	1.01	0.17	5.88
Benzodiazepines and benzodiazepine-related drugs	1.31	0.52	3.26	1.41	0.57	3.49	0.92	0.24	3.48	0.88	0.22	3.56
Narcotic analgesics (N02A)	2.64	1.30	5.38	2.66	1.30	5.43	2.21	0.97	5.05	2.27	0.97	5.34
Any anti-dementia drugs (N06DA+ Ginkgo biloba)	2.21	0.95	5.11	1.85	0.73	4.64	1.91	0.50	7.26	1.92	0.54	6.79
Ginkgo biloba (N06DP01)	2.15	0.89	5.20	1.78	0.67	4.76	1.85	0.45	7.56	1.90	0.51	7.09
Anti-epileptics (N03)	0.74	0.21	2.59	0.85	0.24	2.94	0.47	0.11	1.99	0.42	0.10	1.81
Antiparkinson drugs (N04)	3.51	1.18	10.4	3.80	1.29	11.2	1.44	0.34	6.02	1.43	0.37	5.56
Valerian (N05CP01/51)	0.71	0.16	3.23	0.76	0.17	3.35	1.10	0.21	5.78	0.89	0.16	4.80

Odds ratio (OR) and 95% confidence intervals (95% CI) were derived from logistic regression models.

Model 1: adjusted for sex and age group (65–69, 70–74, 75–79 years)

Model 2: model 1+social status (lower, middle, upper), community size (rural area, small, middle and large city), region (East, north, central, south), living alone (yes, no),

Model 3: Model 2+ frailty (frailty, pre-frailty, no frailty), polypharmacy (yes, no), disability (yes, no), vision impairment (severe, slight, no), daily alcohol drinking (yes, no), BMI <25, 25–30, > = 30), SBP (continuous variables)

Model 4: Model 3+antihypertensives, antidiabetics, nonsteroidal anti-inflammatory drugs (NSAIDs), statins, thyroid therapy (yes, no)

NSMRIs: non selective monoamino reuptake inhibitors

SSRIs: Selective serotonin reuptake inhibitors

Figures in bold denote statistical significance ($p < .050$).

<https://doi.org/10.1371/journal.pone.0182432.t004>

Discussion

Main findings

In the present study, we found that the use of psychotropic drugs overall was associated with increased risks of falls and recurrent falls in the past 12 months among community-dwelling adults aged 65–79 years in Germany. This is particularly true for the use of synthetic psychotropic drugs, mainly synthetic antidepressants, -specifically SSRIs. Our study confirms the association between psychotropic drug use and falls and adds evidence for an increased risk of falls following the use of psychotropic drugs overall and specific subgroups among community-dwelling older adults.

Comparison with other studies

The association between psychotropic drug use and falls has been investigated in previous studies and systematically reviewed or meta-analyzed [10–14]. An earlier meta-analysis covering studies published between 1966–1996 examining the association of drug use with falling in

people aged 60 and older found a pooled odds ratio of 1.73 (95%CI, 1.52–1.97) for any psychotropic use [10]. This finding was confirmed by another meta-analysis including 71 studies published between 1981 and 2007, which reported an odds ratio of 1.78 (1.57–2.01) [13]. Both are well comparable to our finding of an odds ratio of 1.64 (1.14–2.37) for the association between psychotropic drug use and falls among older adults. A recent systematic review of prospective and retrospective studies published between 2008 and 2013 on the association of medication use and falls in older people found that 29 of 36 studies reported a positive association between the risk of falls and use of psychotropic medications including sedatives and hypnotics, antidepressants, and benzodiazepines [14]. Further, a large body of evidence from prospective studies and clinical trials suggests that reduced use of psychotropic medications resulted in reduction of falls [39], strongly supporting the associations between psychotropic drug use and falls.

Our findings on the association between antidepressants use and falls or recurrent falls are also in line with results of other studies. Use of antidepressants, particularly SSRIs, has been consistently found to be associated with increased risks of falls and recurrent falls [40–43]. For example, in a longitudinal analysis of 2948 community-dwelling older adults followed-up for 7 years, antidepressant users, compared with nonusers, were observed to have a 48% greater likelihood of recurrent falls (OR 1.48, 1.12–1.96), particularly those taking SSRIs with an OR of 1.62 (1.15–2.28) [40]. Alike, in a cross-sectional survey including community-dwelling older adults from Australian general practices, antidepressants use was independently associated with multiple falls (OR 1.46, 1.25–1.70). Amongst all psychotropic medications, SSRI use was found to be associated with the highest risk of multiple falls (OR 1.66, 1.36–2.02) [44]. Using data of the Swedish registers with a sample size of more than 1 million persons aged ≥ 65 years, Johnell and colleagues found that among psychotropic drugs, antidepressants had the strongest association with fall-related injuries (adjusted OR 1.42, 1.38–1.45) [45]. These studies suggest that use of antidepressants could result in a 40–70% greater likelihood of falls [40, 44, 45]. In the present study, we found a much higher OR of 2.88 (1.63–5.09) for the association between use of antidepressants and falls, which, however, is comparable to the results of a prospective population-based study reporting an OR of 2.8 (1.9–4.1) among older men aged 60–75 years in Denmark [46]. Nevertheless due to small sample size in our study, the OR for SSRIs use (Table 3) is much higher than that (3.1, 2.0–5.0) found in the Danish study [46].

Interestingly, we found that the use of psychotropic phyto-medicines, either St. Johns' wort, valerian or Gingko biloba, was not associated with any falls or recurrent falls. This may be due to their weak effects compared to their synthetic counterparts. Surprisingly, we did not find an association of falls or recurrent falls with the use of hypnotics and sedatives, antiepileptics, or narcotic analgesics. However, some studies also found no significant association between falls and the use of anxiolytics/hypnotics [46], or a weaker association with the use of hypnotics and sedatives [45] or antiepileptics [47]. In contrast, other studies report positive associations between falls and the use hypnotics and sedatives [12, 14, 47, 48], opioids/narcotics [46, 48] [49] or antiepileptics [46]. There may be several reasons for the inconsistency between findings of our study and of others. First, the sample size for users of hypnotics and sedatives, antiepileptics and narcotic analgesics in our study is relatively small (for example, only 36 persons used synthetic hypnotics and sedatives, Table 2). Second, more than half of all benzodiazepines, hypnotics and sedatives were taken as needed while only a small part of them were taken regularly more than 3 months (appendix Table 1a). Third, evidence of an association between substantially increased risk of falls and use of benzodiazepines (particularly long-acting agents) tends to be found in earlier study. Finally, because of well-known 'after effects' of hypnotics and sedatives that may increase the risk of falls, older adults at high risk of falls may have been cautioned by physicians and thus avoided taking such kind of drugs. As a result, use

of anxiolytics was found to be even associated with a slightly reduced risk of fall (OR 0.97, 95% CI 0.94–1.00) [45]. In our study, although not statistically significant, the adjusted ORs for the use of synthetic hypnotics & sedatives, benzodiazepine and related drugs tend to be <1 (Table 3).

Strength and limitation

The main strength of our study is that we used a nationally representative sample of community-dwelling older adults. The weighted results can be generalized to the German older adults aged 65–79 years. We fitted several logistic regression models to explore the associations between psychotropic drug use and falls controlling for a number of confounding variables including frailty, vision impairment and use of other fall risk-increasing drugs.

Findings of our study are subject to several limitations. First, DEGS1 is not specifically designed for the study of association of psychotropic drug use and falls. Since other sources of linked health administrative data are unavailable, data of national health surveys are a good source that could be used for such kind of studies despite limitations. Persons aged ≥ 80 years and persons who were hospitalized or institutionalized in long-term care facilities were not included in our national health surveys. In addition, community-dwelling older adults with cognitive impairment, depression or other severe mental or physical illnesses (e.g. severe vision impairment and fall injuries) might be less likely to take part in the survey due to the need to travel to the examination sites. These persons might be at psychotropic drug use and high risk of falls. Second, data on falls and drug use were self-reported; recall bias seems unavoidable. In the invitation letter we asked all survey participants to bring the medication packages to examination sites allowing us to verify medication use, which could reduce recall bias greatly. Third, findings of our study might be weakened by the fact that we measured falls in the past 12 months whereas drug use in the last 7 days. Due to this concern, we therefore specifically examined use pattern and use durations of all psychotropic drugs. Results reveal that more than half of all psychotropic drugs have been used for 12 months or longer. This is particularly true for the use of major subgroups of psychotropic drugs such as synthetic antidepressants (72.4%) (S2 Table). Furthermore, sensitivity analyses involving drug use over 12 months demonstrate similar results, further supporting our findings. Fourth, the number of drug users in some subgroups was quite small, which prevented us from detecting weak associations. Fifth, we investigated if any specific psychotropic drug was used in relations to falls. Nevertheless, total daily drug dose exposure, changing doses, interactions between drugs, can all be involved in fall risks, which we could not investigate in the present analysis. Finally and most importantly, subject to observational design, our study did not allow a causality inference between falls and psychotropic drug use and could not avoid indication bias. Indication bias occurs when patients are prescribed drugs for a condition that itself is associated with the outcome of interest [50, 51]. In this analysis, antidepressants are indicated to treat depression, which itself, has been recognized as a key independent risk factor for falls [52–54]. Yet, there are also studies suggesting that antidepressant use (particularly SSRIs) is strongly associated with falls regardless of the presence of depressive symptoms [44].

Conclusions

In summary, we found that the use of psychotropic drugs overall, especially synthetic antidepressants like SSRIs, was associated with a higher risk of falls and recurrent falls among community dwelling older adults aged 65–79 years in Germany. Our study adds new evidence concerning the association of use of psychotropic drugs—mainly synthetic antidepressants like SSRIs— and falls among community-dwelling older adults. Given the severe consequence

of falls and the extensive use of psychotropic drugs among the elderly, a more rational use of such kind of drugs is needed. Due to limitations of our study with small sample sizes for some subgroups of drugs, further studies are required to investigate the associations between the use of other commonly used psychotropic drugs such as hypnotics and sedatives and falls.

Supporting information

S1 Table. List of psychotropic drugs used among people aged 65–79 years. German national health interview and examination survey 2008–2011 (DEGS1). (DOCX)

S2 Table. Use pattern of psychotropic drugs among people aged 65–79 years—by use duration. German national health interview and examination survey 2008–2011 (DEGS1). (DOCX)

Acknowledgments

We thank the survey participants for contributing towards our understanding of psychotropic health.

Author Contributions

Conceptualization: Yong Du, Hildtraud Knopf.

Data curation: Yong Du, Hildtraud Knopf.

Formal analysis: Yong Du.

Funding acquisition: Hildtraud Knopf.

Methodology: Yong Du.

Project administration: Hildtraud Knopf.

Supervision: Hildtraud Knopf.

Writing – original draft: Yong Du.

Writing – review & editing: Yong Du, Ingrid-Katharina Wolf, Hildtraud Knopf.

References

1. WHO Global Report on Falls Prevention in Older Age. http://www.who.int/ageing/publications/Falls_prevention7March.pdf
2. Bergen G, Stevens MR, Burns ER. Falls and fall injuries among adults aged ≥ 65 years—United States, 2014. *Morbidity and Mortality Weekly Report*. 2016; 65(37):938–83.
3. Ambrose AF, Paul G, Hausdorff JM. Risk factors for falls among older adults: A review of the literature. *Maturitas*. 2013; 75(1):51–61. <https://doi.org/10.1016/j.maturitas.2013.02.009> PMID: 23523272
4. Vieira ER, Palmer RC, Chaves PHM. Prevention of falls in older people living in the community. *BMJ (Online)*. 2016; 353. <https://doi.org/10.1136/bmj.i1419> PMID: 27125497
5. Turner S, Kissler R, Rogm W. Falls among older adults in the EU-28: Key facts from the available statistics: EuroSafe, Amsterdam 2015. https://eupha.org/repository/sections/ipsp/Factsheet_falls_in_older_adults_in_EU.pdf.
6. Burns ER, Stevens JA, Lee R. The direct costs of fatal and non-fatal falls among older adults—United States. *J Safety Res*. 2016; 58:99–103. <https://doi.org/10.1016/j.jsr.2016.05.001> PMID: 27620939
7. Scheffer AC, Schuurmans MJ, van Dijk N, van der Hoof T, de Rooij SE. Fear of falling: measurement strategy, prevalence, risk factors and consequences among older persons. *Age and Ageing*. 2008; 37(1):19–24. <https://doi.org/10.1093/ageing/afm169> PMID: 18194967

8. Huang AR, Mallet L, Rochefort CM, Eguale T, Buckeridge DL, Tamblyn R. Medication-related falls in the elderly: causative factors and preventive strategies. *Drugs Aging*. 2012; 29(5):359–76. <https://doi.org/10.2165/11599460-000000000-00000> PMID: 22550966
9. Stubbs B, Brefka S, Denking MD. What Works to Prevent Falls in Community-Dwelling Older Adults? Umbrella Review of Meta-analyses of Randomized Controlled Trials. *Phys Ther*. 2015; 95(8): 1095–110. <https://doi.org/10.2522/ptj.20140461> PMID: 25655877
10. Leipzig RM, Cumming RG, Tinetti ME. Drugs and falls in older people: A systematic review and meta-analysis: I. Psychotropic drugs. *Journal of the American Geriatrics Society*. 1999; 47(1):30–9. PMID: 9920227
11. Hartikainen S, Lonnroos E, Louhivuori K. Medication as a risk factor for falls: critical systematic review. *J Gerontol A Biol Sci Med Sci*. 2007; 62(10):1172–81. PMID: 17921433
12. Woolcott JC, Richardson KJ, Wiens MO, Patel B, Marin J, Khan KM, et al. Meta-analysis of the impact of 9 medication classes on falls in elderly persons. *Archives of Internal Medicine*. 2009; 169(21): 1952–60. <https://doi.org/10.1001/archinternmed.2009.357> PMID: 19933955
13. Bloch F, Thibaud M, Dugue B, Breque C, Rigaud AS, Kemoun G. Psychotropic drugs and falls in the elderly people: updated literature review and meta-analysis. *J Aging Health*. 2011; 23(2):329–46. <https://doi.org/10.1177/0898264310381277> PMID: 20947876
14. Park H, Satoh H, Miki A, Urushihara H, Sawada Y. Medications associated with falls in older people: Systematic review of publications from a recent 5-year period. *European Journal of Clinical Pharmacology*. 2015; 71(12):1429–40. <https://doi.org/10.1007/s00228-015-1955-3> PMID: 26407688
15. Dpa Crews JE, Chou CF, Stevens JA, Saaddine JB. Falls Among Persons Aged \geq 65 Years With and Without Severe Vision Impairment—United States, 2014. *MMWR Morb Mortal Wkly Rep*. 2016; 65(17): 433–7. <https://doi.org/10.15585/mmwr.mm6517a2> PMID: 27148832
16. Hong T, Mitchell P, Burlutsky G, Samarawickrama C, Wang JJ. Visual impairment and the incidence of falls and fractures among older people: Longitudinal findings from the blue mountains eye study. *Investigative Ophthalmology and Visual Science*. 2014; 55(11):7589–93. <https://doi.org/10.1167/iov.14-14262> PMID: 25370514
17. Schultz M, Rosted E, Sanders S. Frailty is associated with a history with more falls in elderly hospitalised patients. *Danish Medical Journal*. 2015; 62(6).
18. Zia A, Kamaruzzaman SB, Tan MP. Polypharmacy and falls in older people: Balancing evidence-based medicine against falls risk. *Postgraduate Medicine*. 2015; 127(3):330–7. <https://doi.org/10.1080/00325481.2014.996112> PMID: 25539567
19. Zia A, Kamaruzzaman SB, Tan MP. The consumption of two or more fall risk-increasing drugs rather than polypharmacy is associated with falls. *Geriatr Gerontol Int*. 2017; 17(3):463–70. <https://doi.org/10.1111/ggi.12741> PMID: 26822931
20. Matsuda PN, Verrall AM, Finlayson ML, Molton IR, Jensen MP. Falls among adults aging with disability. *Archives of Physical Medicine and Rehabilitation*. 2015; 96(3):464–71. <https://doi.org/10.1016/j.apmr.2014.09.034> PMID: 25449191
21. Houry D, Florence C, Baldwin G, Stevens J, McClure R. The CDC Injury Center's response to the growing public health problem of falls among older adults. *Am J Lifestyle Med*. 2016; 10(1). <https://doi.org/10.1177/1559827615600137> PMID: 26688674
22. Do MT, Chang VC, Kuran N, Thompson W. Fall-related injuries among Canadian seniors, 2005–2013: An analysis of the Canadian community health survey. *Health Promotion and Chronic Disease Prevention in Canada*. 2015; 35(7):99–108. PMID: 26378768
23. Pöttsch O, Rößger F. Bevölkerung Deutschlands bis 2060. Ergebnisse der 13. koordinierten Bevölkerungsvorausberechnung. Wiesbaden: Statistisches Bundesamt, Wiesbaden; 2015. p. 20.
24. Rapp K, Freiburger E, Todd C, Klenk J, Becker C, Denking M, et al. Fall incidence in Germany: results of two population-based studies, and comparison of retrospective and prospective falls data collection methods. *BMC Geriatr*. 2014; 14:105. <https://doi.org/10.1186/1471-2318-14-105> PMID: 25241278
25. Du Y, Wolf I-K, Knopf H. Psychotropic drug use and alcohol consumption among older adults in Germany: results of the German Health Interview and Examination Survey for Adults 2008–2011. *BMJ Open*. 2016; 6(10). <https://doi.org/10.1136/bmjopen-2016-012182> PMID: 27855095
26. Scheidt-Nave C, Kamtsiuris P, Gosswald A, et al. German health interview and examination survey for adults (DEGS)—design, objectives and implementation of the first data collection wave. *BMC public health*. 2012; 12:730. <https://doi.org/10.1186/1471-2458-12-730> PMID: 22938722
27. Kamtsiuris P, Lange M, Hoffmann R, Schaffrath Rosario A, Dahm S, Kuhnert R, et al. Die erste Welle der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1): Stichprobendesign, Response, Gewichtung und Repräsentativität. [The first wave of the German Health Interview and Examination Survey for Adults (DEGS1): sample design, response, weighting and representativeness].

- Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz. 2013; 56(5–6):620–30. <https://doi.org/10.1007/s00103-012-1650-9> PMID: 23703478
28. Knopf H, Grams D. Arzneimittelanwendung von Erwachsenen in Deutschland: Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1). [Medication use of adults in Germany: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz. 2013; 56(5–6):868–77. <https://doi.org/10.1007/s00103-013-1667-8> PMID: 23703508
 29. Wolf IK, Du Y, Knopf H. Changes in prevalence of psychotropic drug use and alcohol consumption among the elderly in Germany: results of two National Health Interview and Examination Surveys 1997–99 and 2008–11. BMC Psychiatry. 2017; 17(1):90. <https://doi.org/10.1186/s12888-017-1254-x> PMID: 28279159
 30. Lampert T, Kroll L, Müters S, Stolzenberg H. Measurement of socioeconomic status in the German Health Interview and Examination Survey for Adults (DEGS1) [Messung des sozioökonomischen Status in der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1)]. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz. 2013; 56(5):631–6. <https://doi.org/10.1007/s00103-012-1663-4> PMID: 23703479
 31. Hammond T, Wilson A. Polypharmacy and falls in the elderly: a literature review. Nurs Midwifery Stud. 2013; 2(2):171–5. PMID: 25414854
 32. Klein D, Nagel G, Kleiner A, Ulmer H, Rehberger B, Concin H, et al. Blood pressure and falls in community-dwelling people aged 60 years and older in the VHM&PP cohort. BMC Geriatr. 2013; 13:50. <https://doi.org/10.1186/1471-2318-13-50> PMID: 23692779
 33. Buttery AK, Busch MA, Gaertner B, Scheidt-Nave C, Fuchs J. Prevalence and correlates of frailty among older adults: findings from the German health interview and examination survey. BMC Geriatr. 2015; 15:22. <https://doi.org/10.1186/s12877-015-0022-3> PMID: 25879568
 34. Neuhauser H, Thamm M, Ellert U. [Blood pressure in Germany 2008–2011: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz. 2013; 56(5–6):795–801. <https://doi.org/10.1007/s00103-013-1669-6> PMID: 23703500
 35. Milos V, Bondesson A, Magnusson M, Jakobsson U, Westerlund T, Midlov P. Fall risk-increasing drugs and falls: a cross-sectional study among elderly patients in primary care. BMC Geriatr. 2014; 14:40. <https://doi.org/10.1186/1471-2318-14-40> PMID: 24674152
 36. Tinetti ME, Han L, Lee DSH, McAvay GJ, Peduzzi P, Gross CP, et al. Antihypertensive medications and serious fall injuries in a nationally representative sample of older adults. JAMA Internal Medicine. 2014; 174(4):588–95. <https://doi.org/10.1001/jamainternmed.2013.14764> PMID: 24567036
 37. Findley LR, Bulloch MN. Relationship between nonsteroidal anti-inflammatory drugs and fall risk in older adults. Consultant Pharmacist. 2015; 30(6):346–51. <https://doi.org/10.4140/TCP.n.2015.346> PMID: 26048465
 38. Scott D, Blizzard L, Fell J, Jones G. Statin therapy, muscle function and falls risk in community-dwelling older adults. QJM. 2009; 102(9):625–33. <https://doi.org/10.1093/qjmed/hcp093> PMID: 19633029
 39. Hill KD, Wee R. Psychotropic drug-induced falls in older people: A review of interventions aimed at reducing the problem. Drugs and Aging. 2012; 29(1):15–30. <https://doi.org/10.2165/11598420-000000000-00000> PMID: 22191720
 40. Marcum ZA, Perera S, Thorpe JM, Switzer GE, Castle NG, Strotmeyer ES, et al. Antidepressant Use and Recurrent Falls in Community-Dwelling Older Adults: Findings From the Health ABC Study. Annals of Pharmacotherapy. 2016; 50(7):525–33. <https://doi.org/10.1177/1060028016644466> PMID: 27066988
 41. Naples JG, Kotlarczyk MP, Perera S, Greenspan SL, Hanlon JT. Non-tricyclic and Non-selective Serotonin Reuptake Inhibitor Antidepressants and Recurrent Falls in Frail Older Women. The American Journal of Geriatric Psychiatry. 2016. <http://dx.doi.org/10.1016/j.jagp.2016.08.008>.
 42. Gebara MA, Lipsey KL, Karp JF, Nash MC, Iaboni A, Lenze EJ. Cause or Effect? Selective Serotonin Reuptake Inhibitors and Falls in Older Adults: A Systematic Review. Am J Geriatr Psychiatry. 2015; 23(10):1016–28. <https://doi.org/10.1016/j.jagp.2014.11.004> PMID: 25586602
 43. Stubbs B. A Meta-Analysis Investigating Falls in Older Adults Taking Selective Serotonin Reuptake Inhibitors Confirms an Association but by No Means Implies Causation. Am J Geriatr Psychiatry. 2015; 23(10):1098. <https://doi.org/10.1016/j.jagp.2015.02.004> PMID: 26233930
 44. Kerse N, Flicker L, Pfaff JJ, Draper B, Lautenschlager NT, Sim M, et al. Falls, depression and antidepressants in later life: a large primary care appraisal. PLoS One. 2008; 3(6):e2423. <https://doi.org/10.1371/journal.pone.0002423> PMID: 18560599

45. Johnell K, Jonasdottir Bergman G, Fastbom J, Danielsson B, Borg N, Salmi P. Psychotropic drugs and the risk of fall injuries, hospitalisations and mortality among older adults. *Int J Geriatr Psychiatry*. 2016. <https://doi.org/10.1002/gps.4483> PMID: 27113813
46. Masud T, Frost M, Ryg J, Matzen L, Ibsen M, Abrahamsen B, et al. Central nervous system medications and falls risk in men aged 60–75 years: The study on male osteoporosis and aging (SOMA). *Age and Ageing*. 2013; 42(1):121–4. <https://doi.org/10.1093/ageing/afs118> PMID: 22931902
47. Hartikainen S, Lönnroos E, Louhivuori K. Medication as a Risk Factor for Falls: Critical Systematic Review. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2007; 62(10):1172–81.
48. Kuschel BM, Laflamme L, Möller J. The risk of fall injury in relation to commonly prescribed medications among older people—A Swedish case-control study. *European Journal of Public Health*. 2015; 25(3): 527–32. <https://doi.org/10.1093/eurpub/cku120> PMID: 25085470
49. Pierce JR Jr, Shirley M, Johnson EF, Kang H. Narcotic administration and fall-related injury in the hospital: Implications for patient safety programs and providers. *International Journal of Risk and Safety in Medicine*. 2013; 25(4):229–34. <https://doi.org/10.3233/JRS-130603> PMID: 24305561
50. Salas M, Hofman A, Stricker BH. Confounding by indication: an example of variation in the use of epidemiologic terminology. *Am J Epidemiol*. 1999; 149(11):981–3. PMID: 10355372
51. Signorello LB, McLaughlin JK, Lipworth L, Friis S, Sorensen HT, Blot WJ. Confounding by indication in epidemiologic studies of commonly used analgesics. *Am J Ther*. 2002; 9(3):199–205. PMID: 11941379
52. Stubbs B, Stubbs J, Gnanaraj SD, Soundy A. Falls in older adults with major depressive disorder (MDD): A systematic review and exploratory meta-analysis of prospective studies. *International Psychogeriatrics*. 2016; 28(1):23–9. <https://doi.org/10.1017/S104161021500126X> PMID: 26234532
53. Kvelde T, McVeigh C, Toson B, Greenaway M, Lord SR, Delbaere K, et al. Depressive symptomatology as a risk factor for falls in older people: systematic review and meta-analysis. *J Am Geriatr Soc*. 2013; 61(5):694–706. <https://doi.org/10.1111/jgs.12209> PMID: 23617614
54. Grenier S, Payette MC, Langlois F, Vu TTM, Bherer L. Depressive symptoms are independently associated with recurrent falls in community-dwelling older adults. *International Psychogeriatrics*. 2014; 26(9): 1511–9. <https://doi.org/10.1017/S104161021400074X> PMID: 24758735