



Original Research

Linking EORTC QLQ-C-30 and PedsQL/PEDQOL physical functioning scores in patients with osteosarcoma[☆]



Axel Budde^{a,*}, Katja Baust^a, Leonie Weinhold^b, Mark Bernstein^c, Stefan Bielack^d, Catharina Dhooge^e, Lars Hjorth^f, Katherine A. Janeway^g, Meriel Jenney^h, Mark D. Krailoⁱ, Neyssa Marina^j, Rajaram Nagarajan^k, Sigbjørn Smeland^l, Matthew R. Sydes^m, Patricia De Vosⁿ, Jeremy Whelan^o, Andreas Wiener^p, Gabriele Calaminus^{a,l}, Matthias Schmid^{b,l}

^a Department of Paediatric Haematology and Oncology, University Hospital Bonn, Bonn, Germany

^b Department of Medical Biometry, Informatics, and Epidemiology, University Hospital Bonn, Bonn, Germany

^c IWK Health Centre, Dalhousie University, Halifax, NS, Canada

^d Zentrum für Kinder-, Jugend- und Frauenmedizin, Pädiatrie, Klinikum Stuttgart, Olgahospital, Stuttgart, Germany

^e Department of Internal Medicine and Paediatrics, Faculty of Medicine and Health Sciences, Ghent University Hospital, Ghent, Belgium

^f Department of Clinical Sciences, Department of Paediatrics, Lund University, Skane University Hospital, Lund, Sweden

^g Dana-Farber/Boston Children's Cancer and Blood Disorders Center, Harvard Medical School, Boston, MA, USA

^h Women's Services Clinical Board, University Hospital of Wales, Cardiff, UK

ⁱ Statistics and Data Center, Children's Oncology Group, Monrovia, CA, USA

^j Five Prime Therapeutics, South San Francisco, CA, USA

^k Division of Oncology, Cincinnati Children's Hospital Medical Center, University of Cincinnati College of Medicine, Cincinnati, OH, USA

^l Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway

^m MRC Clinical Trials Unit at UCL, Institute of Clinical Trials and Methodology, University College London, London, UK

ⁿ Department of Paediatric Haematology and Oncology, Ghent University Hospital, Ghent, Belgium

^o Department of Oncology, University College Hospital, London, UK

^p West German Proton Therapy Center Essen, Essen, Germany

Received 18 January 2022; received in revised form 5 March 2022; accepted 15 March 2022

Available online 8 June 2022

Abbreviations: COG, Children's Oncology Group; COSS, Cooperative Osteosarcoma Group; EOI, European Osteosarcoma Intergroup; EORTC QLQ-C30, European Organisation for Research and Treatment of Cancer Core Questionnaire; EURAMOS-1, European American Osteosarcoma Study-1; FACT-G, Functional Assessment of Cancer Therapy - General; IRT, Item Response Theory; LOA, Limit of Agreement; PEDQOL, Paediatric Quality of Life Questionnaire; PedsQL, Paediatric Quality of Life Inventory; PRO, Patient-Reported Outcome; PROMIS, Patient-Reported Outcomes Measurement Information System; QoL, Quality-of-Life; SSG, Scandinavian Sarcoma Group.

[☆] The study includes results from the EURAMOS-1 trial, registered as NCT00134030 and ISRCTN 67613327.

* Corresponding author:

E-mail address: axel.budde@ukbonn.de (A. Budde).

^l These authors contributed equally to this work.

<https://doi.org/10.1016/j.ejca.2022.03.018>

0959-8049/© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

KEYWORDS

Childhood cancer;
EORTC QLQ-C30;
Patient-reported
outcome (PRO);
Paediatric quality of
life inventory
(PedsQL);
Paediatric quality of
life questionnaire
(PEDQOL);
Physical functioning
quality-of-life (QoL);
Score linking

Abstract Purpose: The available questionnaires for quality-of-life (QoL) assessments are age-group specific, limiting comparability and impeding longitudinal analyses. The comparability of measurements, however, is a necessary condition for gaining scientific evidence. To overcome this problem, we assessed the viability of harmonising data from paediatric and adult patient-reported outcome (PRO) measures.

Method: To this end, we linked physical functioning scores from the Paediatric Quality of Life Inventory (PedsQL) and the Paediatric Quality of Life Questionnaire (PEDQOL) to the European Organisation for Research and Treatment of Cancer Core Questionnaire (EORTC QLQ-C30) for adults. Samples from the EURAMOS-1 QoL sub-study of 75 (PedsQL) and 112 (PEDQOL) adolescent osteosarcoma patients were concurrently administered both paediatric and adult questionnaires on 98 (PedsQL) and 156 (PEDQOL) occasions. We identified corresponding scores using the single-group equipercentile linking method.

Results: Linked *physical functioning* scores showed sufficient concordance to the EORTC QLQ-C30: Lin's $\rho = 0.74$ (PedsQL) and Lin's $\rho = 0.64$ (PEDQOL).

Conclusion: Score linking provides clinicians and researchers with a common metric for assessing QoL with PRO measures across the entire lifespan of patients.

© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Quality-of-life (QoL) data are generally collected by self-report questionnaires. Health-related QoL questionnaires can be age-group specific. This age group specificity limits comparability and impedes numerical longitudinal analysis, especially if different instruments are needed to span the age range of the study. Specifically, the motivation for linking scores from paediatric and adult instruments was to make them comparable on a common scale, allowing the study of the QoL developmental trajectory continuously and permitting the analysis with mixed models.

The use of different instruments constitutes a considerable hurdle for the analysis and interpretation of QoL data, since “[t]he comparability of measurements made in differing circumstances by different methods and investigators is a fundamental precondition for all of science” [1]. Therefore, valid methods for linking scores are required.

Dorans provides an overview of applying linking methodology within the realm of patient-reported outcome (PRO) measures [2] (Table 1a).

In the present study, we evaluated the viability of linking *physical functioning* scores of two paediatric PRO questionnaires (the PedsQL and the PEDQOL) to the EORTC QLQ-C30 in a population of survivors of childhood osteosarcoma. We restrict our report to the *physical functioning* domain because we were mainly interested in the viability of linking paediatric and adult instruments. We provide information on linking *emotional functioning*, *cognitive functioning*, *social functioning*, *fatigue* and *pain* domains in the appendix.

2. Materials and methods

The overall study design [25,26] and the methodological specifics of the QoL questionnaire sub-study have been laid out in detail previously [27]. We briefly describe the study design.

2.1. Participants

The EURAMOS-1 trial cohort consisted of 2260 participants who, between the ages 5 and 40 years old, had been diagnosed with a previously untreated resectable high-grade osteosarcoma (at any site, except for craniofacial structures). Among these, 2213 participants were eligible for QoL-assessment (≥ 5 years old) and had a questionnaire in their respective language available (see [27]). Recruitment took place between 2005 and 2011, involving 17 countries and four study groups: the Children's Oncology Group (COG), the Cooperative Osteosarcoma Group (COSS), the European Osteosarcoma Intergroup (EOI), and the Scandinavian Sarcoma Group (SSG). EURAMOS-1 consortium members and their affiliations are listed in Appendix E.1. We obtained demographics from the EURAMOS-1 enrolment survey (sex, date of birth, and study group). Age was stratified as “5 to 15”, “16 to 17” and “18 or older”. As a secondary outcome measure, QoL was assessed prospectively at four time points during and after treatment (Fig. 1).

2.2. Questionnaires

Due to the unavailability of a single questionnaire suited for use across the whole age span of participants and in

Table 1
Publications on linking PRO measures.

Publications
Adults
Health status [3–6]
Physical functioning [7, 8]
Physical and mental health summary scores [9]
Self-regulation [10]
Depression [11–15]
Pain [16]
Pain interference [17]
Anxiety [18–21]
Fatigue [20, 21]
EORTC QLQ-C30 <> FACT-G [22]
Children < > Adults
Emotional distress [23]
Physical functioning in a population of individuals with spinal cord injury [24]

all participating countries, the EURAMOS-1 consortium opted for using different, age- and country-specific instruments (Table 2a).

In the age range 16–18 years old, all patients were asked to complete a paediatric questionnaire (either PedsQL [28] or PEDQOL [29]) and the EORTC QLQ-C30 [30]. We used this sub-sample for score linking. We restricted our study to aggregate scores pertaining to physical functioning, given its significance to QoL in osteosarcoma survivors and the substantial conceptual overlap between instruments in this domain. We linked two sub-sets of participants aged 16–17 years. These sub-sets were administered either the PedsQL or the PEDQOL questionnaire before the EORTC QLQ-C30 on the same day.

2.3. Analyses

2.3.1. Similarity of item content and physical functioning sub-scale structure between instruments

The PedsQL, the PEDQOL and the EORTC QLQ-C30 all contain items that assess the physical functioning domain with multiple items (for details on

scoring, see Table 2b and for verbatim item content see Appendix F).

Item content showed substantial overlap across the three measures. To measure internal consistency of the instruments, we calculated Cronbach’s α . A summary of the results is given in Table 2c.

2.3.2. Summary of physical functioning raw scores, correlation and concordance between instruments

The overall mean physical functioning score, i.e. across all four time points, was 51.6 (SD = 22.7) for the PedsQL and 74.3 (SD = 22.3) for the corresponding EORTC QLQ-C30 (n = 98). The overall mean for physical functioning of the PEDQOL was 46.8 (SD = 25.1) and the corresponding EORTC QLQ-C30 overall mean was 63.5 (SD = 27.2) (n = 156).

The correlations between the EORTC QLQ-C30 physical functioning sub-scale and the corresponding aggregate scores of the two paediatric instruments were both good, but the PedsQL physical functioning raw scores correlated more strongly (r = 0.73; 95% confidence interval (CI): 0.63, 0.81) than those of the PEDQOL (r = 0.64; CI: 0.54, 0.73). The physical functioning raw scores of the paediatric questionnaires showed only moderate agreement with those of the EORTC QLQ-C30 before linking, with similar values for the PedsQL (Lin’s ρ = 0.49; CI: 0.63, 0.81) and the PEDQOL (Lin’s ρ = 0.53; CI: 0.43, 0.63). Given a substantial overlap in item content, we linked the respective aggregate physical

Table 2a
QoL questionnaire by region and age group.

Questionnaire	Region	Age group		
		≥5	16–17	≥18
PedsQL	COG (North America) & EOI (North West Europe)	+	+	+
PEDQOL	COSS (Central Europe) & SSG (Scandinavia)	+	+	+
EORTC QLQ-C30	All	+	+	+




Fig. 1. Timeline for QoL assessments.

Table 2b
QoL questionnaires physical functioning scoring.

Questionnaire	Number of items	Scale points	Period
PedsQL	4	5	Past month
PEDQOL	4	4	Past week
EORTC QLQ-C30	5	4	Past week

functioning scores of the PedsQL and the PEDQOL questionnaires to their EORTC QLQ-C30 equivalent.

2.3.3. Linking design

To produce physical functioning crosswalks (score conversion tables), we linked scores of those participants who had completed one of the two paediatric instruments and the EORTC QLQ-C30 at the same time point. This group consisted of participants who were 16–18 years old. This linking technique, referred to as the single-group design, is akin to a repeated measures design with a single group and two treatments [31]. It is considered the most valid linking design because the scores of identical individuals are linked, thus requiring the smallest sample size to achieve the same level of accuracy as designs with a lesser degree of group equivalency [32].

To ensure that the instruments to be linked showed sufficient conceptual congruity [2], we employed two methods, modelling our approach on Choi *et al.* (2014) and Marrie *et al.* (2020). First, we reviewed the content of the physical functioning items of the three instruments to ensure that they indeed measure approximately the same concept. Second, to assess internal

consistency, we calculated Cronbach's α for the three questionnaires.

2.3.4. Linking function

We performed identity, mean, linear, equipercentile and circle-arc linking procedures (Fig. 2). Previously, we had applied log-linear pre-smoothing to three moments to adjust for potential sampling error introduced by uneven score distributions [33]. Log-linear pre-smoothing is a recommended procedure for small samples such as ours because a smoothed distribution yields more reliable results [33]. We used root mean square error (RMSE) by means of parametric bootstrapping to determine the best linking method (for details see [34], 5.7).

We chose the equipercentile linking method to produce crosswalk tables, as it emerged as the method with the most favourable linking quality parameters, overall.

2.3.5. Evaluation of linking quality

As a first step towards ascertaining the agreement between paediatric and adult QoL instruments, we created Bland–Altman plots [35] (Fig. 3 and Table 2d). We plotted the differences (y-axis) for scores linked from each paediatric questionnaire and those measured by the EORTC QLQ-C30 against subject means (x-axis) to check for patterns and distributions. Following Zhou *et al.* [36], we established that the limits of agreement for linked and measured scores were to be considered "good" if they fell within one standard deviation (SD) of the mean of measured EORTC QLQ-C30 scores, "fair" if they did not extend beyond two SDs, and "poor", otherwise.

Table 2c
Internal consistency reliability of the physical functioning aggregate scores of the three instruments.

Time point	Questionnaire	Linked to	N ¹	Cronbach's α (95% CI)	Item–total correlation ²		
					Min	Mean	Max
E1	PedsQL	EORTC QLQ-C30	38	0.87 (0.80, 0.93)	0.43	0.69	0.82
	PEDQOL		41	0.68 (0.52, 0.85)	0.49	0.60	0.68
	EORTC QLQ-C30	PedsQL	38	0.80 (0.70, 0.89)	0.35	0.65	0.79
E2	PedsQL	EORTC QLQ-C30	24	0.73 (0.57, 0.88)	0.28	0.51	0.82
	PEDQOL		47	0.47 (0.22, 0.72)	0.06	0.48	0.68
	EORTC QLQ-C30	PedsQL	24	0.73 (0.58, 0.87)	0.26	0.61	0.75
E3	PedsQL	EORTC QLQ-C30	20	0.82 (0.73, 0.90)	0.53	0.68	0.77
	PEDQOL		41	0.77 (0.63, 0.92)	0.42	0.57	0.67
	EORTC QLQ-C30	PedsQL	20	0.60 (0.40, 0.80)	0.42	0.54	0.59
E4	PedsQL	EORTC QLQ-C30	20	0.76 (0.61, 0.91)	0.43	0.70	0.85
	PEDQOL		41	0.78 (0.69, 0.87)	0.45	0.67	0.80
	EORTC QLQ-C30	PedsQL	16	0.86 (0.76, 0.95)	0.34	0.68	0.88
E4	PedsQL	EORTC QLQ-C30	27	0.65 (0.44, 0.86)	0.28	0.59	0.74
	PEDQOL		27	0.85 (0.74, 0.95)	0.60	0.79	0.90
	EORTC QLQ-C30	PedsQL	16	0.85 (0.74, 0.95)	0.60	0.79	0.90
		PEDQOL	27	0.74 (0.59, 0.89)	0.31	0.64	0.75

¹ N refers to the number of participants in which Cronbach's α was measured for the instrument in the second column when linked to the instrument in the third column.

² Item–total correlation indicates the correlation between the score on a single item and the aggregate physical functioning sub-scale score.




Fig. 2. Five functions linking physical functioning scores.

Additionally, we calculated Pearson's correlation coefficient r and Lin's concordance correlation coefficient between each of the two paediatric measures and the EORTC QLQ-C30.

We prepared histograms of the differences between measured and linked EORTC QLQ-C30 scores to visually inspect whether the distributions approximate normality (Fig. 4).

Details on software are given in Appendix A.

3. Results

3.1. Participant characteristics

The QoL sub-sample consisted of 2213 osteosarcoma patients. The mean age at registration was 15.1 ($SD = 5.3$) years. Out of the complete sub-sample, 760 participants had completed the PedsQL in the *physical functioning* domain at one or more of the four time points, and 337 had completed the PEDQOL in this domain at one time point or more. Out of these participants, 75 participants between the ages of 16 and 18 had completed both the PedsQL and the EORTC QLQ-C30 in the *physical functioning* domain at the same time point on 98 occasions, and 112 had completed both the PEDQOL and the EORTC QLQ-C30 concurrently on 156 occasions.

Table 3a gives an overview of patient characteristics by linked questionnaire (PedsQL or PEDQOL) for the *physical functioning* domain, including sex, age, and study group, overall and by linked sub-sample.

3.1.1. Bland–Altman plots

We used Bland–Altman plots to compare PedsQL and PEDQOL scores to EORTC QLQ C-30 scores. The interpretation of Bland–Altman plots is premised on normality and homoscedasticity of the distribution. We prepared histograms for the distributions of differences (Fig. 4 and Table 2d) to make a first visual assessment. We then prepared Bland–Altman plots (Fig. 3) displaying the differences in scores between each paediatric instrument and the EORTC QLQ-C30 against the respective means.

To inspect for heteroscedasticity, we prepared quantile–quantile (Q–Q) plots (Fig. 5) for differences between scores linked from the two paediatric questionnaires and EORTC QLQ-C30 scores. We judged that scores linked from the PEDQOL displayed adequate homoscedasticity. However, scores linked from the PedsQL indicated an uneven, left-skewed distribution. Therefore, we log-transformed the score differences, achieving better overall homoscedasticity, albeit with a remaining left skew (Fig. 6). To account for the presence of substantial heteroscedasticity in scores linked from the PedsQL, we prepared a Bland–Altman plot on log-transformed data (Fig. 7a) which indicated a better fit of limits of agreement. Given that log-transformed scores do not lend themselves to easy interpretation for clinical practice, we additionally plotted the score differences in a conventional Bland–Altman plot on the original scale with back-transformed limits of agreement (Fig. 7b) [37,38].

Summarily, we judged agreement for physical functioning scores acceptable, as the limits of agreement did




Fig. 3. Bland–Altman plots for linked vs. observed physical functioning scores.

Table 2d
 Bland–Altman plots: descriptive characteristics.

Bland–Altman analysis					
	Parameter	Count	Value	SD	95% CI
PedsQL	Bias	98	0.53	15.92	(−2.66, 3.72)
	Upper LOA	98	31.73		(26.26, 37.21)
	Lower LOA	98	−30.67		(−36.15, −25.20)
PedQOL	Bias	156	−0.32	22.84	(−3.94, 3.29)
	Upper LOA	156	44.44		(38.25, 50.62)
	Lower LOA	156	−45.09		(−51.27, −38.90)




Fig. 4. Histograms with distributions of differences between *physical functioning* scores.

Table 3a
 Characteristics of participants in the physical functioning domain by paediatric questionnaire, overall and linked.

Characteristics	Physical functioning ¹			
	PedsQL		PEDQOL	
	Overall	Linked to EORTC QLQ-C30	Overall	Linked to EORTC QLQ-C30
Sex, n(%)				
Male	429 (56)	48 (64)	171 (51)	68 (61)
Female	331 (44)	27 (36)	164 (49)	44 (39)
Age (years)²				
Age group, n(%)				
5 to 15	671 (88)	44 (59)	275 (82)	70 (62)
16 to 17	74 (10)	22 (29)	59 (18)	42 (38)
18 or older	15 (2)	9 (12)	1 (0)	0 (0)
Mean (SD)	12.8 (3.0)	15.1 (2.7)	13.4 (2.9)	15.6 (1.3)
Study group, n(%)				
COG	616 (81)	96 (98)	0 (0)	0 (0)
COSS	0 (0)	0 (0)	211 (63)	83 (74)
EOI	144 (19)	2 (2)	59 (18)	22 (20)
SSG	0 (0)	0 (0)	65 (19)	7 (6)

COG: Childrens's Oncology Group; COSS: Cooperative Osteosarcoma Group; EOI: European Osteosarcoma Intergroup; SSG: Scandinavian Sarcoma Group.

¹ The columns pertain to those participants whose PedsQL or PEDQOL scores were linked to their respective EORTC QLQ-C30 scores. Therefore, the table does not contain a separate column for EORTC QLQ-C30 scores.

² Age refers to the age at the time of registration for participation in the study.

not extend beyond two standard deviations of EORTC QLQ-C30 scores for either of the paediatric instruments, and the majority of scores being within one standard deviation of EORTC QLQ-C30 scores.

3.1.2. Correlations between physical functioning aggregate scores of paediatric and adult instruments

Additionally, we calculated Pearson's r and Lin's ρ –[39] concordance correlation coefficients between the EORTC QLQ-C30 and the PedsQL and PEDQOL physical functioning converted scores.

The correlation coefficients for physical functioning scores were good for both the PedsQL and the PEDQOL to EORTC QLQ-C30 conversions, with a Lin's ρ of 0.74 and 0.64, respectively (Table 3b and 3c).

3.1.3. Correlations between other aggregate scores of paediatric and adult instruments

The converted scores of the PedsQL and PEDQOL *fatigue* both correlated well with EORTC QLQ-C30 scores (Lin's $\rho = 0.69$ and Lin's $\rho = 0.71$). Correlation coefficients for *pain* were moderate for the PedsQL

(Lin's $\rho = 0.58$) and good for the PEDQOL (Lin's $\rho = 0.73$). Correlation coefficients for *emotional functioning* were moderate (Lin's $\rho = 0.55$) for the PedsQL and fair for PEDQOL (Lin's $\rho = 0.36$) conversions to EORTC QLQ-C30 scores. The correlation of converted *cognitive functioning* scores with EORTC QLQ-C30 scores was fair for the PedsQL (Lin's $\rho = 0.37$) and moderate for the PEDQOL (Lin's $\rho = 0.47$). Converted *social functioning* scores correlated poorly with EORTC QLQ-C30 scores for both, the PedsQL (Lin's $\rho = 0.17$) and the PEDQOL PedsQL (Lin's $\rho = 0.08$).

4. Discussion

Data harmonisation provides a number of benefits by permitting the pooling of data, such as answering novel research questions or increasing statistical power. Despite a growing interest in harmonising data, retrospective data harmonisation (after data collection) is the rule and prospective harmonisation (before data collection) the exception [3]. While it may be due to a lack of foresight or practicability that retrospective data




Fig. 5. Quantile–quantile plots of differences between physical functioning scores.

harmonisation remains the only option, harmonising data prospectively may also be inherently impossible. This was the case in the international research collaboration the present study grew out of which included longitudinal QoL assessments in adult survivors of childhood osteosarcoma. The use of different PRO measures during childhood and adulthood was unavoidable, as no suitable instrument for both age groups existed.

To obtain harmonised data retrospectively, we linked the scores from two paediatric PRO measures to an

adult PRO measure to assess the quality of life across the lifespan of osteosarcoma survivors. Visual and numerical concordance assessments indicated good agreement between physical functioning aggregate scores. The equipercenile linking method yielded the best overall results for this sample. Sub-sets consisting of 75 (PedsQL) and 112 participants (PEDQOL) yielded 98 (PedsQL) and 156 (PEDQOL) score pairings between paediatric and adult questionnaires and were sufficient to permit score linking for the whole cohort and enabled the analysis of QoL data for a forthcoming publication.

In domains other than *physical functioning*, the concordance estimates obtained with Pearson's r diverged from those obtained with dedicated concordance coefficients (Appendix, Table D.1), thus confirming that Pearson's r is not a useful measure for assessing intra-individual agreement. The Pearson correlation coefficient (Pearson's r) is generally not considered a suitable measure of concordance because it is only informative if the relationship between two variables is linear, thus potentially leading to incorrect conclusions in case of non-linearity. Crucially, Pearson's r only evaluates the extent of a linear relationship on a population level, ignoring intra-individual concordance. Despite its apparent shortcomings, Pearson's r continues to be widely employed in the score linking literature as a measure of agreement between two instruments. This is all the more surprising, given that non-linear score linking methods were presumably developed to specifically account for non-linear agreement between two instruments. Due to its continued popularity and to underscore differences between concordance measures, we nevertheless included Pearson's r alongside Lin's concordance correlation coefficient ρ [39] which we consider more apt. We provide an evaluation according to value ranges to allow a verbal interpretation, similar to the kappa concordance




Fig. 6. Quantile–quantile plot of logarithm of differences between PedsQL and EORTC QLQ-C30 physical functioning scores ($n = 98$).




Fig. 7. Bland–Altman plots for linked vs. observed log-transformed and back-transformed physical functioning scores (n = 98).

coefficient for binary variables [35], with five categories, ranging from “Poor” to “Very Good” (Table 3c).

Building on McNemar’s coefficient of alienation, Dorans [40] defined “Reduction in Uncertainty”. Since a 50% reduction in uncertainty, as measured in score units, requires a Pearson’s *r* of at least 0.866, Dorans recommended a correlation of this magnitude as an appropriate lower bound. This recommendation was made in the context of high-stakes educational testing, as Choi and colleagues [12] have pointed out. For linking health outcome measures, they suggested a correlation of 0.75–0.80 as an appropriate minimum, given that aggregate outcomes are the focus of interest, and in particular when using a single-group design which permits the direct evaluation of accuracy.

A limitation of our study is that our results may not be population invariant, i.e. the linking quality parameters we obtained may not generalise to other populations. Previous studies linking PedsQL or PEDQOL *physical functioning* aggregate scores to the EORTC QLQ-C30 are lacking. Therefore, we were unable to draw comparisons to similar or dissimilar populations and we cannot generalise our findings beyond the highly selective clinical population our sample was drawn from. The aim of our study was to evaluate the feasibility of linking paediatric and adult PRO measures within a population of osteosarcoma survivors. Clearly, our findings are restricted to this narrowly circumscribed area of clinical practice and research. The methodology also does not allow for

harmonisation in completely disparate age groups (e.g. 5–10 year-old with 35–40 year-old).

The use of age-adequate (i.e. age-specific) questionnaires for children seems unavoidable, rendering a direct comparison of paediatric and adult scores in survivors of childhood cancer inherently impossible. Therefore, we see the potential general utility of score linking in this field in offering interoperability of paediatric and adult PRO measures, and the specific value of this study in showing the viability of this approach for the first time. Having established its feasibility, the approach described may be integrated in future study designs involving dissimilar populations. Doing so may yield evidence regarding the population invariance of our results.

Another limitation of our study is that we cannot rule out an order effect, i.e. the relationship of the instruments may have depended on the order of their administration. This point should be addressed in future investigations by randomising the order of administration. In a similar vein, the administration of two questionnaires at the same point may have biased the responses to the second questionnaire. Randomising the order of administration should also reduce fatigue bias, by equalising the directionality of such an effect between the instruments.

We consider the single-group design a major strength of our study, as it provides the firmest methodological grounds for score linking. Its inherent potential

Table 3b
Paediatric questionnaires and EORTC QLQ-C30: Correlation and concordance coefficients post-linking.

Coefficient	Physical functioning	
	PedsQL	PEDQOL
Pearson’s <i>r</i> (95% CI)	0.74 (0.64–0.82)	0.64 (0.54–0.72)
Lin’s ρ (95% CI)	0.74 (0.64–0.82)	0.64 (0.54–0.72)

Table 3c
Interpretation of concordance correlation coefficients [35].

Concordance Correlation coefficient	Strength of agreement
<0.20	Poor
0.21–0.40	Fair
0.41–0.60	Moderate
0.61–0.80	Good
0.81–1.00	Very good

disadvantages should be balanced against its strengths and against the weaknesses of alternative linking designs. Using a single-group design, we obtained actual and linking-derived scores from the same population. This allowed us to evaluate the accuracy of our linking functions directly. As a tangible product, we created crosswalk tables between PedsQL and PEDQOL *physical functioning* aggregate scores (see Appendix, [Table L.1](#)) which will bring forward data harmonisation and will enable us to perform longitudinal analyses within the EURAMOS-1 cohort.

With score linking, it is possible to directly compare scores of osteosarcoma patients obtained with distinct age-group-specific inventories and observe their QoL across the entire lifespan. The approach may create the conditions for conducting longitudinal mixed-model meta-analyses. We consider score linking a promising tool for assuring comparability of intra-individual QoL assessments in studies over time and extending across different stages of life. We anticipate that oncological QoL research may strongly benefit from score linking.

Funding

The study sponsor was the UK Medical Research Council in Europe and the US National Cancer Institute in North America and Australia. Each trial group organised local coordination elements; central coordination and analysis was led from Medical Research Council Clinical Trials Unit at University College of London. Neither the sponsors nor the funders of the trial had a role in trial design, data analysis, or data interpretation. The EURAMOS-1 is an academic clinical trial funded through multiple national and international government agencies and cancer charities: - Children's Oncology Group funding for the EURAMOS-1 trial (AOST0331) was supported by the National Clinical Trial Network (NCTN) Operations Centre Grant U10CA180886, NCTN Statistics and Data Center Grant U10CA180899 and St. Baldrick's Foundation. - European Science Foundation under the European Science Foundation Collaborative Research Programme for Pan-European Clinical Trials, through contract number ERASCT-2003-980409 of the European Commission, DG Research, FP6 (Ref No MM/NG/EMRC/0202) National funding in Europe was provided by the following:

- Belgium: Fonds National de la Recherche Scientifique Belgium FWO (Fonds voor Wetenschappelijk Onderzoek-Vlaanderen)
- Denmark: Danish Medical Research Council
- Finland: Academy of Finland
- Germany: Deutsche Forschungsgemeinschaft ref No: BI 1045/1-1 & 1–2, Deutsche Krebshilfe (DKH) ref No: 50-2723-Bi2
- Hungary: Semmelweis Foundation
- Netherlands: ZonMw (Council for Medical Research)
- Norway: Research Council of Norway

- Sweden: SSG and Swedish Childhood Cancer Fund
- Switzerland: Swiss Paediatric Oncology Group
- United Kingdom: includes funding for the trial coordinating data centre (MRC Clinical Trials Unit at UCL): Cancer Research UK, CRUK/05/013, Medical Research Council: MC_UU_12023/28.

Additional funding to the University of Münster Centre for Clinical Trials, site of the EURAMOS Intergroup Safety Desk: Federal Ministry of Education and Research, Germany, BMBF 01KN1105.

CRedit authorship contribution statement

Axel Budde: Conceptualisation, Data curation, Formal analysis, Methodology, Software, Writing - original draft preparation, Visualisation. **Katja Baust:** Conceptualisation, Data curation, Writing - review and editing, Project administration. **Leonie Weinholt:** Methodology, Validation, Writing - review & editing. **Mark Bernstein:** Investigation, Writing - review and editing. **Stefan Bielack:** Investigation, Writing - review and editing. **Catharina Dhooge:** Investigation, Writing - review and editing. **Lars Hjorth:** Investigation, Writing - review and editing. **Katherine A. Janeway:** Investigation, Writing - review and editing. **Meriel Jenney:** Investigation, Writing - review and editing. **Mark D. Krailo:** Investigation, Writing - review and editing. **Neyssa Marina:** Investigation, Writing - review and editing. **Rajaram Nagarajan:** Investigation, Writing - review and editing. **Sigbjørn Smeland:** Investigation, Writing - review and editing. **Matthew R. Sydes:** Investigation, Methodology, Writing - review and editing. **Patricia DeVos:** Investigation, Writing - review and editing. **Jeremy Whelan:** Investigation, Writing - review and editing. **Andreas Wiener:** Investigation, Writing - review and editing. **Gabriele Calaminus:** Conceptualisation, Investigation, Writing - review and editing, Supervision. **Matthias Schmid:** Conceptualisation, Methodology, Writing - review and editing, Supervision.

Conflict of interest statement

The authors declare the following financial interests/ personal relationships which may be considered as potential competing interests: SB reports grants from Deutsche Krebshilfe, Deutsche Forschungsgemeinschaft, and European Science Foundation during the conduct of the study and personal fees from Lilly, Bayer, Pfizer, Novartis, Isofol, Clinigen, Sensorion, Ipsen, and Roche outside the submitted work. MRS reports grants and nonfinancial support from Astellas, grants from Clovis, grants and nonfinancial support from Janssen, grants and nonfinancial support from Novartis, grants and nonfinancial support from Pfizer, and grants and nonfinancial support from Sanofi, during the conduct of the study and personal fees from Lilly Oncology and personal fees from

Janssen for educational courses and workshops outside the submitted work. NM reports employment by Five Prime Therapeutics, Inc and Sanofi US, outside the submitted work. The remaining authors declare no conflicts of interest.

Acknowledgements

The authors thank all the patients and parents for their contribution and all data managers, especially Ms Eva-Mari Olofsson from the SSG office in Lund, for their support to collect the data from all the patients in timely order.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ejca.2022.03.018>.

A. Software

We conducted all statistical analyses using version 4.1.0 of the *R* platform, version 2.0.7 of *R* package *equate* for score linking, *R* package *blandr* for the calculation of concordance correlation coefficients and the *tidyverse* suite of *R* packages for data preparation and data visualisation.

B. Characteristics of participants by domain, overall and linked.

Table B.1
Characteristics of PedsQL participants by domain, overall and linked.

Characteristics	PedsQL participants by domain											
	Functional								Symptom			
	Physical Functioning		Emotional Functioning		Cognitive Functioning		Social Functioning		Fatigue		Pain	
	Overall	Linked	Overall	Linked	Overall	Linked	Overall	Linked	Overall	Linked	Overall	Linked
Sex, n(%)												
Male	429 (56)	48 (64)	445 (57)	51 (64)	395 (57)	45 (64)	439 (56)	45 (64)	445 (57)	45 (64)	443 (57)	48 (62)
Female	331 (44)	27 (36)	339 (43)	29 (36)	296 (43)	25 (36)	339 (44)	25 (36)	341 (43)	25 (36)	341 (44)	29 (38)
Age (years)¹												
Age group, n(%)												
5 to 15	671 (88)	44 (59)	690 (88)	48 (60)	609 (88)	43 (61)	685 (88)	43 (61)	694 (88)	43 (61)	692 (88)	47 (61)
16 to 17	74 (10)	22 (29)	78 (10)	30 (29)	69 (10)	20 (29)	77 (10)	20 (29)	76 (10)	20 (29)	76 (10)	21 (27)
18 or older	15 (2)	9 (12)	16 (2)	7 (11)	13 (2)	9 (10)	16 (2)	7 (10)	16 (2)	7 (10)	16 (2)	9 (12)
Mean(SD)	12.8 (3.0)	15.1 (2.7)	12.8 (3.0)	15.1 (2.6)	12.8 (3.0)	15.0 (2.6)	12.8 (3.0)	15.0 (2.6)	12.8 (3.0)	15.0 (2.6)	12.8 (3.0)	15.1 (2.7)
Study group, n(%)												
COG	616 (81)	73 (97)	640 (82)	78 (98)	562 (81)	68 (97)	632 (81)	68 (97)	639 (81)	68 (97)	638 (81)	75 (97)
COSS	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
EOI	144 (19)	2 (3)	144 (18)	2 (2)	129 (19)	2 (3)	146 (19)	2 (3)	147 (19)	2 (3)	146 (19)	2 (3)
SSG	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

COG: Children's Oncology Group; COSS: Cooperative Osteosarcoma Group; EOI: European Osteosarcoma Intergroup; SSG: Scandinavian Sarcoma Group.

¹ Age refers to the age at the time of registration for participation in the study.

Table B.2
Characteristics of PEDQOL participants by domain, overall and linked.

Characteristics	PedsQL participants by domain											
	Functional								Symptom			
	Physical Functioning		Emotional Functioning		Cognitive Functioning		Social Functioning		Fatigue		Pain	
	Overall	Linked	Overall	Linked	Overall	Linked	Overall	Linked	Overall	Linked	Overall	Linked
Sex, n(%)												
Male	429 (56)	48 (64)	445 (57)	51 (64)	395 (57)	45 (64)	439 (56)	45 (64)	445 (57)	45 (64)	443 (57)	48 (62)
Female	331 (44)	27 (36)	339 (43)	29 (36)	296 (43)	25 (36)	339 (44)	25 (36)	341 (43)	25 (36)	341 (44)	29 (38)
Age (years)¹												
Age group, n(%)												
5 to 15	671 (88)	44 (59)	690 (88)	48 (60)	609 (88)	43 (61)	685 (88)	43 (61)	694 (88)	43 (61)	692 (88)	47 (61)
16 to 17	74 (10)	22 (29)	78 (10)	30 (29)	69 (10)	20 (29)	77 (10)	20 (29)	76 (10)	20 (29)	76 (10)	21 (27)
18 or older	15 (2)	9 (12)	16 (2)	7 (11)	13 (2)	9 (10)	16 (2)	7 (10)	16 (2)	7 (10)	16 (2)	9 (12)
Mean(SD)	12.8 (3.0)	15.1 (2.7)	12.8 (3.0)	15.1 (2.6)	12.8 (3.0)	15.0 (2.6)	12.8 (3.0)	15.0 (2.6)	12.8 (3.0)	15.0 (2.6)	12.8 (3.0)	15.1 (2.7)
Study group, n(%)												
COG	616 (81)	73 (97)	640 (82)	78 (98)	562 (81)	68 (97)	632 (81)	68 (97)	639 (81)	68 (97)	638 (81)	75 (97)
COSS	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
EOI	144 (19)	2 (3)	144 (18)	2 (2)	129 (19)	2 (3)	146 (19)	2 (3)	147 (19)	2 (3)	146 (19)	2 (3)
SSG	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

COG: Children's Oncology Group; COSS: Cooperative Osteosarcoma Group; EOI: European Osteosarcoma Intergroup; SSG: Scandinavian Sarcoma Group.

¹ Age refers to the age at the time of registration for participation in the study.

C. Internal consistency reliability of the three instruments by domain.

Table C.1
Internal consistency reliability of the *emotional functioning* aggregate scores of the three instruments.

Time point	Questionnaire	Linked to	N	Cronbach's α (95% CI)	Item–total correlation		
					Min	Mean	Max
E1	PedsQL	EORTC QLQ-C30	39	0.74 (0.61, 0.87)	0.36	0.62	0.82
	PEDQOL		47	0.56 (0.70, 0.84)	0.31	0.60	0.79
	EORTC QLQ-C30	PedsQL	39	0.72 (0.82, 0.91)	0.52	0.72	0.91
E2	PedsQL	EORTC QLQ-C30	29	0.72 (0.82, 0.93)	0.44	0.71	0.80
	PEDQOL		51	0.61 (0.73, 0.85)	0.37	0.63	0.77
	EORTC QLQ-C30	PedsQL	29	0.73 (0.82, 0.92)	0.56	0.73	0.85
E3	PedsQL	EORTC QLQ-C30	19	0.60 (0.77, 0.93)	0.55	0.65	0.79
	PEDQOL		43	0.35 (0.56, 0.77)	0.38	0.48	0.75
	EORTC QLQ-C30	PedsQL	19	0.60 (0.77, 0.95)	0.59	0.73	0.87
E4	PedsQL	EORTC QLQ-C30	18	0.66 (0.81, 0.95)	0.29	0.70	0.87
	PEDQOL		32	0.45 (0.65, 0.85)	0.39	0.56	0.76
	EORTC QLQ-C30	PedsQL	18	0.55 (0.73, 0.91)	0.37	0.63	0.77
		PEDQOL	32	0.65 (0.77, 0.90)	0.48	0.67	0.73

Table C.2

Internal consistency reliability of the *cognitive functioning* aggregate scores of the three instruments.

Time point	Questionnaire	Linked to	N	Cronbach's α (95% CI)		Item–total correlation		
						Min	Mean	Max
E1	PedsQL	EORTC QLQ-C30	33	0.47	(0.14, 0.80)	0.47	0.47	0.47
	PEDQOL		36	0.76	(0.63, 0.88)	0.41	0.63	0.76
	EORTC QLQ-C30	PedsQL	33	0.06	(-0.56, 0.68)	0.13	0.13	0.13
E2	PedsQL	EORTC QLQ-C30	18	0.56	(0.18, 0.94)	0.54	0.54	0.54
	PEDQOL		33	0.68	(0.50, 0.85)	0.36	0.54	0.68
	EORTC QLQ-C30	PedsQL	18	0.21	(-0.47, 0.88)	0.27	0.27	0.27
E3	PedsQL	EORTC QLQ-C30	19	0.80	(0.62, 0.97)	0.75	0.75	0.75
	PEDQOL		41	0.84	(0.77, 0.92)	0.53	0.72	0.81
	EORTC QLQ-C30	PedsQL	19	0.62	(0.36, 0.87)	0.66	0.66	0.66
E4	PedsQL	EORTC QLQ-C30	16	0.94	(0.89, 1.00)	0.92	0.92	0.92
	PEDQOL		32	0.74	(0.60, 0.88)	0.54	0.62	0.76
	EORTC QLQ-C30	PedsQL	16	0.04	(-0.88, 0.97)	0.11	0.11	0.11
		PEDQOL	41	0.33	(-0.07, 0.74)	0.69	0.69	0.69

Table C.3

Internal consistency reliability of the *social functioning* aggregate scores of the three instruments.

Time point	Questionnaire	Linked to	N	Cronbach's α (95% CI)		Item–total correlation		
						Min	Mean	Max
E1	PedsQL	EORTC QLQ-C30	38	0.34	(-0.01, 0.69)	0.17	0.42	0.65
	PEDQOL		47	0.24	(-0.09, 0.58)	0.05	0.30	0.53
	EORTC QLQ-C30	PedsQL	38	0.66	(0.45, 0.87)	0.61	0.61	0.61
E2	PedsQL	EORTC QLQ-C30	27	0.62	(0.41, 0.83)	0.39	0.57	0.67
	PEDQOL		40	-0.22	(-0.81, 0.38)	-0.33	0.20	0.81
	EORTC QLQ-C30	PedsQL	27	0.61	(0.32, 0.90)	0.57	0.57	0.57
E3	PedsQL	EORTC QLQ-C30	19	0.48	(0.14, 0.83)	-0.10	0.46	0.71
	PEDQOL		28	-0.27	(-1.05, 0.50)	-0.50	0.15	0.49
	EORTC QLQ-C30	PedsQL	19	0.79	(0.62, 0.96)	0.76	0.76	0.76
E4	PedsQL	EORTC QLQ-C30	17	0.60	(0.36, 0.85)	0.32	0.50	0.68
	PEDQOL		24	-0.02	(-0.68, 0.64)	-0.03	0.21	0.71
	EORTC QLQ-C30	PedsQL	17	0.78	(0.58, 0.98)	0.74	0.74	0.74
		PEDQOL	24	0.89	(0.81, 0.98)	0.85	0.85	0.85

Table C.4

Internal consistency reliability of the *fatigue* aggregate scores of the three instruments.

Time point	Questionnaire	Linked to	N	Cronbach's α (95% CI)		Item–total correlation		
						Min	Mean	Max
E1	PedsQL	EORTC QLQ-C30	40	–	–	–	–	–
	PEDQOL		50	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	40	0.82	(0.71, 0.93)	0.77	0.77	0.77
E2	PedsQL	EORTC QLQ-C30	30	–	–	–	–	–
	PEDQOL		54	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	30	0.64	(0.39, 0.90)	0.59	0.59	0.59
E3	PedsQL	EORTC QLQ-C30	19	–	–	–	–	–
	PEDQOL		45	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	19	0.87	(0.75, 0.99)	0.82	0.82	0.82
E4	PedsQL	EORTC QLQ-C30	17	–	–	–	–	–
			45	0.88	(0.81, 0.95)	0.83	0.83	0.83
			17	–	–	–	–	–

(continued on next page)

Table C.4 (continued)

Time point	Questionnaire	Linked to	N	Cronbach's α (95% CI)		Item–total correlation		
						Min	Mean	Max
	PEDQOL		32	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	17	0.73	(0.50, 0.97)	0.70	0.70	0.70
		PEDQOL	32	0.59	(0.35, 0.83)	0.59	0.59	0.59

Table C.5

Internal consistency reliability of the *pain* aggregate scores of the three instruments.

Time point	Questionnaire	Linked to	N	Cronbach's α (95% CI)		Item–total correlation		
						Min	Mean	Max
E1	PedsQL	EORTC QLQ-C30	40	–	–	–	–	–
	PEDQOL		50	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	40	0.82	(0.71, 0.93)	0.77	0.77	0.77
		PEDQOL	50	0.89	(0.83, 0.95)	0.85	0.85	0.85
E2	PedsQL	EORTC QLQ-C30	30	–	–	–	–	–
	PEDQOL		54	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	30	0.64	(0.39, 0.90)	0.59	0.59	0.59
		PEDQOL	54	0.85	(0.77, 0.93)	0.80	0.80	0.80
E3	PedsQL	EORTC QLQ-C30	19	–	–	–	–	–
	PEDQOL		45	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	19	0.87	(0.75, 0.99)	0.82	0.82	0.82
		PEDQOL	45	0.88	(0.81, 0.95)	0.83	0.83	0.83
E4	PedsQL	EORTC QLQ-C30	17	–	–	–	–	–
	PEDQOL		32	–	–	–	–	–
	EORTC QLQ-C30	PedsQL	17	0.73	(0.50, 0.97)	0.70	0.70	0.70
		PEDQOL	32	0.59	(0.35, 0.83)	0.59	0.59	0.59

D. Paediatric questionnaires and EORTC QLQ-C30: concordance measures.

Table D.1

Paediatric questionnaires and EORTC QLQ-C30: concordance measures.

Concordance Coefficient	Questionnaire	Domain					
		Functional				Symptom	
		Physical Functioning	Emotional Functioning	Cognitive Functioning	Social Functioning	Fatigue	Pain
Pearson's r (95% CI)	PedsQL	0.74 (0.64–0.82)	0.64 (0.52–0.74)	0.37 (0.17–0.54)	0.27 (0.07–0.44)	0.70 (0.58	0.59 (0.45
	PEDQOL	0.64 (0.54–0.72)	0.58 (0.47–0.67)	0.57 (0.45–0.67)	0.16 (–0.01	–0.78)	–0.70)
					–0.32)	0.72 (0.64	0.73 (0.67
						–0.78)	–0.78)
Lin's ρ (95% CI)	PedsQL	0.74 (0.64–0.82)	0.55 (0.42–0.65)	0.37 (0.17–0.54)	0.17 (0.05–0.29)	0.69 (0.58	0.58 (0.45
	PEDQOL	0.64 (0.54–0.72)	0.36 (0.27–0.44)	0.47 (0.35–0.58)	0.08 (–0.01	–0.78)	–0.69)
					–0.16)	0.71 (0.63	0.73 (0.65
						–0.77)	–0.79)

E. EURAMOS-1 consortium.Table E.1
EURAMOS-1 consortium.

Name	Surname	Affiliation Dept/Programme/Centre	Institution Name	City	Country
Sigbjørn	Smeland	Institute for Clinical Medicine	Oslo University Hospital	Oslo	NO
Stefan S	Bielack	Olgahospital Stuttgart	Klinikum Stuttgart	Stuttgart	DE
Jeremy	Whelan		University College Hospital	London	UK
Mark	Bernstein	IWK Health Center	Dalhousie University	Halifax, NS	CA
Kirsten	Sundby Hall	Institute for Clinical Medicine	Oslo University Hospital	Oslo	NO
Catherine	Rechnitzer	Rigshospitalet	University of Copenhagen	Copenhagen	DK
Mikael	Eriksson		Lund University	Lund	SE
Imre	Antal		Semmelweis University	Budapest	HU
Godehard	Friedel	Thoracic surgery	Klinik Schillerhöhe	Gerlingen	DE
Stefanie	Hecker-Nolting	Olgahospital Stuttgart	Klinikum Stuttgart	Stuttgart	DE
Edita	Kabickova		Motol University Hospital	Prague	CZ
Leo	Kager		St. Anna Kinderspital/CCRI	Vienna	AT
Thomas	Kühne		University Hospital Basel	Basel	CH
Susanna	Lang		Medical University of Vienna	Vienna	AT
Regine	Mayer-Steinacker		University Hospital Ulm	Ulm	DE
Peter	Reichardt		HELIOS Klinikum Berlin-Buch	Berlin	DE
Beate	Timmermann		University Hospital Essen	Essen	DE
Thekla	von Kalle	Olgahospital Stuttgart	Klinikum Stuttgart	Stuttgart	DE
Carola	Arndt		Mayo Clinic	Rochester, MN	US
Ching C	Lau		Baylor College of Medicine	Houston, TX	US
Cindy L	Schwartz	M D Anderson Cancer Center	University of Texas	Houston, TX	US
Douglas S	Hawkins		University of Washington	Seattle, WA	US
Holcombe E	Grier		Dana-Farber Cancer Institute	Boston, MA	US
Katherine A	Janeway		Dana-Farber Cancer Institute	Boston, MA	US
Ken L B	Brown		University of British Columbia	Vancouver, BC	CA
Leo	Mascarenhas	Keck School of Medicine	University of Southern California	Los Angeles, CA	US
Lisa	Teot		Boston Children's Hospital	Boston, MA	US
Mark C	Gebhardt		Dana-Farber Cancer Institute	Boston, MA	US
Mark D	Krailo		Children's Oncology Group	Arcadia, CA	US
Michael S	Isakoff		Connecticut Children's Medical Center	Hartford, CT	US
Patrick J	Leavey	Southwestern Children's Medical Center	University of Texas	Dallas, TX	US
Paul A	Meyers		MSKCC	New York, NY	US
R Lor	Randall	Primary Childrens Hospital	The University of Utah	SLC, UT	US
Raj	Nagarajan		Children's Hospital Medical Center	Cincinnati, OH	US
Richard	Gorlick	M D Anderson Cancer Center	The University of Texas	Houston, TX	US
Robert	Goldsby	Paediatric Oncology	UCSF Medical Center—Mission Bay	SF, CA	US
Stephen L	Lessnick		Nationwide Children's Hospital/OSU	Columbus, OH	US
Catherina	Dhooge		University Hospital Ghent	Ghent	BE
Michael	Capra		Our Lady's Children's Hospital	Dublin	IE
Jakob	Anninga				NL
Adrienne M	Flanagan	Cancer Institute	RNOH/UCL	Stanmore/London	UK
Robert	Grimer		Royal Orthopaedic Hospital	Birmingham	UK
Sandra	Strauss		University College Hospital	London	UK
Hans	Gelderblom		Leiden University Medical Center	Leiden	NL
Marleen	Renard		University Hospital Leuven	Leuven	BE
Fiona	Ingleby	MRC Clinical Trials Unit	University College London	London	UK
Gordana	Jovic	MRC Clinical Trials Unit	University College London	London	UK
Trude	Butterfaß-Bahloul		University Hospital Münster	Münster	DE
Gabriele	Calaminus	Paediatric Haematology and Oncology	University Hospital Bonn	Bonn	DE
Pancras	Hogendoorn		Leiden University Medical Center	Leiden	NL
Matthew R	Sydes	MRC Clinical Trials Unit	University College London	London	UK
Neyssa	Marina		Five Prime Therapeutics, Inc	South SF, CA	US

F. Physical functioning items per questionnaire.

F.1 PedsQL physical functioning items.

In the past **ONE month**, how much of a **problem** has this been for you ...

Table F.1

PedsQL physical functioning items.

ABOUT MY HEALTH AND ACTIVITIES (<i>problems with ...</i>)	Never	Almost Never	Sometimes	Often	Always
1. It is hard for me to walk more than one block.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. It is hard for me to run.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. It is hard for me to do sports activity or exercise.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. It is hard for me to lift something heavy.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. It is hard for me to take a bath or shower by myself.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. It is hard for me to do chores around the house.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. I hurt or ache.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. I have low energy	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

F.2 PEDQOL physical functioning items.

Table F.2

PEDQOL physical functioning items.

In der letzten Woche .../In the last week ...	Nie/ Never	Selten/ Rarely	Häufig/ Frequently	Immer/ Always
1.... konnte ich mit meinen Freunden beim Sport mithalten./... I was able to keep up with my friends in sports.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2.... habe ich beim Spielen und beim Sport lieber zugesehen als mitgespielt./ ... I watched rather than played in games and sports.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3.... habe ich mich stark gefühlt./... I felt strong.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4.... fühlte ich mich fit genug, um nach der Schule mit meinen Freunden zu spielen/ ... I felt fit enough to play with my friends after school.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

F.3 EORTC QLQ-C30 physical functioning items.

Table F.3

EORTC QLQ-C30 physical functioning items.

DURING THE PAST WEEK:	Not at All	A Little	Quite a Bit	Very Much
1. Were you short of breath?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Have you had pain?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Did you need to rest?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Have you had trouble sleeping?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Have you felt weak?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

G. Emotional functioning items per questionnaire.

G.1 PedsQL emotional functioning items.

In the past **ONE month**, how much of a **problem** has this been for you ...

Table G.1

PedsQL emotional functioning items.

ABOUT MY FEELINGS (<i>problems with ...</i>)	Never	Almost Never	Sometimes	Often	Always
1. I feel afraid or scared.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. I feel sad or blue.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. I feel angry.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. I have trouble sleeping.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. I worry about what will happen to me.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

G.2 PEDQOL emotional functioning items.

Table G.2

PEDQOL emotional functioning items.

In der letzten Woche .../In the last week ...	Nie/Never	Selten/Rarely	Häufig/Frequently	Immer/Always
1.... fühlte ich mich alleine./ ... I felt alone.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2.... war ich ärgerlich./ ... I have been annoyed.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3.... fühlte ich mich glücklich./ ... I felt happy.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4.... habe ich viel gelacht und Spaßgehabt./ ... I have laughed a lot and had fun.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

G.3 EORTC QLQ-C30 emotional functioning items.

Table G.3

EORTC QLQ-C30 emotional functioning items.

DURING THE PAST WEEK:	Not at All	A Little	Quite a Bit	Very Much
1. Did you feel tense?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Did you worry?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Did you feel irritable?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Did you feel depressed?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

H. Cognitive functioning items per questionnaire.

H.1 PedsQL cognitive functioning items.

In the past **ONE month**, how much of a **problem** has this been for you ...

Table H.1

PedsQL cognitive functioning items.

ABOUT SCHOOL (<i>problems with ...</i>)	Never	Almost Never	Sometimes	Often	Always
1. It is hard to pay attention in class.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. I forget things.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. I have trouble keeping up with my schoolwork.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. I miss school because of not feeling well.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. I miss school to go to the doctor or hospital.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

H.2 PEDQOL cognitive functioning items.

Table H.2
PEDQOL cognitive functioning items.

In der letzten Woche .../In the last week ...	Nie/ Never	Selten/ Rarely	Häufig/ Frequently	Immer/ Always
1.... fiel es mir leicht, neue Dinge zu lernen./ ... I found it easy to learn new things.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2.... fiel es mir schwer, mich zu konzentrieren./ ... I had a hard time concentrating.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3.... war ich genauso schlau wie alle anderen in der Klasse./ ... I was just as smart as everyone else in the class.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4.... konnte ich mir Sachen gut merken./ ... I've been able to remember things well.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5.... brauchte ich sehr lange, um meine Schularbeiten zu machen./ ... it took me a long time to do my schoolwork.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

H.3 EORTC QLQ-C30 cognitive functioning items.

Table H.3
EORTC QLQ-C30 cognitive functioning items.

DURING THE PAST WEEK:	Not at All	A Little	Quite a Bit	Very Much
1. Have you had difficulty concentrating on things, like reading a newspaper or watching television?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Have you had difficulty remembering things?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

I. Social functioning items per questionnaire.

I.1 PedsQL social functioning items.

In the past **ONE month**, how much of a **problem** has this been for you ...

Table I.1
PedsQL social functioning items.

How I Get Along with Others (problems with ...)	Never	Almost Never	Sometimes	Often	Always
1. I have trouble getting along with other kids.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Other kids do not want to be my friend.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Other kids tease me.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. I cannot do things other kids my age can do.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. It is hard to keep up when I play with other kids.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

I.2 PEDQOL social functioning items.

Table I.2
PEDQOL social functioning items.

In der letzten Woche .../In the last week ...	Nie/ Never	Selten/ Rarely	Häufig/ Frequently	Immer/ Always
1... fühlte ich mich in Gruppen von Gleichaltrigen ausgeschlossen./ ... I felt left out in groups of peers.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2... habe ich lieber was alleine gemacht./ ... I preferred to do something on my own.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3... konnte ich mit meinen Freunden über das reden, was mir wirklich Sorgen macht./ ... I was able to talk to my friends about what was really bothering me.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Zum Schluss möchten wir Dich bitten, die folgenden allgemeinen Sätze zu beantworten./Finally, we would like you to answer the following general sentences:				
4. Ich habe es leicht, Freunde zu finden./I have an easy time making friends.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. Ich bin beliebt bei meinen Freunden./I am popular with my friends.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

I.3 EORTC QLQ-C30 social functioning items.

Table I.3

EORTC QLQ-C30 social functioning items.

DURING THE PAST WEEK:	Not at All	A Little	Quite a Bit	Very Much
1. Has your physical condition or medical treatment interfered with your family life?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Has your physical condition or medical treatment interfered with your social activities?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

J. Fatigue items per questionnaire.*J.1 PedsQL fatigue items.**In the past ONE month, how much of a problem has this been for you ...*

Table J.1

PedsQL fatigue items.

ABOUT MY HEALTH AND ACTIVITIES (problems with ...)	Never	Almost Never	Sometimes	Often	Always
1. I have low energy.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

J.2 PEDQOL fatigue items.

Table J.2

PEDQOL fatigue items.

In der letzten Woche .../In the last week ...	Nie/Never	Selten/Rarely	Häufig/Frequently	Immer/Always
1.... fühlte ich mich schlapp und müde./ ... I have felt listless and tired.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

J.3 EORTC QLQ-C30 fatigue items.

Table J.3

EORTC QLQ-C30 fatigue items.

DURING THE PAST WEEK:	Not at All	A Little	Quite a Bit	Very Much
1. Did you need to rest?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Have you felt weak?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3. Have you felt tired?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

K. Pain items per questionnaire.*K.1 PedsQL pain items.**In the past ONE month, how much of a problem has this been for you ...*

Table K.1

PedsQL pain items.

ABOUT MY HEALTH AND ACTIVITIES (problems with ...)	Never	Almost Never	Sometimes	Often	Always
1. I hurt or ache.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

K.2 PEDQOL pain items.

Table K.2
PEDQOL pain items.

In der letzten Woche .../In the last week ...	Nie/Never	Selten/Rarely	Häufig/Frequently	Immer/Always
1.... hatte ich Schmerzen./ ... I've been in pain.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

K.3 EORTC QLQ-C30 pain items.

Table K.3
EORTC QLQ-C30 pain items.

DURING THE PAST WEEK:	Not at All	A Little	Quite a Bit	Very Much
1. Have you had pain?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2. Did pain interfere with your daily activities?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

L. Crosswalks between the PedsQL/the PEDQOL and the EORTC QLQ-C30.

Table L.1
Crosswalk for *physical functioning*.

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
0	20 (5.45)	7 (1.76)	50	73 (2.79)	73 (3.11)
1	20 (5.69)	13 (2.79)	51	73 (2.72)	80 (3.06)
2	20 (5.62)	13 (3.30)	52	73 (2.64)	80 (3.01)
3	20 (5.48)	13 (3.60)	53	80 (2.57)	80 (2.95)
4	20 (5.30)	13 (3.78)	54	80 (2.49)	80 (2.90)
5	20 (5.10)	13 (3.89)	55	80 (2.41)	80 (2.84)
6	20 (4.89)	13 (3.94)	56	87 (2.34)	80 (2.78)
7	20 (4.68)	13 (3.95)	57	87 (2.26)	80 (2.72)
8	20 (4.47)	20 (3.93)	58	87 (2.19)	80 (2.66)
9	27 (4.27)	20 (3.89)	59	87 (2.12)	80 (2.60)
10	30 (4.95)	20 (3.66)	60	87 (2.05)	80 (2.54)
11	30 (4.74)	20 (3.60)	61	87 (1.98)	80 (2.47)
12	33 (4.55)	20 (3.53)	62	87 (1.91)	80 (2.41)
13	33 (4.37)	20 (3.47)	63	87 (1.85)	80 (2.35)
14	33 (4.21)	20 (3.41)	64	87 (1.78)	80 (2.28)
15	33 (4.06)	20 (3.36)	65	87 (1.72)	80 (2.22)
16	33 (3.92)	20 (3.31)	66	87 (1.65)	80 (2.16)
17	33 (3.80)	27 (3.27)	67	93 (1.59)	87 (2.09)
18	33 (3.69)	33 (3.25)	68	93 (1.53)	93 (2.03)
19	33 (3.60)	33 (3.22)	69	93 (1.46)	93 (1.96)
20	40 (3.52)	33 (3.21)	70	93 (1.40)	93 (1.89)
21	40 (3.45)	33 (3.20)	71	93 (1.34)	93 (1.83)
22	44 (3.40)	33 (3.20)	72	93 (1.28)	93 (1.77)
23	47 (3.35)	33 (3.20)	73	93 (1.22)	93 (1.70)
24	47 (3.31)	33 (3.20)	74	93 (1.16)	93 (1.64)
25	47 (3.28)	40 (3.21)	75	100 (1.11)	93 (1.58)
26	47 (3.25)	44 (3.22)	76	100 (1.05)	96 (1.51)
27	47 (3.24)	44 (3.23)	77	100 (1.00)	96 (1.45)
28	53 (3.23)	44 (3.25)	78	100 (0.95)	96 (1.39)
29	53 (3.22)	44 (3.26)	79	100 (0.90)	96 (1.34)
30	53 (3.22)	44 (3.28)	80	100 (0.85)	96 (1.28)
31	53 (3.22)	44 (3.29)	81	100 (0.80)	96 (1.22)
32	56 (3.22)	44 (3.30)	82	100 (0.76)	96 (1.16)
33	56 (3.22)	47 (3.32)	83	100 (0.71)	100 (1.11)

Table L.1 (continued)

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
34	60 (3.23)	53 (3.33)	84	100 (0.67)	100 (1.05)
35	60 (3.24)	53 (3.34)	85	100 (0.63)	100 (1.00)
36	60 (3.24)	53 (3.35)	86	100 (0.59)	100 (0.95)
37	60 (3.24)	53 (3.36)	87	100 (0.55)	100 (0.89)
38	60 (3.24)	53 (3.36)	88	100 (0.51)	100 (0.84)
39	64 (3.24)	53 (3.36)	89	100 (0.47)	100 (0.78)
40	64 (3.23)	53 (3.36)	90	100 (0.43)	100 (0.73)
41	70 (3.22)	53 (3.35)	91	100 (0.40)	100 (0.67)
42	73 (3.20)	60 (3.35)	92	100 (0.36)	100 (0.61)
43	73 (3.17)	67 (3.33)	93	100 (0.32)	100 (0.55)
44	73 (3.14)	67 (3.31)	94	100 (0.28)	100 (0.49)
45	73 (3.09)	67 (3.29)	95	100 (0.24)	100 (0.42)
46	73 (3.05)	67 (3.26)	96	100 (0.20)	100 (0.35)
47	73 (2.99)	67 (3.23)	97	100 (0.16)	100 (0.28)
48	73 (2.93)	67 (3.19)	98	100 (0.12)	100 (0.21)
49	73 (2.86)	67 (3.15)	99	100 (0.07)	100 (0.13)
			100	100 (0.02)	100 (0.04)

Table L.2

Crosswalk for *emotional functioning*.

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
0	0 (4.04)	0 (1.33)	50	58 (4.47)	42 (2.82)
1	0 (4.74)	0 (1.94)	51	62 (4.62)	42 (2.79)
2	0 (5.16)	0 (2.32)	52	62 (4.76)	42 (2.77)
3	0 (5.48)	0 (2.61)	53	62 (4.86)	42 (2.75)
4	0 (5.75)	0 (2.85)	54	62 (4.92)	42 (2.73)
5	0 (5.99)	0 (3.07)	55	67 (4.92)	42 (2.72)
6	0 (6.21)	0 (3.28)	56	67 (4.86)	42 (2.71)
7	0 (6.41)	0 (3.47)	57	67 (4.74)	42 (2.71)
8	0 (6.59)	0 (3.65)	58	67 (4.58)	50 (2.71)
9	0 (6.75)	0 (3.82)	59	67 (4.38)	58 (2.72)
10	0 (6.90)	0 (3.99)	60	75 (4.15)	58 (2.72)
11	0 (7.02)	0 (4.16)	61	83 (3.91)	58 (2.74)
12	0 (7.13)	0 (4.33)	62	83 (3.67)	58 (2.76)
13	0 (7.21)	0 (4.49)	63	83 (3.43)	58 (2.78)
14	0 (7.26)	0 (4.64)	64	83 (3.21)	58 (2.80)
15	17 (7.29)	0 (4.79)	65	83 (2.99)	58 (2.82)
16	17 (7.28)	0 (4.93)	66	83 (2.80)	58 (2.84)
17	17 (7.24)	0 (5.07)	67	83 (2.62)	58 (2.86)
18	17 (7.17)	16 (5.19)	68	83 (2.45)	67 (2.87)
19	17 (7.06)	16 (5.30)	69	83 (2.31)	67 (2.87)
20	21 (6.93)	16 (5.39)	70	83 (2.17)	67 (2.86)
21	29 (6.76)	16 (5.47)	71	92 (2.05)	67 (2.83)
22	29 (6.56)	16 (5.53)	72	92 (1.93)	67 (2.79)
23	29 (6.34)	16 (5.57)	73	92 (1.83)	67 (2.73)
24	29 (6.10)	16 (5.59)	74	92 (1.74)	67 (2.66)
25	29 (5.84)	17 (5.59)	75	92 (1.65)	83 (2.58)
26	29 (5.58)	17 (5.56)	76	92 (1.56)	83 (2.48)
27	29 (5.31)	17 (5.51)	77	92 (1.49)	83 (2.38)
28	29 (5.05)	17 (5.44)	78	92 (1.41)	83 (2.28)
29	29 (4.80)	17 (5.35)	79	92 (1.34)	83 (2.17)
30	33 (4.56)	17 (5.23)	80	100 (1.28)	83 (2.06)
31	33 (4.35)	17 (5.10)	81	100 (1.22)	83 (1.96)
32	33 (4.15)	17 (4.95)	82	100 (1.16)	83 (1.85)
33	33 (3.98)	17 (4.79)	83	100 (1.10)	92 (1.75)

(continued on next page)

Table L.2 (continued)

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
34	33 (3.84)	21 (4.62)	84	100 (1.04)	92 (1.65)
35	33 (3.72)	21 (4.45)	85	100 (0.99)	92 (1.56)
36	33 (3.63)	21 (4.27)	86	100 (0.94)	92 (1.47)
37	33 (3.55)	21 (4.10)	87	100 (0.89)	92 (1.37)
38	33 (3.51)	21 (3.93)	88	100 (0.84)	92 (1.28)
39	33 (3.48)	21 (3.78)	89	100 (0.79)	92 (1.19)
40	42 (3.47)	21 (3.63)	90	100 (0.74)	92 (1.10)
41	50 (3.49)	21 (3.49)	91	100 (0.69)	92 (1.01)
42	50 (3.52)	33 (3.37)	92	100 (0.64)	100 (0.92)
43	50 (3.57)	33 (3.26)	93	100 (0.59)	100 (0.82)
44	50 (3.65)	33 (3.17)	94	100 (0.54)	100 (0.73)
45	50 (3.74)	33 (3.08)	95	100 (0.48)	100 (0.63)
46	58 (3.86)	33 (3.01)	96	100 (0.41)	100 (0.52)
47	58 (3.99)	33 (2.95)	97	100 (0.34)	100 (0.42)
48	58 (4.14)	33 (2.90)	98	100 (0.25)	100 (0.30)
49	58 (4.30)	33 (2.86)	99	100 (0.16)	100 (0.18)
			100	100 (0.06)	100 (0.06)

Table L.3

Crosswalk for *cognitive functioning*.

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
0	0 (10.88)	0 (5.96)	50	83 (5.84)	67 (5.60)
1	0 (11.03)	16 (6.55)	51	83 (5.40)	67 (5.73)
2	0 (11.02)	16 (6.77)	52	83 (4.96)	67 (5.83)
3	0 (10.96)	16 (6.87)	53	83 (4.54)	67 (5.88)
4	0 (10.87)	16 (6.91)	54	83 (4.13)	67 (5.88)
5	0 (10.75)	16 (6.90)	55	83 (3.76)	67 (5.82)
6	0 (10.60)	16 (6.86)	56	83 (3.42)	67 (5.70)
7	0 (10.42)	17 (6.80)	57	83 (3.11)	67 (5.50)
8	0 (10.22)	17 (6.71)	58	83 (2.85)	67 (5.25)
9	0 (9.99)	17 (6.61)	59	83 (2.61)	67 (4.94)
10	0 (9.75)	17 (6.49)	60	83 (2.40)	67 (4.60)
11	0 (9.47)	17 (6.37)	61	83 (2.22)	83 (4.25)
12	0 (9.18)	17 (6.23)	62	83 (2.07)	83 (3.88)
13	0 (8.86)	17 (6.09)	63	83 (1.93)	83 (3.53)
14	0 (8.54)	17 (5.95)	64	83 (1.81)	83 (3.19)
15	0 (8.22)	17 (5.80)	65	83 (1.71)	83 (2.88)
16	0 (7.89)	17 (5.66)	66	83 (1.61)	83 (2.60)
17	0 (7.54)	17 (5.51)	67	83 (1.53)	83 (2.34)
18	0 (7.19)	17 (5.37)	68	83 (1.45)	83 (2.12)
19	0 (6.85)	17 (5.23)	69	83 (1.39)	83 (1.92)
20	0 (6.53)	17 (5.09)	70	83 (1.32)	83 (1.75)
21	0 (6.23)	25 (4.96)	71	83 (1.26)	83 (1.59)
22	0 (5.97)	25 (4.84)	72	83 (1.21)	83 (1.46)
23	0 (5.72)	25 (4.72)	73	83 (1.16)	100 (1.34)
24	0 (5.51)	25 (4.62)	74	83 (1.11)	100 (1.23)
25	42 (5.33)	25 (4.52)	75	100 (1.07)	100 (1.13)
26	50 (5.16)	25 (4.44)	76	100 (1.03)	100 (1.05)
27	50 (5.03)	33 (4.36)	77	100 (0.99)	100 (0.97)
28	50 (4.92)	33 (4.30)	78	100 (0.95)	100 (0.89)
29	50 (4.84)	33 (4.24)	79	100 (0.91)	100 (0.83)
30	50 (4.78)	33 (4.20)	80	100 (0.87)	100 (0.77)
31	50 (4.76)	33 (4.17)	81	100 (0.84)	100 (0.71)
32	50 (4.77)	33 (4.15)	82	100 (0.80)	100 (0.66)
33	50 (4.81)	33 (4.14)	783	100 (0.77)	100 (0.61)
34	50 (4.90)	42 (4.14)	84	100 (0.73)	100 (0.56)

Table L.3 (continued)

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30- equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30- equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
35	50 (5.03)	42 (4.15)	85	100 (0.70)	100 (0.52)
36	50 (5.20)	42 (4.17)	86	100 (0.67)	100 (0.48)
37	50 (5.42)	42 (4.20)	87	100 (0.64)	100 (0.44)
38	67 (5.68)	42 (4.24)	88	100 (0.60)	100 (0.40)
39	67 (5.96)	42 (4.30)	89	100 (0.57)	100 (0.37)
40	67 (6.26)	50 (4.36)	90	100 (0.54)	100 (0.34)
41	67 (6.56)	50 (4.44)	91	100 (0.50)	100 (0.31)
42	67 (6.82)	50 (4.53)	92	100 (0.47)	100 (0.28)
43	67 (7.03)	50 (4.63)	93	100 (0.44)	100 (0.24)
44	67 (7.15)	50 (4.74)	94	100 (0.40)	100 (0.21)
45	67 (7.17)	50 (4.87)	95	100 (0.35)	100 (0.18)
46	67 (7.08)	50 (5.00)	96	100 (0.30)	100 (0.15)
47	67 (6.88)	67 (5.15)	97	100 (0.25)	100 (0.12)
48	67 (6.60)	67 (5.30)	98	100 (0.19)	100 (0.09)
49	67 (6.24)	67 (5.46)	99	100 (0.12)	100 (0.06)
			100	100 (0.05)	100 (0.02)

Table L.4
Crosswalk for social functioning.

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30- equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30- equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
0	0 (0.1)	0 (8e-04)	50	0 (4.5)	0 (1e+00)
1	0 (0.2)	0 (2e-03)	51	0 (4.6)	0 (1e+00)
2	0 (0.2)	0 (3e-03)	52	0 (4.7)	0 (2e+00)
3	0 (0.3)	0 (4e-03)	53	0 (4.7)	0 (2e+00)
4	0 (0.3)	0 (4e-03)	54	0 (4.7)	0 (2e+00)
5	0 (0.3)	0 (5e-03)	55	17 (4.7)	0 (2e+00)
6	0 (0.3)	0 (5e-03)	56	17 (4.7)	0 (2e+00)
7	0 (0.3)	0 (6e-03)	57	17 (4.6)	0 (3e+00)
8	0 (0.3)	0 (6e-03)	58	17 (4.6)	0 (3e+00)
9	0 (0.4)	0 (6e-03)	59	17 (4.5)	0 (3e+00)
10	0 (0.4)	0 (7e-03)	60	33 (4.5)	0 (3e+00)
11	0 (0.4)	0 (7e-03)	61	33 (4.5)	0 (3e+00)
12	0 (0.4)	0 (7e-03)	62	33 (4.5)	0 (3e+00)
13	0 (0.4)	0 (8e-03)	63	33 (4.5)	0 (4e+00)
14	0 (0.4)	0 (8e-03)	64	33 (4.6)	0 (4e+00)
15	0 (0.4)	0 (9e-03)	65	33 (4.7)	0 (4e+00)
16	0 (0.4)	0 (9e-03)	66	33 (4.8)	0 (4e+00)
17	0 (0.4)	0 (1e-02)	67	33 (4.9)	17 (4e+00)
18	0 (0.4)	0 (1e-02)	68	33 (5.1)	17 (5e+00)
19	0 (0.5)	0 (1e-02)	69	33 (5.4)	17 (5e+00)
20	0 (0.5)	0 (1e-02)	70	50 (5.7)	17 (5e+00)
21	0 (0.5)	0 (1e-02)	71	50 (6.0)	17 (5e+00)
22	0 (0.5)	0 (1e-02)	72	50 (6.3)	17 (5e+00)
23	0 (0.5)	0 (2e-02)	73	50 (6.7)	33 (6e+00)
24	0 (0.6)	0 (2e-02)	74	50 (7.1)	50 (6e+00)
25	0 (0.6)	0 (2e-02)	75	50 (7.4)	50 (6e+00)
26	0 (0.7)	0 (2e-02)	76	67 (7.7)	50 (6e+00)
27	0 (0.7)	0 (3e-02)	77	67 (7.8)	50 (7e+00)
28	0 (0.8)	0 (3e-02)	78	67 (7.8)	50 (7e+00)
29	0 (0.8)	0 (4e-02)	79	67 (7.6)	50 (7e+00)
30	0 (0.9)	0 (4e-02)	80	67 (7.3)	50 (7e+00)
31	0 (1.0)	0 (5e-02)	81	75 (6.9)	67 (7e+00)
32	0 (1.1)	0 (6e-02)	82	75 (6.3)	67 (8e+00)
33	0 (1.2)	0 (8e-02)	83	75 (5.7)	67 (8e+00)
34	0 (1.3)	0 (9e-02)	84	75 (5.0)	67 (8e+00)

(continued on next page)

Table L.4 (continued)

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
35	0 (1.5)	0 (1e-01)	85	83 (4.4)	67 (8e+00)
36	0 (1.6)	0 (1e-01)	86	100 (3.8)	67 (8e+00)
37	0 (1.8)	0 (2e-01)	87	100 (3.3)	67 (8e+00)
38	0 (2.0)	0 (2e-01)	88	100 (2.8)	83 (7e+00)
39	0 (2.2)	0 (2e-01)	89	100 (2.5)	83 (7e+00)
40	0 (2.4)	0 (3e-01)	90	100 (2.1)	83 (7e+00)
41	0 (2.6)	0 (3e-01)	91	100 (1.8)	83 (6e+00)
42	0 (2.9)	0 (4e-01)	92	100 (1.6)	83 (6e+00)
43	0 (3.1)	0 (5e-01)	93	100 (1.4)	83 (5e+00)
44	0 (3.4)	0 (6e-01)	94	100 (1.2)	83 (4e+00)
45	0 (3.6)	0 (7e-01)	95	100 (1.0)	83 (4e+00)
46	0 (3.8)	0 (8e-01)	96	100 (0.8)	83 (3e+00)
47	0 (4.1)	0 (9e-01)	97	100 (0.6)	83 (3e+00)
48	0 (4.2)	0 (1e+00)	98	100 (0.5)	83 (2e+00)
49	0 (4.4)	0 (1e+00)	99	100 (0.3)	83 (1e+00)
			100	100 (0.1)	100 (4e-01)

Table L.5

Crosswalk for *fatigue*.

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
0	0 (0.2)	11 (0.7)	50	33 (4.0)	56 (2.7)
1	10 (0.4)	22 (1.4)	51	56 (4.1)	56 (2.7)
2	10 (0.7)	22 (1.8)	52	56 (4.1)	56 (2.8)
3	10 (0.9)	22 (2.0)	53	56 (4.1)	56 (2.8)
4	10 (1.1)	22 (2.1)	54	56 (4.1)	56 (2.8)
5	10 (1.2)	22 (2.2)	55	56 (4.1)	56 (2.9)
6	10 (1.3)	22 (2.2)	56	56 (4.1)	56 (2.9)
7	10 (1.5)	22 (2.2)	57	56 (4.1)	56 (2.9)
8	10 (1.6)	22 (2.2)	58	56 (4.1)	56 (3.0)
9	10 (1.7)	22 (2.2)	59	56 (4.1)	56 (3.0)
10	10 (1.8)	22 (2.3)	60	56 (4.1)	56 (3.0)
11	10 (1.9)	22 (2.3)	61	56 (4.0)	56 (3.1)
12	10 (1.9)	22 (2.3)	62	56 (4.0)	56 (3.1)
13	10 (2.0)	22 (2.3)	63	56 (4.0)	56 (3.1)
14	10 (2.1)	22 (2.3)	64	56 (4.0)	56 (3.1)
15	10 (2.2)	22 (2.3)	65	56 (4.0)	56 (3.2)
16	10 (2.2)	22 (2.3)	66	56 (3.9)	56 (3.2)
17	10 (2.3)	22 (2.3)	67	56 (3.9)	67 (3.2)
18	10 (2.4)	22 (2.3)	68	56 (3.9)	94 (3.2)
19	10 (2.4)	22 (2.3)	69	56 (3.9)	94 (3.2)
20	10 (2.5)	22 (2.3)	70	56 (3.9)	94 (3.2)
21	10 (2.6)	22 (2.3)	71	56 (3.8)	94 (3.2)
22	10 (2.6)	22 (2.3)	72	56 (3.8)	94 (3.2)
23	10 (2.7)	22 (2.4)	73	56 (3.8)	94 (3.2)
24	10 (2.8)	22 (2.4)	74	56 (3.8)	94 (3.2)
25	11 (2.8)	22 (2.4)	75	67 (3.8)	94 (3.1)
26	22 (2.9)	22 (2.4)	76	78 (3.8)	94 (3.1)
27	22 (3.0)	22 (2.4)	77	78 (3.8)	94 (3.1)
28	22 (3.1)	22 (2.4)	78	78 (3.9)	94 (3.1)
29	22 (3.1)	22 (2.4)	79	78 (3.9)	94 (3.0)
30	22 (3.2)	22 (2.4)	80	78 (3.9)	94 (3.0)
31	22 (3.3)	22 (2.4)	81	78 (4.0)	94 (2.9)
32	22 (3.3)	22 (2.4)	82	78 (4.0)	94 (2.8)
33	22 (3.4)	33 (2.5)	83	78 (4.1)	94 (2.8)
34	22 (3.4)	56 (2.5)	84	78 (4.2)	94 (2.7)

Table L.5 (continued)

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
35	22 (3.5)	56 (2.5)	85	78 (4.3)	94 (2.6)
36	22 (3.6)	56 (2.5)	86	78 (4.3)	94 (2.4)
37	22 (3.6)	56 (2.5)	87	78 (4.4)	94 (2.3)
38	22 (3.7)	56 (2.5)	88	78 (4.5)	94 (2.2)
39	22 (3.7)	56 (2.5)	89	78 (4.6)	94 (2.0)
40	22 (3.8)	56 (2.5)	90	78 (4.6)	94 (1.9)
41	22 (3.8)	56 (2.5)	91	78 (4.7)	94 (1.7)
42	22 (3.8)	56 (2.5)	92	78 (4.7)	94 (1.5)
43	22 (3.9)	56 (2.5)	93	78 (4.6)	94 (1.4)
44	22 (3.9)	56 (2.5)	94	78 (4.6)	94 (1.2)
45	22 (3.9)	56 (2.6)	95	78 (4.5)	94 (1.0)
46	22 (4.0)	56 (2.6)	96	78 (4.3)	94 (0.8)
47	22 (4.0)	56 (2.6)	97	78 (4.1)	94 (0.7)
48	22 (4.0)	56 (2.6)	98	78 (3.7)	94 (0.5)
49	22 (4.0)	56 (2.7)	99	78 (3.1)	94 (0.3)
			100	89 (2.1)	100 (0.1)

Table L.6
Crosswalk for *pain*.

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30-equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
0	0 (0.09)	0 (0.07)	50	33 (5.90)	50 (5.84)
1	0 (0.24)	0 (0.20)	51	67 (6.06)	50 (5.94)
2	0 (0.39)	0 (0.32)	52	67 (6.21)	50 (6.01)
3	0 (0.52)	0 (0.42)	53	67 (6.34)	50 (6.04)
4	0 (0.64)	0 (0.52)	54	67 (6.46)	50 (6.04)
5	0 (0.74)	0 (0.60)	55	67 (6.57)	50 (5.99)
6	0 (0.84)	0 (0.68)	56	67 (6.65)	50 (5.92)
7	0 (0.93)	0 (0.76)	57	67 (6.72)	50 (5.81)
8	0 (1.02)	0 (0.83)	58	67 (6.76)	50 (5.68)
9	0 (1.10)	0 (0.89)	59	67 (6.78)	50 (5.53)
10	0 (1.17)	0 (0.96)	60	67 (6.78)	50 (5.37)
11	0 (1.25)	0 (1.02)	61	67 (6.76)	50 (5.20)
12	0 (1.32)	0 (1.08)	62	67 (6.72)	50 (5.02)
13	0 (1.39)	0 (1.14)	63	67 (6.67)	50 (4.86)
14	0 (1.46)	0 (1.21)	64	67 (6.60)	50 (4.69)
15	0 (1.53)	0 (1.27)	65	67 (6.52)	50 (4.53)
16	0 (1.61)	0 (1.34)	66	67 (6.44)	50 (4.39)
17	0 (1.68)	0 (1.40)	67	67 (6.35)	67 (4.25)
18	0 (1.75)	0 (1.47)	68	67 (6.26)	100 (4.12)
19	0 (1.83)	0 (1.54)	69	67 (6.16)	100 (4.00)
20	0 (1.91)	0 (1.62)	70	67 (6.07)	100 (3.89)
21	0 (1.99)	0 (1.70)	71	67 (5.98)	100 (3.80)
22	0 (2.07)	0 (1.77)	72	67 (5.89)	100 (3.72)
23	0 (2.16)	0 (1.86)	73	67 (5.81)	100 (3.65)
24	0 (2.25)	0 (1.94)	74	67 (5.74)	100 (3.59)
25	17 (2.34)	0 (2.03)	75	67 (5.66)	100 (3.53)
26	17 (2.44)	0 (2.13)	76	92 (5.59)	100 (3.49)
27	17 (2.54)	0 (2.22)	77	92 (5.52)	100 (3.45)
28	17 (2.64)	0 (2.33)	78	92 (5.46)	100 (3.41)
29	17 (2.75)	0 (2.43)	79	92 (5.39)	100 (3.37)
30	17 (2.86)	0 (2.54)	80	92 (5.32)	100 (3.33)
31	17 (2.98)	0 (2.66)	81	92 (5.24)	100 (3.28)
32	17 (3.10)	0 (2.78)	82	92 (5.16)	100 (3.23)
33	17 (3.22)	17 (2.91)	83	92 (5.07)	100 (3.16)
34	17 (3.35)	50 (3.05)	84	92 (4.97)	100 (3.08)

(continued on next page)

Table L.6 (continued)

Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30 equivalent score by linked questionnaire (Bootstrap SE)		Original PedsQL score/PEDQOL score	Estimated EORTC QLQ-C30 equivalent score by linked questionnaire (Bootstrap SE)	
	PedsQL	PEDQOL		PedsQL	PEDQOL
35	17 (3.49)	50 (3.19)	85	92 (4.87)	100 (3.00)
36	17 (3.63)	50 (3.34)	86	92 (4.75)	100 (2.89)
37	17 (3.77)	50 (3.50)	87	92 (4.61)	100 (2.78)
38	17 (3.92)	50 (3.67)	88	92 (4.47)	100 (2.65)
39	17 (4.07)	50 (3.84)	89	92 (4.30)	100 (2.51)
40	17 (4.22)	50 (4.02)	90	92 (4.12)	100 (2.36)
41	17 (4.38)	50 (4.21)	91	92 (3.93)	100 (2.19)
42	17 (4.55)	50 (4.40)	92	92 (3.71)	100 (2.01)
43	17 (4.71)	50 (4.60)	93	92 (3.47)	100 (1.83)
44	17 (4.88)	50 (4.80)	94	92 (3.21)	100 (1.63)
45	17 (5.05)	50 (5.00)	95	92 (2.92)	100 (1.42)
46	17 (5.22)	50 (5.20)	96	92 (2.61)	100 (1.21)
47	17 (5.40)	50 (5.38)	97	92 (2.25)	100 (0.98)
48	17 (5.57)	50 (5.56)	98	92 (1.83)	100 (0.74)
49	17 (5.74)	50 (5.71)	99	92 (1.34)	100 (0.48)
			100	100 (0.66)	100 (0.18)

²Score differences are defined as the paediatric instrument as less than the adult instrument.

³The US-English versions of are displayed here in an exemplary fashion.

References

- [1] Dorans Neil J, Holland Paul W. Population invariance and the equatability of tests: basic theory and the linear case. *J Educ Meas* 2000;37(4):281–306. <https://doi.org/10.1111/j.1745-3984.2000.tb01088.x>. ISSN 1745–3984, <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1745-3984.2000.tb01088.x>.
- [2] Neil J. Dorans. Linking scores from multiple health outcome instruments. *Qual Life Res : Int J Qual Life Asp Treat Care Rehabil* 2007;16(Suppl 1):85–94. <https://doi.org/10.1007/s11136-006-9155-3>. ISSN 0962–9343.
- [3] Marrie Ruth Ann, Dufault Brenden, Tyry Tuula, Cutter Gary R, Fox Robert J, Salter Amber. Developing a crosswalk between the rand-12 and the health utilities index for multiple sclerosis. *Mult Scler* 2020;26(9):1102–10. <https://doi.org/10.1177/1352458519852722>.
- [4] Shaw Bronwen E, Syrjala Karen L, Onstad Lynn E, Chow Eric J, Flowers Mary E, Jim Heather, Scott Baker K, Buckley Sarah, Fairclough Diane L, Horowitz Mary M, Lee Stephanie J. Promis measures can be used to assess symptoms and function in long-term hematopoietic cell transplantation survivors. *Cancer* 2018;124(4):841–9. <https://doi.org/10.1002/cncr.31089>. ISSN 0008–543X.
- [5] Kaat Aaron J, Schalet Benjamin D, Rutsohn Joshua, Jensen Roxanne E, Cella David. Physical function metric over measure: an illustration with the patient-reported outcomes measurement information system (promis) and the functional assessment of cancer therapy (fact). *Cancer* 2018;124(1):153–60. <https://doi.org/10.1002/cncr.30981>. ISSN 0008–543X.
- [6] McHorney CA, Cohen AS. Equating health status measures with item response theory: illustrations with functional status items. *Med Care* 2000;38(9 Suppl):II43–59. <https://doi.org/10.1097/00005650-200009002-00008>. ISSN 257079.
- [7] Schalet Benjamin D, Rothrock Nan E, Hays Ron D, Kazis Lewis E, Cook Karon F, Rutsohn Joshua P, Cella David. Linking physical and mental health summary scores from the veterans rand 12-item health survey (vr-12) to the promis® global health scale. *J Gen Intern Med* 2015;30(10):1524–30. <https://doi.org/10.1007/s11606-015-3453-9>. ISSN 1525–1497, <https://link.springer.com/content/pdf/10.1007/s11606-015-3453-9.pdf>.
- [8] Haley Stephen M, Ni Pengsheng, Lai Jin-shei, Tian Feng, Coster Wendy J, Jette Alan M, Straub Donald, Cella David. Linking the activity measure for post-acute care and the quality of life outcomes in neurological disorders. *Arch Phys Med Rehabil* 2011;92(10 Suppl):S37–43. <https://doi.org/10.1016/j.apmr.2011.01.026>. ISSN 0003–9993.
- [9] Schalet Benjamin D, Revicki Dennis A, Cook Karon F, Krishnan Eswar, Fries Jim F, Cella David. Establishing a common metric for physical function: linking the haq-di and sf-36 pf subscale to promis circledR physical function. *J Gen Intern Med* 2015;30(10):1517–23. <https://doi.org/10.1007/s11606-015-3360-0>. ISSN 1525–1497, <https://link.springer.com/content/pdf/10.1007/s11606-015-3360-0.pdf>.
- [10] Måsse Louise C, Allen Diane, Wilson Mark, Williams Geoffrey. Introducing equating methodologies to compare test scores from two different self-regulation scales. *Health Educ Res* 2006;21(suppl_1):i110–20. <https://doi.org/10.1093/her/cyl088>. ISSN 0268–1153, https://academic.oup.com/her/article-pdf/21/suppl_1/i110/1470555/cyl088.pdf.
- [11] Kaat Aaron J, Newcomb Michael E, Ryan Daniel T, Mustanski Brian. Expanding a common metric for depression reporting: linking two scales to promis(r) depression. *Qual Life Res : Int J Qual Life Asp Treat Care Rehabil* 2017;26(5):1119–28. <https://doi.org/10.1007/s11136-016-1450-z>. ISSN 0962–9343.
- [12] Choi Seung W, Schalet Benjamin, Cook Karon F, Cella David. Establishing a common metric for depressive symptoms: linking the bdi-ii, ces-d, and phq-9 to promis depression. *Psychol Assess* 2014;26(2):513–27. <https://doi.org/10.1037/a0035768>. ISSN 1040–3590.
- [13] Tulsky David S, Kisala Pamela A, Kalpakjian Claire Z, Bombardier Charles H, Pohlign Ryan T, Heinemann Allen W, Adam Carle, Choi Seung W. Measuring depression after spinal cord injury: development and psychometric characteristics of the sci-qol depression item bank and linkage with phq-9. *J Spinal Cord Med* 2015;38(3):335–46. <https://doi.org/10.1179/2045772315Y.0000000020>. ISSN 1079–0268.
- [14] Olin Thomas M, Lan Yu, McMakin Dana L, Forbes Erika E, Seeley John R, Lewinsohn Peter M, Pilkonis Paul A. Comparisons across depression assessment instruments in adolescence and young adulthood: an item response theory study using two linking methods. *J Abnorm Child Psychol* 2013;41(8):1267–77. <https://doi.org/10.1007/s11802-013-0000-0>.

- <https://doi.org/10.1007/s10802-013-9756-6>. ISSN 1573–2835, <https://link.springer.com/content/pdf/10.1007/s10802-013-9756-6.pdf>.
- [15] Orlando M, Sherbourne CD, Thissen D. Summed-score linking using item response theory: application to depression measurement. *Psychol Assess* 2000;12(3):354–9. <https://doi.org/10.1037/1040-3590.12.3.354>. ISSN 1040–3590.
- [16] Cook Karon F, Schalet Benjamin D, Kallen Michael A, Rutsohn Joshua P, Cella David. Establishing a common metric for self-reported pain: linking bpi pain interference and sf-36 bodily pain subscale scores to the promis pain interference metric. *Qual Life Res* 2015;24(10):2305–18. <https://doi.org/10.1007/s11136-015-0987-6>. ISSN 1573–2649, <https://link.springer.com/content/pdf/10.1007/s11136-015-0987-6.pdf>.
- [17] Askew Robert L, Kim Jiseon, Chung Hyewon, Cook Karon F, Johnson Kurt L, Amtmann Dagmar. Development of a crosswalk for pain interference measured by the bpi and promis pain interference short form. *Qual Life Res* 2013;22(10):2769–76. <https://doi.org/10.1007/s11136-013-0398-5>. ISSN 1573–2649.
- [18] Kisala Pamela A, Tulsy David S, Kalpakjian Claire Z, Heinemann Allen W, Pohlig Ryan T, Adam Carle, Choi Seung W. Measuring anxiety after spinal cord injury: development and psychometric characteristics of the sci-qol anxiety item bank and linkage with gad-7. *J Spinal Cord Med* 2015;38(3):315–25. <https://doi.org/10.1179/2045772315Y.0000000029>. ISSN 1079–0268.
- [19] Schalet Benjamin D, Cook Karon F, Choi Seung W, Cella David. Establishing a common metric for self-reported anxiety: linking the masq, panas, and gad-7 to promis anxiety. *J Anxiety Disord* 2014;28(1):88–96. <https://doi.org/10.1016/j.janxdis.2013.11.006>. ISSN 0887–6185.
- [20] Lai Jin-shei, Cella David, Yanez Betina, Stone Arthur. Linking fatigue measures on a common reporting metric. *J Pain Symptom Manag* 2014;48(4):639–48. <https://doi.org/10.1016/j.jpainsymman.2013.12.236>. ISSN 0885–3924, <http://www.sciencedirect.com/science/article/pii/S0885392414001444>.
- [21] Noonan Vanessa K, Cook Karon F, Bamer Alyssa M, Choi Seung W, Kim Jiseon, Amtmann Dagmar. Measuring fatigue in persons with multiple sclerosis: creating a crosswalk between the modified fatigue impact scale and the promis fatigue short form. *Qual Life Res* 2012;21(7):1123–33. <https://doi.org/10.1007/s11136-011-0040-3>. ISSN 1573–2649.
- [22] Holzner B, Bode RK, Hahn EA, Cella D, Kopp M, Sperner-Unterweger B, Kemmler G. Equating eortc qlq-c30 and fact-g scores and its use in oncological research. *Eur J Cancer (Oxford, England : 1990)* 2006;42(18):3169–77. <https://doi.org/10.1016/j.ejca.2006.08.016>. ISSN 0959–8049.
- [23] Reeve Bryce B, Thissen David, DeWalt Darren A, Huang I-Chan, Liu Yang, Magnus Brooke, et al. Linkage between the promis(r) pediatric and adult emotional distress measures. *Qual Life Res : Int J Qual Life Asp Treat Care Rehabil* 2016;25(4):823–33. <https://doi.org/10.1007/s11136-015-1143-z>. ISSN 0962–9343.
- [24] Tian Feng, Ni Pengsheng, Mulcahey MJ, Hambleton Ronald K, Tulsy David, Haley Stephen M, Jette Alan M. Tracking functional status across the spinal cord injury lifespan: linking pediatric and adult patient-reported outcome scores. *Arch Phys Med Rehabil* 2014;95(11):2078–85. <https://doi.org/10.1016/j.apmr.2014.05.023>. e15, ISSN 0003–9993.
- [25] Marina N, Bielack S, Whelan J, Smeland S, Krailo M, Sydes MR, Butterfass-Bahloul T, Calaminus G, Bernstein M. International collaboration is feasible in trials for rare conditions: the euramos experience. *Cancer Treat Res* 2009;152:339–53. https://doi.org/10.1007/978-1-4419-0284-9_18. ISSN 0927–3042.
- [26] Whelan JS, Bielack SS, Marina N, Smeland S, Jovic G, Hook JM, et al. Euramos-1, an international randomised study for osteosarcoma: results from pre-randomisation treatment. *Ann Oncol* 2015;26(2):407–14. <https://doi.org/10.1093/annonc/mdu526>. ISSN 0923–7534, <https://academic.oup.com/annonc/article-pdf/26/2/407/24065165/mdu526.pdf>.
- [27] Calaminus Gabriele, Jenney Meriel, Hjorth Lars, Baust Katja, Bernstein Mark, Bielack Stefan, De Vos Patricia, Pancras C, Hogendoorn W, Jovic Gordana, Krailo Mark, Kreitz Kiana, Marina Neyssa, Popoola Babasola O, Sauerland Cristina, Smeland Sigbjørn, Teske Carmen, Schweinitz Clara V, Whelan Jeremy, Wiener Andreas, Sydes Matthew R, Nagarajan Rajaram. Quality of life of patients with osteosarcoma in the european american osteosarcoma study-1 (euramos-1): development and implementation of a questionnaire substudy. *JMIR research protocols* 2019;8(8):e14406. <https://doi.org/10.2196/14406>. ISSN 1929–0748.
- [28] Varni James W, Seid Michael, Rode Cheryl A. The pedsq™: measurement model for the pediatric quality of life inventory. *Med Care* 1999;37(2):126–39. ISSN 00257079.
- [29] Calaminus G, Weinspach S, Teske C, Göbel U. Quality of life in children and adolescents with cancer. first results of an evaluation of 49 patients with the pedqol questionnaire. *Klin Pädiatr* 2000; 212(4):211–5. <https://doi.org/10.1055/s-2000-9679>. ISSN 0300–8630.
- [30] Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, Filiberti A, Flechtner H, Fleishman SB, de Haes JC. The european organization for research and treatment of cancer qlq-c30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993;85(5):365–76. <https://doi.org/10.1093/jnci/85.5.365>. ISSN 0027–8874.
- [31] von Davier A. *Statistical Models for test equating, scaling, and linking*. New York: Springer; 2010. 9780387981383.
- [32] Lord Frederic M. Notes on comparable scales for test scores. *ETS Res Bull Ser* 1950;(2). <https://doi.org/10.1002/j.2333-8504.1950.tb00673.x>. i–21, 1950. ISSN 2333–8504, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/j.2333-8504.1950.tb00673.x>.
- [33] Kolen Michael J, Brennan Robert L. *Test equating, scaling, and linking: Methods and practices. Statistics for Social and Behavioral Sciences*. 3rd ed. New York: Springer; 2014. ISBN 1493903160.
- [34] equate Anthony D Albano. An r package for observed-score linking and equating. *J Stat Software* 2016;74(8):1–36. <https://doi.org/10.18637/jss.v074.i08>.
- [35] Bland JM, Altman DG. *Statistical methods for assessing agreement between two methods of clinical measurement*. *Lancet (London, England)* 1986;1(8476):307–10. ISSN 0140–6736.
- [36] Zhou Xiaoyan, Dibley Michael J, Cheng Yue, Ouyang Xue, Yan Hong. Validity of self-reported weight, height and resultant body mass index in Chinese adolescents and factors associated with errors in self-reports. *BMC Publ Health* 2010;10(1):190. <https://doi.org/10.1186/1471-2458-10-190>. ISSN 1471–2458.
- [37] Euser Anne M, Dekker Friedo W, Le Cessie Saskia. A practical approach to Bland-Altman plots and variation coefficients for log transformed variables. *J Clin Epidemiol* 2008;61(10):978–82. <https://doi.org/10.1016/j.jclinepi.2007.11.003>. ISSN 0895–4356, <https://www.sciencedirect.com/science/article/pii/S0895435607004131>.
- [38] Brehm Merel-Anne, Scholtes Vanessa A, Dallmeijer Annet J, Twisk Jos W, Harlaar Jaap. The importance of addressing heteroscedasticity in the reliability analysis of ratio-scaled variables: an example based on walking energy-cost measurements. *Dev Med Child Neurol* 2012;54(3):267–73. <https://doi.org/10.1111/j.1469-8749.2011.04164.x>. ISSN 1469–8749, <https://onlinelibrary.wiley.com/doi/10.1111/j.1469-8749.2011.04164.x>.
- [39] Lin LI. A concordance correlation coefficient to evaluate reproducibility. *Biometrics* 1989;45(1):255–68. ISSN 0006–341X.
- [40] Neil J. Dorans. Equating, concordance, and expectation. *Appl Psychol Meas* 2004;28(4):227–46. <https://doi.org/10.1177/0146621604265031>.