








ORIGINAL RESEARCH

Cardiovascular Safety Reporting in Contemporary Breast Cancer Clinical Trials

Arsalan Hamid , MD; Markus S. Anker , MD; John C. Ruckdeschel , MD; Muhammad Shahzeb Khan , MD, MSc; Ashal Tharwani, MD; Adebamike A. Oshunbade , MD, MPH; Rodney K. Kipchumba , BS; Samuel C. Thigpen, MD; Stefan D. Anker , MD, PhD; Gregg C. Fonarow , MD; Michael E. Hall , MD, MS; Javed Butler , MD, MPH, MBA

BACKGROUND: Several cancer therapies have been associated with cardiovascular harm in early-phase clinical trials. However, some cardiovascular harms do not manifest until later-phase trials. To limit inter-disease variability, we focused on breast cancer. Thus, we assessed the reporting of cardiovascular safety monitoring and outcomes in phase 2 and 3 contemporary breast cancer clinical trials.

METHODS AND RESULTS: We searched Embase and Medline records for phase 2 and 3 breast cancer pharmacotherapy trials. We examined exclusion criterion as a result of cardiovascular conditions, adverse cardiovascular event reporting, and cardiovascular safety assessment through cardiovascular imaging, ECG, troponin, or natriuretic peptides. Fisher's exact test was utilized to compare reporting. Fifty clinical trials were included in our study. Patients were excluded because of cardiovascular conditions in 42 (84%) trials. Heart failure was a frequent exclusion criterion ($n=31$; 62% trials). Adverse cardiovascular events were reported in 43 (86%) trials. Cardiovascular safety assessments were not reported in 23 (46%) trials, whereas natriuretic peptide and troponin assessments were not reported in any trial. Cardiovascular safety assessments were more frequently reported in industry-funded trials (69.2% versus 0.0%; $P<0.001$), and in trials administering targeted/immunotherapy agents compared with only hormonal/conventional chemotherapy (78.6% versus 22.7%, $P<0.001$).

CONCLUSIONS: Our findings demonstrate significant under-representation of patients with cardiovascular conditions or prevalent cardiovascular disease in contemporary later-phase breast cancer trials. Additionally, cardiovascular safety is not routinely monitored in these trials. Therefore, contemporary breast cancer clinical trials may possibly underestimate the cardiovascular risks of cancer pharmacotherapy agents for use in clinical practice.

Key Words: cancer therapy ■ cardio-oncology ■ cardiovascular disease ■ pharmacotherapy ■ safety

Breast cancer contributes to one quarter of all incident cancers in women and is the most common cause of female cancer-related mortality.¹ Furthermore, breast cancer shares a number of common risk factors with cardiovascular disease (CVD) such as age, obesity, tobacco use, alcohol intake, and hormone replacement therapy.² One third of patients with breast cancer have CVD.³ With greater efficacy and introduction of improved breast cancer therapies,

the 5-year survival of breast cancer patients has risen from 75% to 91%.⁴ Given the increased survival rate, cancer therapy-induced cardiovascular toxicity is a growing cause of morbidity and mortality in patients with breast cancer.⁵ Breast cancer pharmacotherapy agents including conventional chemotherapies (alkylating agents, antimetabolites, and antimicrotubule agents),^{6,7} targeted therapies,⁷ and immunotherapy⁸ are associated with varying degrees of adverse

Correspondence to: Javed Butler, MD, MPH, MBA, Department of Medicine, University of Mississippi Medical Center (L650), 2500 N. State St, Jackson, MS 39216. Email: butljzh@gmail.com

Preprint posted to SSRN November 9, 2021. doi: <https://doi.org/10.2139/ssrn.3959661>

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.121.025206>

For Sources of Funding and Disclosures, see page 9.

© 2022 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](#) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

JAHA is available at: www.ahajournals.org/journal/jaha

CLINICAL PERSPECTIVE

What Is New?

- We highlight that breast cancer pharmacotherapy trials under-represent patients with cardiovascular conditions.
- Cardiovascular safety assessments frequently are not reported in breast cancer pharmacotherapy trials.
- Although patients with cardiovascular disease were often excluded from breast cancer pharmacotherapy trials, included participants commonly experienced adverse cardiovascular events during these trials.

What Are the Clinical Implications?

- Breast cancer trials may underestimate the cardiovascular risks of cancer pharmacotherapy agents for the general population.
- Later-phase breast cancer pharmacotherapy trials may not be able to uncover cardiovascular risks that have not been identified in smaller-scale trials and as a result, subclinical cardiotoxicity may remain undetected in these trials.
- Judiciously conducted cardiovascular safety assessments may allow inclusion of patients with cardiovascular conditions, early identification of adverse cardiovascular events, and demonstrate subclinical cardiovascular harm of pharmacotherapy.

Nonstandard Abbreviations and Acronyms

FDA US Food and Drug Administration

cardiovascular effects. Oncologic pharmacotherapy-induced cardiovascular effects can be serious and possibly fatal, including events such as myocardial infarction, arrhythmias, myocarditis, cardiomyopathy, pericardial disease, stroke, and heart failure.⁷ Recent advances in the field of oncology have led to the rapid introduction of novel therapies, resulting in a rise in the number of agents gaining US Food and Drug Administration (FDA) approval every year.⁹ Rigorous safety and efficacy testing in multiphase clinical trials must be performed before FDA approval. Given the high prevalence of CVD in women with breast cancer, it is important to understand potential cardiovascular harm from novel agents in this population. However, it is unclear whether large contemporary phase 2 and 3 clinical trials adequately assess cardiovascular safety and are able to uncover cardiovascular risks that were not identified in early-phase trials. Therefore, we assessed reporting of cardiovascular safety in breast

cancer trials administering cancer pharmacotherapy through the evaluation of 3 key domains: inclusion/exclusion of patients with cardiovascular conditions, cardiovascular safety assessments (cardiovascular laboratories or imaging assessments), and adverse cardiovascular events.

METHODS

Study Identification

We searched Embase and Medline records before January 4, 2019 for clinical trial articles assessing efficacy of cancer pharmacotherapy in breast cancer until 50 articles that fit inclusion criteria were acquired. The search strategy used was: “Breast” AND “Cancer” with trial filters activated (Figure). The inclusion criteria of studies were as follows: (1) phase 2 and 3 trials; (2) enrolled >50 participants; (3) assessed the use of cancer pharmacotherapy as primary intervention on primary or secondary end points of survival, response, or safety in patients with active breast cancer; and (4) were registered with the national clinical trials registry or other international trial registries (this permits acquisition of further eligibility criteria and adverse event data). Studies were excluded if they were (1) quality of life or cost-analysis studies, (2) expanded access programs, or (3) involved treatment of other (nonbreast) types of cancer (to limit inter-disease variability). Using the above criteria, our search was conducted with an all-inclusive approach to any stage of cancer, pharmacotherapy administered, or length of follow-up for articles published in any journal. This was done to maintain generalizability of our results. Two independent reviewers screened selected studies and abstracted data; in the event of disagreement, a third reviewer resolved discordant assessments. Furthermore, corresponding authors of articles were contacted for queries if needed. If corresponding authors failed to respond, then best judgment and consensus of reviewers was implemented. We acquired 50 published trials (Table S1) after review of 1775 records (Figure). Institutional Review Board approval was waived for this article because it is a review of publicly available data. The data that support the findings of this study are available from the corresponding author upon reasonable request.

Data Abstraction

Data from primary articles, trial databases, protocols, and supplementary files were reviewed as a pooled analysis of trial data (Figure S1). The following data were abstracted from articles, registries, protocols, or supplements: (1) publication year, (2) number of participants, (3) age, (4) race (if specified), (5) median survival duration (if reached/available), (6) trial phase, (7) funding/sponsor, (8) cancer pharmacotherapy intervention, (9) trial end point,




Figure. Trial search.

(10) inclusion/exclusion criteria, (11) safety assessments, and (12) adverse events. Registry data for demographics were utilized when articles did not present clear data. Data were last updated April 25, 2020.

Definition of Reporting Variables

Cardiovascular conditions were defined as any CVD, cardiovascular symptoms (for example, dyspnea, peripheral edema, or chest pain), or any abnormal cardiovascular examination/laboratory parameters (for example, ejection fraction [EF], blood pressure, or heart rate). CVD was classified as heart failure (reported as history of heart failure, symptomatic heart failure, cardiopulmonary failure, or New York Heart Association class >1), angina, ischemic heart disease/coronary artery disease, myocardial infarction (reported as history of myocardial infarction or a recent myocardial infarction), hypertension (reported as history of hypertension or with blood pressure cutoffs), arrhythmia (reported as history of arrhythmias that were uncontrolled, unstable, clinically significant, or other unspecified ECG abnormalities), QT interval prolongation, stroke (ischemic or hemorrhagic), cardiomyopathy, pericardial disease (pericarditis or pericardial effusion), pulmonary hypertension/cor pulmonale, aortic aneurysm, acute

coronary syndrome, cardiac tamponade, hypercholesterolemia/hypertriglyceridemia, cardiac/cardiopulmonary arrest, cardiac infection, structural heart disease, thrombosis (deep venous thrombosis, pulmonary embolism, or thromboembolic disease) or peripheral artery disease/carotid artery disease. Hypertension was divided into a history of hypertension and hypertension. When hypertension was listed without a blood pressure cutoff, this was considered a history of hypertension. Alternatively, hypertension with a blood pressure cutoff (eg, uncontrolled/poorly controlled hypertension [$>140/90$ mmHg under treatment with 2 antihypertensive drugs]) was abstracted as hypertension, because this could possibly indicate hypertension identified upon screening (however, timeline of blood pressure assessment was mostly not specified in trials). Myocardial infarctions were also subdivided into a history of myocardial infarction if the timeline was not specified or a recent myocardial infarction if timeline was specified to have been at most in the past 1 year of study enrollment, entry, randomization, or first dose of therapy.

Cardiovascular safety assessments were defined as planned evaluations at baseline or pre/post therapy administration at any time in the trial. Cardiovascular

safety assessments included ECG, cardiac imaging (echocardiography, multigated acquisition scan, or cardiac magnetic resonance imaging [CMR]), natriuretic peptides (brain natriuretic peptide and N-terminal pro-brain natriuretic peptide), troponins, lipid profile, or creatine phosphokinase. Adverse cardiovascular events were defined as any reported adverse cardiovascular symptom, disease, or condition encountered at any point during the study or that may have resulted in death.

Statistical Analysis

Continuous variables were abstracted as medians or absolute counts and summarized as means and SDs. Categorical variables were summarized as frequencies and percentages. Where demographic data in trials were presented in multiple groups (interventional group or control group), demographic data for the trial overall were summarized as mean of all groups. Where demographic data were available for only 1 study group, it was extrapolated to the entire study population. Fisher's exact test was applied for comparisons of reporting by various trial characteristics (funding, type of agent, and phase of trial). A 2-sided *P* value of <0.05 was considered significant. All statistical analysis was carried out using IBM SPSS Statistics for Windows, Version 23.0 (IBM, Armonk, New York, USA).

RESULTS

General Characteristics

In total, 1775 records were screened and 50 clinical trials that cumulatively enrolled 26 893 participants were included in our study (Figure and Table S1). The majority of these trials were published in 2018 (*n*=39; 78% trials) (Table 1). On average, trials had a population of 538 participants, median age of 55.7 years, median follow-up time of 31.3 months, median progression-free survival of 8.5 months, median overall survival of 24.0 months, and 81.1% of participants were White race (Table 1).

Included trials were most often phase 3 (*n*=27; 54% trials), industry sponsored (*n*=39; 78% trials), and reported a primary end point of therapy response (*n*=23; 46% trials). The most common therapy administration regimen was a combination of targeted therapy with conventional chemotherapy (*n*=20; 40% trials) (Table 1).

Reporting of Cardiovascular Exclusion

The majority of trials (*n*=42; 84% trials) listed at least 1 cardiovascular condition as an exclusion criterion, and 34 (68%) trials listed at least 1 CVD as an exclusion criterion (Table 2). The most frequently reported cause for exclusion was heart failure (*n*=31; 62% trials). Reported exclusion by nonspecific cardiac causes

Table 1. Trial and Participant Characteristics

Characteristic	Trials
Mean number of participants per trial	537.9±880.9
Mean of trial median follow-up, mo	31.3±21.4
Mean of participants median age, y	55.7±7.4
Mean of participants racial distribution, %	
White	81.1±12.9
Black	5.1±8.1
Asian	9.5±11.4
Other*	4.3±3.3
Trials published in year, n (%)	
2019	11 (22.0)
2018	39 (78.0)
Mean of participants median progression-free survival, mo	8.5±4.2
Mean of participants median overall survival, mo	24.0±11.4
Trial phase, n (%)	
2	23 (46.0)
3	27 (54.0)
Trial sponsor/funding, n (%)	
Industry (entirely or partially)	39 (78.0)
University/Research organization/Government	11 (22.0)
Type of agent administered, n (%)	
Targeted+Chemotherapy [†]	20 (40.0)
Targeted+Hormonal	4 (8.0)
Targeted+Chemotherapy [†] +Hormonal	3 (6.0)
Immuno+Chemotherapy [†]	1 (2.0)
Chemotherapy [†] monotherapy	8 (16.0)
Hormonal monotherapy	2 (4.0)
Chemotherapy [†] +Hormonal therapy	12 (24.0)
Primary reported trial end point, n (%)	
Survival (progression free and overall)	20 (40.0)
Response	23 (46.0)
Safety/adverse events	2 (4.0)
Other [‡]	5 (10.0)
Available trial data sources, n (%)	
Included articles	50 (100.0)
Supplementary files	29 (58.0)
Protocols	18 (36.0)

Certain demographic data were not available in trials, including: 25 (50%) trials missing median follow-up duration, 7 (14%) trials missing median age of participants, 30 (60%) trials missing racial distribution of participants, 32 (64%) trials missing median overall survival of participants, and 26 (52%) trials missing median progression-free survival of participants.

*Other includes Native Americans, Alaskans, unknown race, or unspecified as "other race."

[†]Chemotherapy here includes cytotoxic chemotherapy agents, antimetabolites, alkylating agents, platinum agents, and microtubule-inhibiting agents.

[‡]Other includes clinical benefit rate, time to treatment failure, time to progression, resumption of menstruation, and disease control rate.

was encountered in 26 (52%) trials (for example, listed as "uncontrolled heart disease" or "history of cardiac disease"). Other common causes of exclusion were

Table 2. Reporting of Cardiovascular Exclusion Criteria

Exclusion criteria	Number of trials reported exclusion, n (%)
Any cardiovascular condition	42 (84)
Any CVD	34 (68)
Heart failure	31 (62)
Arrhythmia*	29 (58)
Low ejection fraction	27 (54)
Angina	26 (52)
Nonspecific cardiac cause [†]	26 (52)
Recent myocardial infarction [‡]	25 (50)
Hypertension [§]	11 (22)
History of hypertension	10 (20)
Structural disease	8 (16)
Ischemic/Coronary heart disease	8 (16)
History of myocardial infarction	6 (12)
QT interval prolongation	6 (12)
Pericardial disorders (effusion or pericarditis)	5 (10)
Stroke (ischemic or hemorrhagic)	4 (8)
Dyspnea	4 (8)
Pro-arrhythmic medication use	4 (8)
Cardiomyopathy	3 (6)
Thrombosis/thromboembolism	2 (4)
Hypotension	2 (4)
Hypercholesterolemia/ hypertriglyceridemia	2 (4)
Aortic aneurysm	1 (2)
Peripheral artery disease	1 (2)
Elevated creatine phosphokinase levels	1 (2)

CVD indicates cardiovascular disease.

*Includes arrhythmia, atrioventricular block, atrial fibrillation, bradycardia, high-risk arrhythmia, brugada, torsades, and tachycardia. These arrhythmias were largely labeled as unstable or uncontrolled.

[†]Nonspecific reporting such as “Uncontrolled cardiac/cardiovascular/heart disease, history of heart/cardiac disease or dysfunction, significant or clinically significant cardiac disorders/disease, clinically significant cardiac disease, cardiac/cardiovascular disease precluding study, nonmalignant systemic disease (cardiac), major cardiovascular comorbidity, severe heart disease with life expectancy from disease <2years, active cardiac disease, other cardiac disease, serious cardiac disease/illness; inclusion of participants with normal cardiopulmonary/cardiac function as assessed by echocardiography, normal/adequate cardiac function, or functionally preserved heart”. The nonspecificity terms were summarized in the interest of brevity.

[‡]Recent myocardial infarction includes myocardial infarctions before specific timeline of study entry, randomization, enrollment, or first dose therapy (8 studies did not specify baseline from which recent myocardial infarctions will be excluded). Different trials listed exclusions at different timelines, 18 trials excluded patients with myocardial infarctions within 6 months, 5 trials excluded patients with myocardial infarctions within 12 months, and 2 trials excluded patients with myocardial infarctions within the past 3 months.

[§]Hypertension as classified as an elevated blood pressure cutoff. Blood pressure cutoffs varied, eg, >140/90 (while on 2 antihypertensive medication) or if systolic blood pressure was >150, >160, >180, <90mmHg or if diastolic blood pressures were >90 or >100mmHg.

arrhythmias in 29 (58%) trials, low EF in 27 (54%) trials, and a recent myocardial infarction in 25 (50%) trials (Table 2). Exclusion criteria set for EF varied from trial to trial, ranging from EF <40% to EF <55%.

Of note, 10 (20%) trials excluded patients with a history of hypertension and 11 (22%) trials excluded patients with hypertension with various blood pressure cutoffs (Table 2). Altogether, 20 (40%) trials excluded patients with a history of hypertension or hypertension (with listed blood pressure cutoffs).

Reporting of mCardiovascular Safety Assessments

Cardiovascular safety assessments were rarely reported by the included trials. Nearly half of these trials (n=23; 46% trials) did not report any cardiovascular safety assessments either at baseline or throughout treatment cycles (Table 3). The most frequently reported cardiovascular safety assessments were ECGs (n=23; 46% trials) and cardiac imaging (n=20; 40% trials) (Table 3). The 20 (40%) trials that conducted cardiac imaging assessments all utilized echocardiograms and/or multigated acquisition scans, while only 1 (2%) trial conducted some cardiac imaging assessments with CMR. Cardiac biomarker measurements were rarely reported. Assessment of creatine kinase or creatine phosphokinase levels was reported in only 2 (4.3%) trials, and no trial reported assessment of natriuretic peptides or troponins (Table 3).

Although 27 (54%) trials reported exclusion of patients based on low EF, only 20 (40%) trials reported cardiac imaging as a safety assessment (Tables 2 and 3). Further assessment of this discordance demonstrated that one quarter (n=7; 26% trials) of these 27 trials, which reported exclusion of patients with low EF, did not report any cardiac imaging (by echocardiography or myocardial gated uptake assessment) as a safety assessment performed during the trial.

Reporting of Adverse Cardiovascular Events

Adverse cardiovascular events were commonly encountered, as 43 (86%) trials reported at least 1 cardiovascular adverse event and 36 (72%) trials reported at least 1 CVD event (Table 4). The most frequently reported adverse events were hypertension (n=23; 46% trials), thrombosis/thromboembolisms (n=23; 46% trials), dyspnea (n=22; 44% trials), edema (n=22; 44%), heart failure (n=19; 38% trials), and arrhythmia (n=19; 38% trials) (Table 4). The majority of trials (n=46; 92% trials) reported adverse events via a standardized grading system (ie, the National Cancer Institute common terminology criteria for adverse events/common toxicity criteria). Death caused by CVD was reported in one third of these trials (n=18; 36% trials). Common causes of death were heart failure, stroke, myocardial infarctions, thromboembolic events, arrhythmias, and cardiac arrest.

Table 3. Reporting of Cardiovascular Safety Assessments

Safety assessment	Number of trials reported assessment, n (%)
Any cardiovascular assessment	27 (54)
ECG	23 (46)
Cardiac imaging*	20 (40)
Lipid profile	5 (10)
Creatine kinase/creatinine phosphokinase	2 (4)
Troponin	0 (0)
BNP/NT-proBNP	0 (0)

BNP indicates brain natriuretic peptide; and NT-proBNP, N-terminal pro-brain natriuretic peptide.

*Cardiac imaging was conducted in trials through echocardiograms, multigated acquisition scan, or cardiac magnetic resonance imaging (CMR). Only 1 (2%) trial reported use of CMR, while 20 (40%) trials reported use of echocardiograms and/or multigated acquisition scans as cardiac imaging modalities.

Symptoms such as dyspnea (n=22; 44% trials), peripheral edema (n=22; 44% trials), and chest pain (n=6; 12% trials) were commonly reported; however, they were primarily reported with no specified cause (Table 4). Only 1 trial reported chest pain specifically as cardiac chest pain. Furthermore, reporting of cardiovascular events in a nonspecific manner was encountered in 4 (8%) trials, with events reported as a “cardiac/cardiovascular adverse event” (Table 4).

Association of Trial Factors With Reporting

Industry-funded trials more frequently reported cardiovascular safety assessments, compared with non-industry-funded trials (69.2% versus 0.0%, respectively, $P<0.001$). Furthermore, industry-funded trials more frequently reported cardiovascular adverse events, compared with non-industry-funded trials (97.4% versus 45.5%, respectively; $P<0.001$) (Table 5).

Trials administering targeted/immunotherapy agents more frequently reported cardiovascular safety assessments, compared with trials administering conventional chemotherapy or hormonal therapy (78.6% versus 22.7%, respectively, $P<0.001$). We also observed more frequent exclusion of participants with cardiovascular conditions in trials administering targeted/immunotherapy agents, compared with trials administering conventional chemotherapy or hormonal therapy (96.4% versus 68.2%, respectively; $P=0.015$) (Table 5).

DISCUSSION

This study provides unique insight as the first report assessing cardiovascular safety reporting within contemporary later-phase breast cancer pharmacotherapy clinical trials by 3 major criteria: eligibility, cardiovascular safety assessment monitoring, and adverse event reporting.

Table 4. Reporting of Adverse Cardiovascular Events

Adverse cardiovascular event reporting	Number of trials reported adverse events, n (%)
Reported any cardiovascular adverse event	43 (86)
Reported any CVD adverse event	36 (72)
Hypertension	23 (46)
Thrombosis/thromboembolism	23 (46)
Dyspnea	22 (44)
Edema	22 (44)
Heart failure	19 (38)
Arrhythmia*	19 (38)
Stroke	14 (28)
Low/decreased ejection fraction	12 (24)
Pericardial disease (effusion or pericarditis)	9 (18)
Myocardial infarction	8 (16)
Hypotension	8 (16)
Hypercholesterolemia/hypertriglyceridemia	7 (14)
Cardiac/cardiopulmonary arrest	7 (14)
LV dysfunction/failure	7 (14)
Chest pain	6 (12)
Ischemic heart/coronary artery disease/coronary artery stenosis	5 (10)
Cardiomyopathy	4 (8)
Nonspecific cardiac event†	4 (8)
QT interval prolongation	3 (6)
Cardiac decompensation/insufficiency/circulatory collapse	3 (6)
Acute coronary syndrome	3 (6)
Creatine phosphokinase/troponin elevation	3 (6)
Cardiac tamponade	2 (4)
Palpitations	2 (4)
Pulmonary hypertension/cor pulmonale	2 (4)
Cardiac infection	2 (4)
Angina	1 (2)
Structural heart disease	1 (2)
Vasovagal episode	1 (2)
Peripheral artery disease	1 (2)

CVD indicates cardiovascular disease; and LV, left ventricle.

*Includes atrial fibrillation, atrial flutter, bradycardia, heart block, premature ventricular contraction, sinus arrhythmia, supraventricular tachycardia, tachycardia, torsades, ventricular tachycardia, ventricular arrhythmia, ventricular extrasystole, or ECG abnormality.

†Nonspecific reporting such as “cardiac/cardiovascular adverse events, cardiac disorder, other cardiovascular adverse events, cardiac complications, any cardiovascular disease, cardiac events not otherwise specified, cardiac other”. Reporting terms were summarized in the interest of brevity.

Our findings demonstrate significant underrepresentation of patients with CVD (by exclusion of cardiovascular conditions or prevalent CVD) in these clinical trials. Although patients with CVD were often excluded from

Table 5. Association of Cardiovascular Safety Reporting and Trial Characteristics

Trial characteristics	Trials reported, n (%)*					
	Reported any cardiovascular exclusion		Reported any cardiovascular safety assessment		Reported any cardiovascular event	
	Reporting	P value	Reporting	P value	Reporting	P value
Funding		0.174		<0.001		<0.001
Industry	31 (79.5)		27 (69.2)		38 (97.4)	
University/Research organization/ Government	11 (100.0)		0 (0.0)		5 (45.5)	
Use of targeted/immunotherapy		0.015		<0.001		1.000
Included in regimen	27 (96.4)		22 (78.6)		24 (85.7)	
Not included in regimen	15 (68.2)		5 (22.7)		19 (86.4)	
Use of anthracyclines		0.087		1.000		1.000
Included in regimen	14 (100.0)		8 (57.1)		12 (85.7)	
Not included in regimen	28 (77.8)		19 (52.8)		31 (86.1)	
Phase		1.000		0.087		0.225
2	19 (82.6)		9 (39.1)		18 (78.3)	
3	23 (85.2)		18 (66.7)		25 (92.6)	

*Percentage is reported as fraction of trial characteristic group (for example, 79.5% of industry-funded trials reported any cardiovascular exclusion).

breast cancer pharmacotherapy clinical trials, included participants commonly experienced adverse cardiovascular events during these trials. Furthermore, our findings highlight that cardiovascular safety may not be routinely monitored in all breast cancer pharmacotherapy studies. Therefore, potential cardiovascular harm from these cancer pharmacotherapy agents might be underestimated and subclinical cardiotoxicity may remain undetected in later-phase breast cancer clinical trials. Lastly, other factors such as trial funding source and type of pharmacotherapy agent were associated with discrepant cardiovascular safety monitoring.

The underrepresentation of patients with CVD in cancer pharmacotherapy trials linked to FDA drug approval has previously been described by Bonsu et al. They observed that 34% of pharmacotherapy trials for any cancer type and ~50% for breast cancer exclude patients with CVD.¹⁰ This was lower compared with our observation that 68% of breast cancer trials exclude patients with CVD and was far lower compared with our finding that 84% of trials exclude patients with cardiovascular conditions. Therefore, we highlight that in addition to direct exclusion of patients with CVD, breast cancer pharmacotherapy trials may further exclude patients with CVD by using cardiovascular symptoms (dyspnea) or cardiovascular parameters (left ventricular EF) as cardiovascular exclusion criterion. As a result, patients with cardiovascular conditions are excluded in the majority of these trials.

Similar to our observations, Bonsu et al also encountered nonspecific reported exclusion of patients with CVD in more than one third of trials compared with approximately half of trials in our study.¹⁰ The use of nonspecific/ill-defined terminology (eg, “history of cardiac

disease” or “uncontrolled heart disease”) possibly indicates the use of subjective exclusion criteria, resulting in the possibility of either the inclusion of patients with real risk factors for cardiovascular toxicity or exclusion of patients with minimal to no added risk of toxicity.

An evaluation of cancer trials by Bonsu et al suggested that cancer pharmacotherapy trials linked with FDA drug approval underreport adverse CVD events in comparison to population-based estimates.¹¹ Furthermore, they demonstrated that 45% of breast cancer trials do not report adverse CVD events, compared with 28% in our observation.¹¹ Our study demonstrates that cancer trials sometimes report events as symptoms (dyspnea) or abnormal parameters (decreased left ventricular ejection fraction), rather than diagnosed CVD; this may also have reflected on the findings of Bonsu et al on the underreporting of CVD. Furthermore, our study highlights a possibility that this observation of adverse event underreporting may be because of underdetection as a result of poor conduct of cardiovascular safety assessments, because nearly half of trials in our observation did not report any form of cardiovascular safety assessment. Therefore, improved conduct of cardiovascular safety assessments may possibly permit better adverse cardiovascular event detection and identification at earlier stages.

Preclinical studies offer an opportunity to study cardiovascular risks of cancer pharmacotherapy before introduction in human models. These are largely conducted in animal models, based on studied cardiovascular pharmacology and physiology. However, preclinical studies on their own are not sufficient to uncover cardiovascular risks of cancer pharmacotherapy in humans because of several limitations including

size of the model (smaller animals usually utilized) and species-dependent sensitivity to myocardial injury.¹² This highlights the importance of clinical trials in uncovering cardiovascular toxicity in humans that otherwise cannot be observed in preclinical models.

Large, later-phase clinical trials present an opportunity to uncover cardiovascular risks (particularly subclinical cardiovascular risks) not previously demonstrated in smaller trials, and ultimately these later-phase trials are the primary drivers of decision making in clinical practice. Currently, guidelines on the management of patients with high cardiovascular risk receiving cancer pharmacotherapy are nonexistent. However, studies suggest the use of less cardiotoxic agents (or at minimal dose), risk class allocation based on risk scores from initial evaluation, then appropriate monitoring (with periodic echocardiography, ECG, troponin, natriuretic peptides), and management.¹³ Patients with cardiovascular risks still receive cancer pharmacotherapy but the criteria for their inclusion are not standardized. Therefore, exclusion of patients who otherwise could be candidates to receive therapy in clinical practice may impact the generalizability of results garnered from trials.

Cancer trials are not the only type of trials that exclude patients with comorbidities/multimorbidity. Patients included in clinical trials have half as many medical comorbidities as compared with the general population.¹⁴ The percentage of included patients with comorbidities varies depending on the patient condition under treatment.¹⁴ While a more inclusive approach to clinical trials may generate more generalizable data, this raises several issues with primary trial outcomes. Clinical trials are often created with narrowly defined populations (for example, by excluding patients with CVD), with the intention of generating results specific to the primary outcome.¹⁵ Factors such as regimen compliance¹⁶ and patient survival^{17,18} are directly impacted by the presence of comorbidities (such as CVD) in patients with cancer. These factors are therefore likely to affect trial success and may alter results, particularly with respect to cancer trials where more than one third of all trials in our observation used survival end points to determine therapy efficacy. Because of these reasons, breast cancer trials may opt to include healthier populations when assessing pharmacotherapy in the interest of primary outcomes.

Rather than exclude patients with CVD, cancer trials may possibly consider improvement in protocol-driven safety assessments (baseline, pretreatment, and posttreatment), and identification of high-risk participants who require more frequent assessment or dose adjustments. Early changes in left ventricular strain and effective arterial elastance in patients with breast cancer receiving pharmacotherapy have been indicative of future heart failure symptoms.¹⁹ Furthermore, brain

natriuretic peptide²⁰ and troponin²¹ are proven predictors of cancer pharmacotherapy-related cardiotoxicity and cardiac events. However, reporting of cardiac imaging as a safety assessment was observed in less than half of trials, and no trials we studied reported using troponin, brain natriuretic peptide, or N-terminal pro-brain natriuretic peptide for screening or safety assessments.

The European Society of Cardiology supports the use of natriuretic peptides and cardiac troponins in patients receiving cancer pharmacotherapy.²² Biomarkers can help risk stratify patients at baseline and uncover early cardiac injury or strain. Elevations in cardiac biomarkers can indicate patients requiring closer monitoring, initiation of cardioprotective therapy, and do not necessarily mean cessation of therapy.²² However, baseline troponin levels at times may not always correlate with increased risks of cancer pharmacotherapy-related cardiotoxicity.^{23,24} As a result, their consistent use may not be widely adopted as yet. The European Society of Cardiology recommendations on cardiac biomarkers have evolved to more clearly defined timelines of collection from the 2016 to 2020 position statements.^{22,25} Therefore, in the evolving field of cardio-oncology, the consensus on their use in routine surveillance in pharmacotherapy-induced cardiotoxicity is also evolving. While biomarker assessments offer better surveillance of cardiotoxicity in patients receiving cancer pharmacotherapy, there are still no established standards with respect to assay selection, cut-off values for clinically significant changes, and the ideal timing of sampling.²² This may suggest why biomarker analyses were not conducted in any of the contemporary breast cancer trials assessed in our study. Cardiac biomarkers offer an opportunity for cancer pharmacotherapy trials in the future to identify and prevent cancer pharmacotherapy-induced cardiotoxicity.

While 20 (40%) trials reported use of cardiac imaging modalities, we observed only 1 (2%) trial that reported use of CMR. CMR as a modality offers the benefit of early identification of cancer pharmacotherapy-induced cardiotoxicity compared with echocardiograms or multigated acquisition scans with the utility of T1/T2 mapping, the ability to detect subtle decreases in left ventricular ejection fraction, myocardial edema (in T2 images), early inflammation, and early fibrosis.²⁶ In the future, we suggest that cancer pharmacotherapy trials may utilize CMR in higher-risk populations to allow inclusion of participants with cardiovascular conditions because of the comfort of identification of early cardiotoxicity, before the development of overt cardiotoxicity. However, because of cost constraints, CMR is largely used when echocardiograms and multigated acquisition scans are indeterminate,²⁶ which may explain why we observed its negligible use in cancer pharmacotherapy trials. Lack of use of cardiac biomarkers as well as cardiac imaging (particularly CMR) may result

in a missed opportunity for the identification of newly uncovered cancer pharmacotherapy-induced cardiotoxicity. Improved conduct in cardiovascular safety assessments may permit better detection of incident early CVD, patients requiring dose adjustments, and may uncover subclinical risks of therapy. This would permit safer inclusion of participants with comorbidities and would generate data that could lead to better-informed decision making in clinical practice.

Trials frequently reported adverse cardiovascular events despite most of them excluding patients with prevalent CVD. Hypertension was the most commonly reported adverse event among trials and can be caused by a variety of agents.²⁷ Specifically, it is one of the most common adverse effects of targeted therapies.^{25,28} Targeted therapies are relatively new and were evaluated in a majority of trial regimens in our study. An analysis of clinical trials with patients receiving targeted therapy reported cardiotoxic adverse events more often than any other type of adverse event.²⁹ In our study, clinical trials excluded patients with CVD more frequently and took greater caution in the conduct and reporting of cardiovascular safety assessments when targeted therapy or immunotherapy were administered (as the sole therapy or in combination with other agents). This may be because targeted therapies are relatively newer agents and exhibit a large variety of cardiotoxic effects, ranging from cardiac pump dysfunction (HER2 monoclonal antibodies/HER2 inhibitors) to prolonged QT interval (tyrosine kinase inhibitors), which is why more stringent monitoring is required.⁷ In contrast, while anthracyclines are known to cause cardiovascular dysfunction, we observed no significant difference in reporting irrespective of anthracycline use.²⁵

Industry funding/sponsorship may play a role in the presentation of positive results and outcomes; however, for the most part, this has no effect on study quality.^{30,31} Our study similarly demonstrated better study practices in industry-funded trials with greater reporting of safety assessments and adverse cardiovascular events. Industry-funded studies are often thoroughly scrutinized; thus, they may take further precautions to maintain safety and transparency in trials.³¹

Our study has some limitations that must be taken into consideration. Detailed safety assessments are often reported in study protocols, and protocols were unavailable for two thirds of trials (Table 1). We thoroughly searched for protocols of all trials, but many trials do not upload trial protocols. Hence, throughout our study we can only comment on what trials opted to report (in terms of safety assessments and eligibility criteria). Furthermore, this study retrieved trials conducted in patients with breast cancer; therefore, results cannot be generalized to other cancer types.

CONCLUSIONS

The study populations of contemporary later-phase breast cancer trials administering pharmacotherapy are underrepresentative of patients with CVD. This possibly results in a lack of generalizability of generated data. Before the initiation of pharmacotherapy and during its course, cardiovascular safety assessments are rarely reported, suggesting room for improvement in the conduct and reporting of cardiovascular safety assessments. Reporting of cardiovascular safety assessment data may depend on the source of trial funding and the type of agent administered. Judiciously conducted cardiovascular safety assessments may allow inclusion of patients with CVD, early identification of adverse cardiovascular events, and demonstrate subclinical cardiovascular harm of pharmacotherapy. Thus, later-phase breast cancer clinical trials may not be able to uncover cardiovascular risks that have not been identified in smaller-scale trials. These findings suggest that contemporary later-phase breast cancer clinical trials may underestimate the cardiovascular risks of cancer pharmacotherapy. Trials conducted in populations more inclusive of CVD, with increased scrutiny in monitoring of cardiovascular harm related to novel cancer therapies, could possibly lead to more informed clinical decision making for oncology patients with CVD in clinical practice.

ARTICLE INFORMATION

Received December 25, 2021; accepted April 29, 2022.

Affiliations

Department of Medicine (A.H., M.S.K., R.K.K., S.C.T., J.B.), Division of Hematology/Oncology, Department of Medicine, Cancer Center and Research Institute, University of Mississippi Medical Center, Jackson, MS (J.C.R.) and Division of Cardiology, Department of Medicine, University of Mississippi Medical Center, Jackson, MS (A.A.O., M.E.H.), University of Mississippi Medical Center, Jackson, MS; Department of Cardiology (CBF), Charite University Medicine Berlin (M.S.A.); Department of Cardiology (CVK), Charite Universitätsmedizin Berlin (S.D.A.); Berlin Institute of Health Center for Regenerative Therapies (BCRT) (M.S.A., S.D.A.); DZHK (German Centre for Cardiovascular Research), partner site Berlin (M.S.A., S.D.A.), Berlin, Germany; Department of Medicine, Cleveland Clinic Foundation, Cleveland, OH (A.T.); and Division of Cardiology, David Geffen School of Medicine at UCLA, Los Angeles, CA (G.C.F.).

Sources of Funding

None.

Disclosures

Dr Markus Anker reports personal fees from Servier, outside the submitted work. Dr John Ruckdeschel serves on the Boards of Genyoux-Omnitura and Real-Time Micro Spectroscopy, and he has received research support from the National Cancer Institute, Novartis, Immunomedics, and EpicentRx. Dr Gregg Fonarow serves as a consultant for Abbott, Amgen, AstraZeneca, Bayer, Janssen, Medtronic, Merck, and Novartis. Dr Javed Butler serves as a consultant for Abbott, Adrenomed, Amgen, Array, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol Myers Squibb, CVRx, G3 Pharmaceutical, Impulse Dynamics, Innolife, Janssen, LivaNova, Luitpold, Medtronic, Merck, Novartis, NovoNordisk, Relypsa, Roche, V-Wave Limited, and Vifor. The remaining authors have no disclosures to report.

Supplemental Material

Table S1
Figure S1

REFERENCES

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68:394–424. doi: 10.3322/caac.21492
- Mehta LS, Watson KE, Barac A, Beckie TM, Bittner V, Cruz-Flores S, Dent S, Kondapalli L, Ky B, Okwuosa T, et al. Cardiovascular disease and breast cancer: where these entities intersect: a scientific statement from the American Heart Association. *Circulation*. 2018;137:e30–e66. doi: 10.1161/CIR.0000000000000556
- Ng HS, Vitry A, Koczwara B, Roder D, McBride ML. Patterns of comorbidities in women with breast cancer: a Canadian population-based study. *Cancer Causes Control*. 2019;30:931–941. doi: 10.1007/s10552-019-01203-0
- American Cancer Society. Cancer Facts & Figures 2019. *Cancer*. 2019. Available from: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2019/cancer-facts-and-figures-2019.pdf> Accessed May 2020
- Shakir DK, Rasul KI. Chemotherapy induced cardiomyopathy: pathogenesis, monitoring and management. *J Clin Med Res*. 2009;1:8–12. doi: 10.4021/jocmr2009.02.1225
- Pai VB, MCJDS N. Cardiotoxicity of chemotherapeutic agents. *Drug Saf*. 2000;22:263–302. doi: 10.2165/00002018-200022040-00002
- Moslehi JJ. Cardiovascular toxic effects of targeted cancer therapies. *N Engl J Med*. 2016;375:1457–1467. doi: 10.1056/NEJMra1100265
- Asnani A. Cardiotoxicity of immunotherapy: incidence, diagnosis, and management. *Curr Oncol Rep*. 2018;20:44. doi: 10.1007/s11912-018-0690-1
- Centerwatch.com. FDA Approved Drugs for Oncology. 2019. Available from: <https://www.centerwatch.com/drug-information/fda-approved-drugs/therapeutic-area/12/oncology>. Accessed May 2020.
- Bonsu J, Charles L, Guha A, Awan F, Woyach J, Yildiz V, Wei L, Jneid H, Addison D. Representation of patients with cardiovascular disease in pivotal cancer clinical trials. *Circulation*. 2019;139:2594–2596. doi: 10.1161/CIRCULATIONAHA.118.039180
- Bonsu JM, Guha A, Charles L, Yildiz VO, Wei L, Baker B, Brammer JE, Awan F, Lustberg M, Reinbolt R, et al. Reporting of cardiovascular events in clinical trials supporting FDA approval of contemporary cancer therapies. *J Am Coll Cardiol*. 2020;75:620–628. doi: 10.1016/j.jacc.2019.11.059
- Madonna R, Cadeddu C, Deidda M, Mele D, Monte I, Novo G, Pagliaro P, Pepe A, Spallarossa P, Tocchetti CG, et al. Improving the preclinical models for the study of chemotherapy-induced cardiotoxicity: a position paper of the Italian working group on drug cardiotoxicity and cardioprotection. *Heart Fail Rev*. 2015;20:621–631. doi: 10.1007/s10741-015-9497-4
- Herrmann J, Lerman A, Sandhu NP, Villarraga HR, Mulvagh SL, Kohli M. Evaluation and management of patients with heart disease and cancer: cardio-oncology. *Mayo Clin Proc*. 2014;89:1287–1306. doi: 10.1016/j.mayocp.2014.05.013
- Hanlon P, Hannigan L, Rodriguez-Perez J, Fischbacher C, Welton NJ, Dias S, Mair FS, Guthrie B, Wild S, McAllister DA. Representation of people with comorbidity and multimorbidity in clinical trials of novel drug therapies: an individual-level participant data analysis. *BMC Med*. 2019;17:201. doi: 10.1186/s12916-019-1427-1
- Mosenifar Z. Population issues in clinical trials. *Proc Am Thorac Soc*. 2007;4:185–188. doi: 10.1513/pats.200701-009GC
- Sogaard M, Thomsen RW, Bossen KS, Sorensen HT, Norgaard M. The impact of comorbidity on cancer survival: a review. *Clin Epidemiol*. 2013;5:3–29. doi: 10.2147/CLEP.S47150
- Iachina M, Green A, Jakobsen E. The direct and indirect impact of comorbidity on the survival of patients with non-small cell lung cancer: a combination of survival, staging and resection models with missing measurements in covariates. *BMJ Open*. 2014;4:e003846. doi: 10.1136/bmjopen-2013-003846
- Saussele S, Krauss MP, Hehlmann R, Lauseker M, Proetel U, Kalmanti L, Hanfstein B, Fabarius A, Kraemer D, Berdel WE, et al. Impact of comorbidities on overall survival in patients with chronic myeloid leukemia: results of the randomized CML study IV. *Blood*. 2015;126:42–49. doi: 10.1182/blood-2015-01-617993
- Narayan HK, French B, Plappert T, Hyman D, Smith AM, Margulies KB, Ky B. Detailed echocardiographic phenotyping in breast cancer patients. *Circulation*. 2017;135:1397–1412. doi: 10.1161/CIRCULATIONAHA.116.023463
- Skovgaard D, Hasbak P, Kjaer A. BNP predicts chemotherapy-related cardiotoxicity and death: Comparison with gated equilibrium radionuclide ventriculography. *PLoS One*. 2014;9:e96736. doi: 10.1371/journal.pone.0096736
- Cardinale D, Sandri MT, Colombo A, Colombo N, Boeri M, Lamantia G, Civelli M, Peccatori F, Martinelli G, Fiorentini C, et al. Prognostic value of troponin I in cardiac risk stratification of cancer patients undergoing high-dose chemotherapy. *Circulation*. 2004;109:2749–2754. doi: 10.1161/01.CIR.0000130926.51766.CC
- Pudil R, Mueller C, Celutkienė J, Henriksen PA, Lenihan D, Dent S, Barac A, Stanway S, Moslehi J, Suter TM, et al. Role of serum biomarkers in cancer patients receiving cardiotoxic cancer therapies: a position statement from the cardio-oncology study group of the heart failure association and the cardio-oncology council of the European Society of Cardiology. *Eur J Heart Fail*. 2020;22:1966–1983. doi: 10.1002/ehf.2017
- Feola M, Garrone O, Occelli M, Francini A, Biggi A, Visconti G, Albrile F, Bobbio M, Merlano M. Cardiotoxicity after anthracycline chemotherapy in breast carcinoma: effects on left ventricular ejection fraction, troponin i and brain natriuretic peptide. *Int J Cardiol*. 2011;148:194–198. doi: 10.1016/j.ijcard.2009.09.564
- Leya J, Hayek S. American college of cardiology. Role of troponin and bnp in the management of patients with cancer. 2017. Available from: <https://www.acc.org/latest-in-cardiology/articles/2019/12/10/15/15/role-of-troponin-and-bnp-in-the-management-of-patients-with-cancer>. Accessed February 2022
- Zamorano JL, Lancellotti P, Rodriguez Munoz D, Aboyans V, Asteggiano R, Galderisi M, Habib G, Lenihan DJ, Lip GYH, Lyon AR, et al. 2016 ESC position paper on cancer treatments and cardiovascular toxicity developed under the auspices of the ESC committee for practice guidelines: the task force for cancer treatments and cardiovascular toxicity of the European Society of Cardiology (ESC). *Eur Heart J*. 2016;37:2768–2801. doi: 10.1093/eurheartj/ehw211
- de Baat EC, Naaktgeboren WR, Leiner T, Teske AJ, Habets J, Grotenhuis HB. Update in imaging of cancer therapy-related cardiac toxicity in adults. *Open Heart*. 2021;8:e001506. doi: 10.1136/openhrt-2020-001506
- Chang HM, Okwuosa TM, Scarabelli T, Moudgil R, Yeh ETH. Cardiovascular complications of cancer therapy: best practices in diagnosis, prevention, and management: part 2. *J Am Coll Cardiol*. 2017;70:2552–2565. doi: 10.1016/j.jacc.2017.09.1095
- Chen ZI, Ai DI. Cardiotoxicity associated with targeted cancer therapies. *Mol Clin Oncol*. 2016;4:675–681. doi: 10.3892/mco.2016.800
- Seruga B, Sterling L, Wang L, Tannock IF. Reporting of serious adverse drug reactions of targeted anticancer agents in pivotal phase III clinical trials. *J Clin Oncol*. 2011;29:174–185. doi: 10.1200/JCO.2010.31.9624
- Riaz H, Raza S, Khan MS, Riaz IB, Krasuski RA. Impact of funding source on clinical trial results including cardiovascular outcome trials. *Am J Cardiol*. 2015;116:1944–1947. doi: 10.1016/j.amjcard.2015.09.034
- Bekelman JE, Li Y, Gross CP. Scope and impact of financial conflicts of interest in biomedical research: a systematic review. *JAMA*. 2003;289:454–465. doi: 10.1001/jama.289.4.454

SUPPLEMENTAL MATERIAL

Table S1. List of all included studies.

Serial Number	Title	Citation
1	Sorafenib in combination with docetaxel as first-line therapy for HER2-negative metastatic breast cancer: Final results of the randomized, double-blind, placebo-controlled phase II MADONNA study	Mavratzas A, Baek S, Gerber B, Schmidt M, Moebus V, Foerster F, Grischke EM, Fasching P, Strumberg D, Solomayer E, et al. Sorafenib in combination with docetaxel as first-line therapy for HER2-negative metastatic breast cancer: Final results of the randomized, double-blind, placebo-controlled phase II MADONNA study. <i>Breast.</i> 2019;45:22-28
2	Randomized phase II study evaluating weekly oral vinorelbine versus weekly paclitaxel in estrogen receptor-positive, HER2-negative patients with advanced breast cancer (NorBreast-231 trial)	Aapro M, Ruiz-Borrego M, Hegg R, Kukielka-Budny B, Morales S, Cinieri S, Freitas-Junior R, Garcia-Estevez L, Szombara E, Borges GS, et al. Randomized phase ii study evaluating weekly oral vinorelbine versus weekly paclitaxel in estrogen receptor-positive, HER2-negative patients with advanced breast cancer (NorBreast-231 trial). <i>Breast.</i> 2019;45:7-14

3	<p>Concurrent neoadjuvant chemotherapy and estrogen deprivation in patients with estrogen receptor-positive, human epidermal growth factor receptor 2-negative breast cancer (CBCSG-036): A randomized, controlled, multicenter trial</p>	<p>Yu KD, Wu SY, Liu GY, Wu J, Di GH, Hu Z, Hou YF, Chen CM, Fan L, Tang LC, et al. Concurrent neoadjuvant chemotherapy and estrogen deprivation in patients with estrogen receptor-positive, human epidermal growth factor receptor 2-negative breast cancer (CBCSG-036): A randomized, controlled, multicenter trial. <i>Cancer</i>. 2019;125:2185-2193</p>
4	<p>A multi-national, randomised, open-label, parallel, phase III non-inferiority study comparing NK105 and paclitaxel in metastatic or recurrent breast cancer patients</p>	<p>Fujiwara Y, Mukai H, Saeki T, Ro J, Lin YC, Nagai SE, Lee KS, Watanabe J, Ohtani S, Kim SB, et al. A multi-national, randomised, open-label, parallel, phase III non-inferiority study comparing NK105 and paclitaxel in metastatic or recurrent breast cancer patients. <i>Br J Cancer</i>. 2019;120:475-480</p>
5	<p>Efficacy and safety of everolimus plus exemestane in postmenopausal women with hormone receptor-positive, human epidermal growth factor receptor 2-</p>	<p>Tesch H, Stoetzer O, Decker T, Kurbacher CM, Marme F, Schneeweiss A, Mundhenke C, Distelrath A, Fasching PA, Lux MP, et al. Efficacy and safety of everolimus plus exemestane in postmenopausal women with hormone</p>

	<p>negative locally advanced or metastatic breast cancer:</p> <p>Results of the single-arm, phase IIIB 4EVER trial</p>	<p>receptor-positive, human epidermal growth factor receptor 2-negative locally advanced or metastatic breast cancer:</p> <p>Results of the single-arm, phase IIIB 4EVER trial. <i>Int J Cancer</i>. 2019;144:877-885</p>
6	<p>PF-05280014 (a trastuzumab biosimilar) plus paclitaxel compared with reference trastuzumab plus paclitaxel for HER2-positive metastatic breast cancer: a randomised, double-blind study</p>	<p>Pegram MD, Bondarenko I, Zorzetto MMC, Hingmire S, Iwase H, Krivorotko PV, Lee KS, Li RK, Pikiel J, Aggarwal R, et al. PF-05280014 (a trastuzumab biosimilar) plus paclitaxel compared with reference trastuzumab plus paclitaxel for HER2-positive metastatic breast cancer: A randomised, double-blind study. <i>Br J Cancer</i>. 2019;120:172-182</p>
7	<p>Randomized Phase II Study Evaluating Palbociclib in Addition to Letrozole as Neoadjuvant Therapy in Estrogen Receptor–Positive Early Breast Cancer: PALLET Trial</p>	<p>Johnston S, Puhalla S, Wheatley D, Ring A, Barry P, Holcombe C, Boileau JF, Provencher L, Robidoux A, Rimawi M, et al. Randomized phase II study evaluating palbociclib in addition to letrozole as neoadjuvant therapy in estrogen receptor-positive early breast cancer: PALLET Trial. <i>J Clin Oncol</i>. 2019;37:178-189</p>

8	<p>Safety, activity, and molecular heterogeneity following neoadjuvant non-pegylated liposomal doxorubicin, paclitaxel, trastuzumab, and pertuzumab in HER2-positive breast cancer (Opti-HER HEART): an open-label, single-group, multicenter, phase 2 trial</p>	<p>Gavila J, Oliveira M, Pascual T, Perez-Garcia J, Gonzalez X, Canes J, Pare L, Calvo I, Ciruelos E, Munoz M, et al. Safety, activity, and molecular heterogeneity following neoadjuvant non-pegylated liposomal doxorubicin, paclitaxel, trastuzumab, and pertuzumab in HER2-positive breast cancer (Opti-HER HEART): An open-label, single-group, multicenter, phase 2 trial. <i>BMC Med.</i> 2019;17:8</p>
9	<p>Intense dose-dense epirubicin, paclitaxel, cyclophosphamide versus weekly paclitaxel, liposomal doxorubicin (plus carboplatin in triple-negative breast cancer) for neoadjuvant treatment of high-risk early breast cancer (GeparOcto—GBG 84): A randomised phase III trial</p>	<p>Schneeweiss A, Mobus V, Tesch H, Hanusch C, Denkert C, Lubbe K, Huober J, Klare P, Kummel S, Untch M, et al. Intense dose-dense epirubicin, paclitaxel, cyclophosphamide versus weekly paclitaxel, liposomal doxorubicin (plus carboplatin in triple-negative breast cancer) for neoadjuvant treatment of high-risk early breast cancer (GeparOcto—GBG 84): A randomised phase III trial. <i>Eur J Cancer.</i> 2019;106:181-192</p>

10	A randomized and open-label phase II trial reports the efficacy of neoadjuvant lobaplatin in breast cancer	Wu X, Tang P, Li S, Wang S, Liang Y, Zhong L, Ren L, Zhang T, Zhang Y. A randomized and open-label phase ii trial reports the efficacy of neoadjuvant lobaplatin in breast cancer. <i>Nat Commun.</i> 2018;9:832
11	Neoadjuvant chemotherapy with or without anthracyclines in the presence of dual HER2 blockade for HER2-positive breast cancer (TRAIN-2): a multicentre, open-label, randomised, phase 3 trial	van Ramshorst MS, van der Voort A, van Werkhoven ED, Mandjes IA, Kemper I, Dezentje VO, Oving IM, Honkoop AH, Tick LW, van de Wouw AJ, et al. Neoadjuvant chemotherapy with or without anthracyclines in the presence of dual her2 blockade for HER2-positive breast cancer (TRAIN-2): A multicentre, open-label, randomised, phase 3 trial. <i>Lancet Oncol.</i> 2018;19:1630-1640
12	A Randomized Phase II Study of Eribulin/Cyclophosphamide or Docetaxel/Cyclophosphamide as Neoadjuvant Therapy in	Yardley DA, Shipley D, Zubkus J, Wright GL, Ward PJ, Mani A, Shastry M, Finney L, DeBusk L, Hainsworth JD. A randomized phase ii study of eribulin/cyclophosphamide or docetaxel/cyclophosphamide as neoadjuvant therapy in operable HER2-

	Operable HER2-negative Breast Cancer	negative breast cancer. <i>Clin Breast Cancer</i> . 2019;19:1-9
13	Atezolizumab and nab-paclitaxel in advanced triple-negative breast cancer	Schmid P, Adams S, Rugo HS, Schneeweiss A, Barrios CH, Iwata H, Dieras V, Hegg R, Im SA, Shaw Wright G, et al. Atezolizumab and nab-paclitaxel in advanced triple-negative breast cancer. <i>N Engl J Med</i> . 2018;379:2108-2121
14	Combination versus sequential paclitaxel plus gemcitabine as first-line chemotherapy for women with metastatic breast cancer: a prospective randomized phase II study.	Shao B, Song G, Li H, Dil L, Jiang H, Liang X, Yan Y, Zhang R, Ran R, Wang J, et al. Combination versus sequential paclitaxel plus gemcitabine as first-line chemotherapy for women with metastatic breast cancer: A prospective randomized phase II study. <i>J BUON</i> . 2018;23:1583-1590
15	Fulvestrant plus goserelin versus anastrozole plus goserelin versus goserelin alone for hormone receptor-positive, HER2-negative tamoxifen-pretreated premenopausal women with	Kim JY, Im SA, Jung KH, Ro J, Sohn J, Kim JH, Park YH, Kim TY, Kim SB, Lee KS, et al. Fulvestrant plus goserelin versus anastrozole plus goserelin versus goserelin alone for hormone receptor-positive, her2-negative tamoxifen-pretreated premenopausal women with

	recurrent or metastatic breast cancer (KCSG BR10-04): a multicentre, open-label, three-arm, randomised phase II trial (FLAG study)	recurrent or metastatic breast cancer (KCSG BR10-04): A multicentre, open-label, three-arm, randomised phase II trial (FLAG study). <i>Eur J Cancer</i> . 2018;103:127-136
16	Intermittent versus continuous first-line treatment for HER2-negative metastatic breast cancer: the Stop & Go study of the Dutch Breast Cancer Research Group (BOOG)	Claessens AKM, Bos M, Lopez-Yurda M, Bouma JM, Rademaker-Lakhai JM, Honkoop AH, de Graaf H, van Druten E, van Warmerdam LJC, van der Sangen MJC, et al. Intermittent versus continuous first-line treatment for her2-negative metastatic breast cancer: The stop & go study of the dutch breast cancer research group (BOOG). <i>Breast Cancer Res Treat</i> . 2018;172:413-423
17	Paclitaxel With Inhibitor of Apoptosis Antagonist, LCL161, for Localized Triple-Negative Breast Cancer, Prospectively Stratified by Gene Signature in a Biomarker-Driven Neoadjuvant Trial	Bardia A, Parton M, Kummel S, Estevez LG, Huang CS, Cortes J, Ruiz-Borrego M, Telli ML, Martin-Martorell P, Lopez R, et al. Paclitaxel with inhibitor of apoptosis antagonist, LCL161, for localized triple-negative breast cancer, prospectively stratified by gene signature in a biomarker-

		driven neoadjuvant trial. <i>J Clin Oncol.</i> 2018;JCO2017748392
18	Multicenter Phase II Study of Lurbinectedin in BRCA-Mutated and Unselected Metastatic Advanced Breast Cancer and Biomarker Assessment Substudy	Cruz C, Llop-Guevara A, Garber JE, Arun BK, Perez Fidalgo JA, Lluch A, Telli ML, Fernandez C, Kahatt C, Galmarini CM, et al. Multicenter phase II study of lurbinectedin in BRCA -mutated and unselected metastatic advanced breast cancer and biomarker assessment substudy. <i>J Clin Oncol.</i> 2018;36:3134-3143
19	Everolimus Plus Exemestane vs Everolimus or Capecitabine Monotherapy for Estrogen Receptor–Positive, HER2-Negative Advanced Breast CancerThe BOLERO-6 Randomized Clinical Trial	Jerusalem G, de Boer RH, Hurvitz S, Yardley DA, Kovalenko E, Ejlersen B, Blau S, Ozguroglu M, Landherr L, Ewertz M, et al. Everolimus plus exemestane vs everolimus or capecitabine monotherapy for estrogen receptor-positive, HER2-negative advanced breast cancer: The BOLERO-6 randomized clinical trial. <i>JAMA Oncol.</i> 2018;4:1367-1374
20	UCBG 2-08: 5-year efficacy results from the UNICANCER-PACS08	Campono M, Lacroix-Triki M, Roca L, Spielmann M, Wildiers H, Cottu P, Kerbrat P, Levy C, Desmoulins I, Bachelot T, et al.

	randomised phase III trial of adjuvant treatment with FEC100 and then either docetaxel or ixabepilone in patients with early-stage, poor prognosis breast cancer	UCBG 2-08: 5-year efficacy results from the UNICANCER-PACS08 randomised phase iii trial of adjuvant treatment with FEC100 and then either docetaxel or ixabepilone in patients with early-stage, poor prognosis breast cancer. <i>Eur J Cancer.</i> 2018;103:184-194
21	Adjuvant dose-dense doxorubicin-cyclophosphamide versus docetaxel-doxorubicin-cyclophosphamide for high-risk breast cancer: First results of the randomised MATADOR trial (BOOG 2004-04)	van Rossum AGJ, Kok M, van Werkhoven E, Opdam M, Mandjes IAM, van Leeuwen-Stok AE, van Tinteren H, Imholz ALT, Portielje JEA, et al. Adjuvant dose-dense doxorubicin-cyclophosphamide versus docetaxel-doxorubicin-cyclophosphamide for high-risk breast cancer: First results of the randomised MATADOR trial (BOOG 2004-04). <i>Eur J Cancer.</i> 2018;102:40-48
22	Results From the First Multicenter, Open-label, Phase IIIb Study Investigating the Combination of Pertuzumab With Subcutaneous Trastuzumab and a Taxane in Patients	Woodward N, De Boer RH, Redfern A, White M, Young J, Truman M, Beith J. Results from the first multicenter, open-label, phase IIIb study investigating the combination of pertuzumab with subcutaneous trastuzumab and a taxane in patients with HER2-positive metastatic

	With HER2-positive Metastatic Breast Cancer (SAPPHIRE)	breast cancer (SAPPHIRE). <i>Clin Breast Cancer</i> . 2019;19:216-224
23	Effect of Adjuvant Trastuzumab for a Duration of 9 Weeks vs 1 Year With Concomitant Chemotherapy for Early Human Epidermal Growth Factor Receptor 2-Positive Breast Cancer: The SOLD Randomized Clinical Trial	Joensuu H, Fraser J, Wildiers H, Huovinen R, Auvinen P, Utriainen M, Nyandoto P, Villman KK, Halonen P, Granstam-Bjorneklett H, et al. Effect of adjuvant trastuzumab for a duration of 9 weeks vs 1 year with concomitant chemotherapy for early human epidermal growth factor receptor 2-positive breast cancer: The SOLD randomized clinical trial. <i>JAMA Oncol</i> . 2018;4:1199-1206
24	Double-Blind Phase III Trial of Adjuvant Chemotherapy With and Without Bevacizumab in Patients With Lymph Node-Positive and High-Risk Lymph Node-Negative Breast Cancer (E5103)	Miller KD, O'Neill A, Gradishar W, Hobday TJ, Goldstein LJ, Mayer IA, Bloom S, Brufsky AM, Tevaarwerk AJ, Sparano JA, et al. Double-blind phase III trial of adjuvant chemotherapy with and without bevacizumab in patients with lymph node-positive and high-risk lymph node-negative breast cancer (E5103). <i>J Clin Oncol</i> . 2018;36:2621-2629

25	Talazoparib in patients with advanced breast cancer and a germline BRCA mutation	Litton JK, Rugo HS, Ettl J, Hurvitz SA, Goncalves A, Lee KH, Fehrenbacher L, Yerushalmi R, Mina LA, Martin M, et al. Talazoparib in patients with advanced breast cancer and a germline brca mutation. <i>N Engl J Med.</i> 2018;379:753-763
26	Phase III Randomized Study of Ribociclib and Fulvestrant in Hormone Receptor–Positive, Human Epidermal Growth Factor Receptor 2–Negative Advanced Breast Cancer: MONALEESA-3	Slamon DJ, Neven P, Chia S, Fasching PA, De Laurentiis M, Im SA, Petrakova K, Bianchi GV, Esteva FJ, Martin M, et al. Phase iii randomized study of ribociclib and fulvestrant in hormone receptor-positive, human epidermal growth factor receptor 2-negative advanced breast cancer: MONALEESA-3. <i>J Clin Oncol.</i> 2018;36:2465-2472
27	A randomized, double-blind, phase 2 study of ruxolitinib or placebo in combination with capecitabine in patients with advanced HER2-negative breast cancer and elevated	O'Shaughnessy J, DeMichele A, Ma CX, Richards P, Yardley DA, Wright GS, Kalinsky K, Steis R, Diab S, Kennealey G, et al. A randomized, double-blind, phase 2 study of ruxolitinib or placebo in combination with capecitabine in patients with advanced her2-negative breast

	C-reactive protein, a marker of systemic inflammation	cancer and elevated c-reactive protein, a marker of systemic inflammation. <i>Breast Cancer Res Treat.</i> 2018;170:547-557
28	A randomized phase II trial of trastuzumab plus capecitabine versus lapatinib plus capecitabine in patients with HER2-positive metastatic breast cancer previously treated with trastuzumab and taxanes: WJOG6110B/ELTOP	Takano T, Tsurutani J, Takahashi M, Yamanaka T, Sakai K, Ito Y, Fukuoka J, Kimura H, Kawabata H, Tamura K, et al. A randomized phase ii trial of trastuzumab plus capecitabine versus lapatinib plus capecitabine in patients with her2-positive metastatic breast cancer previously treated with trastuzumab and taxanes: WJOB6110B/ELTOP. <i>Breast.</i> 2018;40:67-75
29	Neoadjuvant PF-05280014 (a potential trastuzumab biosimilar) versus trastuzumab for operable HER2+ breast cancer	Lammers PE, Dank M, Masetti R, Abbas R, Hilton F, Coppola J, Jacobs I. Neoadjuvant PF-05280014 (a potential trastuzumab biosimilar) versus trastuzumab for operable HER2+ breast cancer. <i>Br J Cancer.</i> 2018;119:266-273
30	Efficacy and safety of ABP 980 compared with reference trastuzumab in women with HER2-positive	von Minckwitz G, Colleoni M, Kolberg HC, Morales S, Santi P, Tomasevic Z, Zhang N, Hanes V. Efficacy and safety of ABP 980 compared with reference trastuzumab

	early breast cancer (LILAC study): a randomised, double-blind, phase 3 trial	in women with HER2-positive early breast cancer (LILAC study): A randomised, double-blind, phase 3 trial. <i>Lancet Oncol.</i> 2018;19:987-998
31	Bi-weekly eribulin therapy for metastatic breast cancer: a multicenter phase II prospective study (JUST-STUDY)	Ohtani S, Nakayama T, Yoshinami T, Watanabe KI, Hara F, Sagara Y, Kawaguchi H, Higaki K, Matsunami N, Hasegawa Y, et al. Bi-weekly eribulin therapy for metastatic breast cancer: A multicenter phase II prospective study (JUST-STUDY). <i>Breast Cancer.</i> 2018;25:438-446
32	Efficacy and safety of trastuzumab, lapatinib, and paclitaxel neoadjuvant treatment with or without prolonged exposure to anti-HER2 therapy, and with or without hormone therapy for HER2-positive primary breast cancer: a randomised, five-arm, multicentre, open-label phase II trial	Masuda N, Toi M, Yamamoto N, Iwata H, Kuroi K, Bando H, Ohtani S, Takano T, Inoue K, Yanagita Y, et al. Efficacy and safety of trastuzumab, lapatinib, and paclitaxel neoadjuvant treatment with or without prolonged exposure to anti-HER2 therapy, and with or without hormone therapy for HER2-positive primary breast cancer: A randomised, five-arm, multicentre, open-label phase II trial. <i>Breast Cancer.</i> 2018;25:407-415

33	<p>Ribociclib plus endocrine therapy for premenopausal women with hormone-receptor-positive, advanced breast cancer (MONALEESA-7): a randomised phase 3 trial</p>	<p>Tripathy D, Im SA, Colleoni M, Franke F, Bardia A, Harbeck N, Hurvitz SA, Chow L, Sohn J, Lee KS, et al. Ribociclib plus endocrine therapy for premenopausal women with hormone-receptor-positive, advanced breast cancer (MONALEESA-7): A randomised phase 3 trial. <i>Lancet Oncol.</i> 2018;19:904-915</p>
34	<p>Neoadjuvant endocrine therapy with exemestane followed by response-guided combination therapy with low-dose cyclophosphamide in postmenopausal patients with estrogen receptor-positive breast cancer: A multicenter, open-label, phase II study</p>	<p>Sato N, Masuda N, Morimoto T, Ueno T, Kanbayashi C, Kaneko K, Yasojima H, Saji S, Sasano H, Morita S, et al. Neoadjuvant endocrine therapy with exemestane followed by response-guided combination therapy with low-dose cyclophosphamide in postmenopausal patients with estrogen receptor-positive breast cancer: A multicenter, open-label, phase II study. <i>Cancer Med.</i> 2018</p>
35	<p>Induction therapy with paclitaxel and bevacizumab followed by switch maintenance therapy with eribulin in Japanese patients</p>	<p>Inoue K, Ninomiya J, Saito T, Kimizuka K, Kurosumi M. Induction therapy with paclitaxel and bevacizumab followed by switch maintenance therapy with eribulin in japanese patients with HER2-negative</p>

	with HER2-negative metastatic breast cancer: A multicenter, collaborative, open-label, phase II clinical study for the SBCCSG 35 investigators	metastatic breast cancer: A multicenter, collaborative, open-label, phase II clinical study for the SBCCSG 35 investigators. <i>BMC Cancer</i> . 2018;18:671
36	The efficacy and feasibility of dose-dense sequential chemotherapy for Japanese patients with breast cancer	Takabatake D, Kajiwara Y, Ohtani S, Itano Y, Yamamoto M, Kubo S, Ikeda M, Takahashi M, Hara F, Aogi K, et al. The efficacy and feasibility of dose-dense sequential chemotherapy for japanese patients with breast cancer. <i>Breast Cancer</i> . 2018;25:717-722
37	Randomized phase II trial of fulvestrant plus everolimus or placebo in postmenopausal women with hormone receptor-positive, human epidermal growth factor receptor 2-negative metastatic breast cancer resistant to aromatase	Kornblum N, Zhao F, Manola J, Klein P, Ramaswamy B, Brufsky A, Stella PJ, Burnette B, Telli M, Makower DF, et al. Randomized phase ii trial of fulvestrant plus everolimus or placebo in postmenopausal women with hormone receptor-positive, human epidermal growth factor receptor 2-negative metastatic breast cancer resistant to aromatase

	inhibitor therapy: Results of PrE0102	inhibitor therapy: Results of PrE0102. <i>J Clin Oncol.</i> 2018;36:1556-1563
38	A randomized Phase III trial of neoadjuvant recombinant human endostatin, docetaxel and epirubicin as first-line therapy for patients with breast cancer (CBCRT01)	Chen J, Yao Q, Huang M, Wang B, Zhang J, Wang T, Ming Y, Zhou X, Jia Q, Huan Y, et al. A randomized phase iii trial of neoadjuvant recombinant human endostatin, docetaxel and epirubicin as first-line therapy for patients with breast cancer (CBCRT01). <i>Int J Cancer.</i> 2018;142:2130-2138
39	Carboplatin in BRCA1/2-mutated and triple-negative breast cancer BRCAness subgroups: The TNT Trial	Tutt A, Tovey H, Cheang MCU, Kernaghan S, Kilburn L, Gazinska P, Owen J, Abraham J, Barrett S, Barrett-Lee P, et al. Carboplatin in brca1/2-mutated and triple-negative breast cancer BRCAness subgroups: The TNT Trial. <i>Nat Med.</i> 2018;24:628-637
40	Open-label randomised phase III trial of vinflunine versus an alkylating agent in patients with heavily pretreated metastatic breast cancer	Cortes J, Perez-Garcia J, Levy C, Gomez Pardo P, Bourgeois H, Spazzapan S, Martinez-Janez N, Chao TC, Espie M, Nabholz JM, et al. Open-label randomised phase III trial of vinflunine versus an alkylating agent in patients with heavily

		pretreated metastatic breast cancer. <i>Ann Oncol.</i> 2018;29:881-887
41	Everolimus Plus Letrozole for Treatment of Patients With HR(+), HER2(-) Advanced Breast Cancer Progressing on Endocrine Therapy: An Open-label, Phase II Trial	Safra T, Kaufman B, Kadouri L, Efrat Ben-Baruch N, Ryvo L, Nisenbaum B, Evron E, Yerushalmi R. Everolimus plus letrozole for treatment of patients with HR(+), HER2(-) advanced breast cancer progressing on endocrine therapy: An open-label, phase II trial. <i>Clin Breast Cancer.</i> 2018;18:e197-e203
42	Randomized phase II study of anastrozole plus tegafur-uracil as neoadjuvant therapy for ER-positive breast cancer in postmenopausal Japanese women (Neo-ACET BC)	Nakayama T, Sagara Y, Takashima T, Matsunami N, Masuda N, Miyoshi Y, Taguchi T, Aono T, Ito T, Kagimura T, et al. Randomized phase II study of anastrozole plus tegafur-uracil as neoadjuvant therapy for ER- positive breast cancer in postmenopausal japanese women (Neo-ACET BC). <i>Cancer Chemother Pharmacol.</i> 2018;81:755-762
43	Phase III, randomized, double-blind study comparing the efficacy, safety, and immunogenicity of SB3	Pivot X, Bondarenko I, Nowecki Z, Dvorkin M, Trishkina E, Ahn JH, Vinnyk Y, Im SA, Sarosiek T, Chatterjee S, et al. Phase III, randomized, double-blind study comparing


	(trastuzumab biosimilar) and reference trastuzumab in patients treated with neoadjuvant therapy for human epidermal growth factor receptor 2-positive early breast cancer	the efficacy, safety, and immunogenicity of SB3 (trastuzumab biosimilar) and reference trastuzumab in patients treated with neoadjuvant therapy for human epidermal growth factor receptor 2-positive early breast cancer. <i>J Clin Oncol.</i> 2018;36:968-974
44	Adjuvant anastrozole versus exemestane versus letrozole, upfront or after 2 years of tamoxifen, in endocrine-sensitive breast cancer (FATA-GIM3): a randomised, phase 3 trial	De Placido S, Gallo C, De Laurentiis M, Bisagni G, Arpino G, Sarobba MG, Riccardi F, Russo A, Del Mastro L, Cogoni AA, et al. Adjuvant anastrozole versus exemestane versus letrozole, upfront or after 2 years of tamoxifen, in endocrine-sensitive breast cancer (FATA-GIM3): A randomised, phase 3 trial. <i>Lancet Oncol.</i> 2018;19:474-485
45	Addition of the PARP inhibitor veliparib plus carboplatin or carboplatin alone to standard neoadjuvant chemotherapy in triple-negative breast cancer (BrighTNess): a randomised, phase 3 trial	Loibl S, O'Shaughnessy J, Untch M, Sikov WM, Rugo HS, McKee MD, Huober J, Golshan M, von Minckwitz G, Maag D, et al. Addition of the PARP inhibitor veliparib plus carboplatin or carboplatin alone to standard neoadjuvant chemotherapy in triple-negative breast cancer

		(BrighTNess): A randomised, phase 3 trial. <i>Lancet Oncol.</i> 2018;19:497-509
46	Phase III, Randomized study of dual human epidermal growth factor receptor 2 (HER2) blockade with lapatinib plus trastuzumab in combination with an aromatase inhibitor in postmenopausal women with HER2-positive, hormone receptor-positive metastatic breast cancer: ALTERNATIVE	Johnston SRD, Hegg R, Im SA, Park IH, Burdaeva O, Kurteva G, Press MF, Tjulandin S, Iwata H, Simon SD, et al. Phase III, randomized study of dual human epidermal growth factor receptor 2 (HER2) blockade with lapatinib plus trastuzumab in combination with an aromatase inhibitor in postmenopausal women with her2-positive, hormone receptor-positive metastatic breast cancer: ALTERNATIVE. <i>J Clin Oncol.</i> 2018;36:741-748
47	Pertuzumab and trastuzumab with or without metronomic chemotherapy for older patients with HER2-positive metastatic breast cancer (EORTC 75111-10114): an open-label, randomised, phase 2 trial from the Elderly	Wildiers H, Tryfonidis K, Dal Lago L, Vuylsteke P, Curigliano G, Waters S, Brouwers B, Altintas S, Touati N, Cardoso F, et al. Pertuzumab and trastuzumab with or without metronomic chemotherapy for older patients with HER2-positive metastatic breast cancer (EORTC 75111-10114): An open-label, randomised, phase 2 trial from the Elderly Task Force/Breast

	Task Force/Breast Cancer Group	Cancer Group. <i>Lancet Oncol.</i> 2018;19:323-336
48	Comparing neoadjuvant nab-paclitaxel vs paclitaxel both followed by anthracycline regimens in women with ERBB2/ HER2-negative breast cancer-the evaluating treatment with neoadjuvant abraxane (ETNA) trial a randomized phase 3 clinical trial	Gianni L, Mansutti M, Anton A, Calvo L, Bisagni G, Bermejo B, Semiglazov V, Thill M, Chacon JI, Chan A, et al. Comparing neoadjuvant nab-paclitaxel vs paclitaxel both followed by anthracycline regimens in women with ERBB2/ HER2-negative breast cancer-the evaluating treatment with neoadjuvant abraxane (ETNA) trial: A randomized phase 3 clinical trial. <i>JAMA Oncol.</i> 2018;4:302-308
49	Sequential versus simultaneous use of chemotherapy and gonadotropin-releasing hormone agonist (GnRHa) among estrogen receptor (ER)-positive premenopausal breast cancer patients: Effects on ovarian function, disease-free survival, and overall survival	Zhang Y, Ji Y, Li J, Lei L, Wu S, Zuo W, Jia X, Wang Y, Mo M, Zhang N, et al. Sequential versus simultaneous use of chemotherapy and gonadotropin-releasing hormone agonist (GnRHa) among estrogen receptor (ER)-positive premenopausal breast cancer patients: Effects on ovarian function, disease-free survival, and overall survival. <i>Breast Cancer Res Treat.</i> 2018;168:679-686

50	Neoadjuvant letrozole for postmenopausal estrogen receptor-positive, HER2-negative breast cancer patients, a study from the Danish Breast Cancer Cooperative Group (DBCG)	Skriver SK, Laenkholm AV, Rasmussen BB, Handler J, Grundtmann B, Tvedskov TF, Christiansen P, Knoop AS, Jensen MB, Ejlersen B. Neoadjuvant letrozole for postmenopausal estrogen receptor-positive, HER2-negative breast cancer patients, a study from the Danish Breast Cancer Cooperative Group (DBCG). <i>Acta Oncol.</i> 2018;57:31-37
----	---	--

Figure S1. Data collection strategy.



*The search string applied was “breast:ti,ab,kw AND cancer:ti,ab,kw AND ([controlled clinical trial]/lim OR [randomized controlled trial]/lim)”.