



MEETING ABSTRACT

Open Access

# Maternal proviral load and vertical transmission of Human T cell Lymphotropic Virus type 1 in Guinea-Bissau

Carla van Tienen<sup>1,2\*</sup>, Samuel J McConkey<sup>1</sup>, Thushan I de Silva<sup>1,3</sup>, Matthew Cotten<sup>1</sup>, Steve Kaye<sup>1</sup>, Ramu Sarge-Njie<sup>1</sup>, Carlos da Costa<sup>1,4</sup>, Nato Gonçalves<sup>1,4</sup>, Julia Parker<sup>1</sup>, Tim Vincent<sup>1</sup>, Assan Jaye<sup>1</sup>, Peter Aaby<sup>5</sup>, Hilton Whittle<sup>1</sup>, Maarten Schim van der Loeff<sup>1,6</sup>

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses  
Leuven and Gembloux, Belgium. 5-8 June 2011

## Background

The relative importance of routes of transmission of Human T cell Lymphotropic Virus type 1 (HTLV-1) in Guinea-Bissau is largely unknown; vertical transmission is thought to be important, but there are very few existing data. We aimed to examine factors associated with transmission in mothers and children in Guinea-Bissau, where HTLV-1 is endemic (prevalence of 5%).

## Methods

A cross-sectional survey was performed among mothers and their children (aged <15 years) in a rural community in Guinea-Bissau. A questionnaire to identify risk factors for infection and a blood sample were obtained. HTLV-1 proviral load in peripheral blood was determined and PCR was performed to compare Long Terminal Repeat (LTR) sequences in mother-child pairs.

## Results

Fourteen out of 55 children (25%) of 31 HTLV-1 infected mothers were infected versus none of 70 children of 30 uninfected mothers. The only factor significantly associated with HTLV-1 infection in the child was the proviral load of the mother; the risk of infection increased significantly with the log<sub>10</sub> proviral load in the mother's peripheral blood (OR 5.5, 95% CI 2.1-14.6, per quartile), adjusted for weaning age and maternal income. HTLV-1 sequences of the LTR region obtained

from mother-child pairs were identical within pairs but differed between the pairs.

## Conclusions

Vertical transmission plays an important role in HTLV-1 transmission in this community in Guinea-Bissau. The risk of transmission increases with the mother's proviral load in the peripheral blood. Identical sequences in mother-child pairs give additional support to the maternal source of the children's infection.

## Author details

<sup>1</sup>Viral diseases program, Medical Research Council, Fajara, Gambia.

<sup>2</sup>Department of Medical Microbiology and Infectious Diseases, Erasmus Medical Centre, Rotterdam, Netherlands. <sup>3</sup>Centre for Medical Molecular Virology, Division of Infection and Immunity, University College London, London, UK. <sup>4</sup>Projeto de Saúde de Bandim, Bissau, Guinea-Bissau. <sup>5</sup>Projeto de Saúde de Bandim/Indepth Network, Bissau, Guinea-Bissau. <sup>6</sup>Municipal Health Services, Amsterdam, Netherlands.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A72

Cite this article as: van Tienen et al.: Maternal proviral load and vertical transmission of Human T cell Lymphotropic Virus type 1 in Guinea-Bissau. *Retrovirology* 2011 **8**(Suppl 1):A72.

\* Correspondence: carlavantienen@gmail.com

<sup>1</sup>Viral diseases program, Medical Research Council, Fajara, Gambia  
Full list of author information is available at the end of the article