

MEETING ABSTRACT

Open Access

Basic and clinical immunology – 3032. 18FDG-PET/CT findings (SUV-MAX) are correlated with survival in stage-4 squamous cell carcinoma

Arzu Didem Yalcin^{1*}, Metin Erkilic², Bulent Kargi³, Aysegul Kargi⁴, Burhan Savas⁵, Antonia Dimitrakopoulou-Strauss⁶, Ludwig Strauss⁶

From 2nd WAO International Scientific Conference (WISC 2012) Hyderabad, India. 6-9 December 2012

Background

Gene promoter hypermethylation is now regarded as a promising biomarker for the risk and progression of lung cancer. Vit-D is a steroid hormone that is now widely accepted that it exerts several extraskelatal actions, including anti-tumorigenic and immunomodulatory effects in vitro and in vivo. Its potential in cancer prevention and treatment is currently under detailed investigation. With the prospect of more effective therapeutic options for advanced stage disease, there are still insufficient current follow-up procedures for lung cancer. Therefore, in the present study, we investigated the concentrations of sCD200, sTRAIL, Vit-D, Hcy in the peripheral blood of metastatic non-small cell lung cancer (MNSCLC) patients. As well as the corresponding F-18-Deoxyglucose positron emission tomography (PET/CT) results were evaluate

Methods

Consecutive 22 adenocarcinoma and 15 squamous cell carcinoma patients of MNSCLC referred to our institute were included in this study. The sCD200, sTRAIL, Hcy and Vit-D concentrations in the serum samples from the 37 NSCLC patients were analyzed using a TRAIL/APO2L ELISA kit (Diaclone, France) and Roche kit for Vit-D and Hcy according to the manufacturer's instructions. The absorbance of each patient on a spectrophotometer using 450 nm and the concentration of sTRAIL (pg/mL), sCD200 (pg/mL), Hcy (mmol/L) and Vit-D (ng/mL) was measured.

Results

Levels of sCD200, sTRAIL, Vit-D, Hcy are not to be correlated with survival in patients with MNSCLC. In patients with squamous cell carcinoma SUVMAX and survival are found to be correlated. The correlation is $r = -0.6954$, $p = 0.004$, $n = 15$. The data were evaluated using the StataMP software package, version 12.1 (StataCorp, College Station, TX, USA) on a Mac Pro 2x2.93 GHz, 2*6 Core Intel Xeon system with 24GB RAM using Mac OS X 10.7.4 (Apple, Cupertino, CA, USA). An error level of $p < 0.05$ was generally used for the statistical analysis. The Wilcoxon matched pairs signed rank test was used to assess differences in the variables prior and three months after onset of treatment. Correlation tables were calculated for all variables. Furthermore, a multivariate correlation/regression analysis was applied to the data, using the survival as the dependent variable.

Conclusion

F-18-Deoxyglucose (FDG), a glucose analogue, is an advanced imaging technique and allows a highly sensitive whole body search for malignant foci, which are detected by their increased glucose metabolism versus benign tissues, and successful scanning has been performed in a wide variety of cancers.

Author details

¹Internal Medicine, Allergy and Immunology, Education and Research Hospital, Turkey. ²Department of Nuclear Medicine, Akdeniz University Hospital, Turkey. ³Medical Park Hospital, Turkey. ⁴Department of Internal Medicine, Antalya Education and Research Hospital, Antalya, Turkey. ⁵Department of Internal Medicine, Oncology Unit, Akdeniz University Hospital, Antalya, Turkey. ⁶Clinical Cooperation Unit Nuclear Medicine, German Cancer Research Center, Im Neuenheimer Feld 280, 69120, Heidelberg, Germany.

¹Internal Medicine, Allergy and Immunology, Education and Research Hospital, Turkey
Full list of author information is available at the end of the article

Published: 23 April 2013

doi:10.1186/1939-4551-6-S1-P207

Cite this article as: Yalcin *et al.*: Basic and clinical immunology – 3032. 18FDG-PET/CT findings (SUV-MAX) are correlated with survival in stage-4 squamous cell carcinoma. *World Allergy Organization Journal* 2013 6(Suppl 1):P207.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

