

POSTER PRESENTATION

Open Access

Clinical scale zinc finger nuclease (ZFN)-driven gene-editing of PD-1 in tumor infiltrating lymphocytes (TIL) for the potential treatment of metastatic melanoma

Joal D Beane^{1*}, Gary K Lee², Zhili Zheng¹, Nimisha Gandhi², Daniel Abate-Daga¹, Mini Bharathan¹, Mary Black¹, Matthew Mendel², Zhiya Yu¹, Sadik H Kassim¹, Smita Chandran¹, Martin Giedlin², Dale Ando², Ed Rebar², Andreas Reik², Michael Holmes², Philip D Gregory², Nicholas P Restifo³, Steven A Rosenberg⁴, Richard A Morgan⁵, Steven A Feldman¹

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Multiple inhibitory pathways exist to block the immune response to cancer potentially limiting the effectiveness of adoptive cell transfer (ACT). Programmed cell death-1 (PD-1) is a member of the CD28 superfamily and is expressed on activated T cells. Its ligands, PDL-1 and PDL-2 are expressed on a variety of tumor cells, including melanoma. The binding of PD-1 to PDL-1 inhibits T cell effector function, and represents an important mechanism for PDL-1 expressing tumors to evade the host immune response to cancer. PD-1 thus represents an attractive target for gene-editing of tumor-targeted T cells prior to ACT. To this end, our aim was to eliminate PD-1 expression in tumor infiltrating lymphocytes (TIL) by genome-editing using zinc finger nucleases (ZFNs) directed against the PD-1 gene at a scale sufficient for patient treatment. Using the MaxCyte GT Flow Transfection System to deliver mRNA encoding the PD-1 ZFNs, we show that our clinical scale TIL production process yielded efficient modification of the PD-1 gene locus, with an average modification frequency of 74.8% (n = 3, range 69.9 - 84.1%) of the alleles in a bulk TIL population, which resulted in a 76% reduction in PD-1 surface-expression. Importantly, the PD-1 gene-edited TIL product displayed an effector memory phenotype and expanded approximately 500 - 2000 fold during a rapid cell expansion *in vitro* while retaining T cell effector function. Thus further

study to determine the safety of adoptive cell transfer using PD-1 gene-edited TIL for the treatment of metastatic melanoma is warranted.

Authors' details

¹Surgery Branch, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA. ²Sangamo Biosciences, Richmond, CA, USA. ³National Cancer Institute, Bethesda, MD, USA. ⁴US National Institutes of Health (NIH), Bethesda, MD, USA. ⁵bluebird bio, Cambridge, MA, USA.

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P2

Cite this article as: Beane et al.: Clinical scale zinc finger nuclease (ZFN)-driven gene-editing of PD-1 in tumor infiltrating lymphocytes (TIL) for the potential treatment of metastatic melanoma. *Journal for ImmunoTherapy of Cancer* 2014 2(Suppl 3):P2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Surgery Branch, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

Full list of author information is available at the end of the article