

RESEARCH ARTICLE

Open Access

Movement disorders in neuroleptic-naïve patients with schizophrenia spectrum disorders

Moges Ayehu¹, Teshome Shibre^{2,3}, Barkot Milkias² and Abebaw Fekadu^{2,4*}

Abstract

Background: Spontaneous Movements Disorders (SMDs) or dyskinetic movements are often seen in patients with schizophrenia and other psychotic disorders, and are widely considered to be adverse consequences of the use of antipsychotic medications. Nevertheless, SMDs are also observed in the pre-neuroleptic era and among patients who were never exposed to antipsychotic medications. The aim of this study was to determine the extent of SMDs among antipsychotic-naïve patients in a low income setting, and to evaluate contextually relevant risk factors.

Methods: The study was a cross-sectional facility-based survey conducted at a specialist psychiatric hospital in Addis Ababa, Ethiopia. Consecutive consenting treatment-naïve patients with a diagnosis of schizophrenia, schizoaffective disorder and schizophreniform disorder contacting services for the first time were assessed using the Simpson-Angus Rating Scale (SAS) and the Abnormal Involuntary Movement Scale (AIMS) to evaluate the presence of SMDs. Scale for the Assessment of Negative Symptoms (SANS) and Scale for the Assessment of Positive Symptoms (SAPS) were administered to evaluate negative and positive symptom profiles respectively. Body mass index (BMI) was used as a proxy measure for nutritional status.

Result: Sixty-four patients, 67.2% male (n = 43), with first contact psychosis who met the DSM-IV-TR criteria for schizophrenia (n = 47), schizophreniform disorder (n = 5), and schizoaffective disorder (n = 12) were assessed over a two month study period. Seven patients (10.9%) had SMDs. BMI (OR = 0.6, 95% CI = 0.40, 0.89; p = 0.011) and increasing age (OR = 1.10; 95% CI = 1.02, 1.20; p = 0.017) were associated with SMD.

Conclusions: This finding supports previous suggestions that abnormal involuntary movements in schizophrenia and other psychotic disorders may be related to the pathophysiology of psychotic disorders and therefore cannot be attributed entirely to the adverse effects of neuroleptic medication.

Keywords: Parkinsonism, Dyskinesia, Spontaneous movement disorders, Dyskinetic movements, Antipsychotics, Psychotic disorder, Schizophrenia, Schizophreniform disorder, Schizoaffective disorder, Ethiopia, Developing country

Background

Different neurological motor abnormalities, especially dyskinesia and parkinsonism, have been reported in patients with schizophrenia [1-5]. With the introduction of antipsychotic medications and their numerous neurological adverse effects, these neurological abnormalities were generally presumed to be consequences of treatment with these newly introduced medications [6]. However, a wide range of neurological motor disturbances has

been reported in patients with schizophrenia who have never been treated with antipsychotic drugs, pre-dating the antipsychotic-era [7-10]. For example, research in the past two decades has identified spontaneous dyskinesia in a proportion of drug-naïve patients with schizophrenia [11-14]. Spontaneous movement disorders (SMD), including tardive dyskinesia-like syndrome, were reported in 4% to 11% of patients with no prior treatment history with antipsychotic medications [10-17]. In a recent review of 13 studies, the median rate of spontaneous dyskinesia was 9%, with spontaneous parkinsonism being the commoner abnormality [5]. Muscle rigidity and bradykinesia, often referred to together as akinetic type signs, are reported to be the most common parkinsonian signs in drug-naïve

* Correspondence: Abe.wassie@kcl.ac.uk

²Department of Psychiatry, Addis Ababa University, College of Health Sciences, School of Medicine, Addis Ababa, Ethiopia

⁴Department of Psychological Medicine, Centre for Affective Disorders, King's College London, Institute of Psychiatry, London, UK

Full list of author information is available at the end of the article

patients and occur more often than non-akinetic type signs such as tremor, glabellar tap, and salivation [12,18-22]. Non-akinetic parkinsonism becomes more severe following treatment with antipsychotics than the akinetic type and it has been suggested that it may be primarily a drug-induced phenomenon [21,22]. The most commonly reported dyskinesic symptoms are involuntary orofacial movements [8,14]. SMDs have also been reported in relatives of those with schizophrenia and dyskinesia, as well as in subjects with schizophrenia spectrum personality conditions, especially in schizotypal personality [17,23,24].

The main factors reported to have significant association with SMD were patients' age, psychopathology, pre-morbid functioning, educational achievement, and IQ [12,14,19-22,25-31]. In addition, there is preliminary evidence that pre-existing abnormal movements increase the risk of subsequent extrapyramidal side effects.

Based on the above observations, it has been proposed that idiopathic extrapyramidal disturbances may be intrinsic to the pathophysiology of schizophrenia and that antipsychotic medications may act by modifying the expression of disease-based motor dysfunctions [32-34], perhaps through specific alterations in subcortical dopamine neuronal activity due to basal ganglia dysregulation and dysfunction in cortical-basal, ganglia-cortical circuitry [18]. It is unclear how much 'neuroleptic-induced' motor disturbance should actually be attributed to the natural history of psychosis, aging, or other nondrug causes. Identifying this may be relevant to reducing negative attitude towards the side effects of antipsychotic medications and enhancing our understanding of the pathophysiology of psychosis. At present, nearly all patients with psychotic disorders in high income countries are exposed to antipsychotic medications [25]. In low income countries like Ethiopia, up to 90% of patients may not be exposed to antipsychotic medications [35,36], providing an opportunity to explore the prevalence and determinants of SMD among patients not exposed to antipsychotic treatments.

There are no published data from Ethiopia reporting on the prevalence of extrapyramidal symptoms or side effects; however, the incidence and prevalence of these symptoms are likely to be comparable to what has been reported in the broader literature. There are at least three outstanding issues that need addressing, which may be potentially relevant to our knowledge of extrapyramidal symptoms: 1) The contribution of antipsychotics above and beyond the natural risk of movement disorders is unclear; 2) the prevalence of movement disorders in treatment-naïve patients is not well documented in low income settings; 3) it has not yet been established whether there are unique risk factors more relevant for low income settings, such as malnutrition, infection and trauma.

Objectives

The primary objective of the study was to assess the extent of SMDs among antipsychotic-naïve patients with schizophrenia spectrum disorders (schizophrenia, schizoaffective and schizophreniform disorder) attending the outpatient department of a psychiatric hospital in Ethiopia. We additionally aimed to determine the association of socio-demographic and clinical factors with SMDs, including the occurrence of potential risk factors that may be contextually relevant, such as nutritional status.

Methods

Study design and setting

The study was a facility-based cross-sectional survey conducted at a psychiatric hospital located in the capital city, Addis Ababa. The hospital is the oldest and largest public psychiatric hospital in Ethiopia. Patients nationwide come to the hospital to receive treatment for severe mental disorders. The hospital service is organized under specialist programs or specialist case teams and each case team is expected to have expert knowledge about the disorders it treats. Patients with psychotic disorders are treated by psychoses case teams. These case teams manage all individuals with psychosis referred or seen by the hospital.

Participants

Participants of this study were patients aged between 18 and 60 years with a clinical diagnosis of schizophrenia, schizoaffective and schizophreniform disorders with no previous antipsychotic treatment or current treatment with medications that could induce extrapyramidal symptoms such as metoclopramide or other antiemetics. Other drugs that may induce extrapyramidal symptoms such as amoxapine and buspirone were not considered because they were not available in the Ethiopia market. Persons above the age of 60 were excluded because of the concern that older age may increase the risk of movement disorders as well as the potential for somatic and neurological diseases. Additionally, patients with catatonia were excluded as catatonia may be confused with other movement abnormalities.

Assessment

Successive patients were screened from the triage unit by a final-year psychiatry resident and consenting patients fulfilling inclusion criteria were offered a detailed assessment. Assessments consisted of an evaluation of basic socio-demographic characteristics, diagnosis, symptom severity, physical health status and occurrence of SMDs. Nutritional status, history of serious head injury and birth trauma were evaluated as contextually relevant risk factors. All assessments were made by the same final-year

psychiatry resident. A diagnosis of schizophrenia, schizoaffective disorder or schizophreniform disorder was confirmed using a diagnostic check list from the Text Revision of the Diagnostic and Statistical Manual of Mental Disorders [37]. The presence of extrapyramidal symptoms was evaluated using Simpson-Angus Scale (SAS) [38] and Abnormal Involuntary Movement Scale (AIMS) [39]. The SAS is a 10-item scale and has been shown to have good sensitivity [38]. The extrapyramidal symptoms assessed by the SAS are rigidity, hypersalivation, glabellar reflex, and tremor. On most items, severity is scored from 0 (none) to 4 (severe). A score of 1 usually indicates the presence of the extrapyramidal symptoms in a mild form. The commonly used threshold value to diagnose parkinsonism is 0.3 or more, which is used as a cut-off score to confirm occurrence of parkinsonism in the current report. The AIMS is commonly used to evaluate dyskinesic symptoms. The 12 questions of the AIMS cover anatomical locations of the dyskinesic movements (face, extremities, and trunk), global severity of the dyskinesia, global judgment of incapacitation and awareness of abnormality. The AIMS is administered through a clinician interview and physical examination. Scores on AIMS range from 0 (absent) to 5 (severe). Conventionally, dyskinesia is present, according to the AIMS scale, when there are mild movements in at least two areas or moderate movements in at least one area [40].

Patient symptoms profile and severity were assessed using the Scale for the Assessment of Positive Symptoms (SAPS) and Scale for the Assessment of Negative Symptoms (SANS) [41,42]. Both scales were designed to provide a detailed assessment of positive and negative symptoms of schizophrenia and can be used separately or in tandem. The domains assessed in the SAPS include hallucinations, delusions, bizarre behavior, and thought disorders, while the domains assessed in the SANS include affective flattening, poverty of speech, apathy, anhedonia, and inattentiveness. Nutritional status of participants was measured by the Quetelet's body mass index (BMI), a measure often used to define nutritional status in adults. It is estimated as weight in kilograms divided by height in metres squared. The BMI is easy to measure and closely relates to individuals' food consumption levels. The BMIs of the participants were categorized according to the recommendations of the WHO Expert Committee for chronic energy deficiency [43]. According to this WHO classification, participants with BMI values of less than 16 were considered to have severe energy deficiency; those with scores of 16–16.9 to have moderate deficiency; and those with scores of 17–18.4 to have mild deficiency. Those with BMI values of 18.5–24.9 were considered to be normal. Mid-arm circumference was also assessed as an additional measure of nutritional status. Finally, use of substances of

potential abuse (alcohol, khat, cigarettes and cannabis) was assessed. The assessment considered initially whether the participant had lifetime or current use of these substances and those reporting use in the past year were questioned in more detail about the frequency and quantity of the substances used.

Data management

Data were collected by a final-year psychiatry resident who received a half-day of training in addition to the extensive residency training he had received. Data were entered into the Statistical Package for Social Sciences, Version 15 for Windows for analysis. The focus of the analyses was on simple descriptive outputs (frequencies, percentages, means and standard deviations). Normality of continuous variables was checked visually using histogram and tested using Kolmogorov-smirnov test. For normally distributed data, equality of variance was checked using Leven's test before applying T-test. The few analyses for comparison and associations used the Independent Sample T-test and logistic regression analysis. Because of the small number of cases, only bivariate associations were explored.

Ethical considerations

The proposal was approved by the Scientific Committee of the Department of Psychiatry, School of Medicine, College of Health Sciences, Addis Ababa University and the Ethics Committee of Amanuel Specialised Mental Hospital. Interviews were conducted in private after obtaining informed consent. Analysis was conducted on anonymised data. Any new clinically relevant information generated in the research interview was passed on to the clinical team for the benefit of the participant.

Result

Socio-demographic and clinical characteristics of the participants

Sixty-four patients, mostly men ($n = 43/64$; 67.2%), with first contact psychosis were assessed over a 3 month period (April 01- June 30, 2013) prior to exposure to anti-psychotic or other psychotropic medications. The age of participants was between 18 and 60 years and the mean (SD) age was 29.2 (9.5). Most participants were single (44/64; 68.8%) and had a diagnosis of schizophrenia ($n = 47$; 73.4%). The duration of illness prior to presentation (duration of untreated psychosis) was between 1 month and 156 months, with a median (IQR) of 24 months (12, 57 months). Details are provided in Table 1.

Spontaneous movement disorders or dyskinesic movements

Based on standard cut-off points used to make a diagnosis of movement disorders, seven cases (10.9%) were

Table 1 Distribution of patients by socio-demographic characteristics

Characteristics		Number	Percent
Sex	Male	43	67.2
	Female	21	32.8
Age (in years)	18-24	28	43.8
	25-34	13	20.3
	35 and above	23	35.9
Educational status	Not able to read and write	10	15.6
	Basic writing and reading upto grade 6	21	32.8
	Above grade 6	33	51.5
Religion	Orthodox Christian	37	57.8
	Muslims	22	34.4
	Others	5	7.8
Marital status	Married	11	27.3
	Single	44	68.8
	Divorced or separated	9	14.1
Ethnicity	Amhara	20	31.3
	Oromo	16	25.0
	Guragae	16	25.0
	Others	12	18.8
Occupation	Unemployed	27	42.2
	Self-employed	14	21.9
	Farmer	14	21.9
	Student/house wife	9	14.1
Living arrangement	Living with parent or sibling	52	81.3
	Living alone or with own family	12	18.7
Diagnosis	Schizophrenia	47	73.4
	Schizophreniform disorder	5	7.8
	Schizoaffective disorder	12	18.8
Duration of illness	≤1 year	23	35.9
	1-2 years	11	17.2
	2-4 years	14	21.8
	4-13 years	16	25.0

found to have spontaneous movement disorders. Four participants had parkinsonism while one had dyskinesia and two had both parkinsonism and dyskinesia. Four of those with SMD were female and three male. Although a proportionately higher number of women had SMD, this difference was not statistically significant (Fisher's exact $p = 0.204$). Those with SMD were older, with a mean (SD) age of 38.3 (12.5) years compared with participants without SMD who had a mean age of 28.1 (8.6) years. This difference was statistically significant (mean difference = 10.2; $t = 2.8$; $p = 0.006$). These differences are reflected in the mean scores for both the SAS

and AIMS scales, which were relatively higher among older participants, women and those with more chronic illness (Table 2). The topography of dyskinetic movements was primarily orofacial, while the parkinsonian symptoms were found in the elbow, shoulder and arm regions, followed by wrist and leg regions.

Nutritional status and substance use behavior

Based on the WHO Expert Committee classification for Chronic Energy Deficiency in adults, close to half of the participants ($n = 29$; 45.3%) were in the underweight range with a BMI value of less than 18.5. Among those with low BMI values, 12 (41.4%) had BMI values of less than 16, which is classified as severely underweight. These details are presented in Figure 1. Thirty six (56.3%) participants had either current or past history of substance use (alcohol, khat, cigarettes or cannabis) and among those 52.4% of those used substances daily.

Exploratory analysis of associated factors

Because of the relatively small sample size and the small number of cases with SMD, results presented are based on a bivariate model. Increasing age was associated with SMD (OR = 1.10; 95% CI = 1.02, 1.20; $p = 0.017$). An inverse association was found between patients' nutritional status and the presence of spontaneous movement disorders; therefore increasing body weight (a proxy measure of dietary health), assessed with BMI, was inversely associated with SMD (OR = 0.6; 95% CI = 0.4, 0.8; $p = 0.01$). The mean score on the Simpson-Angus scale was also marginally associated with the SANS score ($\beta = .25$; 95% CI = 0.0, 0.06; $p = 0.049$).

Discussion

This is the first study on this subject in Ethiopia and is also one of few research reports on this subject from Africa. We evaluated a total of sixty-four patients with first contact psychosis who met the DSM-IV-TR criteria for schizophrenia, schizophreniform, and schizoaffective psychosis over the period of the study. The principal finding of this study is the confirmation that movement disorders occur in persons with schizophrenia spectrum disorders prior to exposure to antipsychotic medications, which are often presumed to be the cause of movement disorders in those with psychotic disorders. In this study at least one in ten treatment-naïve individuals (10.9%; 7/64) had spontaneous movement disorders. Most of those with SMD (6/7) had parkinsonism, while nearly half (3/7) had dyskinesia either alone or with parkinsonian movement disorders. The prevalence of spontaneous movement disorders in our study (10.9%) is consistent with the prevalence finding of SMDs in previous studies [15], particularly that of spontaneous parkinsonism [12,17,18], which lies between 4% and 21%. Rigidity and motor

Table 2 The mean, minimum and maximum Simpson-Angus Rating Scale and the Abnormal Involuntary Movement Scale scores of participants by demographic and clinical characteristics

Characteristics		Simpson-Angus (SAS) score			Abnormal Involuntary Movements (AIMS) score		
		Mean	Minimum	Maximum	Mean	Minimum	Maximum
Gender	Male	0.4	0.0	8.0	0.3	0.0	12.0
	Female	1.43	0.0	11.0	0.8	0.0	10.0
Age	<24	0.4	0.0	11.0	0.2	0.0	6.0
	25-34	0.3	0.0	4.0	0.9	0.0	12.0
	≥ 35	1.4	0.0	11.0	0.4	0.0	10.0
BMI	≥ 18.5	0.2	0.0	4.0	0.0	0.0	0.0
	<18.5	1.4	0.0	11.0	1.0	0.0	12.0
Duration of illness (in months)	1-12	0.4	0.0	6.0	0.0	0.0	0.0
	13-24	0.2	0.0	2.0	0.0	0.0	0.0
	25-48	0.3	0.0	4.0	0.0	0.0	0.0
	49-156	2.0	0.0	11.0	1.8	0.0	12.0
Diagnosis	Schizophrenia	0.9	0.0	11.0	0.6	0.0	12.0
	Schizophreniform disorder	1.2	0.0	6.0	0.0	0.0	0.0
	Schizoaffective disorder-bipolar type	0.0	0.0	0.0	0.0	0.0	0.0
	Schizoaffective disorder- depressive type	0.0	0.0	0.0	0.0	0.0	0.0

slowness were the common manifestations of parkinsonism in our study, consistent with previous studies [12,18-22] which reported akinetic symptoms, such as rigidity and bradykinesia, much more frequently than non-akinetic type movement disorders, such as tremor, glabellar tap, and salivation in antipsychotic-naïve patient with parkinsonian movement disorders. Consistent with previous studies [8,14], the commonly found dyskinetic symptoms in our study were orofacial movements. Thus, the findings in our study support the assertion that spontaneous movement disorders may be partly a disease-

induced state-dependent change, independent of the effect of antipsychotic medications.

One of the incidental findings of this study is the high rate of protein energy malnutrition: 45.5% (n = 29) of participants had BMI value under 18.5, which was significantly higher among those with spontaneous movement disorders, where 6/7 of those with SMD had low BMI. Alternatively, this could suggest that higher BMI was protective of SMD. The small number of cases prohibits in-depth speculation about the relevance of this finding. However, one postulation may be whether




Figure 1 Proportion of participants with evidence of malnutrition according to the BMI-based WHO classification.

this finding may have relevance to the aetiology of schizophrenia.

The aetiology of schizophrenia and other psychotic disorders remains largely undefined despite extensive research on possible biological mechanisms. Genetic constitution is important, but environmental factors like an unhealthy lifestyle with a poor diet may be involved. One area of focus regarding environmental risk factors has been around the role of polyunsaturated fatty acids. Low maternal fish and seafood consumption during pregnancy has been reported to increase the risk for a low IQ and suboptimal neuro-developmental outcomes in childhood [44], factors that in turn are associated with an increased risk for adult mental disorders like schizophrenia [45]. Research done by Horrobin [46] and Fenton [47] reported lower levels of polyunsaturated fatty acids (PUFA) in brain content, red blood cells and skin fibroblasts among patients with schizophrenia, compared with a healthy population. Hedelin et al. [48] also found that a high dietary intake of fish, omega-3 and omega-6 PUFA, as well as vitamin D, may have a protective effect against positive psychotic-like symptoms. However, we did not find studies reporting association between poor nutrition and SMD among persons with schizophrenia. This association may be attributed to specific alterations in subcortical dopamine neuronal activity [18]. Further studies are required to determine whether poor nutrition is a risk factor for SMDs.

The other factor significantly associated with SMD was increasing age. Although some previous studies did not find such an association [4], increasing age is often considered an important risk factor for SMD [9,49]. Additionally, there was weak association between negative symptoms and parkinsonian movement disorder. Exploratory analysis did not find an association between duration of illness, gender and SMD, which is consistent with previous reports [4].

Limitation of the study

The limitations of cross-sectional studies hold true also for this study. The number of recruited patients was relatively small and only patients who came to the hospital were recruited, limiting generalisability. Because of the small sample size, the analysis for association between potential risk factors and SMD should be considered exploratory.

Conclusions

This study supports the argument that involuntary movements (and parkinsonian symptoms) in schizophrenia may be, at least in part, intrinsic to the pathophysiology of the illness rather than a side effect of its treatment. Nutrition may simply be a contextual risk factor or represent a factor linked to the pathophysiology of the illness. Further

studies need to confirm these observations and systematically examine the relationship between SMDs and these risk factors with a larger sample size and more robust methodology. Additionally, determining the course of SMDs and their impact on treatment would also be of interest.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MA, TS and AF conceived of the research idea. BM participated in the initial case screening. MA and AF analysed the data and wrote the initial draft. All authors contributed to the writing of the final draft and read and approved the manuscript.

Acknowledgements

The study was conducted by a research grant awarded by Addis Ababa University to MA. The funding body had no role in developing the study design; in the collection, analysis, and interpretation of data; in the writing of the manuscript; and in the decision to submit the manuscript for publication. We are grateful to Ms. Meghan Davis for editing the English.

Author details

¹Department of Psychiatry, Hawassa University, College of Medicine and Health Sciences, Hawassa, Ethiopia. ²Department of Psychiatry, Addis Ababa University, College of Health Sciences, School of Medicine, Addis Ababa, Ethiopia. ³University of Toronto, Ontario Shores Center for Mental Health Sciences, Toronto, Canada. ⁴Department of Psychological Medicine, Centre for Affective Disorders, King's College London, Institute of Psychiatry, London, UK.

Received: 23 November 2013 Accepted: 27 September 2014

Published online: 09 October 2014

References

1. Gerlach J, Casey D: **Tardive dyskinesia.** *Acta Psychiatr Scand* 1988, **77**:369–378.
2. Kane J, Woerner M, Lieberman J: *Tardive dyskinesia: rate incidence and risk factors, in Dyskinesia Research and Treatment.* Berlin: Springer; 1985.
3. Kane J, Smith J: **Tardive dyskinesia: rate and risk factors.** *Arch Gen Psychiatry* 1982, **39**:473–481.
4. McCreddie R, Srinivasan T, Padmavati R, Thara R: **Extrapyramidal Symptoms in un medicated schizophrenia.** *J Psychiatr Research* 2005, **39**:261–266.
5. Pappa S, Dazzan P: **Spontaneous movement disorders in antipsychotic-naïve patients with first-episode psychoses: a systematic review.** *Psychol Med* 2009, **39**:1065–1076.
6. Whitty P, Owoeye O, Waddington J: **Neurological Signs and Involuntary Movements in Schizophrenia: Intrinsic To and Informative on Systems Pathobiology.** *Schizophr Bull* 2009, **35**:415–424.
7. Kraepelin E: *Dementia praecox and paraphrenia.* E & S Livingstone: Edinburgh; 1919.
8. Turner T: **Rich and mad in Victorian England.** *Psychol Med* 1989, **19**:29–44.
9. Liddle P: **Tardive dyskinesia in schizophrenia.** *Br J Psychiatry* 2013, **203**:6–7.
10. Sarro S, Pomarol-Clotet E, Canales-Rodriguez EJ, Salvador R, Gomar JJ, Ortiz-Gil J, Landin-Romero R, Vila-Rodriguez F, Blanch J, McKenna PJ: **Structural brain changes associated with tardive dyskinesia in schizophrenia.** *Br J Psychiatry* 2013, **203**:51–57.
11. Caligiuri M, Lohr J: **A disturbance in the control of muscle force in neuroleptic-naïve schizophrenic patients.** *Biol Psychiatry* 1994, **35**:104–111.
12. Chatterjee A, Chakos M, Koreen A: **Prevalence and clinical correlates of extrapyramidal signs and spontaneous dyskinesia in never-medicated schizophrenic patients.** *Am J Psychiatry* 1995, **152**:1724–1729.
13. McCreddie R, Thara R, Kamath S: **Abnormal movements in never-medicated Indian patients with schizophrenia.** *Br J Psychiatry* 1996, **168**:221–226.
14. Fenton W, Wyatt R, McGlashan T: **Risk factors for spontaneous dyskinesia in schizophrenia.** *Arch Gen Psychiatry* 1994, **51**:643–650.
15. Guy W: *ECDEU assessment manual for psychopharmacology.* Washington, DC: US Department of Health, Education and Welfare; 1976.

16. Gervin M, Browne S, Lane A: Spontaneous abnormal involuntary movements in first-episode schizophrenia and schizophreniform disorder: baseline rate in a group of patients from an Irish catchment area. *Am J Psychiatry* 1998, **155**:1202–1206.
17. Puri B, Barnes T, Chapman M, Hutton S, Joyce E: Spontaneous dyskinesia in first episode schizophrenia. *J Neurol Neurosurg Psychiatry* 1999, **66**:76–78.
18. Caligiuri M, Lohr J, Jeste D: Parkinsonism in neuroleptic-naive schizophrenic patients. *Am J Psychiatry* 1993, **150**:1343–1348.
19. Fenn D, Moussaoui D, Hoffman W: Movements in never-medicated schizophrenics. *Psychopharmacology* 1996, **123**:206–210.
20. Kopala L, Good K, Honer W: Extrapyramidal signs and clinical symptoms in first-episode schizophrenia: response to low-dose risperidone. *J Clin Psychopharmacology* 1997, **17**:308–313.
21. Peralta V, Cuesta M, Martinez-Larrea A, Serrano J: Differentiating primary from secondary negative symptoms in schizophrenia. *Am J Psychiatry* 2000, **157**:1461–1466.
22. Honer W, Kopala L, Rabinowitz J: Extrapyramidal symptoms and signs in first-episode, antipsychotic exposed and non-exposed patients with schizophrenia or related psychotic illness. *J Psychopharmacology* 2005, **19**:277–285.
23. Shawn L, Cassidy M, Helene A: Spontaneous Dyskinesia in Subjects with Schizophrenia Spectrum Personality. *Am J Psychiatry* 1998, **155**:70–7563.
24. Jeroen P, Diederik E, Jim V, Andr A, Renr S, Peter N: Dyskinesia and Parkinsonism in Antipsychotic-Native Patients With Schizophrenia, First-Degree Relatives and Healthy Controls. A Meta-analysis. *Schizophr Bull* 2010, **36**:723–731.
25. Fenton W: Prevalence of spontaneous dyskinesia in schizophrenia. *J Clin Psychiatry* 2000, **61**(Suppl. 4):10–14.
26. Cortese L, Caligiuri M, Malla A, Manchanda R, Takhar J, Haricharan R: Relationship of neuromotor disturbances to psychosis symptoms in first-episode neuroleptic-naive schizophrenia patients. *Schizophr Res* 2005, **75**:65–75.
27. Pareek P, Bhojraj T, Montrose D, Rajarethinam R, Sweeney J, Keshava M: Involuntary movements and their correlates in first-episode psychoses. *Acta Neuropsychiatr* 2010, **22**(5):262–263.
28. McCreadie R, Latha S, Thara R, Padmavati R, Ayankaran J: Poor memory, negative symptoms and abnormal movements in never-treated Indian patients with schizophrenia. *Br J Psychiatry* 1997, **171**:360–363.
29. Sandyk R, Kay S: The relationship of tardive dyskinesia to positive schizophrenia. *Int J Neurosci* 1991, **56**:107–139.
30. White T, Brown K, Woods J: Tardive dyskinesia and positive symptoms of schizophrenia. *Acta Psychiatrica Scand* 1991, **83**:377–379.
31. Waddington J, Youssef H, Dolphin C, Kinsella A: Cognitive dysfunction, negative symptoms, and tardive dyskinesia in schizophrenia. Their association in relation to topography of involuntary movements and criterion of their abnormality. *Arch Gen Psychiatry* 1987, **44**:907–912.
32. Yarden P, DiScipio W: Abnormal movements and prognosis in schizophrenia. *Am J Psychiatry* 1971, **128**:317–323.
33. Crown T, Cross A, Johnstone E, Owen F, Owens D, Waddington J: Abnormal involuntary movements in schizophrenia: are they related to the disease process or its treatment? Are they associated with changes in dopamine receptors. *J Clin Psychopharmacology* 1982, **2**:336–340.
34. Rogers D: The motor disorders of severe psychiatric illness: a conflict of paradigms. *Br J Psychiatry* 1985, **147**:221–232.
35. Fekadu A, Kebede D, Alem A, Fekadu D, Mogga S, Negash A, Medhin G, Beyero T, Shibre T: Clinical outcome in bipolar disorder in a community-based follow-up study in Butajira, Ethiopia. *Acta Psychiatr Scand* 2006, **114**(6):426–434.
36. Kebede D, Alem A, Shibre T, Negash A, Fekadu A, Fekadu D, Deyassa N, Jacobsson L, Kullgren G: Onset and clinical course of schizophrenia in Butajira-Ethiopia—a community-based study. *Soc Psychiatry Psychiatr Epidemiol* 2003, **38**(11):625–631.
37. American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, Text revision*. Washington, DC: American Psychiatric Association; 2000.
38. Simpson G, Angus J: A rating scale for extrapyramidal side effects. *Acta Psychiatrica Scand* 1970, **212**:S11–S19.
39. Guy W: *ECDEU Assessment Manual for Psychopharmacology. Revised DHEW Pub. (ADM)*. National Institute for Mental Health. Psychopharmacology Research: Rockville, MD; 1976.
40. Schooler N, Kamath J: Research diagnoses for tardive dyskinesia(letter). *Arch Gen Psychiatry* 1982, **39**:486–487.
41. Andreasen N: *Scale for the Assessment of Positive Symptoms (SAPS)*. University of Iowa: Iowa City; 1984.
42. Andreasen N: *Modified Scale for the Assessment of Negative Symptoms (SANS)*. Iowa City: University of Iowa; 1984.
43. WHO Expert Committee on Physical Status: The use and interpretation of anthropometry: report of a WHO expert committee. In *WHO Technical Report Series 854*. Geneva: World Health Organization; 1995.
44. Hibbeln J, Davis J, Steer C: Maternal seafood consumption in pregnancy and neurodevelopmental outcomes in childhood (ALSPAC study): an observational cohort study. *Lancet* 2007, **369**(9561):578–585.
45. Koenen K, Moffitt T, Roberts A: Childhood IQ and adult mental disorders: a test of the cognitive reserve hypothesis. *Am J Psychiatry* 2009, **166**(1):50–57.
46. Horrobin D: The membrane phospholipid hypothesis as a biochemical basis for the neurodevelopmental concept of schizophrenia. *Schizophr Res* 1998, **30**(3):193–208.
47. Fenton W, Hibbeln J, Knable M: Essential fatty acids, lipid membrane abnormalities, and the diagnosis and treatment of schizophrenia. *Biol Psychiatry* 2000, **47**(1):8–21.
48. Hedelin M, Luf M: Dietary intake of fish, omega-3, omega-6 polyunsaturated fatty acids and vitamin D and the prevalence of psychotic-like symptoms in a cohort of 33000 women from the general population. *BMC Psychiatry* 2010, **10**:38.
49. Merrill RM, Lyon JL, Maticco PM: Tardive and spontaneous dyskinesia incidence in the general population. *BMC Psychiatry* 2013, **13**:152.

doi:10.1186/s12888-014-0280-1

Cite this article as: Ayehu et al.: Movement disorders in neuroleptic-naive patients with schizophrenia spectrum disorders. *BMC Psychiatry* 2014 **14**:280.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

