

## Review

# Suicide by poisoning in Pakistan: review of regional trends, toxicity and management of commonly used agents in the past three decades

Maria Safdar\*, Khalid Imran Afzal\*, Zoe Smith, Filza Ali, Pervaiz Zarif and Zahid Farooq Baig

## Background

Suicide is one of the leading mental health crises and takes one life every 40 seconds. Four out of every five suicides occur in low- and middle-income countries. Despite religion being a protective factor against suicide, the estimated number of suicides is rapidly increasing in Pakistan.

## Aims

Our review focuses on the trends of suicide and means of self-poisoning in the past three decades, and the management of commonly used poisons.

## Method

We searched two electronic databases (PubMed and PakMediNet) for published English-language studies describing agents used for suicide in different regions of Pakistan. A total of 46 out of 85 papers ( $N = 54\,747$  cases) met our inclusion criteria.

## Results

Suicidal behaviour was more common among individuals younger than 30 years. Females comprised 60% of those who attempted suicide in our study sample, although the ratio of completed suicides favoured males. There were regional trends in the choice of agent for overdose. Organophosphate poisoning was reported across the nation, with a predominance of cases

from the agricultural belt of South Punjab and interior Sindh. Aluminium phosphide ('wheat pills') was a preferred agent in North Punjab, whereas parphenylenediamine ('*kala pathar*') was implicated in deaths by suicide from South Punjab. Urban areas had other means for suicide, including household chemicals, benzodiazepines, kerosene oil and rat poison.

## Conclusions

Urgent steps are needed, including psychoeducational campaigns on mental health and suicide, staff training, medical resources for prompt treatment of self-poisoning and updated governmental policy to regulate pesticide sales.

## Keywords

Low- and middle-income countries; suicide; mortality; epidemiology; self-harm.

## Copyright and usage

© The Author(s), 2021. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Suicide is the second leading cause of death in 15- to 29-year-olds globally, and 10- to 34-year-olds in the USA.<sup>1,2</sup> The World Health Organization (WHO) estimates that 800 000 people die by suicide every year, which translates into one death every 40 seconds, and 79% of global suicides occur in low- and middle-income countries (LMICs).<sup>3</sup> The World Bank Atlas defines low-income countries as having a gross national income (GNI) per capita of \$1025 or less in 2018, and lower-middle-income countries as having a GNI per capita of \$1026–\$3995.<sup>4</sup> Although pesticide ingestion, hanging and firearms are among the most common methods of suicide worldwide,<sup>1</sup> trends vary between nations regarding the age groups, access and availability of the means.<sup>5–8</sup> Mirroring global studies, the three most common methods for suicides in Pakistan are poisoning, firearms and hanging.<sup>9,10</sup>

## Pakistan

Pakistan is the fifth most populous country in the world.<sup>11</sup> It is predominantly an agricultural country and, according to the 2017 National Census, around 64% of its population of 207 million is considered rural.<sup>12,13</sup> The population ratio favours males (51.23%), with a male:female ratio of 1.05.<sup>12</sup> About 50% of the population is under 20 years of age, and 35% is under 15 years of age. The literacy rate of Pakistan, as measured by the ability of people aged ≥15 years to read and write, is around 59%, which is lower than

the average literacy rate in other South Asian countries (71.70%) and for LMICs overall (75%).<sup>14</sup> Men have a literacy rate of 71%, whereas women have a literacy rate of <47%.<sup>12,14,15</sup> The literacy rate in large urban centres such as Karachi and Lahore, the two largest cities in the country, is close to 75%, whereas the average literacy rate in rural areas is <50%.<sup>16</sup> Along with other factors, terrorism has negatively affected sustained economic growth in Pakistan over the past two decades, leading to a high unemployment rate. The health indicators of the country continue to remain poor.<sup>17–20</sup>

## Geography and demography

Geographically, the country is composed of four provinces – Punjab, Sindh, Balochistan, and Khyber Pakhtunkhwa (KPK) – and Gilgit–Baltistan, a newly created province in the north (Fig. 1).<sup>21,22</sup> The Punjab and Sindh are fertile plains with agriculture-based economies. Balochistan and KPK are bound by strong tribal traditions. Gun ownership is a shared pride between the two provinces. Balochistan is rugged, rich in minerals and mostly barren.<sup>23,24</sup> In the north of Pakistan, Gilgit–Baltistan is home to three large mountainous ranges: the Himalayas, the Karakoram and the Hindu Kush. The scenic region has beautiful valleys and river-irrigated lands.<sup>21,25</sup> Shah and Amjad<sup>26</sup> measured the cultural diversity of different regions of Pakistan. They found a high masculinity index score in all provinces, indicating a difference in social genders, with clear-cut roles. Uncertainty avoidance index scores were low in all provinces, mainly because a firm belief in Allah

\* Joint first authors.



**Fig. 1** Geographical map of Pakistan. AJK, Azad Jammu and Kashmir; KPK, Khyber Pakhtunkhwa.

(God Almighty) led to most people not feeling threats or uncertainty about the future. Individualism index scores were low in all provinces, especially in KPK and Balochistan, signifying collectivism as a national culture. The people of Pakistan possessed a strong urge toward group cohesiveness and the expectation of loyalty.<sup>26</sup>

Approximately 96% of the population of Pakistan is Muslim.<sup>27,28</sup> Like other major religions, Islam condemns suicide, declaring it an unforgivable sin.<sup>29–31</sup> This could be a significant deterrent to suicide, evidenced by the traditionally low rates reported in Muslim countries compared with non-Muslim countries.<sup>32</sup> Based on religious tenets, both suicide and self-harm are illegal and punishable by imprisonment and fines under Pakistani law, adding another deterrent to suicide.<sup>33–35</sup> Studies from other LMICs and higher-income (GNI per capita of  $\geq$ \$12 376) Muslim-majority countries also show a lower suicide rate than non-Muslim-majority countries.<sup>36,37</sup> Arya et al describe the geographical heterogeneity of suicide rates in the neighbouring LMIC of India, focusing on religion, caste, tribe, etc. The authors found that the rate of suicide was lowest for Sikhs and Muslims, and highest for Hindus and Christians.<sup>38</sup>

### Suicide statistics

Pakistan has no vital registrations and lacks accurate figures for death by suicide.<sup>39</sup> As compared with the 2017 global suicide death rate per 100 000 people for both genders of 9.98,<sup>40</sup> the estimated age-standardised suicide rate in Pakistan is 4.4 per 100 000 people.<sup>41</sup> The suicide death rates in neighbouring India,

Bangladesh and Sri Lanka are 13.33, 5.73 and 7.55 per 100 000 people, respectively. Despite the low estimated rate, recent data suggest that suicide is becoming a significant public health problem in Pakistan.<sup>42–45</sup> The WHO published a report showing an increase in the reported suicide rate of 2.6% from the year 2000.<sup>1</sup> Because of the social, legal and religious factors noted above, suicide and self-harm are not reported or are underreported. Recent reports have shown rapidly increasing rates for suicide and self-harm across the country.<sup>34,42,46</sup> Shekhani et al noted a stigmatisation of suicidal behaviour contributing toward a lack of research on the subject.<sup>10</sup> We did not find literature on suicide or self-harm that compared different regions of Pakistan or differentiated between urban and rural populations.

To address the gap in current knowledge, this is the first study to map the regional trends of suicide by poisoning in Pakistan, and detail urban versus rural differences. We also aim to provide a detailed account of the pathophysiology and management strategies of agents used in suicide attempts, to give readers a comprehensive review on the subject. Our analysis will provide future research directions and inform policy for suicide prevention in Pakistan, focusing on regional and urban versus rural differences in suicide attempts.

### Method

We searched two electronic databases (PubMed and PakMediNet) for studies describing agents used for suicide in different regions

of Pakistan, using the following terms: suicide, death, poisoning, drugs, overdose and Pakistan. We considered studies published in the English language within the past 30 years, and conducted the search from October to December 2019. Our null hypothesis was that there is no regional or urban versus rural difference in suicide by poisoning in Pakistan. We included primary research, case series and case reports, focusing on different agents used by adults of both genders, aged  $\geq 18$  years, who attempted suicide. Studies involving ex-pat Pakistanis and those using means of suicide other than overdose were excluded. We did not include single case reports as most focused on uncommon means of death or unusual clinical presentations that were not the focus of our study. The Postgraduate Medical Institute at Lahore, Pakistan, approved all of the data collection for this research project according to its policies regarding studies involving human patients.

After retrieving 85 articles from both databases, two independent reviewers screened the titles and abstracts for relevance. Sixty-two papers met the inclusion criteria; however, sixteen were case reports and were not included. Most studies were descriptive, with only three that used a case-control design. The majority of the studies were from urban areas (74%) and addressed determinants rather than risk factors. The WHO defines determinants as a range of behavioural, biological and socioeconomic factors that influence the health of populations.<sup>47</sup> The risk factors are characteristics or attributes within an individual that influence the likelihood of disease.<sup>10</sup> Most studies reported gender (95.3%) and age (93.0%) differences. We identified eight distinctive regions, including North and South Punjab, North and South KPK, interior Sindh (all cities except Karachi), urban Sindh (represented by studies from the largest city of Karachi), Balochistan and Gilgit-Baltistan (Table 1). The four predominant agents used in the attempted and completed suicides were organophosphates, aluminium phosphide (or 'wheat pills'), parphenylenediamine (or 'kala pathar') and others (including over-the-counter medications and household chemicals). We describe the clinical presentation, pathophysiological mechanism, morbidity, mortality and available treatments in the Discussion section. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

## Results

Table 1 shows the distribution of studies according to the regions, with details on study design, cohort size, gender, mean age, geographical region, city, suicide attempts or completion methods, and mortality. The exact doses of agents used in the suicide attempts were inconsistently reported and were not statistically meaningful for our study. The majority of the studies were from urban areas (74%) and addressed determinants rather than risk factors. Most studies reported the gender (95.3%) and age (93.0%) of the individuals. We identified eight distinctive regions, including North and South Punjab, North and South KPK, interior Sindh (all cities except Karachi) and urban Sindh (represented by studies from the largest city Karachi), Balochistan and Gilgit-Baltistan (Table 1). The four predominant agents used in the attempted and completed suicides were organophosphates, aluminium phosphide, parphenylenediamine and others (including over-the-counter medications and household chemicals). Of the 47 studies, 53.2% examined organophosphates ( $n = 25$ ), 36.2% examined over-the-counter agents and household chemicals ( $n = 17$ ), 23.4% examined *kala pathar* ( $n = 11$ ), 19.1% examined wheat pills ( $n = 9$ ) and 4.3% examined 'intoxication' without indicating the agent used ( $n = 2$ ). Note that some studies examined multiple agents, so the total exceeds the

number of studies included. With the exception of two studies, all papers were published in the past two decades.

The total number of cases across the 53 studies was 54 747 (see Table 2). A total of 60% of overall study participants were female and 40% were male. Suicidal behaviour was more common among individuals aged  $< 30$  years, with a mean age of 27.9 years. See Table 3 for more comprehensive demographic information. Urban Sindh had the most publications (14 studies;  $n = 25\ 458$ ), followed by North Punjab (12 studies;  $n = 2319$ ), South Punjab (7 studies;  $n = 1901$ ), interior Sindh (6 studies;  $n = 1027$ ), North KPK (4 studies;  $n = 438$ ), South KPK (2 studies;  $n = 541$ ) and one study each from Balochistan ( $n = 46$ ) and Gilgit-Baltistan ( $n = 46$ ) (see Table 4 for demographic information by region). The overall mortality rate, regardless of the method, was 24.5%. Organophosphates were the most widely reported agent (25 studies;  $n = 35\ 479$ ), with an average mortality rate of 13.9% (11 studies;  $n = 2364$ ). The highest average mortality rate was for wheat pills, at 44.7% (9 studies;  $n = 2070$ ). The lowest average mortality rate was for over-the-counter agents and household chemicals (17 studies;  $n = 20\ 911$ ), at 12.1%. For *kala pathar*, the average mortality rate was 38.6% (11 studies;  $n = 2364$ ). See Table 5 for more demographic information by different agents.

Studies from Karachi (i.e. urban Sindh) included 25 458 individuals, of whom 55.1% were women aged 20–43 years (mean age 27.5 years). The average overall mortality rate for this region was 7.46 and ranged from 0 to 42%. Most studies from Karachi (73.3%) found organophosphates as the agent chosen for death by suicide, with an average mortality rate of 9.33 (range 0–20%). Other agents were also examined, including benzodiazepines, off-label agents, pesticides, corrosives, kerosene oil, rat poison, non-steroidal anti-inflammatory drugs (NSAIDs)/analgesics, and antidepressants. Two studies found that 55–91% of 771 people chose benzodiazepines as the agent of choice for attempting suicide. However, benzodiazepine overdose was associated with a 0% mortality rate in these studies. One study found that 18% of 2546 individuals chose off-label agents, whereas another study found that 15% of 705 individuals chose pesticides. Two studies of 3708 individuals found that 13.5% used corrosives. Kerosene oil was examined in two studies, with 2–14% of 15 259 individuals using it to commit suicide. Finally, rat poison (11% of 2546 individuals), NSAIDs/analgesics (11% of 324 individuals) and antidepressants (10% of 324 individuals) were all examined in one paper.

Interior Sindh included six studies from three cities: Hyderabad, Jamshoro and Nawabshah. The latter two cities are rural. There were 1027 individuals aged 16–43 years (mean age 32.5 years), of whom 53.1% were female. The most commonly studied agent was organophosphates (66.6% of studies, 987 individuals), whereas the other two studies examine *kala pathar* (40 individuals). Mortality rates for organophosphates ranged from 17 to 27% (mean 20.5%), whereas aluminium phosphide (two studies;  $n = 40$ ) was higher at 38–42% (mean 40%). Overall mortality rates for this region averaged at 27%.

Within North Punjab, a total of 2319 cases were noted in 12 studies, with a male:female gender ratio of 50.5%:49.5% favouring males. The age range was 20–40 years (mean age 26.6 years) across six cities (Kharian, Lahore, Mianwali, Rawalpindi, Sahiwal and Wah Cantt). Of these cities, Mianwali and Sahiwal are considered rural, and the other four are urban. The overall mortality rates range from 2.5 to 87%, with a mean percentage of 43.8%. In North Punjab, almost half of individuals who ingested wheat pills died by suicide, indicating the high lethality of the agent. The overall mortality rate for wheat pills ranged from 33 to 87%, with an average of 52%. Other agents examined in the region included organophosphates (four studies), corrosives (two studies), benzodiazepines (one study), generic agents (one study), medicine (one study),

**Table 1** Studies on commonly used agents for poisoning in Pakistan, by region

Region	City	Reference	Total cases	Age (mean + s.d./range/ median)^	Gender, male: female	Poison used	Mortality	Study type
Organophosphate (25 articles published)								
North Punjab (5 articles)	Wah Cantt <sup>a</sup>	Bhatti N, et al. <sup>48</sup>	126	25.9 ± 9.75	48:52	Drugs 58%, organophosphates 18%, wheat pills 10%, corrosives 6%	–	Descriptive
	Rawalpindi	Maqbool F, et al. <sup>49</sup>	62	23.3 ± 6.1	53:47	Organophosphates	–	Descriptive
	Rawalpindi <sup>a</sup>	Khurram M, et al. <sup>50</sup>	85	24.35 ± 7.69	41:59	Medicine 53%, organophosphates 21%, corrosives 10%	2.5%	Descriptive
	Mianwali <sup>a</sup>	Tahir MN, et al. <sup>51</sup>	108	11–40	78:22	Toxic substance 36%, pesticides 31%, drug overdose 11%	–	Descriptive
	Lahore <sup>a</sup>	Naheed T, et al. <sup>52</sup>	114	25.89 ± 11.48	53:47	Urban: household toxins and drugs; rural: wheat pills and organophosphates	–	Descriptive
North KPK (3 articles)	Peshawar <sup>a</sup>	Ali Z, et al. <sup>53</sup>	128	25.79 ± 11.23	40:60	Organophosphates 31%, benzodiazepine 13%, wheat pills 11%	Total 12–44%	Descriptive
	Peshawar <sup>a</sup>	Rahim F, et al. <sup>54</sup>	92	26.8 ± 13.9	40:60	Medicine 53%, organophosphates 36%, wheat pills 11%	25%	Descriptive
	Peshawar	Bilal M, et al. <sup>55</sup>	50	30.88 ± 15.72	54:46	Organophosphates	10%	Descriptive
Interior Sindh (4 articles)	Hyderabad	Shaikh MA. <sup>56</sup>	100	37.5 ± 9.5 (43 median)	78:22	Organophosphates	18%	Descriptive
	Nawabshah	Imran S, et al. <sup>57</sup>	387	26.14 ± 10.09	63:37	Organophosphates	27%	Descriptive
	Nawabshah	Faiz MS, et al. <sup>58</sup>	300	32 ± 5.2	17:83	Organophosphates	17%	Descriptive
Urban Sindh (12 articles)	Jamshoro	Shaikh MA, et al. <sup>59</sup>	200	38.4 ± 3.5 (45 median)	81:19	Organophosphates	20%	Descriptive
	Karachi <sup>a</sup>	Amir A, et al. <sup>60</sup>	2546	26.57 ± 11.82	51:49	Organophosphates 46%, off-label products 18%, rat poison 11%	4%	Descriptive
	Karachi <sup>a</sup>	Khan NU, et al. <sup>61</sup>	705	4–32 (21 median)	–	Drugs 24%, pesticides 15%, household toxins 9%	–	Descriptive
	Karachi	Ahmed A, et al. <sup>62</sup>	248	27.28 ± 11.5	27:73	Organophosphates	14%	Descriptive
	Karachi <sup>a</sup>	Imtiaz F, et al. <sup>63</sup>	11 925	14–22 (20 median)	34:66	Organophosphates 62%, kerosine oil 14%, drugs 23%	18%	Descriptive
	Karachi <sup>a</sup>	Asghar SP, et al. <sup>64</sup>	40	12–56 (30 median)	20:80	Organophosphates	Nil	Descriptive
	Karachi <sup>a</sup>	Bashir F. <sup>65</sup>	374	25 ± 10.1	62:38	Organophosphates 47%, corrosives 14%, drugs 23%	–	Descriptive
	Karachi	Ali P, et al. <sup>66</sup>	100	28.6 ± 9.8	68:32	Organophosphates	20%	Case Series
	Karachi	Ather N, et al. <sup>67</sup>	2708	–	51:49	Organophosphates	5%	Descriptive
	Karachi <sup>a</sup>	Turabi A, et al. <sup>68</sup>	3334	8–50 (30 median)	50:50	Organophosphates 60%, corrosives 13%, rodenticides 7%, kerosine oil 2%	–	Descriptive
	Karachi	Hussain AM, et al. <sup>69</sup>	52	–	–	Organophosphates	8%	Descriptive
	Karachi <sup>a</sup>	Jamil H. <sup>70</sup>	1900	11–30	47:53	Organophosphates 40%, tranquilisers 21%, sedatives/hypnotics 10%	5.6%	Descriptive
Karachi	Jamil H. <sup>71</sup>	755	11–30	40:60	Organophosphates	–	Descriptive	
Balochistan (1 article)	Quetta	Khan NK, et al. <sup>72</sup>	46	15–35	Nil:100	Organophosphates	–	Descriptive
Wheat pills (9 articles published)								
North Punjab (7 articles)	Rawalpindi	Hassan A, et al. <sup>73</sup>	77	16–60	47:53	Wheat pills	33%	Descriptive
	Kharian	Iftikhar R, et al. <sup>74</sup>	52	25.10 ± 5.35	48:52	Wheat pills	87%	Descriptive
	Lahore	Ghazi MA. <sup>75</sup>	–	25 ± 5	–	Wheat pills	70%	Review
	Lahore <sup>a</sup>	Naheed T, et al. <sup>52</sup>	114	25.89 ± 11.48	53:47	Urban: household toxins and drugs; rural: wheat pills and organophosphates	–	Descriptive
	Lahore <sup>a</sup>	Shoaib S, et al. <sup>76</sup>	107	11–60	55:45	Wheat pills 33%, bleach 26%, benzodiazepines 19%	26%	Descriptive
Lahore <sup>a</sup>	Asif A, et al. <sup>77</sup>	1390	11–90	69:31	Poison unknown 83%, rat poison pills 4%, wheat pills 3%	–	Descriptive	
Sahiwal	Qureshi MA, et al. <sup>78</sup>	110	12–40	41:59	Wheat pills	44%	Descriptive	
North KPK (2 articles)	Peshawar <sup>a</sup>	Ali Z, et al. <sup>53</sup>	128	25.79 ± 11.23	40:60	Organophosphates 31%, benzodiazepines 13%, wheat pills 11%	Total 12–44%	Descriptive
	Peshawar <sup>a</sup>	Rahim F, et al. <sup>54</sup>	92	26.8 ± 13.9	40:60	Medicine 53%, organophosphates 36%, wheat pills 11%	25%	Descriptive
<i>Kala pathar</i> (11 articles published)								

South Punjab (7 articles)	Sahiwal	Akram A, et al. <sup>79</sup>	88	>14 (92%)	22:78	<i>Kala pathar</i>	–	Descriptive	
	Multan	Tanweer S, et al. <sup>80</sup>	122	23.21 ± 8.2	20:80	<i>Kala pathar</i>	28%	Descriptive	
	Multan	Haider SH, et al. <sup>81</sup>	32	21.06 ± 3.25	34:66	<i>Kala pathar</i>	28%	Descriptive	
	Bahawalpur	Khan MA, et al. <sup>82</sup>	1258	5–63 (21 median)	65:35	<i>Kala pathar</i>	24%	Descriptive	
	Bahawalpur	Ishtiaq R, et al. <sup>83</sup>	109	22 ± 3.4	29:71	<i>Kala pathar</i>	39%	Descriptive	
	Bahawalpur	Qasim AP, et al. <sup>84</sup>	109	11–60	11:89	<i>Kala pathar</i>	21%	Descriptive	
	Rahim Yar Khan	Akbar K, et al. <sup>85</sup>	65	24.35 ± 9.8	28:72	<i>Kala pathar</i>	–	Descriptive	
South KPK (2 articles)	Dera Ismail Khan	Ansari RZ, et al. <sup>86</sup>	503	12–39	21:79	<i>Kala pathar</i>	80%	Descriptive	
	Dera Ismail Khan	Khan H, et al. <sup>87</sup>	38	22.08 ± 6.42	5:95	<i>Kala pathar</i>	47%	Descriptive	
Interior Sindh (2 articles)	Hyderabad	Kazi MA et al. <sup>88</sup>	24	15–35	29:71	<i>Kala pathar</i>	42%	Descriptive	
	Nawabshah	Khuhro BA, et al. <sup>89</sup>	16	25.87 ± 5.59	13:87	<i>Kala pathar</i>	38%	Descriptive	
Over-the-counter drugs and household chemicals (17 articles published)									
North Punjab (6 articles)	Wah Cantt <sup>a</sup>	Bhatti N, et al. <sup>48</sup>	126	25.9 ± 9.75	48:52	Drugs 58%, organophosphates 18%, wheat pills 10%, corrosives 6%	–	Descriptive	
	Rawalpindi <sup>a</sup>	Khurram M, et al. <sup>50</sup>	85	24.35 ± 7.69	41:59	Medicine 53%, organophosphates 21%, corrosives 10%	2.5%	Descriptive	
	Mianwali <sup>a</sup>	Tahir MN, et al. <sup>51</sup>	108	11–40	78:22	Toxic substances 36%, pesticides 31%, drug overdose 11%	–	Descriptive	
	Lahore <sup>a</sup>	Naheed T, et al. <sup>52</sup>	114	25.89 ± 11.48	53:47	Urban: household toxins and drugs; rural: wheat pills and organophosphates	–	Descriptive	
Lahore <sup>a</sup>		Shoaib S, et al. <sup>76</sup>	107	11–60	55:45	Wheat pills 33%, bleach 26%, benzodiazepines 19%	26%	Descriptive	
		Asif A, et al. <sup>77</sup>	1390	11–90	69:31	Poison unknown 83%, rat poison pills 4%, wheat pills 3%	–	Descriptive	
		Hashmi MU, et al. <sup>90</sup>	206	23.44 ± 7.19	34:66	Corrosives	–	Descriptive	
South Punjab (1 article)	Multan								
North KPK (2 articles)	Peshawar <sup>a</sup>	Ali Z, et al. <sup>53</sup>	128	25.79 ± 11.23	40:60	Organophosphates 31%, benzodiazepines 13%, wheat pills 11%	Total 12–44%	Descriptive	
	Peshawar <sup>a</sup>	Rahim F, et al. <sup>54</sup>	92	26.8 ± 13.9	40:60	Medicine 53%, organophosphates 36%, wheat pills 11%	25%	Descriptive	
Urban Sindh (8 articles)	Karachi <sup>a</sup>	Amir A, et al. <sup>60</sup>	2546	4–32 (21 median)	51:49	Organophosphates 46%, off-label products 18%, rat poison 11%	4%	Descriptive	
	Karachi <sup>a</sup>	Khan NU, et al. <sup>61</sup>	705	14–22	–	Drugs 24%, pesticides 15%, household toxins 9%	–	Descriptive	
	Karachi <sup>a</sup>	Imtiaz F, et al. <sup>63</sup>	11 925	25 + 10.1	34:66	Organophosphates 62%, kerosine oil 14%, drugs 23%	18%	Descriptive	
	Karachi <sup>a</sup>	Bashir F, et al. <sup>65</sup>	374	25 + 10.1	62:38	Organophosphates 47%, corrosives 14%, drugs 23%	–	Descriptive	
	Karachi <sup>a</sup>	Patel MJ, et al. <sup>91</sup>	324	36.2 ± 17.0	33:67	Benzodiazepines 55%, NSAIDs/analgesics 11%, antidepressants 10%	Nil	Descriptive	
	Karachi <sup>a</sup>	Turabi A, et al. <sup>68</sup>	3334	8–50	50:50	Organophosphates 60%, corrosives 13%, rodenticides 7%, kerosine oil 2%	–	Descriptive	
Karachi <sup>a</sup>		Khan MM, et al. <sup>92</sup>	447	27.50 + 10.62	40:60	Drugs 73% (benzodiazepines 91%)	Nil	Descriptive	
		Jamil H. <sup>70</sup>	1900	11–30	47:53	Organophosphates 40%, tranquilisers 21%, sedatives/hypnotics 10%	5.6%	Descriptive	
Miscellaneous (2 articles published)									
North KPK (1 article)	Chitral	Ahmed Z, et al. <sup>93</sup>	168	10–50	38:62	Drowning 52%, hanging 26%, gunshot 17%, intoxication 5%	–	Descriptive	
Gilgit–Baltistan (1 article)	Ghizer	Khan MM, et al. <sup>94</sup>	49	16–>70 Majority 16–35	Nil:100	Jumping into a river/lake 40%, intoxication 30%, strangulation 11%, firearm 5%	–	Descriptive	

KPK, Khyber Pakhtunkhwa; NSAID, non-steroidal anti-inflammatory drug.  
a. Studies cited more than once because of multiple toxins/drugs involved.



**Table 2** Overall demographic information for included studies

Demographics	Mean/percentage
Participants	N = 54 747
Studies	N = 53
Female:male	60%:40%
Age	27.9 years
Urban:rural area	74%:26%
Mortality rate	24.5%
<b>Regions</b>	<b>Number of participants</b>
Balochistan	46
Gilgit–Baltistan	49
Interior Sindh	1027
North Khyber Pakhtunkhwa	438
North Punjab	2319
South Khyber Pakhtunkhwa	541
South Punjab	1901
Urban Sindh	25 458
<b>Regions</b>	<b>Number of studies</b>
Balochistan	1
Gilgit–Baltistan	1
Interior Sindh	6
North Khyber Pakhtunkhwa	4
North Punjab	12
South Khyber Pakhtunkhwa	2
South Punjab	7
Urban Sindh	14

**Table 3** Demographic information of included studies by region

Region/demographics	Mean/percentage
<b>Balochistan</b>	
Participants	n = 46
Female:male	100%:0%
Age	20.5 years
Mortality rate	Not reported
<b>Gilgit–Baltistan</b>	
Participants	n = 49
Female:male	100%:0%
Age	25.5 years
Mortality rate	Not reported
<b>Interior Sindh</b>	
Participants	n = 1027
Female:male	53.1%:46.9%
Age	32.5 years
Mortality rate	27%
<b>North Khyber Pakhtunkhwa</b>	
Participants	n = 438
Female:male	57%:43%
Age	28.5 years
Mortality rate	21%
<b>North Punjab</b>	
Participants	n = 2319
Female:male	49.5%:50.5%
Age	26.6
Mortality rate	43.8%
<b>South Khyber Pakhtunkhwa</b>	
Participants	n = 541
Female:male	87%:13%
Age	23.8 years
Mortality rate	63.5%
<b>South Punjab</b>	
Participants	n = 1901
Female:male	68.4%:31.6%
Age	23.4 years
Mortality rate	28%
<b>Urban Sindh</b>	
Participants	n = 25 458
Female:male	55.1%:46.9%
Age	27.5 years
Mortality rate	7.46%

**Table 4** Demographic information of included studies by agent

Agent/demographics	Mean/percentage
<b>Organophosphates</b>	
Participants	n = 35 479
Studies	n = 25
Urban:rural	16%:84%
Female:male	52.3%:47.7%
Age	28.9 years
Mortality rate	13.9%
<b>Over-the-counter agents and household chemicals</b>	
Participants	n = 20 911
Studies	n = 17
Urban:rural	6%:94%
Female:male	51.6%:48.4%
Age	26.0 years
Mortality rate	12.1%
<b>Kala pathar</b>	
Participants	n = 2364
Studies	n = 11
Urban:rural	36%:64%
Female:male	74.8%:25.2%
Age	24.5 years
Mortality rate	38.6%
<b>Wheat pills</b>	
Participants	n = 2070
Studies	n = 9
Urban:rural	11%:89%
Female:male	50.9%:49.1%
Age	27.7 years
Mortality rate	44.7%
<b>Miscellaneous</b>	
Participants	n = 217
Studies	n = 2
Urban:rural	0%:100%
Female:male	81%:19%
Age	27.8 years
Mortality rate	Not reported

‘toxic substance’ (one study), pesticides (one study), household toxins (one study), bleach (one study), *kala pathar* (one study) and rat poison (one study). Mortality rates were not reported for these agents.

There were 1901 cases in 7 studies from three cities in South Punjab (Bahawalpur, Multan and Rahim Yar Khan). This region consisted of all urban cities, although the healthcare facilities’ catchment area extends into vast agricultural lands. Women comprised 68.4% of the samples, with an age range of 21–30 years (mean age 23.4 years). All seven studies examined parphenylenediamine (*kala pathar*) poisoning, with a mortality rate of 28% (ranging from 21 to 39%). Only one study examined corrosives as the substance of choice for overdose, but this study did not report mortality.

North KPK included four studies with 438 cases from two cities: Peshawar (urban) and Chitral (rural). Women comprised 57% of the reported cases, with an age range of 26–31 years (mean 28.5 years). Mortality rates ranged from 10 to 44%, with an average overall mortality rate of 21%. No clear choice of agent for overdose emerged; however, similar to urban Sindh, organophosphates were included in three of the four studies, with a prevalence rate of 31–36%. Aluminium phosphide and benzodiazepines were the agents of choice 11% and 13% of the time, respectively. Interestingly, one study included methods outside of poisoning, finding that only 5% of individuals preferred an overdose by agents compared with other methods (drowning 52%, hanging 26%, firearms 17%).

For South KPK, there were two studies, both from Dera Ismail Khan, which is a rural area. There were 541 participants across the two studies, of whom 87% were female, and the average age was 23.8 years (range 12–39 years). Both studies only examined *kala pathar*,

**Table 5** Overview of commonly used poisons in Pakistan

Poison	Symptoms	Diagnosis	Management	Lethal dose	Mortality	Regional prevalence	Cost
Organophosphates	Bronchorrhea, bronchoconstriction, excessive sweating, constricted pupils, abdominal cramps, involuntary defaecation and urination, tachycardia, QT prolongation, headaches, dizziness, drowsiness, confusion, anxiety, slurred speech, ataxia, psychosis, convulsions, coma, hypotension and respiratory depression	Clinical	Supportive care; decontaminate the patient and prevent further absorption via the gut, eyes, skin or lungs; administer atropine followed by enzyme reactivation by pralidoxime	Depends on many factors	10–27%	All over Pakistan, but more prevalent in North Punjab and Sindh	1200–1600 PKR per litre
Aluminium phosphide	Epigastric pain, vomiting, diarrhoea, dizziness and dyspnoea, multiorgan failure involving the heart, kidneys, lungs and liver	Silver nitrate test	Supportive care as no antidote is available; gastric lavage with potassium permanganate and mineral or coconut oil; renal replacement therapy in the early stage may be helpful	150–500 mg	33–87%	North Punjab	700–1000 PKR per 500 mg
Paraphenylenediamine	Angioneurotic oedema, rhabdomyolysis causing myoglobinuria, cola-coloured urine, oliguria and acute tubular necrosis leading to acute renal failure	Clinical	Supportive care as no antidote is available; early tracheostomy; intravenous fluids to prevent renal failure; renal replacement therapy in cases where ATN develops	7–10 g	21–47%	South Punjab and South KPK	500 PKR per 10 g
Over-the-counter agents and household chemicals	May present with CNS depression, CNS stimulation or a mixed picture; the heart rate, blood pressure, body temperature, respiratory rate, skin clamminess, pupillary reaction and neuromuscular abnormalities would provide clues to the correct diagnosis	Clinical	Supportive care; decontamination and gastric lavage with activated charcoal antidote (flumazenil for benzodiazepines) should be used if available; haemodialysis, haemofiltration and exchange transfusion could facilitate the removal of some agents	Depends on agent used	2.5–25%	Urban areas	Diazepam is one of the most commonly used benzodiazepines; it is 37 PKR for 30 10-mg tablets

ATN, acute tubular necrosis; KPK, Khyber Pakhtunkhwa; CNS, central nervous system.

finding a high overall mortality rate of 63.5% (range 47–80%). One study included only 38 participants, 95% of whom were female, with a mortality rate of 47%. The second study confirmed the findings of the first paper, with a much higher number of reported cases (503 cases). The number of men in the second study rose to 21%, and the mortality rate rose to around 80%.

In Balochistan, there was only one study examining agents used by people attempting suicide. This sample included only 46 female participants in Quetta, an urban centre and the largest city in the province. This study only examined organophosphates but did not report mortality rates.

The Gilgit–Baltistan region included only one study, in the city of Ghizer (a rural town). This study included 49 individuals, all of whom were female. The means of suicide included jumping into a body of water (40%), ingesting a poisonous agent (30%), strangulation (11%) and the use of a firearm (5%). Mortality rates were not reported in this study.

In summary, organophosphate poisoning was reported from all four provinces. However, organophosphates played a more substantial role in the cases of suicide reported from North Punjab and interior Sindh, where it accounted for up to 60% of reported cases. Aluminium phosphide (wheat pill) poisoning was noted in agent overdoses reported mainly from North Punjab and North KPK, whereas paraphenylenediamine (*kala pathar*) was primarily used in suicide from South Punjab, with some reports from South KPK and interior Sindh. Compared with the rural population (where pesticides and paraphenylenediamine were most common), the urban population chose more varied agents for overdose, including household chemicals (bleach, corrosives), medicines (sedatives, tranquilisers, NSAIDs, antidepressants), rat poison pills and other toxic substances. Other means of suicide, such as hanging (asphyxiation), gunshot and drowning, were not the focus of our paper. Some studies in our analysis reported the reason for the suicide attempt. Five themes emerged, including financial problems, family conflicts, illicit spousal relationships, serious medical illness and failed romance. Studies did not report risk factors for suicide consistently enough to allow for a complete analysis of regional or urban versus rural differences in these risk factors.

## Discussion

To our knowledge, this is the first study to focus on the regional difference in suicide by poisoning in Pakistan. The results also suggest urban versus rural differences in the choice of poison. We discuss determinants of suicide behaviour and comprehensive management strategies for commonly used agents, to address existing gaps in suicide literature.

Our study found that pesticides (organophosphates and aluminium phosphide) are the most frequently used agents for suicide across Pakistan. As noted above, agriculture is the backbone of Pakistan's economy. The main crops include cotton, wheat, rice, maize and sugarcane, in addition to a large variety of regional fruits and vegetables.<sup>13,95</sup> The need to meet the ever-increasing demand is one of the driving forces of the phenomenal rise in pesticide use in farming and agriculture. It does not spare even the remote areas of Pakistan.<sup>96,97</sup> Pesticides are regulated in Pakistan by the Agriculture Pesticide Ordinance of 1971 (amended up to 1997) and Agriculture Pesticides Rules of 1973.<sup>98</sup> Pakistan's Agriculture and Research Council detailed several elements regarding registration, production, procurement, transportation, distribution, sale, storage, usage and the safe disposition of empty containers.<sup>98</sup> There are also institutional arrangements for pesticide monitoring and research.<sup>99</sup> However, pesticides are readily available, and their

unrestricted use continues to be widespread.<sup>100</sup> A sobering study from the Khoj Foundation in 2009 reported that Pakistan used 14 times more pesticides for wheat and rice crops than India. Furthermore, the researchers found:

'Pesticides are often stored in living rooms, among cookware and plates, and the bags in which they are sold are sometimes reused and sewn into quilts or floor covering. Utensils used to mix pesticides are often also used for cooking. They found that because women are not involved in the decision making around pesticide use and work both in the fields and in the home where pesticides are stored, they are at increased risk of poisoning.'<sup>101</sup>

Corresponding to these findings, several studies have investigated suboptimal or a complete lack of knowledge and awareness of pesticide hazards in these regions.<sup>102–104</sup> Although unintentional poisoning is beyond the scope of this paper, this information is crucial in providing a glimpse of the problem and how it relates to easy accessibility and means for self-harm and suicide.

In our analysis, organophosphate overdose was reported in studies from across Pakistan, with the highest number of cases from the Punjab and Sindh regions (Table 5). Twelve studies were from Karachi, representing urban Sindh. We believe that, being the largest city of the province and Pakistan, Karachi receives patients with suicide overdose from all over Sindh, to receive care in its well-equipped medical institutions.<sup>105</sup> Thus, the number of organophosphate poisonings from Karachi likely represents rural rather than urban Sindh. Similarly, studies from other metropolitan cities, such as Lahore or Rawalpindi in Punjab, treated patients with poisoning who were transferred from the surrounding rural areas to receive treatments. In the wheat-growing regions of North Punjab, aluminium phosphide or wheat pills are more readily available and were the most common agents to attempt suicide. North KPK also reported a high incidence of aluminium phosphide use.

Interestingly, there was no report of aluminium phosphide overdose from urban or interior Sindh, indicating that availability could be the critical factor in the choice of agent in suicide. As opposed to inhalational or skin contact in unintentional poisoning, ingestion was the most common method for suicide by pesticide.<sup>106–109</sup> The chemical structure and management of organophosphates and aluminium phosphide poisoning are discussed later in the paper.

Paraphenylenediamine is an ingredient in a compound commonly known as *kala pathar* (Black Stone) in Urdu. It is used as a chemical ingredient in temporary tattoo ink, fabrics, dark makeup, photocopying inks, printing, rubber products and gasoline. In the Indian subcontinent and North Africa, paraphenylenediamine is an ingredient of black henna, which is used for hair dye and tattoo ink.<sup>110–112</sup> Paraphenylenediamine was noted as the agent of choice for suicide in South Punjab, South KPK and interior Sindh. Its easy availability, unrestricted sale as a hair dye and the associated low cost of 10 PKR for a single dose (1 USD = 160.36 PKR (at the time of publication)) are the likely reasons behind the increasing number of cases in recent years.<sup>113</sup> The ease of preparing the suicide concoction by mixing *kala pathar* in water increases the probability of its use in poisoning.<sup>114</sup> Following the increasing number of cases, a unified social and print media campaign against the rapidly rising number of suicides with *kala pathar* led Punjab's government to issue a temporary ban on its open trading in September 2017 in South Punjab. In April 2018, the Punjab government expanded the temporary ban on *kala pathar* throughout the whole province.<sup>114–116</sup> The management of paraphenylenediamine poisoning is discussed later.

We found significant differences in the choice of agents for suicide in urban versus rural populations (see Table 4). *Kala*



*pathar* was used in 36% of overdose cases in the urban areas as opposed to 64% from the rural regions. More than 85% of the poisoning cases choosing organophosphates, aluminium phosphide and miscellaneous agents were from rural areas, whereas 94% of over-the-counter poisoning cases were from urban areas. Over-the-counter agents included drugs/medicines (benzodiazepines, tranquilisers, NSAIDs/analgesics, antidepressants, etc.), household toxins (bleach, rat poison pills or rodenticides, insecticides) and kerosene oil. The availability, accessibility and ease of use appeared to be significant factors influencing the choice of agents for suicide in our study.

Interestingly, the gender distribution was relatively similar for all agents except *kala pathar*, which favoured females (74% female *v.* 25% male). The category of ‘miscellaneous agents’ was mostly reported in males (19% female *v.* 81% male). Drowning or jumping into a lake or a river was a preferred method for suicide in North KPK and Gilgit–Baltistan, where there is ready access to rivers, lakes and streams. Except for North Punjab, where the female:male suicide ratio is almost equal, all other reported regions showed a higher incidence of reported suicide in females compared with males (see Table 3). The average age of suicide in our data was 27.9 years, with the youngest reported age of 20.5 years in Balochistan.

Suicide is a complex phenomenon, and its identity is often shrouded in mystery. Unspoken religious and cultural factors, especially in LMICs, may contribute toward its inadequate understanding, and Pakistan is no exception.<sup>116</sup> Our study highlights social determinants such as financial problems, gender and cultural stressors influencing suicide. Although not reported in all of the studies, we identified economic issues, family conflicts, illicit spousal relationships, serious illness and failed romance as commonly identified reasons for suicide. Pakistan is an economically strained country with a high unemployment rate.<sup>19</sup> Previous reports from the region similarly found a range of socially and culturally specific family problems, typically involving spouses, in-laws, parent–child conflicts, unfulfilled expectations at work or failure in school, and mental turmoil to be factors in suicide attempts.<sup>116</sup> Pakistan’s regional differences influence the execution of cultural norms. As discussed earlier, a low individualism index promotes collective culture, and a high masculinity index defines boundaries and gender roles.<sup>26</sup> A deviation from tribal tradition could lead to a sense of betrayal among other clan members that can incite violence, especially against women.<sup>10</sup> We found that all reported cases of suicide from Balochistan and Gilgit–Baltistan were females. In a recent news report, female suicides in the region were associated with the lack of freedom in choosing potential husbands.<sup>117</sup>

Ali et al suggest domestic and social issues as the most common reason for overdose accounting for up to 70% of the cases.<sup>118</sup> In comparison, prior psychiatric history of suicide was possibly linked with suicide attempts in only 10% of the patients.<sup>54</sup> As opposed to high-income countries, where primary psychiatric disorders such as major depression are often reported to be present in 80%–90% of deaths by suicide, in Pakistan, a premorbid mental health diagnosis is often absent.<sup>119</sup> Treatment could potentially be delayed, as the patient’s history, although very important, is often unreliable in suicide attempts.<sup>120,121</sup> Fear of persecution, stigma and confidentiality around such a sensitive issue may lead to the concealment of facts, both by the patient and the family.<sup>122</sup>

Gender inequality and discrimination are significant issues both globally and in Pakistan.<sup>123</sup> The country has a deep-rooted patriarchal culture with unequal gender role expectations.<sup>124–126</sup> Women are expected to do household chores for the extended family. Men are the primary authority figures and considered the traditional breadwinners, which gives them a superior position to women. Although an increasing number of women are

economically active, both in rural and urban areas, society has yet to recognise their contribution.<sup>23</sup> Women are seldom included in decision-making and continue to be victims of abuse.<sup>125,127</sup> Lack of gender-sensitive policies seems to hinder equitable political and economic status, birth gender ratios, illiteracy rates, maternal mortality rates and other health indicators in South Asian women.<sup>128</sup> As opposed to the West, marriage does not seem to be a protective factor against suicide in Pakistan. This likely indicates the high level of marital stress married women face compared with single women.<sup>129</sup> Ali et al identified the pursuit of higher education as an agent toward change for all genders in Pakistan. The authors also recognised the role of mass media in supporting women’s empowerment.<sup>125</sup>

With the increasing availability of handheld devices and internet access in both urban and rural areas of Pakistan, the influence of social media on suicide behaviour cannot be disregarded. In a recent study, Cheng et al identified the role of social media and Facebook in depicting suicide and having an intended effect of similar choice of agent in other suicides.<sup>130,131</sup> Others have focused on local newspapers and the impact of reporting suicide on the front pages.<sup>132</sup>

Religious beliefs can provide a series of effective coping strategies (e.g. prayer, rituals, religious services and social networks) that are considered as protective factors against suicide.<sup>133</sup> A strong belief in God and that whatever happens is by Allah’s will may create an atmosphere of acceptance rather than desperation in Muslims.<sup>26</sup> Rezaeian argues that Islam attempts to address the underlying factors contributing to the suicidal state, such as promoting mental health by the remembrance of the creator (*Zikr*), decreasing poverty by the distribution of wealth through mandatory charity (*Zakat*), and forbidding alcohol and other intoxicants.<sup>134</sup> Although religious beliefs and laws against suicide may be a deterrent, inadvertent negative consequences, such as a delay in help-seeking, fear of prosecution by the police and legal authorities, stigma and a lack of reliable statistics, can also occur.<sup>30</sup> For religious families, suicide is viewed as a sin and a failure rather than an illness. It may dictate family reactions, treatment-seeking behaviours, explanations of disease and adherence to treatment.<sup>39</sup> It is important to note that the clinicians’ own religious view of suicidal behaviour may lead to unconscious biases in delivering clinical care, and could lead to moral and ethical dilemmas when treating such patients.<sup>30</sup>

Recent literature has challenged the notion of outright faith-based protection. Pritchard et al explored ‘hidden’ or missed suicides in Islamic countries. They suggested that the official records seemed to be at odds with the study results purporting a higher number of suicides in Muslim-majority countries than previously reported.<sup>39,135</sup> The authors identified the risk of the unrecognised or denied extent of suicidal behaviour, undermining the necessary steps to support the individual and prevent fatal outcomes. Similarly, Jordans et al found a higher reported suicide rate in South Asia, mainly driven by Bangladesh (a Muslim-majority country), India and Sri Lanka, compared with the global average.<sup>136</sup>

Our data did not report on individual risk factors for suicide. However, we include a brief overview to emphasise its importance in the study of suicide. Previous analyses showed poor impulse control, premorbid depression, a history of physical/sexual or emotional abuse, high risk-taking behaviour and low self-esteem as contributing toward self-harm and suicide.<sup>41</sup> Cognitive factors such as low IQ and limited education; poor problem-solving or inadequate communication skills; lack of distress tolerance; and the timing of the attempt, such as after a similar attempt in the family or neighbourhood, may also have a significant effect on the choice of agent in self-poisoning.<sup>19,137,138</sup> Copycat suicides or Werther’s syndrome have long been identified as drivers of cluster suicides.<sup>139–143</sup>

Although we did not look for the timing of cluster suicides in our analysis, it should be explored in future studies.<sup>113</sup> In short, the prevalence, characteristics and methods of suicidal behaviour vary widely between different communities, across other demographic groups and over time.<sup>144</sup>

Structural determinants of health account for some of the regional variations noted in our study.<sup>145,146</sup> We argue that the easy and unrestricted availability of drugs/medicines could be one reason for these regional variations. Ali et al have raised concern about the lack of regulation for over-the-counter drugs in Pakistan, leading to misuse and overuse.<sup>118</sup> The authors did not consider the risk of suicide overdose with uncontrolled access to medications, which we believe should be factored into future regulations. Pakistan's growing income inequality and increase in poverty are concerning.<sup>146,147</sup> Li and Katikireddi emphasised the urban-rural inequalities as a driver of suicide trends.<sup>148</sup> The efforts to decriminalise suicide in Pakistan gained momentum likely after India decriminalised suicide in 2015.<sup>10,149,150</sup> Although Islam condemns those who commit suicide, no legal or societal punishment is mentioned for suicide survivors in the Quran.<sup>122</sup> In February of 2018, the Pakistan Senate passed a bill for treatment of those who attempt suicide and survive, rather than punishment under Section 325 of the Pakistan Penal Code.<sup>151</sup>

With the alarming rise of suicide rates in Pakistan, we must emphasise urgent steps to halt and gradually reverse the suicide trends. It is imperative to initiate mental health literacy and psychoeducational campaigns in vulnerable communities, to identify high-risk individuals and the hazardous effects of agricultural chemicals.<sup>152</sup> Furthermore, increasing the availability of resources for timely and prompt treatment of overdose may prevent dire consequences. The role of partnership with local leaders and utilisation of existing resources in such endeavours, such as governmental or non-governmental organisations, especially in rural areas, cannot be overemphasised.<sup>153,154</sup> In a recent article, Eddleston and Gunnell focused on preventing suicide through regulating pesticides, especially in LMICs.<sup>155</sup> Chowdhury et al reported the promising effects of a ban on class I pesticides in Bangladesh and a corresponding overall decrease in suicide rate in the region.<sup>156</sup>

Similarly, Sri Lanka and South Korea have achieved success through governmental regulations in the availability of pesticides and insecticides.<sup>5,157,158</sup> With our collective effort, there is no reason that Pakistan could not achieve the same. After the next three decades, a strikingly different review focusing on suicide rate reduction success may be reported. As noted above, the Punjab Government has taken the first steps to ban potentially harmful agents.<sup>115</sup> The Federal Government of Pakistan must follow suit in steering the campaign against suicide in the right direction. In a recent paper, Zia emphasised the need for clear warning labels, phrases in local languages and symbols on pesticides and other hazardous chemicals. The author suggested that the advertisement must include safety warnings as for cigarettes, and a strict following of Food and Agriculture Organization of the United Nations guidelines should be implemented.<sup>159</sup> The need for systemic media campaigns for awareness and safe pesticide is necessary. We believe that despite the recent step of passing the decriminalisation of suicide bill in the Senate, it will take a concerted effort to decrease stigma against suicide survivors.

## Management of individual agents

A summary of the management of agents is as follows (see Table 5):

### Organophosphates

Organophosphate compounds are a diverse group of chemicals used in domestic, industrial and agricultural settings. Examples include

insecticides and pesticides (malathion, parathion, etc.), herbicides (glyphosate, atrazine, etc.) and nerve gases (sarin, tabun, VX).<sup>160</sup> Organophosphate poisoning is one of the most common methods used for suicide, and is a leading cause of death in young people in Pakistan, China, India, Sri Lanka and other Asian countries.<sup>161-163</sup> It is recognised as the principal mode of poisoning in southern Punjab, and accounts for 47-60% of instances reported in Sindh.<sup>164-166</sup> Data from other parts of the country suggest organophosphates as a cause of poisoning in 20 to 40% of cases.<sup>50,54,165</sup>

Inhalation, ingestion or skin contact can lead to organophosphate poisoning. The organophosphate molecule binds and inactivates acetylcholinesterase enzyme in red blood cells. This leads to an overabundance of acetylcholine within both nicotinic and muscarinic synapses and the neuromuscular junctions.<sup>166</sup> The nicotinic effects are rapid in onset and may include twitching of fine muscles, fasciculations and hyperreflexia, which may progressively lead to flaccid paralysis. Muscarinic receptors are located in both the sympathetic and parasympathetic nervous systems, and are usually slower in onset because of their action via G-protein-coupled receptors. Symptoms such as bronchorrhea, bronchoconstriction, excessive sweating, constricted pupils, abdominal cramps, involuntary defaecation and urination, tachycardia, QT prolongation, headaches, dizziness, drowsiness, confusion, anxiety, slurred speech, ataxia, psychosis, convulsions, coma, hypotension and respiratory depression can occur.<sup>167-169</sup>

The diagnosis of organophosphate poisoning is clinical and based on the presenting history, collateral information from the attendants and the clinical signs. Confirmation of organophosphate poisoning can be obtained by measuring plasma butyrylcholinesterase activity or acetylcholinesterase in whole blood; however, such assays are not readily available to inform clinical decision-making.<sup>170</sup> The first step is to decontaminate the patient and prevent further absorption via the eyes, skin or lungs. Personal protective equipment must be used to avoid exposure. The standard treatment of organophosphate poisoning is the reversal of muscarinic manifestations using atropine, followed by enzyme reactivation by pralidoxime. Frequent atropine doses or continuous infusion are used to clear excessive respiratory secretions and to treat bradycardia.<sup>171</sup> Atropine should be continued for 1-3 days after successful atropinisation. Pralidoxime facilitates the recovery of neuromuscular transmission at the nicotinic synapses. It significantly reduces atropine consumption in organophosphate poisoning, and signs of atropinisation may occur earlier with its use than without its administration.<sup>172</sup>

In our analysis, mortality ranged from 10 to 27%. It was dependent on the amount of substance ingested, the time to reach an emergency department or time to initiation of treatment, and the use of a ventilator for assisted breathing.<sup>173</sup> Other predictors of mortality include age >40 years, bradycardia, low pH, high glucose, high lactate dehydrogenase and low Glasgow Coma Scale score.<sup>44,174</sup>

### Aluminium phosphide

Aluminium phosphide is a highly toxic, solid fumigant insecticide and rodenticide used for grain conservation.<sup>175</sup> It is referred to as wheat pills in Pakistan, and is also known as rice pills or rice tablets in other countries.<sup>176,177</sup> It is not regulated by the government and is available for over-the-counter purchase without any restriction, making it an ideal agent for self-poisoning in the wheat-growing areas of northern and central Punjab.<sup>178</sup> Studies have reported its use as an agent of suicide by ingestion from Rawalpindi,<sup>178</sup> Kharian,<sup>75</sup> Lahore,<sup>179</sup> Sahiwal<sup>78</sup> and Peshawar in KPK.<sup>54</sup> In these areas, domestic conflicts or petty quarrels are a frequent cause of overdose, resulting in fatal outcomes.<sup>78</sup> The lack of an antidote makes it a prevalent and particularly lethal suicide agent.<sup>175</sup>

When exposed to moisture in the stomach after ingestion, phosphine gas is produced. This toxic gas inhibits cytochrome c oxidase and other vital cellular enzymes, disrupting several metabolic pathways and destabilising cell membranes. Disruption of mitochondrial function produces reactive hydroxyl radicals, leading to cellular hypoxia, free-radical-mediated injury and eventual cell death.<sup>176,180</sup> The presenting symptoms of aluminium phosphide poisoning may include epigastric pain, vomiting, diarrhoea, dizziness and dyspnoea.<sup>178</sup> Multiorgan failure involving the heart, kidneys, lungs and liver later ensues, with metabolic acidosis, hepatic necrosis, renal failure, cardiac arrhythmia, congestive heart failure and hypotensive shock.<sup>180,181</sup>

A silver nitrate test can be performed to confirm the diagnosis. Paper impregnated with silver nitrate turns black after exposure to the patient's breath or gastric contents, as a result of the reaction between phosphides and silver nitrate. The sensitivity of the test strip is 50% with a breath test and 100% with gastric contents.<sup>182</sup>

The treatment is supportive because of the absence of an antidote. Gastric lavage with potassium permanganate and mineral or coconut oil has been shown to reduce morbidity.<sup>183</sup> Besides symptomatic treatment, renal replacement therapy in the early stage is also recommended.<sup>184</sup>

Aluminium phosphide is termed 'agent of sure death',<sup>185</sup> and the mortality rate ranged from 33 to 87% in our data.<sup>186</sup> The lethal dose for an adult is 150–500 mg. The presence of vomiting, exposure of tablets before ingestion and early availability of supportive care can help decrease mortality.

#### Paraphenylenediamine

Paraphenylenediamine is an ingredient of a compound commonly known as *kala pathar* or 'Black Stone' in Urdu. It is used as a chemical ingredient in temporary tattoo ink, fabrics, dark makeup, photocopying inks, printing, rubber products and gasoline. In the Indian subcontinent and North Africa, it is an ingredient of black henna for hair dye and tattoo ink.<sup>84–86</sup> Paraphenylenediamine is metabolised into benzoquinone diamine by cytochrome P450 peroxidase, and further oxidation results in the formation of Brandowaski's base. Both of these by-products are responsible for their toxicity.<sup>187,188</sup> Paraphenylenediamine ingestion is another conventional means to commit suicide in southern Punjab.<sup>189</sup>

The most common clinical presentations after paraphenylenediamine intoxication include cervicofacial oedema, rhabdomyolysis causing myoglobinuria, cola-coloured urine, oliguria and acute tubular necrosis leading to renal failure.<sup>190</sup> A study of 150 cases of paraphenylenediamine poisoning from Sudan revealed angioneurotic oedema and conjunctival discolouration in 100% of cases, and acute kidney injury requiring haemodialysis in 60% of cases.<sup>191</sup>

There is no antidote available for paraphenylenediamine poisoning. As the chemical is nondialysable, the mainstay of management remains supportive.<sup>192</sup> The patient must be observed in the intensive care unit. Management includes early tracheostomy for cervicofacial oedema and intravenous fluids, with aggressive diuresis and urine alkalisation for renal failure.<sup>193–195</sup> Rhabdomyolysis may lead to acute tubular necrosis, requiring haemodialysis.

The outcome of paraphenylenediamine ingestion depends on the dose taken. The lethal dose of paraphenylenediamine is unknown, and estimates vary from 7 to 10 g.<sup>196,197</sup> A large quantity (>7 g) might cause death within the first 6–24 h from angioneurotic oedema or cardiotoxicity.<sup>198</sup> The mortality ranges from 21 to 47%.

#### Others

This group included over-the-counter agents, prescription medicine, agents of abuse and household chemicals. This type of poisoning was more common in young patients (15–35 years) from urban

backgrounds.<sup>50,76,77,91,199</sup> Males overdosed at a higher rate than females.<sup>91,199</sup> Benzodiazepines were the most common agent used for overdose;<sup>91,116,197</sup> however, other agents used were NSAIDs, analgesics, sedatives, tricyclics, anti-emetics, antiallergics, anti-epileptics, oral hypoglycaemics, warfarin, digoxin, methamphetamine and cocaine.<sup>50,91,197</sup> Corrosives, kerosene oil, rubbing alcohol, copper sulphate, bleach, rat poison pills and home insecticide sprays were also used.<sup>63,76,77,91,199</sup> Most patients taking an overdose had an intention to commit suicide; however, other reasons for overdose were to gain attention, express distress or get revenge.<sup>50,200</sup> The researchers interviewed a total of 80 individuals admitted after suicide overdose, to determine their intention to die, and noted that the patients with such an intention chose organophosphates because of its known high lethality.

The most common presentation was drowsiness owing to central nervous system depression; others presented with central nervous system stimulation or a mixed picture.<sup>91</sup> Heart rate, blood pressure, body temperature, respiratory rate, skin clamminess, pupillary reaction and neuromuscular abnormalities provided clues to the correct diagnosis.

Treatment includes decontamination and gastric lavage with activated charcoal. The use of the benzodiazepine antidote flumazenil remains controversial as it could precipitate withdrawal seizures in individuals who have developed tolerance from chronic use.<sup>201</sup> Flumazenil use in paediatric benzodiazepine overdose may be used as young children are unlikely to be tolerant to benzodiazepines.<sup>202</sup> Haemodialysis, haemofiltration and exchange transfusion could facilitate removing the agents or chemicals from circulation. Supportive care is indicated for strict airway monitoring, gastrointestinal protection and the treatment of hypo- or hypertension.

Mortality varied from 2.5 to 25%, depending on the place of study. General medical wards reported lower death rates than intensive care units, likely related to the severity of the patient's condition.<sup>50,54</sup> Mortality was also dependant on the level of care available in the centre where the patient was under treatment.

There are several limitations to this analysis. We considered papers in the English language, from only two electronic databases, and excluded single case reports in this retrospective analysis. Significant variations in the reported information in descriptive studies make it difficult to analyse or present the data in a meta-analysis. Limited data were available from Balochistan and Gilgit-Baltistan, and studies from other provinces also represented only larger cities. Risk factors were not available for extensive analysis. More comprehensive studies are required to explore how individual differences influence regional trends of suicide and other means of suicide that were not addressed in our review.

## Summary and future directions

Our study found that there are striking regional and urban versus rural differences in the choice of agents used for suicide. As the suicide rate in Pakistan is rapidly increasing, we must take several steps to reverse the trend of the past three decades. We should launch customised mental health literacy and public health awareness campaigns across the country, to address the stigma against suicide and mental health. The success and acceptance of such endeavours will depend on partnership with local authorities, tribal or clan leaders, religious leaders and influential community figures. Print (newspapers, magazines, etc.), electronic (network television, radio, etc.) and social media (Facebook, Twitter, Instagram, YouTube, etc.) may enhance the dissemination of the message. Efforts are needed to enforce the existing national pesticide policy. There is a need to have regulations to restrict over-the-counter sales of potentially dangerous medications,



such as benzodiazepines, opiates and opioid derivatives. Finally, our hospitals need consistent medical supplies and specialised equipment, along with training of medical staff, to manage victims adequately. These interventions are necessary to reduce morbidity and mortality related to suicide poisoning in this time of crisis.

### Data availability

The authors confirm that the data supporting the findings of this study are available within the article.

**Maria Safdar** , Department of Forensic Medicine, Postgraduate Medical Institute, Pakistan; **Khalid Imran Afzal** , Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Illinois, USA; **Zoe Smith** , Department of Psychology, Loyola University, Illinois, USA; **Filza Ali**, Department of Forensic Medicine, CMH Multan Institute of Medical Sciences, Pakistan; **Pervaiz Zarif** , Department of Forensic Medicine, Postgraduate Medical Institute, Pakistan; **Zahid Farooq Baig** , Department of Medicine, CMH Lahore Medical College and Institute of Dentistry, Pakistan

**Correspondence:** Maria Safdar. Email: mariasafdar79@gmail.com

First received 12 Jul 2020, final revision 26 Apr 2021, accepted 7 May 2021

### Author contributions

M.S., F.A., Z.F.B. and P.Z. identified review articles and planned the review. Z.S. analysed the data and wrote the Results section and the associated tables. M.S. and K.I.A. wrote the manuscript. Z.F.B. wrote the management section of the Discussion section and created the associated table. All authors contributed to and have approved the final manuscript.

### Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

### Declaration of interest

None.

### References

- World Health Organization (WHO). *Preventing Suicide: A Global Imperative*. WHO, 2014 ([https://apps.who.int/iris/bitstream/handle/10665/131056/9789241564878\\_eng.pdf;jsessionid=3D97FDA8D92F043F43C07B007DD13D14?sequence=%208](https://apps.who.int/iris/bitstream/handle/10665/131056/9789241564878_eng.pdf;jsessionid=3D97FDA8D92F043F43C07B007DD13D14?sequence=%208)).
- World Health Organization (WHO). *10 Leading Causes of Death by Age Group, United States – 2017*. Centers for Disease Control and Prevention, 2019 ([https://www.cdc.gov/injury/images/lc-charts/leading-causes\\_of\\_death\\_by\\_age\\_group\\_2017\\_1100w850h.jpg](https://www.cdc.gov/injury/images/lc-charts/leading-causes_of_death_by_age_group_2017_1100w850h.jpg)).
- World Health Organization (WHO). *Mental Health and Substance Use: Suicide Data*. WHO, 2019 ([https://www.who.int/mental\\_health/prevention/suicide/suicideprevent/en/](https://www.who.int/mental_health/prevention/suicide/suicideprevent/en/)).
- Prydz EB, Wadhwa D. *Classifying Countries by Income*. The World Bank, 2019 (<https://datatopics.worldbank.org/world-development-indicators/stories/the-classification-of-countries-by-income.html#:~:text=As%20of%201%20July%202019,between%20%243%2C996%20and%20%2412%2C375%3B%20high%2D>).
- Cha ES, Chang SS, Choi Y, Lee WJ. Trends in pesticide suicide in South Korea, 1983–2014. *Epidemiol Psychiatr Sci* 2019; **29**: e25.
- Demir M. Trends in suicide methods by age group. *Asia Pac Psychiatry* 2018; **10**(4): e12334.
- Ohberg A, Lonnqvist J, Sarna S, Vuori E, Penttila A. Trends and availability of suicide methods in Finland. Proposals for restrictive measures. *Br J Psychiatry* 1995; **166**(1): 35–43.
- Snowdon J. Differences between patterns of suicide in East Asia and the West. The importance of sociocultural factors. *Asian J Psychiatr* 2018; **37**: 106–11.
- Abdullah M, Khalili MT, Ahmad I, Hallahan B. Psychological autopsy review on mental health crises and suicide among youth in Pakistan. *Asia Pac Psychiatry* 2018; **10**(4): e12338.
- Shekhani SS, Perveen S, Hashmi DE, Akbar K, Bachani S, Khan MM. Suicide and deliberate self-harm in Pakistan: a scoping review. *BMC Psychiatry* 2018; **18**: 44.
- United States Census Bureau. *U.S. Census Bureau Current Population*. United States Census Bureau, 2020 (<https://www.census.gov/popclock/print.php?component=counter&image=https://www.census.gov/images/census-logo-whiteBG.png>).
- Pakistan Bureau of Statistics. *6th Population and Housing Census 2017*. Pakistan Bureau of Statistics, 2017 (<https://www.pbs.gov.pk/content/final-results-census-2017>).
- Rehman A, Jingdong L, Shahzad B, Chandio AA, Hussain I, Nabi G, et al. Economic perspectives of major field crops of Pakistan: an empirical study. *Pac Sci Rev B: Humanit Soc Sci* 2015; **1**(3): 145–58.
- UNESCO Institute for Statistics. *Literacy Rate, Adult Total (% of People Ages 15 and Above) - Pakistan 2017*. The World Bank, 2020 (<https://data.worldbank.org/indicator/SE.ADT.LITR.ZS?contextual=aggregate&locations=PK>).
- Hunter R. *Education in Pakistan*. World Education Services, 2020 (<https://wenr.wes.org/2020/02/education-in-pakistan>).
- Rehman A, Jingdong L, Hussain I. The province-wise literacy rate in Pakistan and its impact on the economy. *Pac Sci Rev B: Humanit Soc Sci* 2015; **1**(3): 140–4.
- Zakaria M, Jun W, Ahmed H. Effect of terrorism on economic growth in Pakistan: an empirical analysis. *Econ Res Ekonomika Istraživanja* 2019; **32** (1): 1794–812.
- Khan NH, Ju Y, Hassan ST. Modeling the impact of economic growth and terrorism on the human development index: collecting evidence from Pakistan. *Environ Sci Pollut Res* 2018; **25**(34): 34661–73.
- World Economic Outlook. *Unemployment Rate Percent - Pakistan 2021*. International Monetary Fund, 2021 (<https://www.imf.org/external/datamapper/LUR@WEO/PAK>).
- Shah Z. *Underperforming on Most Social Development Indicators*. The Express Tribune, 2017 (<https://tribune.com.pk/story/1410585/underperforming-social-development-indicators>).
- Anjum A, Saeed Ali T, Akber Pradhan N, Khan M, Karmaliani R. Perceptions of stakeholders about the role of health system in suicide prevention in Ghizer, Gilgit-Baltistan, Pakistan. *BMC Public Health* 2020; **20**: 991.
- Rej A. *Gilgit-Baltistan to Become a New Province of Pakistan, Announces Khan Government*. The Diplomat, 2020 (<https://thedi diplomat.com/2020/11/gilgit-baltistan-to-become-a-new-province-of-pakistan-announces-khan-government/>).
- UKessays. *Culture Diversity In Pakistan Cultural Studies Essay*. UKessays, 2018 (<https://www.ukessays.com/essays/cultural-studies/culture-diversity-in-pakistan-cultural-studies-essay.php?vref=1>).
- Finlayson C. Chapter 9.3: Pakistan and Bangladesh. In *World Regional Geography: People, Places and Globalization*. University of Minnesota Libraries Publishing, 2014.
- Bukhari SM. *The Serenity of Ghanche: Of Mountains, Rivers and Valleys*. Dawn, 2017 (<https://www.dawn.com/news/1176091>).
- Shah SAM, Amjad S. Cultural diversity in Pakistan: national vs provincial. *Mediterr J Soc Sci* 2011; **2**: 331–44.
- Pakistan Bureau of Statistics. *Population by Religion*. Pakistan Bureau of Statistics, [updated 9 Jan 2021; cited 9 Jan 2021]. (<http://www.pbs.gov.pk/sites/default/files/tables/POPULATION%20BY%20RELIGION.pdf>).
- Hussain R. "Pakistan." In *The Oxford Encyclopedia of the Islamic World*. Oxford Islamic Studies Online, 2021.
- Gearing RE, Lizardi D. Religion and suicide. *J Relig Health* 2009; **48**(3): 332–41.
- Khan MM, Mian AI. 'The one truly serious philosophical problem': ethical aspects of suicide. *Int Rev Psychiatry* 2010; **22**(3): 288–93.
- Chaleby KS. Issues in forensic psychiatry in Islamic jurisprudence. *Bull Am Acad Psychiatry Law* 1996; **24**(1): 117–24.
- Simpson ME, Conklin GH. Socioeconomic development, suicide and religion: a test of Durkheim's theory of religion and suicide. *Soc Forces* 1989; **67**(4): 945–64.
- Faruqui R, Afghan S. P03-445 - suicide risk awareness in Pakistan: influence of religious, cultural, legal, socio-economic and interpersonal factors. *Eur Psychiatry* 2011; **26**: 1615.
- Khan MM, Hyder AA. Suicides in the developing world: case study from Pakistan. *Suicide Life Threat Behav* 2006; **36**(1): 76–81.
- Lester D. Suicide and Islam. *Arch Suicide Res* 2006; **10**(1): 77–97.
- Eskin M, AlBuhairan F, Rezaeian M, Abdel-Khalek AM, Harlak H, El-Nayal M, et al. Suicidal thoughts, attempts and motives among university students in 12 Muslim-majority countries. *Psychiatr Q* 2019; **90**(1): 229–48.
- Karamouzin M, Rostami M. Suicide statistics in Iran: let's get specific. *Am J Mens Health* 2019; **13**(1): 1557988318807079.
- Arya V, Page A, Dandona R, Vijayakumar L, Mayer P, Armstrong G. The geographic heterogeneity of suicide rates in India by religion, caste, tribe, and other backward classes. *Crisis* 2019; **40**(5): 370–4.

- 39 Pritchard C, Iqbal W, Dray R. Undetermined and accidental mortality rates as possible sources of underreported suicides: population-based study comparing Islamic countries and traditionally religious Western countries. *BJPsych Open* 2020; **6**(4): e56.
- 40 Institute for Health Metrics and Evaluation. *GBD Results Tool: Suicide by Gender 2018*. Global Health Data Exchange, 2018 (<http://ghdx.healthdata.org/gbd-results-tool>).
- 41 Bachmann S. Epidemiology of suicide and the psychiatric perspective. *Int J Environ Res Public Health* 2018; **15**(7): 1425.
- 42 Khan MM, Mahmud S, Karim MS, Zaman M, Prince M. Case-control study of suicide in Karachi, Pakistan. *Br J Psychiatry* 2008; **193**(5): 402–5.
- 43 Yousafzai AW, Yousafzai S, Khan SA. Rising suicide rates in Pakistan: is it about time to break the silence? *J Ayub Med Coll Abbottabad* 2020; **32**(2): 153–4.
- 44 Khan MM, Ahmed A, Khan SR. Female suicide rates in Ghizer, Pakistan. *Suicide Life Threat Behav* 2009; **39**(2): 227–30.
- 45 Anjum A, Saeed Ali T, Akber Pradhan N, Khan M, Karmaliani R. Perceptions of stakeholders about the role of health system in suicide prevention in Ghizer, Gilgit-Baltistan, Pakistan. *BMC Public Health* 2020; **20**: 991.
- 46 Hassan T. *Why Are More Pakistanis Taking Their Own Lives?* Dawn, 2019 (<https://www.dawn.com/news/1481826>).
- 47 World Health Organization (WHO). *The World Health Report 2004 - Changing History*. WHO, 2004 (<https://www.who.int/whr/2004/en/>).
- 48 Bhatti N, Khan DA, Saleem S, Ijaz A, Amir M. Frequency of drug poisoning in adults at tertiary care hospital. *Wah Cantt. Pak J Pathol* 2015; **26**(1): 27–34.
- 49 Maqbool F, Satti AI, Jeelani RA, Baqai HZ. Organophosphate poisoning – clinical profile. *J Rawal Med College (KRMC)* 2015; **19**(1): 15–9.
- 50 Khurram M, Mahmood N. Deliberate self-poisoning: experience at a medical unit. *J Pak Med Assoc* 2008; **58**(8): 455–7.
- 51 Tahir MN, Akbar AH, Naseer R, Khan QO, Khan F, Yaqub I. Suicide and attempted suicide trends in Mianwali, Pakistan: social perspective. *East Mediterr Health J* 2014; **19**(3): S111–4.
- 52 Naheed T, Akbar N, Akbar N, Munir R. Acute poisoning in the city of Punjab - how can we help these souls? *J Fatima Jinnah Med Coll Lahore* 2007; **1**(3–4): 56–8.
- 53 Ali Z, Afridi MAR, Muhammad R, Rahim A, Rahman SKU, Ullah N, et al. Outcome and predictors of in-hospital mortality in patients presenting with acute poisoning to a teaching hospital. *J Postgrad Med Inst* 2018; **32**(2): 155–61.
- 54 Rahim F, Ullah F, Haroon M, Ashfaq M, Afridi AK. Acute poisoning treated in medical intensive care unit. *Gomal J Med Sci* 2016; **14**: 129–32.
- 55 Bilal M, Khan Y, Saad A, Awais N. The pattern of organophosphorus poisoning and its short term outcomes in various socioeconomic groups. *Khyber Journal of Medical Sciences (KJMS)* 2014; **7**(1): 11–16. ([https://www.researchgate.net/profile/Awais-Naeem/publication/335230875\\_THE\\_PATTERN\\_OF\\_ORGANOPHOSPHORUS\\_POISONING\\_AND\\_ITS\\_SHORT\\_TERM\\_OUTCOMES\\_IN\\_VARIOUS\\_SOCIOECONOMIC\\_GROUPS/links/5d59ab22299bf151badea9b7/THE-PATTERN-OF-ORGANOPHOSPHORUS-POISONING-AND-ITS-SHORT-TERM-OUTCOMES-IN-VARIOUS-SOCIOECONOMIC-GROUPS.pdf](https://www.researchgate.net/profile/Awais-Naeem/publication/335230875_THE_PATTERN_OF_ORGANOPHOSPHORUS_POISONING_AND_ITS_SHORT_TERM_OUTCOMES_IN_VARIOUS_SOCIOECONOMIC_GROUPS/links/5d59ab22299bf151badea9b7/THE-PATTERN-OF-ORGANOPHOSPHORUS-POISONING-AND-ITS-SHORT-TERM-OUTCOMES-IN-VARIOUS-SOCIOECONOMIC-GROUPS.pdf)).
- 56 Shaikh MA. Mortality in patients presenting with organophosphorus poisoning at Liaquat university of medical and health sciences. *Pak J Med Sci* 2011; **27**(5): 1022–4.
- 57 Imran S, Awan EA, Memon MIS, Memon A. Frequency and outcomes of organophosphate poisoning at tertiary care hospital in Nawabshah. *J Liaquat Uni Med Health Sci* 2017; **16**(2): 118–20.
- 58 Faiz MS, Mughal S, Memon AQ. Acute and late complications of organophosphate poisoning. *J Coll Physicians Surg Pak* 2011; **21**(5): 288–90.
- 59 Shaikh MA, Ujjan ID, Memon SH. Evaluation of patients with organophosphorus poisoning at a tertiary care hospital of Sindh. *Med Channel* 2011; **17**(3): 51–3.
- 60 Amir A, Haleem F, Mahesar G, Sattar RA, Qureshi T, Syed JG, et al. Epidemiological, poisoning characteristics and treatment outcomes of patients admitted to the national poisoning control centre at Karachi, Pakistan: a six month analysis. *Cureus* 2019; **11**(11): e6229.
- 61 Khan NU, Khan UR, Feroze A, Khan SA, Ali N, Ejaz K, et al. Trends of acute poisoning: 22 years experience from a tertiary care hospital in Karachi, Pakistan. *J Pak Med Assoc* 2016; **66**(10): 1237–42.
- 62 Ahmed A, Ali L, Shehbaz L, Nasir S, Rizvi SRH, Aman MZ, et al. Prevalence and characteristics of organophosphate poisoning at a tertiary care centre in Karachi, Pakistan. *Pak J Surg* 2016; **32**(4): 269–73.
- 63 Imtiaz F, Ali M, Ali L. Prevalence of chemical poisoning for suicidal attempts in Karachi, Pakistan. *Emerg Med (Los Angel)* 2015; **5**: 247.
- 64 Asghar SP, Ather N, Farooq M, Sidra S, Asghar S, Ijaz A. Presentation and management of organophosphate poisoning in an intensive care unit. *Pak Armed Forces Med J* 2014; **64**(1): 134–8.
- 65 Bashir F, Ara J, Kumar S. Deliberate self poisoning at national poisoning control centre. *J Liaquat Univ Med Health Sci* 2014; **13**(1): 3–8.
- 66 Ali P, Anwer A, Bashir B, Jabeen R, Haroon H, Makki K. Clinical pattern and outcome of organophosphorus poisoning. *J Liaquat Univ. Med. Health Sci* 2012; **11**(1): 15–8.
- 67 Ather NA, Ara J, Khan EA, Sattar RA, Durrani R. Acute organophosphate insecticide poisoning. *Journal of Surgery Pakistan (International)* 2008; **13**(2): 71–4.
- 68 Turabi A, Danyal A, Hasan S, Durrani A, Ahmed M. Organophosphate poisoning in the urban population; study conducted at National Poison Control Center Karachi. *Biomedica* 2008; **24**: 124–9.
- 69 Hussain AM, Sultan ST. Organophosphorus insecticide poisoning: management in surgical intensive care unit. *J Coll Physicians Surg Pak* 2005; **15**(2): 100–2.
- 70 Jamil H. Acute poisoning - A review of 1900 cases. *J Pak Med Assoc* 1990; **40**: 131.
- 71 Jamil H. Organophosphorus insecticide poisoning. *J Pak Med Assoc* 1989; **39**: 27–31.
- 72 Khan NK, Shamim H. Deliberate self harm due to organophosphates. *J Pak Inst Med Sci. (JPIMS)* 2003; **14**(2): 784–9.
- 73 Hassan A, Manzoor MS, Shafique M, Adil. Wheat pill poisoning: clinical manifestation and its outcome. *J Rawal Med Univ* 2014; **18**(1): 49–51.
- 74 Iftikhar R, Tariq KM, Saeed F, Khan MB, Babar NF. Wheat pill: characteristics and outcome. *Pak Armed Forces Med J* 2011; **61**(3): 350–3.
- 75 Ghazi MA. Wheat pill [aluminum phosphide] poisoning; commonly ignored dilemma. A comprehensive clinical review. *Professional Med J -Q* 2013; **20**(6): 855–63.
- 76 Shoaib S, Nadeem M, Khan Z. Causes and outcome of suicidal cases presented to a medical ward. *Ann King Edward Med Univ* 2016; **11**(1): 30–2.
- 77 Asif A, Yusuf F, Haider K, Gul H, Usman S, Akbar S, et al. Epidemiology of attempted suicides in emergency of Mayo Hospital in 2004. *Ann King Edward Med Univ* 2005; **11**(4): 384–6.
- 78 Qureshi MA, Nadeem S, Ahmed T, Tariq F, Rehman H, Qasim AP. Aluminium phosphide poisoning: clinical profile and outcome of patients admitted in a tertiary care hospital. *Ann Punjab Med Coll* 2018; **12**(3): 191–4.
- 79 Akram A, Shahid RA, Tariq M. Kala Pathar (Paraphenylene Diamine) poisoning; Role of tracheostomy: our experience at DHQ hospitals. *Pak J Med Health Sci* 2018; **12**(2): 865–6.
- 80 Tanweer S, Saeed M, Zaidi S, Aslam W. Clinical profile and outcome of Paraphenylene Diamine. *J Coll Physicians Surg Pak*. 2018; **28**(5): 374–7.
- 81 Haider SH, Sultan A, Salman Z, Waris S, Bandesha Y. Paraphenylenediamine poisoning: clinical presentations and outcomes. *Anaesth Pain & Intensive Care* 2017; **21**(4): 43–7.
- 82 Khan MA, Akram S, Shah HBU, Hamdani SAM, Khan M. Epidemic of Kala Pathar (Paraphenylene Diamine) poisoning: an emerging threat in Southern Punjab. *J Coll Physicians Surg Pak*. 2018; **28**(1): 44–7.
- 83 Ishtiaq R, Shafiq S, Imran A, Masroor Ali Q, Khan R, Tariq H, et al. Frequency of acute hepatitis following acute paraphenylene diamine intoxication. *Cureus* 2017; **9**(4): e1186.
- 84 Qasim AP, Ali AM, Baig A, Moazzam MS. Emerging trend of self harm by using 'kala pathar' hair dye (paraphenylene diamine): an epidemiological study. *Ann Punjab Med Coll* 2016; **10**(1): 26–30.
- 85 Akbar K, Iqbal J, Rehman H, Iqbal R. Acute renal failure among kala pathar poisoning. *J Sheikh Zayed Med Coll*. 2017; **8**(2): 1153–6.
- 86 Ansari RZ, Khosa AH, Yadain SM, Shafi S, Haq AU, Khalil ZH. Incidence of paraphenylene diamine poisoning in three district headquarter hospitals of Pakistan. *J Ayub Med Coll Abbottabad* 2019; **31**(4): 544–7.
- 87 Khan H, Khan N, Khan N, Ahmad I, Shah F, Rahman AU, et al. Clinical presentation and outcome of patients with paraphenylenediamine (kala-pathar) poisoning. *Gomal J Med Sci* 2016; **14**: 3–6.
- 88 Kazi MA, Shaikh AR, Samad A, Bibi I, Khan M. Kala pathar (Paraphenylene Diamine) poisoning: an ICU based observational study at Hyderabad, Pakistan. *Indo Am. J. P. Sci* 2018; **5**(9): 9334–7.
- 89 Khuhro BA, Khaskheli MS, Shaikhet AA. Paraphenylene diamine poisoning: our experience at PMC hospital Nawabshah. *Anaesth Pain & Intensive Care* 2012; **16**(3): 243–6.
- 90 Hashmi MU, Ali M, Ullah K, Aleem A, Khan IH. Clinico-epidemiological characteristics of corrosive ingestion: a cross sectional study at a tertiary care hospital of Multan, South Punjab Pakistan. *Cureus* 2018; **10**(5): e2704.
- 91 Patel MJ, Shahid M, Riaz M, Kashif W, Ayaz SI, Khan MS, et al. Drug overdose: a wake up call! Experience at a tertiary care centre in Karachi, Pakistan. *J Pak Med Assoc* 2008; **58**(6): 298–301.
- 92 Khan MM, Reza H. Benzodiazepine self-poisoning in Pakistan: implications for prevention and harm reduction. *J Pak Med Assoc* 1998; **48**: 293–5.



- 93 Ahmed Z, QNisa A, Yousafzai SK, Chaudhry J. Trends and patterns of suicide in people of Chitral, Khyber Pakhtunkhwa, Pakistan. *Khyber Med Univ J* 2016; **8** (2): 72–7.
- 94 Khan MM, Ahmed A, Khan SR. Female suicide rates in Ghizer, Pakistan. *Suicide Life Threat Behav* 2009; **39**(2): 227–30.
- 95 Miller V, Yusuf S, Chow CK, Dehghan M, Corsi DJ, Lock K, et al. Availability, affordability, and consumption of fruits and vegetables in 18 countries across income levels: findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet Glob Health* 2016; **4**(10): e695–703.
- 96 Mohammad N, Mohammad Rasul J, Hizbullah K. Pesticide use in Swat Valley, Pakistan. *Mt Res Dev* 2008; **28**(3): 201–4.
- 97 Ullah R, Asghar R, Baqar M, Mahmood A, Ali SN, Sohail M, et al. Assessment of organochlorine pesticides in the Himalayan riverine ecosystems from Pakistan using passive sampling techniques. *Environ Sci Pollut Res Int* 2019; **26**(6): 6023–37.
- 98 Ali MA. *The Pesticides Registered with Recommendations for Safe Handling and Use in Pakistan*. Pakistan Agricultural Research Council, 2018.
- 99 Jabbar A, Mallick S. *Institutional arrangements for pesticide monitoring and research*. In *Pesticides and Environment Situation in Pakistan: 6–8*. Sustainable Development Policy Institute, 1994.
- 100 Mughal FH. *Environmental Impact of Pesticide Overuse*. Dawn, 2018 (<https://www.dawn.com/news/1406013>).
- 101 Global Greengrants Funds. *Poisoned by Pesticides in Pakistan*. Global Greengrants Funds, 2011 (<https://www.greengrants.org/2011/11/10/poisoned-by-pesticides-in-pakistan/>).
- 102 Ahmad A, Shahid M, Khalid S, Zaffar H, Naqvi T, Pervez A, et al. Residues of endosulfan in cotton growing area of Vehari, Pakistan: an assessment of knowledge and awareness of pesticide use and health risks. *Environ Sci Pollut Res Int* 2019; **26**(20): 20079–91.
- 103 Saeed MF, Shaheen M, Ahmad I, Zakir A, Nadeem M, Chishti AA, et al. Pesticide exposure in the local community of Vehari District in Pakistan: an assessment of knowledge and residues in human blood. *Sci Total Environ* 2017; **587–8**: 137–44.
- 104 Khan M, Mahmood HZ, Damalas CA. Pesticide use and risk perceptions among farmers in the cotton belt of Punjab, Pakistan. *Crop Protection* 2015; **67**: 184–90.
- 105 Medical and Health Services, Departments, Karachi Metropolitan Corporation 2021. (<http://www.kmc.gos.pk/Departments.aspx>).
- 106 Amir A, Raza A, Qureshi T, Mahesar GB, Jaffer S, Haleem F, et al. Organophosphate poisoning: demographics, severity scores and outcomes from national poisoning control centre, Karachi. *Cureus* 2020; **12** (5): e8371.
- 107 Khan M, Damalas CA. Occupational exposure to pesticides and resultant health problems among cotton farmers of Punjab, Pakistan. *Int J Environ Health Res* 2015; **25**(5): 508–21.
- 108 Bakhsh K, Ahmad N, Kamran MA, Hassan S, Abbas Q, Saeed R, et al. Occupational hazards and health cost of women cotton pickers in Pakistani Punjab. *BMC Public Health* 2016; **16**: 961.
- 109 Boedeker W, Watts M, Clausing P, Marquez E. The global distribution of acute unintentional pesticide poisoning: estimations based on a systematic review. *BMC Public Health* 2020; **20**: 1875.
- 110 Abdelraheem MB, El-Tigani MA, Hassan EG, Ali MA, Mohamed IA, Nazik AE. Acute renal failure owing to paraphenylene diamine hair dye poisoning in Sudanese children. *Ann Trop Paediatr* 2009; **29**(3): 191–6.
- 111 Abdelraheem M, Ali ET, Hussien R, Zijlstra E. Paraphenylene diamine hair dye poisoning in an adolescent. *Toxicol Ind Health* 2011; **27**(10): 911–3.
- 112 Ismaeel T. *Kala Pathar Rains Black Death on DG Khan*. The Express Tribune, 2019 (<https://tribune.com.pk/story/1921981/kala-pathar-rains-black-death-dg-khan>).
- 113 Birmani TS. *156 Women Attempted Suicide by Consuming 'Kala Pathar' in Dera Ghazi Khan Last Year*. Dawn, 2018 (<https://www.dawn.com/news/1388360>).
- 114 Ahmed S. *Kala Pathar Trade Banned to Stem Suicide Cases*. Dawn, 2017 (<https://www.dawn.com/news/1360913>).
- 115 INP. *Kala Pathar Banned to Stop Suicides in S Punjab*. The Nation, 2018 (<https://nation.com.pk/10-Apr-2018/kala-pathar-banned-to-stop-suicides-in-s-punjab>).
- 116 Institute of Medicine Committee on Pathophysiology Prevention of Adolescent Adult Suicide. *Reducing Suicide: A National Imperative* (eds Goldsmith SK, Pellmar TC, Kleinman AM, Bunney WE). National Academies Press (US), 2002.
- 117 Shah N. *Suicide: A Plague in Balochistan*. Daily Times, 2018 (<https://dailylimes.com.pk/303073/suicide-a-plague-in-balochistan/>).
- 118 Ali M, Abbasi BH, Ahmad N, Fazal H, Khan J, Ali SS. Over-the-counter medicines in Pakistan: misuse and overuse. *Lancet* 2020; **395**(10218): 116.
- 119 Kripipe D, Williams AJ, Hannam-Swain S, Upton S, Brown K, Bandara P, et al. Psychiatric morbidity and suicidal behaviour in low- and middle-income countries: a systematic review and meta-analysis. *PLoS Med* 2019; **16**(10): e1002905-e.
- 120 Hom MA, Stanley IH, Duffy ME, Rogers ML, Hanson JE, Gutierrez PM, et al. Investigating the reliability of suicide attempt history reporting across five measures: a study of US military service members at risk of suicide. *J Clin Psychol* 2019; **75**(7): 1332–49.
- 121 Hom MA, Joiner TE, Bernert RA. Limitations of a single-item assessment of suicide attempt history: implications for standardized suicide risk assessment. *Psychol Assess* 2016; **28**(8): 1026–30.
- 122 Naveed S, Qadir T, Afzaal T, Waqas A. Suicide and its legal implications in Pakistan: a literature review. *Cureus* 2017; **9**(9): e1665-e.
- 123 Hausmann R, Tyson LDA, Zahidi S (eds). *The Global Gender Gap Report 2012*. World Economic Forum Geneva, 2012 ([http://www3.weforum.org/docs/WEF\\_GenderGap\\_Report\\_2012.pdf](http://www3.weforum.org/docs/WEF_GenderGap_Report_2012.pdf)).
- 124 Chauhan K. Patriarchal Pakistan: women's representation, access to resources, and institutional practices. In *Gender Inequality in the Public Sector in Pakistan: Representation and Distribution of Resources* (ed. Chauhan K): 57–87. Palgrave Macmillan, 2014.
- 125 Ali TS, Krantz G, Gul R, Asad N, Johansson E, Mogren I. Gender roles and their influence on life prospects for women in urban Karachi, Pakistan: a qualitative study. *Glob Health Action* 2011; **4**: 7448.
- 126 Georgas JE, Berry JW, Van de Vijver FJ, Kağıtçıbaşı ÇE, Poortinga YH. *Families Across Cultures: A 30-Nation Psychological Study*. Cambridge University Press, 2006.
- 127 Rabbani F, Qureshi F, Rizvi N. Perspectives on domestic violence: case study from Karachi, Pakistan. *East Mediterr Health J* 2008; **14**(2): 415–26.
- 128 Gill R, Stewart DE. Relevance of gender-sensitive policies and general health indicators to compare the status of South Asian women's health. *Womens Health Issues* 2011; **21**(1): 12–8.
- 129 Shahid M, Iqbal R, Khan MM, Khan MZ, Shamsi US, Nakeer R. Risk factors for deliberate self-harm in patients presenting to the emergency departments of Karachi. *J Coll Physicians Surg Pak* 2015; **25**(1): 50–5.
- 130 Cheng Q, Chen F, Yip PS. Media effects on suicide methods: a case study on Hong Kong 1998–2005. *PLoS One* 2017; **12**(4): e0175580.
- 131 Ruder TD, Hatch GM, Ampanozi G, Thali MJ, Fischer N. Suicide announcement on Facebook. *Crisis* 2011; **32**(5): 280–2.
- 132 Sun FK, Lu CY, Tseng YS, Chiang CY. Factors predicting recovery from suicide in attempted suicide patients. *J Clin Nurs* 2017; **26**(23–4): 4404–12.
- 133 Gearing RE, Alonzo D. Religion and suicide: new findings. *J Relig Health* 2018; **57**(6): 2478–99.
- 134 Rezaeian M. Islam and suicide: a short personal communication. *Omega (Westport)* 2008; **58**(1): 77–85.
- 135 Pritchard C, Amanullah S. An analysis of suicide and undetermined deaths in 17 predominantly Islamic countries contrasted with the UK. *Psychol Med* 2007; **37**(3): 421–30.
- 136 Jordans MJ, Kaufman A, Brenman NF, Adhikari RP, Luitel NP, Tol WA, et al. Suicide in South Asia: a scoping review. *BMC Psychiatry* 2014; **14**: 358.
- 137 Khezeli M, Hazavehei SM, Ariapooran S, Ahmadi A, Soltanian A, Rezapur-Shahkolai F. Individual and social factors related to attempted suicide among women: a qualitative study from Iran. *Health Care Women Int* 2019; **40**(3): 295–313.
- 138 Cole AB, Littlefield AK, Gauthier JM, Bagge CL. Impulsivity facets and perceived likelihood of future suicide attempt among patients who recently attempted suicide. *J Affect Disord* 2019; **257**: 195–9.
- 139 Celik M, Kalendaroglu A, Almis H, Turgut M. Copycat suicides without an intention to die after watching TV programs: two cases at five years of age. *Noro Psikiyatri Arsivi* 2016; **53**(1): 83–4.
- 140 Ladwig KH, Kunrath S, Lukaschek K, Baumert J. The railway suicide death of a famous German football player: impact on the subsequent frequency of railway suicide acts in Germany. *J Affect Disord* 2012; **136** (1–2): 194–8.
- 141 Koburger N, Mergl R, Rummel-Kluge C, Ibelshausen A, Meise U, Postuvan V, et al. Celebrity suicide on the railway network: can one case trigger international effects? *J Affect Disord* 2015; **185**: 38–46.
- 142 Wang W, Chen X, Li S, Yan H, Yu B, Xu Y. Cusp catastrophe modeling of suicide behaviors among people living with HIV in China. *Nonlinear Dynamics Psychol Life Sci* 2019; **23**(4): 491–515.
- 143 Mirza I, Jenkins R. Risk factors, prevalence, and treatment of anxiety and depressive disorders in Pakistan: systematic review. *BMJ* 2004; **328** (7443): 794.
- 144 Hansen H, Braslow J, Rohrbaugh RM. From cultural to structural competency-training psychiatry residents to act on social determinants of health and institutional racism. *JAMA Psychiatry* 2018; **75**(2): 117–8.

- 145 Metz J, Hansen H. Structural competency: theorizing a new medical engagement with stigma and inequality. *Soc Sci Med* 2014; **103**: 126–33.
- 146 Kirmani N. *The Spectre of Inequality*. The News International, 2020 (<https://www.thenews.com.pk/tns/detail/610312-the-spectre-of-inequality>).
- 147 Shams K, Kadow A. Income inequality, remittances and economic wellbeing in rural Pakistan: linkages and empirical evidence. *Asia-Pac J Reg Sci* 2020; **4**(2): 499–519.
- 148 Li M, Katikireddi SV. Urban-rural inequalities in suicide among elderly people in China: a systematic review and meta-analysis. *Int J Equity Health* 2019; **18**(1): 2.
- 149 Varshney M, Gupta R, Balhara YPS. Yes, India has done it: decriminalization of suicide in India. *Asian J Psychiatry* 2015; **17**: 103.
- 150 Majeed MH, Amir Sherazi SA, Afzal MY. Decriminalization of suicide in Pakistan - treatment not punishment. *Asian J Psychiatr* 2018; **35**: 6–7.
- 151 Our Correspondent. Dunya News TV. *Treatment, Not Punishment: Senate Passes Bill to Decriminalise Attempted Suicide*. 2018 (<https://dunyanews.tv/en/Pakistan/428389-Treatment-punishment-Senate-bill-decriminalise-attempted-suicide>).
- 152 Munawar K, Abdul Khayom JH, Bokhary IZ, Park MS, Choudhry FR. A systematic review of mental health literacy in Pakistan. *Asia Pac Psychiatry* 2020; **12**(4): e12408.
- 153 Cha ES, Chang SS, Gunnell D, Eddleston M, Khang YH, Lee WJ. Impact of paraquat regulation on suicide in South Korea. *Int J Epidemiol* 2016; **45**(2): 470–9.
- 154 Gunnell D, Fernando R, Hewagama M, Priyangika WD, Konradsen F, Eddleston M. The impact of pesticide regulations on suicide in Sri Lanka. *Int J Epidemiol* 2007; **36**(6): 1235–42.
- 155 Eddleston M, Gunnell D. Preventing suicide through pesticide regulation. *Lancet Psychiatry* 2020; **7**(1): 9–11.
- 156 Chowdhury FR, Dewan G, Verma VR, Knipe DW, Isha IT, Faiz MA, et al. Bans of WHO class I pesticides in Bangladesh-suicide prevention without hampering agricultural output. *Int J Epidemiol* 2018; **47**(1): 175–84.
- 157 Gunnell D, Knipe D, Chang SS, Pearson M, Konradsen F, Lee WJ, et al. Prevention of suicide with regulations aimed at restricting access to highly hazardous pesticides: a systematic review of the international evidence. *Lancet Glob Health* 2017; **5**(10): e1026–37.
- 158 Knipe DW, Gunnell D, Eddleston M. Preventing deaths from pesticide self-poisoning-learning from Sri Lanka's success. *Lancet Glob Health* 2017; **5**(7): e651–2.
- 159 Zia S. Pesticide Policy. *eSocialSciences* 2016; Working papers: **id9077** (<https://ideas.repec.org/p/ess/wpaper/id9077.html>).
- 160 Lee EC. Clinical manifestations of sarin nerve gas exposure. *JAMA* 2003; **290**(5): 659–62.
- 161 Arshad S, Abid F, Aziz HA, Anjum R, Nadeem I. Specific poisons in Pakistan: a mini review. *Res Pharm Health Sci* 2016; **2**(3): 179–86.
- 162 Najeeb K. Pattern of suicide. Causes and methods employed. *J Coll Physicians Surg Pak* 2001; **11**: 759–61.
- 163 Yimaer A, Chen G, Zhang M, Zhou L, Fang X, Jiang W. Childhood pesticide poisoning in Zhejiang, China: a retrospective analysis from 2006 to 2015. *BMC Public Health* 2017; **17**: 602.
- 164 Ahmad R, Ahad K, Rashid I, Ashiq M. Acute poisoning due to commercial pesticides in Multan. *Pak J Med Sci* 2002; **18**(3): 227–31.
- 165 Ali Z AM, Muhammad R, Rahim A, Rahman SKU, Ullah N, et al. Outcome and predictors of in-hospital mortality in patients presenting with acute poisoning to a teaching hospital. *J Postgrad Med Inst*. 2018; **32**(2): 155–61.
- 166 Robb EL, Baker MB. *Organophosphate Toxicity*. StatPearls Publishing, 2020 (<https://www.ncbi.nlm.nih.gov/books/NBK470430/>).
- 167 Eddleston M, Singh S, Buckley N. Organophosphorus poisoning (acute). *Clin Evid* 2004; **12**: 1941–53.
- 168 Eddleston M, Singh S, Buckley N. Organophosphorus poisoning (acute). *Clin Evid* 2005; **13**: 1744–55.
- 169 Asghar SP AN, Farooq M, Sidra, Asghar S, Ijaz A. Presentation and management of organophosphate poisoning in an intensive care unit. *Pak Armed Forces Med J*. 2014; **64**(1): 134–8.
- 170 Eddleston M, Buckley NA, Eyer P, Dawson AH. Management of acute organophosphorus pesticide poisoning. *Lancet* 2008; **371**(9612): 597–607.
- 171 Clark RF. Insecticides: organic phosphorus compounds and carbamates. In *Goldfrank's Toxicologic Emergencies* (eds Goldfrank LR, Flomenbaum NE, Lewin NA, Howland MA, Hoffman RS, Nelson LS): 1346–60. McGraw-Hill, 2002.
- 172 Eyer P. The role of oximes in the management of organophosphorus pesticide poisoning. *Toxicol Rev* 2003; **22**(3): 165–90.
- 173 Calvert GM, Plate DK, Das R, Rosales R, Shafey O, Thomsen C, et al. Acute occupational pesticide-related illness in the US, 1998–1999: surveillance findings from the SENSOR-pesticides program. *Am J Ind Med* 2004; **45**(1): 14–23.
- 174 Ahmed SM, Das B, Nadeem A, Samal RK. Survival pattern in patients with acute organophosphate poisoning on mechanical ventilation: a retrospective intensive care unit-based study in a tertiary care teaching hospital. *Indian J Anaesth* 2014; **58**(1): 11–7.
- 175 Karimani A, Mohammadpour AH, Zirak MR, Rezaee R, Megarbane B, Tsatsakis A, et al. Antidotes for aluminum phosphide poisoning - an update. *Toxicol Rep* 2018; **5**: 1053–9.
- 176 Navabi SM, Navabi J, Aghaei A, Shaahmadi Z, Heydari R. Mortality from aluminium phosphide poisoning in Kermanshah Province, Iran: characteristics and predictive factors. *Epidemiol Health* 2018; **40**: e2018022.
- 177 Hosseinian A, Pakravan N, Rafiei A, Feyzbakhsh SM. Aluminum phosphide poisoning known as rice tablet: a common toxicity in North Iran. *Indian J Med Sci* 2011; **65**(4): 143–50.
- 178 Hassan A MM, Shafique M, Adil. Wheat pill poisoning: clinical manifestation and its outcome. *Journal of Rawalpindi Medical College*. 2014; **18**(1): 49–51.
- 179 Khan ZU. *Rise in Wheat Pill Poisoning: Study*. Dawn, 2008 (<https://www.dawn.com/news/884170/rise-in-wheat-pill-poisoning-study>).
- 180 Chugh SN, Chugh K, Ram S, Malhotra KC. Electrocardiographic abnormalities in aluminium phosphide poisoning with special reference to its incidence, pathogenesis, mortality and histopathology. *J Indian Med Assoc* 1991; **89**(2): 32–5.
- 181 Hena Z, McCabe ME, Perez MM, Sharma M, Sutton NJ, Peek GJ, et al. Aluminium phosphide poisoning: successful recovery of multiorgan failure in a pediatric patient. *Int J Pediatr Adolesc Med* 2018; **5**(4): 155–8.
- 182 Chugh SN, Ram S, Chugh K, Malhotra KC. Spot diagnosis of aluminium phosphide ingestion: an application of a simple test. *J Assoc Physicians India* 1989; **37**(3): 219–20.
- 183 Shadnia S, Rahimi M, Pajoumand A, Rasouli MH, Abdollahi M. Successful treatment of acute aluminium phosphide poisoning: possible benefit of coconut oil. *Hum Exp Toxicol* 2005; **24**(4): 215–8.
- 184 Nasa P, Gupta A, Mangal K, Nagrani SK, Raina S, Yadav R. Use of continuous renal replacement therapy in acute aluminium phosphide poisoning: a novel therapy. *Ren Fail* 2013; **35**(8): 1170–2.
- 185 Mahajan VV, Pargal L. Aluminium phosphide poisoning: an agent of sure death. *Indian J Forensic Med Toxicol* 2012; **6**: 231–5.
- 186 Bogle RG, Theron P, Brooks P, Dargan PI, Redhead J. Aluminium phosphide poisoning. *Emergency Med J* 2006; **23**(1): e3.
- 187 Prabhakaran AC. Paraphenylene diamine poisoning. *Indian J Pharmacol* 2012; **44**(3): 423–4.
- 188 Hill RM, Hatkevich CE, Kazimi I, Sharp C. The Columbia-Suicide Severity Rating Scale: associations between interrupted, aborted, and actual suicide attempts among adolescent inpatients. *Psychiatry Res* 2017; **255**: 338–40.
- 189 Akbar MA, Khaliq SA, Malik NA, Shahzad A, Tarin SM, Chaudhry GM. Kala pathar (paraphenylene diamine) intoxication: a study at Nishtar Hospital Multan. *Nishtar Med J* 2010; **2**: 111–5.
- 190 Suliman SM, Fadlalla M, Nasr Mel M, Beliel MH, Fesseha S, Babiker M, et al. Poisoning with hair-dye containing paraphenylene diamine: ten years experience. *Saudi J Kidney Dis Transpl* 1995; **6**(3): 286–9.
- 191 Kondle R, Pathapati RM, Saginela SK, Malliboina S, Makineed VP. Clinical profile and outcomes of hair dye poisoning in a teaching hospital in Nellore. *Int Scho Res Not* 2012; **5**: 624253.
- 192 Umair SF, Amin I, Urrehman A. Hair dye poisoning: "an early intervention". *Pak J Med Sci* 2018; **34**(1): 230–2.
- 193 Daga MK, Sinha N, Mahapatra HS, Kumar R, Lalmaisawma R, Nayak HK, et al. Paraphenylene diamine poisoning. *J Indian Med Assoc* 2011; **109**(1): 49.
- 194 Chaudhary SC, Sawlani KK, Singh K. Paraphenylenediamine poisoning. *Niger J Clin Pract* 2013; **16**(2): 258–9.
- 195 Punjani NS. Paraphenylene diamine (hair dye) poisoning leading to critical illness neuropathy. *J Neurol Disord* 2014; **2**: 1–2.
- 196 Bowen DA. A case of phenylenediamine poisoning. *Med Sci Law* 1963; **3**: 216–9.
- 197 Jan A, Khan MJ, Khan MTH, Khan MTM, Fatima S. Poisons implicated in homicidal, suicidal and accidental cases in North-West Pakistan. *J Ayub Med Coll Abbottabad* 2016; **28**(2): 308–11.
- 198 Gude D, Bansal DP, Ambegaonkar R, Prajapati J. Paraphenylenediamine: blackening more than just hair. *J Res Med Sci*. 2012; **17**(6): 584–6.
- 199 Parkar Khan N, Perez-Nunez R, Shamim N, Khan U, Naseer N, Feroze A, et al. Intentional and unintentional poisoning in Pakistan: a pilot study using the Emergency Departments surveillance project. *BMC Emerg Med* 2015; **15**: S2.

- 200 Parkar SR, Dawani V, Weiss MG. Clinical diagnostic and sociocultural dimensions of deliberate self-harm in Mumbai, India. *Suicide Life Threat Behav* 2006; **36**(2): 223–38.
- 201 Weinbroum AA, Flaishon R, Sorkine P, Szold O, Rudick V. A risk-benefit assessment of flumazenil in the management of benzodiazepine overdose. *Drug Safety* 1997; **17**(3): 181–96.
- 202 Shalansky SJ, Naumann TL, Englander FA. Effect of flumazenil on benzodiazepine-induced respiratory depression. *Clin Pharm* 1993; **12**(7): 483–7.

