

## Potential Effects of Pomegranate on Lipid Peroxidation and Pro-inflammatory Changes in Daunorubicin-induced Cardiotoxicity in Rats

Hayder M. Al-Kuraishy, Ali I. Al-Gareeb

Department of Pharmacology, Toxicology and Medicine, College of Medicine, Al-Mustansiriya University, Baghdad, Iraq

### Correspondence to:

Dr. Hayder M. Al-Kuraishy, Department of Pharmacology, Toxicology and Medicine, College of Medicine, Al-Mustansiriya University, P.O. Box 14132, Baghdad, Iraq. E-mail: hayderm36@yahoo.com

**How to cite this article:** Al-Kuraishy HM, Al-Gareeb AI. Potential effects of pomegranate on lipid peroxidation and pro-inflammatory changes in daunorubicin-induced cardiotoxicity in rats. *Int J Prev Med* 2016;7:85.

### ABSTRACT

**Background:** Daunorubicin-induced acute cardiotoxicity caused by oxidative stress and free radical formation. Pomegranate possessed a significant *in vitro* free radical scavenging activity. Therefore, the aim of this study was estimations of the role of pomegranate effects in daunorubicin-induced cardiotoxicity.

**Methods:** A total of 21 Sprague male rats were allocated into three groups, seven animals in each group. Group A: Control group received distilled water. Group B: Treated group with daunorubicin 20 mg/kg via intraperitoneal injection daily for the 12<sup>th</sup> day for total cumulative dose of 240 mg/kg. Group C: Pretreatment group with pomegranate 25 mg/kg for 6 days orally, then daunorubicin 20 mg/kg administrated concomitantly for the next 6 days with a cumulative dose of 120 mg/kg. Cardiac troponin I ([cTn I] pg/ml), malondialdehyde (MDA) (ng/ml), interleukin 17 (IL-17 pg/ml), and cardiac lactate dehydrogenase (LDH) (pm/ml), all these biomarkers were used to measure the severity of cardiotoxicity.

**Results:** Daunorubicin at a dose of 20 mg/kg lead to pronounced cardiac damage that reflected on through elevations of serum cTn and serum LDH levels significantly  $P < 0.01$ , it induced lipid peroxidation during cardiotoxicity that reflected through an elevation in the serum MDA significantly  $P < 0.01$ , moreover, daunorubicin induces pro-inflammatory changes in cardiotoxicity; it raises the IL-17 serum level significantly  $P < 0.01$  as compared with control. Pomegranate pretreatment demonstrated a significant cardioprotection from daunorubicin-induced cardiotoxicity; it attenuated the cardiac damage through reduction of cTn, LDH, MDA, and serum IL-17 level significantly  $P < 0.01$  as compared with daunorubicin-treated group.

**Conclusions:** Pomegranate demonstrated significant cardioprotection in daunorubicin-induced cardiotoxicity through reduction of oxidative stress, lipid peroxidation, pro-inflammatory, and cardiac injury biomarkers.

**Keywords:** Cardiotoxicity, daunorubicin, pomegranate

### Access this article online

#### Quick Response Code:



Website: [www.ijpvmjournal.net/www.ijpm.ir](http://www.ijpvmjournal.net/www.ijpm.ir)

DOI:  
10.4103/2008-7802.184314

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: [reprints@medknow.com](mailto:reprints@medknow.com)

## INTRODUCTION

Daunorubicin is an anthracycline antibiotic that mainly used as an antineoplastic agent against various tumor types; its value was limited due to their perilous side effects like cardiotoxicity.<sup>[1]</sup>

Daunorubicin-induced acute cardiotoxicity is caused by oxidative stress and free radical formation that induced by a high daunorubicin dose more than 15 mg/kg, which provoking oxidative myocardial injury. Furthermore, daunorubicin is converted by intracellular endogenous enzymes into semiquinone that generating a free radical.<sup>[2]</sup> The heart is more vulnerable to oxidative damage due to less antioxidant enzyme activity (superoxide dismutase, glutathione, and catalase enzymes) consequently, accumulated free radicals leading to lipid peroxidation, mitochondrial membrane damage, injury of nucleic acid, and destruction of myocardium endoplasmic reticulum.<sup>[3]</sup> Moreover, the daunorubicin-induced release of cytokine-like tumor necrosis factor and interleukin 17 (IL-17) that lead to dilated cardiomyopathy and ventricular dysfunction.<sup>[4]</sup>

Daunorubicin-semiquinone radical binds iron and form complex called anthracycline-iron free radical complex, which reduces oxygen to superoxide that converted into oxygen and hydrogen peroxide, all of these free radicals lead to oxidative stress and lipid peroxidation in daunorubicin-induced cardiotoxicity.<sup>[5]</sup>

Thus, antioxidant agents and medicinal plants demonstrated a cardioprotective effect on daunorubicin-induced acute cardiotoxicity, one of these medicinal plants named pomegranate (*punica grantum*) which is belongs to the family of Punicaceae, there are two species in this family *punica granatum* and *punica protopunica*, the second one found only in socotra (Indian ocean Island). The most significant and beneficial effects of pomegranate are related to their active constituents such as punicic acid, anthocyanins, flavones, and flavonols.<sup>[6]</sup>

Pomegranate possessed a significant *in vitro* free radical scavenging activity and is being studied for its potential biological property in human being; numerous studies demonstrated a reliable anti-oxidant effect of pomegranate in attenuation of risk factors of ischemic heart disease through inhibitions of low-density lipoprotein oxidation, foam cell formation, and macrophage oxidation.<sup>[7,8]</sup>

Furthermore, pomegranate is regarding as angiotensin converting enzymes inhibitor that inhibits (ACEIs), thus, it is regarded as a natural ACEI in reduction of blood pressure and affording a cardioprotective effect.<sup>[9]</sup>

Vascular and cardiac protection effects of pomegranate may be produced through enhancement of natural anti-oxidant system or act as direct anti-oxidant.<sup>[10]</sup>

Therefore, the aim of this study was estimations the role of pomegranate effects on the oxidative stress, lipid peroxidation, and pro-inflammatory changes in daunorubicin-induced cardiotoxicity.

## METHODS

This experimental study was done in Department of Clinical Pharmacology, College of Medicine, Al-Mustansiriya University, Baghdad, Iraq, from May to June 2015, this study was approved by Scientific Juries and Moral Committee of Animal Experimentation. Humane care for animals was done according to the guide to the care and use of laboratory animal.

### Study design

A total of 21 Sprague male rats were allocated into three groups, seven animals in each group. The animal body weight was 250–300 g, they were attained from Iraqi National Center, the animals were housed and fed an ordinary chow and distilled water was provided *ad libitum*, then the animals were kept at room temperature and 12-h light-dark cycle. All drugs were purchased from private pharmacy pomegranate extract capsule 250 mg (apeiron, Germany) and daunorubicin 200 mg (rubex vial, santus pharma), then dilution was done with specific solutes to reach the final concentration of 25 mg/kg and 20 mg/kg of pomegranate and daunorubicin, respectively.

- Group A: Control group received distilled water 6 ml/kg orally for 12 days
- Group B: Treated group with daunorubicin 20 mg/kg via intraperitoneal injection daily for 12 days for total cumulative dose of 240 mg/kg
- Group C: Pretreatment group with pomegranate 25 mg/kg for 6 days orally, then daunorubicin 20 mg/kg administrated concomitantly for the next 6 days with cumulative doses of 120 mg/kg.

After 24 hrs of the last dose of drugs, animal decapitated under light anesthesia, and 3 ml of blood sample was obtained. Serum was taken by centrifugation at 3500 rpm for 15 min.

### Assessment of cardiac biomarkers

Cardiac troponin I ([cTn I] pg/ml), malondialdehyde (MDA) (ng/ml), IL-17 pg/ml and cardiac lactate dehydrogenase (LDH) (pmol/ml), all of these biomarkers were used to measure the severity of cardiotoxicity, they were estimated by ELISA kit method according to the kit instructions (ELISA kit Catalog NO: e-el-ro566).

### Statistical methods

The results were presented as mean  $\pm$  standard error ANOVA Bonferroni test was used for evaluating the significance of differences regarding  $P < 0.01$ .

## RESULTS

There were 14 rats received daunorubicin with a cumulative dose of 240 mg/kg in daunorubicin-treated group and 120 mg/kg in pretreatment groups as part of the protocol, none of these treated animals with daunorubicin were died through the experimental period.

Daunorubicin at a dose of 20 mg/kg leads to pronounced cardiac damage that reflected on elevation of serum cTn I and serum LDH from  $25.09 \pm 0.57$  pg/ml and  $88.74 \pm 0.30$  pm/ml in control rats to  $211.20 \pm 0.615$  pg/ml and  $301.04 \pm 0.989$  pm/ml, respectively with significance differences value of  $P = 0.00001$  for cTn and  $P = 0.00009$  for LDH respectively, also; it induced lipid peroxidation during cardiotoxicity that reflected on an elevation in the serum MDA level from  $25.25 \pm 0.49$  ng/ml in control rats to  $282.75 \pm 2.75$  ng/ml in daunorubicin-treated rats significantly  $P = 0.00003$ . Moreover, daunorubicin induces pro-inflammatory changes in cardiotoxicity; it raises the IL-17 serum levels significantly from  $56.245 \pm 0.84$  pg/ml in control rats to the  $231.04 \pm 0.707$  pg/ml in daunorubicin-treated rats  $P = 0.000026$ , [Table 1].

Pomegranate pretreatment demonstrated a significant cardioprotection from daunorubicin-induced cardiotoxicity; it attenuated the cardiac damage through reduction of serum cTn I and serum LDH with  $P = 0.00012$  and  $P = 0.00037$ , respectively.

In addition, pomegranate decreases lipid peroxidation significantly through reduction of serum level of MDA level from  $282.75 \pm 2.75$  ng/ml in daunorubicin-treated rats to the  $133.175 \pm 0.671$  ng/ml in pomegranate pretreated rats significantly  $P = 0.000103$ .

Regarding the pro-inflammatory amelioration in pomegranate, it reduced serum IL-17 levels from  $231.04 \pm 0.707$  pg/ml in daunorubicin treated rats to the  $165.85 \pm 0.120$  pg/ml in pomegranate pretreated rats significantly  $P = 0.00021$ , [Table 2].

Furthermore, the intergroup differences demonstrated significant differences  $P < 0.01$  among daunorubicin and daunorubicin with pomegranate as compared with control. Therefore, cardiac injury, lipid peroxidation, and pro-inflammatory biomarkers were higher in daunorubicin-treated rats and lower in the pomegranate pretreated rats, [Table 3].

## DISCUSSION

Daunorubicin-induced cardiotoxicity has been widely reviewed, and it has been reported that daunorubicin and other anthracyclines antibiotic cause heart failure, cardiac electrical abnormalities, and cardiomyopathy.<sup>[11]</sup>

The probable mechanism for proposed cardiotoxic effects of daunorubicin includes myocardial oxidative stress, mitochondrial damage, lipid peroxidation, and dysregulation of adrenergic signaling.<sup>[12]</sup>

In this study, daunorubicin at a dose of 20 mg/kg significantly leads to a cardiotoxic injury that reflected off an elevation in serum level of cTn I and LDH, cTn I has been accounted as a highly sensitive marker of myocardial injury and can be of value in the detection of anthracycline-induced cardiotoxicity.<sup>[13]</sup> Therefore, elevation of both LDH and cTn I during daunorubicin chemotherapy pointed out the occurrence of cardiotoxicity as found in this study.

Moreover, daunorubicin induces oxidative stress, free radical formation and finally induction of lipid peroxidation since; cardiac tissues are rich in mitochondria and its metabolism mainly depend on oxidative pathway, thus, intracellular is buildups of daunorubicin provoked mitochondrial free radicals productions which lead to lipid peroxidation that reflected off an increasing in the serum level of MDA.<sup>[14]</sup> Xiao, 2015 study showed that MDA is a stable end product of lipid peroxidation and regarded as a reliable marker for estimation of free radical induced tissue damage and lipid peroxidation<sup>[15]</sup> as in daunorubicin-induced cardiotoxicity of present study.

Moreover, the current study demonstrated that daunorubicin induces inflammatory consequences of cardiotoxicity through an increment in the serum level of IL-17.

IL-17 is a pro-inflammatory cytokine linking adaptive and innate immunity, secreted from a subset of CD4 called Th17, also, it is secreted from natural killer cell,<sup>[16]</sup> it is responsible for the development of dilated cardiomyopathy in viral myocarditis through enhancement of lymphocyte and neutrophil infiltrations, induction of IL-6 and interferon secretions and cyclooxygenase-2 (COX-2) inhibition.<sup>[17]</sup>

Amador, *et al.* study pointed out that IL-17 blocking antibodies decreased cardiac inflammatory cytokine

**Table 1: Effects of daunorubicin on cardiac, pro-inflammatory and lipid peroxidation biomarkers during experimental cardiotoxicity compared with control**

Biomarkers	Control (n=7)	Daunorubicin (n=7)	95% CI Upper-Lower	t	P value
MDA ng/ml	25.25±0.49	282.75±2.75*	-230.75-284.24	-34.84	0.00003*
cTnI pg/ml	25.09±0.57	211.20±0.615*	-179.32-192.89	-83.88	0.00001*
LDH pm/ml	88.74±0.30	301.04±0.989*	-202.77-221.8	-77.6409	0.00009*
IL-17 pg/ml	56.245±0.84	231.04±0.707*	-165.87-183.71	-60.173	0.000026*

Data are expressed as mean±SE \*P<0.01 as compared with control, MDA=Maondialdehyde, cTnI=Cardiac troponin I, LDH=lactate dehydrogenase, IL-17=Interleukin 17

**Table 2: Effects of pre-treatments with pomegranate on cardiac, pro-inflammatory and lipid peroxidation biomarkers in daunorubicin-induced cardiotoxicity**

Biomarkers	Pomegranate (n=7)	Daunorubicin (n=7)	95% CI Upper-Lower	t	P value
MDA ng/ml	133.175±0.671*	282.75±2.75	-122.98-176.17	-19.97	0.000103*
cTnl pg/ml	82.815±0.247*	211.20±0.615	-123.02-133.74	-73.21	0.00012*
LDH pm/ml	177.205±0.657*	301.04±0.989	-113.97-133.70	-39.42	0.00037*
IL-17 pg/ml	165.85±0.120*	231.04±0.707	-58.31-72.06	-34.35	0.00021*

Data are expressed as mean±SE \*P<0.01 as compared with daunorubicin group. MDA=Maondialdehyde, cTnl=Cardiac troponin I, LDH=Lactate dehydrogenase, IL-17=Interleukin 17

**Table 3: Effects of daunorubicin and daunorubicin with pomegranate on cardiac, pro-inflammatory and lipid peroxidation biomarkers during experimental cardiotoxicity compared with control**

Biomarkers	Control (n=7)	Daunorubicin (n=7)	Daunorubicin + Pomegranate (n=7)
MDA ng/ml	25.25±0.49	282.75±2.75*	133.175±0.671**
cTnl pg/ml	25.09±0.57	211.20±0.615*	82.815±0.247**
LDH pm/ml	88.74±0.30	301.04±0.989*	177.205±0.657**
IL-17 pg/ml	56.245±0.84	231.04±0.707*	165.85±0.120**

Data are expressed as mean±SE \*P<0.01 as compared with control, P value calculating via ANOVA Bonferrony test, \*\*P<0.01 as compared with daunorubicin group. MDA=Maondialdehyde, cTnl=Cardiac troponin I, LDH=Lactate dehydrogenase, IL-17=Interleukin 17

injury via down-regulation of tumor growth factor and up-regulation of cardiac COX-2/prostaglandin E2 pathway which is responsible for cardioprotection and coronary vasodilatations.<sup>[18]</sup>

These studies correspond with the present findings, which showed that daunorubicin-induced inflammatory changes in cardiotoxicity that reflected on through an elevation in serum level of IL-17 was significantly during daunorubicin-induced cardiotoxicity.

Thus, oxidative stress, lipid peroxidation, inflammatory induction, and direct myocardial damage all were interconnected in provoking and augmenting of cardiotoxicity induced by daunorubicin.

Regarding, the cardioprotection from daunorubicin-induced cardiotoxicity Conway, *et al.* study revealed that a numerous of synthetic drugs are currently used to overcome the cardiotoxicity such as  $\beta$ -blockers, ACEI inhibitors, statins, and cyclosporine,<sup>[19]</sup> these agents leads to various side effects such as dry coughs with ACEI inhibitors, fatigue with  $\beta$ -blockers, and muscle damage with statins, which are not appearing with the use of herbal medicine, thus the utilize of these supplements had to turn out to be well-liked in latest years.<sup>[20]</sup>

Therefore, pretreatment with pomegranate significantly attenuates the cardiotoxic effects of daunorubicin through significant reduction of cardiac injury biomarkers, lipid peroxidation, and pro-inflammatory cytokine; this effect may relate to the polyphenolic compounds which have significant biological activity including antioxidant effect.<sup>[21]</sup>

The antioxidant activity of pomegranate is related to ellagic acid, which has *in vitro* a prominent free radical scavenging activity and potentiates different antioxidant pharmacological and biochemical pathways thus; it may be regarded as a cardioprotective agent but not prevent other toxicity like cisplatin-induced nephrotoxicity in rats.<sup>[22]</sup>

Moreover, pomegranate contained other antioxidant constituents such as quercetin, gallic acid, cyanide-3-glucose, and myricetin that are potent agents in attenuations the reduction of heart weight/body weight ratio in doxorubicin cardiotoxicity via a protective effect on cardiac myofibrils from free radical oxidative stress.<sup>[23,24]</sup>

In addition, pomegranate significantly reduced the level of caspase-3 protease which is a marker for apoptosis and DNA fragmentation, thus; pomegranate like other antioxidants prevents daunorubicin-induced apoptosis in experimental cardiotoxicity.<sup>[25]</sup> However, unfortunately; this marker was not measured in present study.

Furthermore, pomegranate regarded as natural ACEI and through this effect it demonstrated a potential cardioprotective action since; ACEIs have a potent antioxidant activity, free radical scavenging activity, mitochondrial protection, and inhibition of free radical generations, all of these mechanisms may explain the role of ACEI in the prevention of anthracycline-induced cardiotoxicity.<sup>[26]</sup> Mohan, *et al.* research demonstrated that pomegranate in animal model study, lead to potential cardioprotection from doxorubicin-induced cardiotoxicity through reduction in serum levels of cardiac creatine kinase-MB and LDH with significant elevation in the antioxidant glutathione,<sup>[27]</sup> which *per se* explain the attenuation in daunorubicin-induced cardiotoxicity in the present study.

Furthermore, pomegranate boosts HDL effect by stimulation of paraoxonase enzyme (PON-1), an enzyme attached to the surface of HDL that show an age-dependent reduction in its activity, therefore, pomegranate revealed a significant cardioprotection regardless of age unlike statins which boost HDL but not PON-1 activity.<sup>[28]</sup>

In addition, puniceic acid of pomegranate led to inhibition of vascular adhesion molecule in cardiac and vascular endothelium, hence pomegranate regarded as anti-inflammatory agent<sup>[29]</sup> which may explain the reduction effect of pomegranate on IL-17 in the

present study, also, a numerous studies demonstrated the anti-inflammatory property of pomegranate through inhibition of nuclear factor E2-related factor during carcinogenesis which plays an important role in initiation of inflammatory pathways to cardiotoxicity.<sup>[30]</sup> Finally, co-administration of pomegranate with daunorubicin leads to significant attenuation in the induced cardiotoxicity through reduction of lipid peroxidation, antioxidant activity, and anti-inflammatory effect.

## CONCLUSIONS

Pomegranate demonstrated significant cardioprotection in daunorubicin-induced cardiotoxicity through reduction of oxidative stress, lipid peroxidation, pro-inflammatory, and cardiac injury biomarkers.

## Acknowledgements

We would like to thank professor Dr. Sadiq Alhamash Dean, College of Medicine, Al-Mustansiriyah University for his great supports.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

**Received:** 31 Jul 15 **Accepted:** 03 May 16

**Published:** 20 Jun 16

## REFERENCES

- Su H, Gorodny N, Gomez LF, Gangadharmath U, Mu F, Chen G, et al. Noninvasive molecular imaging of apoptosis in a mouse model of anthracycline-induced cardiotoxicity. *Circ Cardiovasc Imaging* 2015;8:e001952.
- Saleem MT, Chetty MC, Kavimani S. Antioxidants and tumor necrosis factor alpha-inhibiting activity of sesame oil against doxorubicin-induced cardiotoxicity. *Ther Adv Cardiovasc Dis* 2014;8:4-11.
- Savitha S, Panneerselvam C. Mitigation of age-dependent oxidative damage to DNA in rat heart by carnitine and lipoic acid. *Mech Ageing Dev* 2007;128:206-12.
- De Giorgi U, Rosti G, Zaniboni A, Ballardini M, Minzi MR, Baioni M, et al. High-dose epirubicin, preceded by dexrazoxane, given in combination with paclitaxel plus filgrastim provides an effective mobilizing regimen to support three courses of high-dose dense chemotherapy in patients with high-risk stage II-III breast cancer. *Bone Marrow Transplant* 2003;32:251-5.
- Saprin AN, Kalina EV, Serezhenkov VA, Kotova IaN, Solomka VS, Shcherbak NP. Changes in the free-radical state and level of free iron during the formation of drug resistance in tumor cells. *Biofizika* 2006;51:485-90.
- Soucek P, Kondrova E, Hermanek J, Stopka P, Boumendjel A, Ueng YF, et al. New model system for testing effects of flavonoids on doxorubicin-related formation of hydroxyl radicals. *Anticancer Drugs* 2011;22:176-84.
- Minaiyan M, Zolfaghari B, Taheri D, Gomarian M. Preventive effect of three pomegranate (*Punica granatum* L.) seeds fractions on cerulein-induced acute pancreatitis in mice. *Int J Prev Med* 2014;5:394-404.
- Qabaha KI. Antimicrobial and free radical scavenging activities of five Palestinian medicinal plants. *Afr J Tradit Complement Altern Med* 2013;10:101-8.
- Aviram M, Dornfeld L. Pomegranate juice consumption inhibits serum angiotensin converting enzyme activity and reduces systolic blood pressure. *Atherosclerosis* 2001;158:195-8.
- Dassprakash MV, Arun R, Abraham SK, Premkumar K. *In vitro* and *in vivo* evaluation of antioxidant and antigenotoxic potential of *Punica granatum* leaf extract. *Pharm Biol* 2012;50:1523-30.
- Lian JB, Wu ZJ, Fang QJ, Yu J, He RL. Sodium ferulate protects against daunorubicin-induced cardiotoxicity in juvenile rats. *Zhongguo Ying Yong Sheng Li Xue Za Zhi* 2015;31:54-8.
- Hanna AD, Lam A, Tham S, Dulhunty AF, Beard NA. Adverse effects of doxorubicin and its metabolic product on cardiac RyR2 and SERCA2A. *Mol Pharmacol* 2014;86:438-49.
- Desai VG, C Kwekel J, Vijay V, Moland CL, Herman EH, Lee T, et al. Early biomarkers of doxorubicin-induced heart injury in a mouse model. *Toxicol Appl Pharmacol* 2014;281:221-9.
- Govender J, Loos B, Marais E, Engelbrecht AM. Mitochondrial catastrophe during doxorubicin-induced cardiotoxicity: A review of the protective role of melatonin. *J Pineal Res* 2014;57:367-80.
- Xiao NN. Effects of resveratrol supplementation on oxidative damage and lipid peroxidation induced by strenuous exercise in rats. *Biomol Ther (Seoul)* 2015;23:374-8.
- Mombelli S, Cochaud S, Merrouche Y, Garbar C, Antonicelli F, Laprevotte E, et al. IL-17A and its homologs IL-25/IL-17E recruit the c-RAF/S6 kinase pathway and the generation of pro-oncogenic LMW-E in breast cancer cells. *Sci Rep* 2015;5:11874.
- Konya C, Paz Z, Apostolidis SA, Tsokos GC. Update on the role of interleukin 17 in rheumatologic autoimmune diseases. *Cytokine* 2015;75:207-15.
- Amador CA, Barrientos V, Peña J, Herrada AA, González M, Valdés S, et al. Spironolactone decreases DOCA-salt-induced organ damage by blocking the activation of T helper 17 and the downregulation of regulatory T lymphocytes. *Hypertension* 2014;63:797-803.
- Conway A, McCarthy AL, Lawrence P, Clark RA. The prevention, detection and management of cancer treatment-induced cardiotoxicity: A meta-review. *BMC Cancer* 2015;15:366.
- Khan G, Haque SE, Anwer T, Ahsan MN, Safhi MM, Alam MF. Cardioprotective effect of green tea extract on doxorubicin-induced cardiotoxicity in rats. *Acta Pol Pharm* 2014;71:861-8.
- Hassanpour Fard M, Ghule AE, Bodhankar SL, Dikshit M. Cardioprotective effect of whole fruit extract of pomegranate on doxorubicin-induced toxicity in rat. *Pharm Biol* 2011;49:377-82.
- Jilanchi S, Nematbakhsh M, Mazaheri S, Talebi A, Zolfaghari B, Pezeshki Z, et al. Pomegranate flower extract does not prevent cisplatin-induced nephrotoxicity in female rats. *Int J Prev Med* 2014;5:1621-5.
- Tavasoli S, Zarnani AH, Vafa M, Moradi-Lakeh M, Pazoki-Toroudi H, Eghtesadi S. The effect of pomegranate extract on survival and peritoneal bacterial load in cecal ligation and perforation model of sepsis in rats. *Int J Prev Med* 2014;5:104-9.
- Bishayee A, Thoppil RJ, Darvesh AS, Ohanyan V, Meszaros JG, Bhatia D. Pomegranate phytoconstituents blunt the inflammatory cascade in a chemically induced rodent model of hepatocellular carcinogenesis. *J Nutr Biochem* 2013;24:178-87.
- Upaganlawar A, Balaraman R. Cardioprotective effects of co-administration of pomegranate extract and Vitamin E on electrocardiographic, biochemical and apoptotic changes in isoproterenol induced myocardial infarction in rats. *Pharmacologia* 2015;6:178-85.
- Donnini S, Terzuoli E, Ziche M, Morbidelli L. Sulfhydryl angiotensin-converting enzyme inhibitor promotes endothelial cell survival through nitric-oxide synthase, fibroblast growth factor-2, and telomerase cross-talk. *J Pharmacol Exp Ther* 2010;332:776-84.
- Mohan M, Patankar P, Ghadi P, Kasture S. Cardioprotective potential of *Punica granatum* extract in isoproterenol-induced myocardial infarction in Wistar rats. *J Pharmacol Pharmacother* 2010;1:32-7.
- Parsaeyan N, Mozaffari-Khosravi H, Mozayan MR. Effect of pomegranate juice on paraoxonase enzyme activity in patients with type 2 diabetes. *J Diabetes Metab Disord* 2012;11:111.
- Banerjee N, Kim H, Talcott S, Mertens-Talcott S. Pomegranate polyphenolics suppressed azoxymethane-induced colorectal aberrant crypt foci and inflammation: Possible role of miR-126/VCAM-1 and miR-126/PI3K/AKT/mTOR. *Carcinogenesis* 2013;34:2814-22.
- González-Trujano ME, Pellicer F, Mena P, Moreno DA, García-Viguera C. Antinociceptive and anti-inflammatory activities of a pomegranate (*Punica granatum* L.) extract rich in ellagitannins. *Int J Food Sci Nutr* 2015;66:395-9.