

DOI: 10.1093/femsre/fuac011 Advance access publication date: 23 February 2022 Review Article

Proteome expansion in the *Potyviridae* evolutionary radiation

Fabio Pasin^{(1,2,*}, José-Antonio Daròs⁽¹⁾, Ioannis E. Tzanetakis⁽²⁾

¹Instituto de Biología Molecular y Celular de Plantas (IBMCP), Consejo Superior de Investigaciones Científicas-Universitat Politècnica de València (CSIC-UPV), 46011 Valencia, Spain

²School of Science, University of Padova, 35121 Padova, Italy

³Department of Entomology and Plant Pathology, Division of Agriculture, University of Arkansas System, 72701 Fayetteville, AR, USA

*Corresponding author: Instituto de Biología Molecular y Celular de Plantas (IBMCP), Consejo Superior de Investigaciones Científicas-Universitat Politècnica de València (CSIC-UPV), UPV Building 8E, Ingeniero Fausto Elio, 46011 Valencia, Spain. E-mail: f.pasin@csic.es

One sentence summary: *Potyviridae* represents one of the most notable evolutionary radiations among RNA viruses; its species richness is accompanied by large genomic layout diversity and expansion of the encoded protein modules.

Editor: Guo Hui-Shan

Abstract

Potyviridae, the largest family of known RNA viruses (realm Riboviria), belongs to the picorna-like supergroup and has important agricultural and ecological impacts. Potyvirid genomes are translated into polyproteins, which are in turn hydrolyzed to release mature products. Recent sequencing efforts revealed an unprecedented number of potyvirids with a rich variability in gene content and genomic layouts. Here, we review the heterogeneity of non-core modules that expand the structural and functional diversity of the potyvirid proteomes. We provide a family-wide classification of P1 proteinases into the functional Types A and B, and discuss pretty interesting sweet potato potyviral ORF (PISPO), putative zinc fingers, and alkylation B (AlkB)—non-core modules found within P1 cistrons. The atypical inosine triphosphate pyrophosphatase (ITPase/HAM1), as well as the pseudo tobacco mosaic virus-like coat protein (TMV-like CP) are discussed alongside homologs of unrelated virus taxa. Family-wide abundance of the multitasking helper component proteinase (HC-pro) is revised. Functional connections between non-core modules are highlighted to support host niche adaptation and immune evasion as main drivers of the *Potyviridae* evolutionary radiation. Potential biotechnological and synthetic biology applications of potyvirid leader proteinases and non-core modules are finally explored.

Keywords: Potyviridae, virus comparative genomics, non-core proteome module, evolutionary radiation, host adaptation, immune evasion

Introduction

Understanding the origin and evolution of viruses is complex, yet it is fundamental to fully realize the ecological, agricultural and medical impact of the virosphere (Jones and Naidu 2019, Zimmerman *et al.* 2020, Holmes *et al.* 2021, Liang and Bushman 2021). Plant virus diseases are major threats to food security; they occur worldwide and greatly affect developing countries (Jones and Naidu 2019, Savary *et al.* 2019). Conceptual frameworks rationalize the polyphyletic origins and evolution of the plant virome, as well as its ecological impact on crops and wild species (Lefeuvre *et al.* 2019, Dolja, Krupovic and Koonin 2020). Genomic resources for plant viruses have increased in the past four decades (Pasin, Menzel and Daròs 2019), but our knowledge of plant virus evolution and host adaptation mechanisms is nonetheless incomplete.

The plant-infecting Potyviridae is the largest RNA virus family (realm *Riboviria*) (Fig. 1A). The most recent virus taxonomy based on phylogenomic analyses places the family within the phylum Pisuriviricota (Fig. 1B), which comprises of members of the former picorna-like supergroup (Koonin *et al.* 2020). Potyvirid genomes are a mosaic of modules with polyphyletic origins that can be linked to multiple unrelated viruses, either within and outside *Pisuriviricota* (Dolja, Krupovic and Koonin 2020, Gibbs *et al.* 2020). Despite their complex origin, emergence and diversification of modern po-

tyvirids have been traced to plant-associated astro-like viruses (plastroviruses) and protopotyviruses, groups of viruses identified in plant transcriptomes and aquatic samples (Lauber *et al.* 2019, Wolf *et al.* 2020) (Fig. 1B).

Potyviridae includes > 200 plant virus species currently assigned to the twelve genera Arepavirus, Bevemovirus, Brambyvirus, Bymovirus, Celavirus, Ipomovirus, Macluravirus, Poacevirus, Roymovirus, Rymovirus, Tritimovirus, and Potyvirus, with this last being the most speciose (Gibbs et al. 2020, International Committee on Taxonomy of Viruses 2020). Potyvirids have positive single-stranded RNA genomes of 8-11 kb that are translated into polyproteins, which are in turn hydrolyzed by viral proteinases to release a set of mature products (Adams, Antoniw and Beaudoin 2005, Revers and García 2015). First studies of potyvirid genomes identified a basic layout with conserved gene abundance and order. Yet recent discoveries spurred by sequencing technological advances have revealed a large variability in the genomic structures and gene content. Potyvirid polyproteins indeed show a common core led by diversified leaders that are enriched in non-core modules which expand the proteome structural and functional heterogeneity.

Here, we present a pan-family survey of the structural and functional diversity of the *Potyviridae* proteomes by delineating core and non-core modules (see Supporting Information). We pro-

Received: July 15, 2021. Revised: February 2, 2022. Accepted: February 8, 2022

[©] The Author(s) 2022. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.



Figure 1. Potyviridae within the known RNA virosphere. **(A)** Species and genus abundance in the recognized families of the realm Riboviria; families with > 100 species are labeled (see Table S1 of Supporting Information). **(B)** Schematic phylogenetic tree of the RNA virus RNA-dependent RNA polymerases (RdRp). RNA virus phyla are indicated, and the branches of Pisuviricota are expanded; reverse transcriptases of group II introns and non-long-terminal-repeat transposons were used as an outgroup; scale bar = 0.5. The overall tree topology was taken from Wolf *et al.* (2018), and updated to include the virus lineages (red text) with reported ancestral status to Potyviridae (Lauber *et al.* 2019, Wolf *et al.* 2020).

vide a family-wide classification of P1 proteinases, and review knowledge of non-core domains. We examine abundance of the leader helper component proteinase (HC-pro) within the family, and uncover a putative papain-like protease domain in polyprotein leaders of known and putative *Celavirus* members. Using non-core module evolution as a case study, we summarize main molecular mechanisms that have acted in the *Potyviridae* radiation.

We also discuss the finding that common immune evasion roles can be identified in potyvirid leader cistrons and those of plant, fungal and animal viruses; pointing to host adaptation as a main



Figure 2. Layout and leader diversity of Potyviridae polyproteins. RNA molecules and encoded polyproteins are represented as lines and arrowed boxes, respectively; the hypervariable N termini (leaders), and the conserved middle and C-terminal (core) regions are indicated. Representative layouts are as follows: I, Type-A P1 and HC-pro in full-length polyproteins; II, Type-A P1 and HC-pro in truncated polyproteins generated by P3 frameshifting (P3N-PIPO, and P3N-ALT); III, PISPO production by frameshifting in Type-A P1 cistrons; IV, tandem of Type-A and Type-B P1; V, Type-B P1 and HC-pro; VI, single or tandem copies of HC-pro, and P1 absence; VII, leader-less RNA1, and additional RNA2; VIII, Type-A or Type-B P1 including alkylation B (AlkB) or DUF3725; IX, inosine triphosphate pyrophosphatase (HAM1) in polyproteins with Type-B P1 and no HC-pro; X, atypical leader with non-conserved domains. Diagrams are for illustrative purposes and not to scale; PPV, plum pox virus; SPFMV, sweet potato feathery mottle virus; CVYV, cucumber vein yellowing virus; WSMV, wheat streak mosaic virus; ANSSV, areca palm necrotic spindle-spot virus; CYNMV, Chinese yam necrotic mosaic virus; BaYMV, barley yellow mosaic virus; ENMV, endive necrotic mosaic virus; BlVY, blackberry virus Y; UCBSV, Ugandan cassava brown streak virus; CeLV, celery latent virus.

driver of their evolution. A perspective on the applications of potyvirid leader proteinases and other non-core modules in biotechnology and synthetic biology is also presented.

Core and non-core modules of Potyviridae proteomes

Genera of Potyviridae have a common polyprotein core which is expanded by a heterogeneous array of non-core modules (Fig. 2). A set of eight mature proteins is conserved in the middle and carboxy (C) terminus of the polyproteins, namely P3, 6 kDa protein 1 (6K1), cytoplasmic inclusion (CI) protein, 6 kDa protein 2 (6K2), viral genome-linked protein (VPg), nuclear inclusion protein A proteinase (NIa-pro), nuclear inclusion protein B (NIb), and coat protein (CP) (Revers and García 2015). P3N-PIPO and P3N-ALT are generated by a frameshifting mechanism in the P3 cistron, and are conserved (Yang, Li and Wang 2021, Choi et al. 2022). These conserved proteins have a common but polyphyletic origin (Gibbs et al. 2020). NIa-pro and NIb are homologous to picorna-like signature genes, being, respectively, a chymotrypsin-like cysteine proteinase and an RNA-dependent RNA polymerase (RdRp) with phylogenetic affinity to animal-infecting Astroviridae and other *Pisuriviricota* members. CP was likely acquired from other filamentous RNA viruses, whereas CI is a superfamily 2 helicase most closely related to flavivirid homologs (Koonin et al. 2008, Zamora et al. 2017, Dolja, Krupovic and Koonin 2020).

Organization of polyprotein amino (N) termini (leaders) is highly variable and bears distinctive genus- or even speciesspecific features (Fig. 2). Protein hidden Markov model (HMM) profiles allow for sensitive homology detection and have been applied to infer evolution of viral proteomes, as well as virus identification in metatranscriptomic datasets and taxonomic assignment (Nasir and Caetano-Anollés 2015, Wolf et al. 2018, Bin Jang et al. 2019, Callanan et al. 2020). A combination of HMM and protein profile scans was applied here to quantitatively survey the diversity and abundance of the Potyviridae non-core modules. P1 and HC-pro are the most common, but not universal, leader cistrons (Yang, Li and Wang 2021). Other non-core modules identified in few potyvirid species include the pretty interesting sweet potato potyviral ORF (PISPO), putative zinc fingers and DUF3725, alkylation B (AlkB), inosine triphosphate pyrophosphatase (ITPase/HAM1), as well as a pseudo tobacco mosaic virus-like coat protein (TMV-like CP) domain

Diversity and evolution of non-core modules

P1 proteinases—two phylogenetically and biochemically distinct lineages

P1 is the least abundant among the potyvirid proteinases (Figs 2 and 3A). The C terminus includes a well-conserved chymotrypsinlike serine protease domain, a common module of RNA viruses, which autocatalytically releases P1 from the polyprotein (Rodamilans *et al.* 2018, Mann and Sanfaçon 2019). The N terminus is hypervariable, intrinsically disordered and dispensable for P1 proteolysis (Valli, López-Moya and García 2007, Pasin, Simón-Mateo and García 2014). It can tolerate sequence insertions and diverse atypical domains and functional motifs can be found within it (Fig. 2). *Potyvirus* P1 is active *in planta* and in plant-based translation systems but its proteolysis is very low or absent in animal systems (Rohožková and Navrátil 2011). This supports the hypothesis that activation of potyviral P1 requires a plant co-factor, the identity of which is yet unknown.

Family-wide phylogenesis of the conserved protease domain supports the presence of two distinct lineages—Types A and B (Fig. 3B). Type A is predominant (88%; Fig. 3C), and includes homologs that display plant co-factor dependency in *in vitro* cleavage assays; it is found in all members of Potyvirus and Rymovirus, and in 3/7 of ipomoviruses (Fig. 3D). Type-B proteinases do not need plant co-factors, displaying robust self-processing in multiple translation systems including bacteria (Rodamilans, Valli and García 2013, Shan *et al.* 2018). This lineage is found in all *Ipomovirus*, *Roymovirus*, *Poacevirus*, *Tritimovirus*, and *Brambyvirus* members (Fig. 3D). A tandem of both lineages is found in the ipomoviruses cucumber vein yellowing virus (CVYV), squash vein yellowing virus (SqVYV), and *Coccinia* mottle virus (CocMoV) (Figs 2, 3B and D) (Dombrovsky, Reingold and Antignus 2014, Desbiez *et al.* 2016). Type-A P1 acts as a viral accessory factor, since deletion mutants are infectious and capable of replication and systemic movement (Rohožková and Navrátil 2011, Pasin, Simón-Mateo and García 2014). Consistent with its dispensability, ~10% of the recognized potyvirid species lack P1 (see Arepavirus, Bevemovirus, Bymovirus, Celavirus, and Macluravirus of Fig. 3D).

Pretty interesting sweet potato potyviral ORF (PISPO)

The potyvirus sweet potato feathery mottle virus (SPFMV) has a large Type-A P1 and defective HC-pro (Yang, Li and Wang 2021). Transcriptional slippage takes place within P1 with the derived transcripts coding for a truncated P1 and the frameshift protein PISPO, which participates in RNA silencing suppression (Fig. 2, and see below) (Mingot *et al.* 2016, Untiveros *et al.* 2016). Besides SPFMV, PISPO is present in sweet potato virus 2, C, and G (Clark *et al.* 2012).

Zinc fingers and DUF3725

Zinc-finger domains mediate interaction with DNA, RNA, and proteins, and have a variety of cellular functions that include antiviral immunity regulation. A divalent cation coordinates two cysteines and histidines in CCHH zinc-fingers, but different cysteine/histidine compositions are found in the non-canonical CCHC, CCCH, and CCCC zinc-fingers, all of which have reported RNA interacting ability (Cassandri *et al.* 2017, Corley, Burns and Yeo 2020, Wang and Zheng 2021).

Putative CCCC or CCHC zinc fingers are present in all known Type-B P1s, and are found with varying degrees of conservation in the N terminus of many Type-A orthologs (Fig. 4). Duplicated P1 zinc fingers can be identified in blackberry virus Y (BlVY; Brambyvirus) and sweet potato mild mottle virus (SPMMV; Ipomovirus). Conserved CCCC and CCHC motifs of Type-B P1s were functionally linked to RNA silencing suppression (Valli, Dujovny and García 2008, Kenesi et al. 2017, Gupta and Tatineni 2019a). Consistently, zinc fingers are known in diverse RNA silencing suppressors of plant viruses (Csorba, Kontra and Burgyán 2015, Sōmera, Sarmiento and Truve 2015).

Type-A P1s lack silencing suppression activity and biological roles of their putative zinc fingers remain unknown. The CCCC motif in some of them partially overlaps DUF3725 (Pfam: PF12523) (Fig. 4). DUF3725 is found in *Streptomyces* bacteriophage proteins (ATE85218.1) that share similarities to the zinc-binding domain of DnaG-like primases, which coordinates template binding and RNA primer synthesis in the replication of double-stranded DNA viruses (Gao et al. 2019).

Alkylation B (AlkB)

AlkB domains are ubiquitous among prokaryotes and eukaryotes; having iron(II)- and α -ketoglutarate-dependent dioxygenase activity that reverses nucleic acid methylation damage (Fedeles *et al.* 2015).

AlkB is found in atypical P1s of endive necrotic mosaic virus (ENMV; Potyvirus) and BlVY (Brambyvirus; Fig. 5). AlkB of the latter catalyzes the *in vitro* removal of RNA methyl groups (van den Born *et al.* 2008). Hypermethylation of viral RNA genomes negatively affects plant and animal cell infection (Martínez-Pérez *et al.* 2017, Zhang, Qian and Jia 2021). AlkB was suggested to safeguard viral genomic integrity through repair of methylation damage and promote long-term infection of perennial hosts (van den Born *et al.* 2008, Martínez-Pérez *et al.* 2017). A plant AlkB domain inserted within the tobacco etch virus (TEV) genome was rapidly lost and it did not confer any fitness benefit (Willemsen *et al.* 2017), detailed

4 | FEMS Microbiology Reviews, 2022, Vol. 46, No. 4



Figure 3. Family-wide phylogeny and abundance of Type-A and Type-B P1 lineages. **(A)** Abundance of the potyvirid-encoded proteinases P1, HC-pro and NIa-pro. Ratio of species with at least one domain of the indicated proteinase vs. total species is shown in parentheses (Table S2). **(B)** P1 phylogeny. Protease domain sequences were aligned (Figure S1), and phylogeny was inferred (Supporting Methods); numbers beside branches indicate bootstrap support values; scale bar = 1. All Potyvirus branches are unlabeled and are included in the Type-A lineage; branches of remaining genera are labeled. The ipomoviruses cucumber vein yellowing virus (CYYV), *Coccinia* mottle virus (CocMoV), squash vein yellowing virus (SqVYV) encode a Type-B homolog and an additional Type-A copy (branch II). Common reed chlorotic stripe virus (CRCSV), *Spartina* mottle virus (SpMoV), and longan witches' broom-associated virus (LWBD) are orphans. **(C)** Family counts of Types A and B. **(D)** Abundance of Types A and B across genera of *Potyviridae*. Absolute numbers (Counts) and counts per species (%) are shown. Unassigned includes CRCSV, SpMoV, and LWBD.

characterization of AlkB roles in potyvirid infection remains to be addressed.

AlkB is embedded within replication-associated proteins of plant RNA viruses in the families Alphaflexiviridae, Betaflexiviridae, Closteroviridae, and Secoviridae (Fig. 5). Phylogenomic analysis of plant viruses has highlighted a divergent evolutionary history for AlkB compared to other viral protein domains; it was concluded that AlkB probably emerged by multiple independent acquisition events (Bratlie and Drabløs 2005). For example, the divergent genomic organization and significant phylogenetic separation of BlVY and ENMV suggests that the two viruses acquired the domain independently (Fig. 5B). BlVY and ENMV have been identified in plants of the Rosaceae and Asteraceae, respectively (Susaimuthu et al. 2008, Desbiez et al. 2017), known to host several AlkB-encoding viruses (Fig. 5C). Mixed infections are common in plants, and the AlkB origin in potyvirids can possibly be traced to independent events of inter-family gene transfer.

AlkB is distributed across divergent taxonomic groups of RNA and DNA viruses that include invertebrate RNA viruses (Shi et al. 2016), DNA bacteriophages (Yoshikawa et al. 2018), and giant DNA viruses (Fig. 5). A complex DNA methylation landscape was observed in genomes of the last of these (Jeudy *et al.* 2020), and viral AlkB may have roles in its regulation.

Inosine triphosphate pyrophosphatase (ITPase/HAM1)

ITPase is widespread in cellular organisms, hydrolyzing triphosphates of non-canonical purine nucleotides to prevent their incorporation in nucleic acids and preserve genome integrity (Simone, Pavlov and Borgstahl 2013).

A viral ITPase, also known as HAM1, was first identified in Ugandan cassava brown streak virus (UCBSV) and cassava brown streak virus (CBSV) (Figs 2 and 6). The two are atypical ipomoviruses that lack canonical Type-A P1 and HC-pro, encoding a single Type-B P1 with RNA silencing suppressor activity (Mbanzibwa et al. 2009, Dombrovsky, Reingold and Antignus 2014, Alicai et al. 2016, Shan et al. 2018). ITPase was later identified in *Euphorbia* ringspot virus (EuRSV; *Potyvirus*), encoding Type-A P1 and HC-pro (Fig. 6A) (Knierim, Menzel and Winter 2017). CBSV ITPase, although not essential for infection of experimental hosts, was involved in viral

Type-A P1	🔲 Туре-В	3 P1 🔲 DUF3725			• •
Brambyvirus	BIVY	433 E A K A L L N G L M Y		PSCSDAAVLPEAL LTLE	HKR-SC
, .	BrSMV	116 I RAATRRGLAY	- DIAAQLYMC	PKCCSASDKVLYF DTN	HND-SC
	WSMV	66 LYKATAY <mark>G</mark> LG <mark>Y</mark> -	· - DLAANVFV <mark>C</mark>	GK <mark>C</mark> RSSCTQYRYF IED	HF AC
Tritimovirus	ONMV	66 LYKAMSL <mark>G</mark> LG <mark>Y</mark> -	· - DLAADVFV <mark>C</mark>	CGM <mark>C</mark> RSSCAHYRYF IED	HF A <mark>C</mark>
Thurnovirus	TOgMV	85 L Y I A R A H G L A Y -	· - DLAADVF I C	TQ <mark>C</mark> SSSCANYRYF TED	HE AC
	YOgMV	66 L Y I A V E H G L G Y -	- DLAAETFVC	KQCRSSCRRYKYF IDE	HA AC
1	WEqMV		- DLAAELFVC	GSCRSASPNISYF KI	
Poacevirus	Caiva				
/ oucornuo	TriMV	911 L RATRMNI DY			
i.	TMMoV	75 L K RA I Q NGL L Y		CETCWEWATHVELL YEGEVK	CK AC
	SPMMV	483 L L K A T K V Q V F Y	- DGIKDIFEC	PYCHVSSNELEGL NGDN	CEKC
	CBSV	105 L R R A T S Y <mark>G</mark> F M Y -	· - <mark>D</mark> PVLKAFE <mark>C</mark>	PK <mark>C</mark> RTKATELEAF CSD	CDHC
Ipomovirus	UCBSV	109 L R R A T S Y <mark>G</mark> F M <mark>Y</mark> -	· - DPVACAFE <mark>C</mark>	CPV <mark>C</mark> RTKATALEAF TSD	CDHC
	SqVYV	607 L R R A A R Y G L A F -	· - <mark>D</mark> CALEAYE <mark>C</mark>	CPM <mark>C</mark> GMQCTYLESF YFD	CD FC
	CocMoV	597 L R K A A K F G V G Y -	· - DIVMNHYVC	PVCKMRATDISLL GSN	C D F C
I	CVYV	592 L RKATKYGVGF		CRCHVRCTDISLM NTM	CN EC
	SPFMV	80 L KGAIEHGEVY -	- KWESELTFC		
Potyvirus	SPVG				
	SPV2	801 SNALOYGEVE			
Inomovirus i	SPMMV	691 RIALARGTIY		CPDCDVIVDSEFGWF	CF - DC
ipoinovii do i	WMVBV	1 83 T R K AML S G L K H		PECDETEDVFF KK	C S C
	SuWMV	83 T R K A M L S G L K H	- DYTANQWIC	PECDETEDTFF KK	C S C
Potyvirus	ZTMV	83 C S K A L I S G L T Y -	- DFNSEVWIC	PE <mark>C</mark> GECEDKFF KK	C E C
	PRSV	83 I RKTMVGRLG <mark>Y</mark>	· - DFESELWV <mark>C</mark>	RN <mark>C</mark> DKTSEKYF KK	C D C
I	ZSV	83 A R K A L C G N F S Y	· - DPEMDAFQC	- E <mark>C</mark> GEQSDSSV SA	CPEC
(CocMoV	83 VRKSMCSGLHY	- NPNMDEFEC	- ECGFMSEQFS KV	CE DC
Ipomovirus	CVYV	83 VRRALCSALYY -	- NPNEEDFEC	C-ECGNSNTSWKRV	CE DC
Unassigned I	Sqviv				
	PeSV	821 KEALENGEDY	- FIFTGTYFC	PYCSLORDS I KEMKKAVGSLM	
Roymovirus	RYMV	56 L L E A F R NG V SY	- NPERL TYG	CPFCSVEAANIPEALAKINSEG	
Brambyvirus	BIVY	25 A D A A K R R G L L F	-DHGSETYEC	PRCGEIWRNLDDYM - AEGGKM	HPK-KC
, .	NeYSV	8 V L A A K R R G I E F -	- DLEDEVYW <mark>C</mark>	PR <mark>C</mark> GEVFNGLQEV DEE	HRH-G <mark>C</mark>
	CeMV	32 A R A A N R R S L H Y -	- DCEDDVYL C	VK <mark>C</mark> SESFDSVQQF KDE	H E <mark>C</mark>
	ApVY	32 ALAAKKRNLRY -	· - <mark>D</mark> AEDDVYH <mark>C</mark>	LD <mark>C</mark> DETFDTVEQF RME	H D <mark>C</mark>
	CalMoV	32 A V A A H K M G W V Y -	· - DVEMDDYQC	PR <mark>C</mark> GEIYNSKEEI RNY	H E C
	LMV	32 YLAAQRIGFYY -	- DWDDDVYEC	PTCEAIYHSLDEIKNW	H E C
		32 ALCIENNGFEY		YKCKHAFNNKEEIHE	
	LYSV				
	SWW				
_	NDV	24 VSAAARRDI TW		NKCGNQYRSI HDF FFN	H SC
Potyvirus	OYDV	14 L T L AQRKRACY	SEDDAVHYHC	TKCNFAFESLRMVR PVN	H DC
	BICMV	41 P I SARKRNL CF	DDDEDGHYHC	HFCDCECDSKNHL EE	HERDIC
	CAI23781.1	29 R I SRRERL V C <mark>H</mark> O	DGDEEGHFH	TH <mark>C</mark> GRECASLTMLKQKN	H T <mark>C</mark>
	LuMV	18 K Y T R K I R	DEDGEYR <mark>C</mark>	TL <mark>C</mark> GLGCDSMTMAK PVN	H N C
	PSbMV	18 <mark>A</mark> R T R R I R	DADGEYR <mark>C</mark>	TQ <mark>C</mark> DMGFDSMTMAR PVN	H C <mark>C</mark>
	QED42798.1	36 T K T P E V L <mark>G</mark> K L <mark>Y</mark> C	СК <mark>С</mark>	PICDTPCPTKSFYRK	HMATAC
	QHB15168.1	36 EY I KENIGNLYC	2 K	PECGAPCPTKSFMS R	
			, RC		
	QFZ79297.1	33 FRI IVREKKVYO	3 K (GVCGDEMFIKSMLKK	HLASKC

Figure 4. Putative zinc-finger motifs and DUF3725 in P1. Alignment of a conserved cysteine-rich region of Type-A and Type-B P1s is shown; inverted triangles indicate putative zinc-finger residues described to be involved in RNA silencing suppression activity of Type-B homologs (Valli, Dujovny and García 2008, Kenesi *et al.* 2017, Gupta and Tatineni 2019a). For each virus, position of the first aligned polyprotein residue is indicated, and colored in red to label duplications; accession numbers are shown or given in Table S2 alongside virus complete names.

accumulation and symptom development. Contrary to the predicted antimutagenic activity of ITPase, viral mutation rates were not reduced in transgenic plants overexpressing CBSV ITPase, nor they were increased in CBSV clones lacking ITPase (Tomlinson *et al.* 2019). Use of improved sequencing approaches and alternative experimental systems could help shed light on the ITPase roles in potyvirid infection.

Metagenomics surveys have uncovered ITPase across diverse RNA and DNA virus taxa (Fig. 6). The ITPase fold is found in plant and invertebrate RNA viruses (Shi *et al.* 2016, Le Lay *et al.* 2020,



Figure 5. AlkB in Potyviridae and divergent virus taxa. (A) Diagrams of representative ORF bearing AlkB; relevant domains are colored. Species and taxonomic groups (left) are labeled; BIVY, blackberry virus Y (genus *Brambyvirus*); ENMV, endive necrotic mosaic virus (genus Potyvirus); Escherichia coli alkB as a standard. (B) AlkB phylogeny. Protein sequences in plant viruses were aligned (Figure S2), and phylogeny was inferred; numbers beside branches indicate bootstrap support values; scale bar = 1; E. coli AlkB was included as a reference (black); Potyviridae accessions are labeled (red). (C) Conserved residues in viral AlkB proteins. Alignment blocks show regions of *E. coli* AlkB that participate in catalysis (inverted triangles) or α -ketoglutarate binding (diamonds) (Yu *et al.* 2006). Position of the first residue is indicated (left), and the (poly)protein size is shown in parentheses. Right, virus taxonomic groups and plant host families are indicated (R, Rosaceae; A, Asteraceae).

Leiva *et al.* 2022), as well as in bacteriophages and giant DNA viruses (Kiljunen *et al.* 2005, Deeg, Chow and Suttle 2018, Sun and Ku 2021).

Tobacco mosaic virus-like coat protein (TMV-like CP)

Bymovirus is the only potyvirid genus whose member transmission is mediated by soil-borne plasmodiophorids (Jiang *et al.* 2020). Bymoviruses have bipartite genomes with RNA1 encoding the potyvirid polyprotein core, and RNA2, which encodes a second polyprotein processed in P2-1 and P2-2 (You and Shirako 2010). P2-1 is closely related to HC-pro (see below). P2-2 shares no similarity with other potyvirid proteins, and bymoviruses with its truncation or complete deletion are able to replicate and systemically move, but could not be transmitted by the natural vector (You and Shirako 2010).

Plasmodiophorid-transmitted viruses include Virgaviridae and Benyviridae members (Tamada and Kondo 2013), whose capsid proteins show homology with bymovirus P2-2 (Dessens and Meyer 1996) (Fig. 7A). HMM-profile scans detect a 'pseudo' TMV-like CP domain conserved in P2-2 of all full-length bymovirus accessions and absent in oat mosaic virus, whose reference sequence is of a mechanically propagated isolate which lacks most of P2-2 (You and Shirako 2010). TMV-like CP sequences of bymoviruses cluster



Figure 6. HAM1/ITPase in Potyviridae and divergent virus taxa. (A) Diagrams of representative viral ORF including the inosine triphosphate pyrophosphatase (ITPase/HAM1) fold; relevant domains are colored. Left, taxonomic groups and species are shown; UCBSV, Ugandan cassava brown streak virus (*Ipomovirus*); EuRSV, *Euphorbia* ringspot virus (Potyvirus); E. coli ITPase rgdB is included as a standard. (B) Conserved residues in ITPase sequences. Alignment blocks show regions of E. coli RdgB that participate in substrate binding or catalysis (inverted triangle) (Savchenko et al. 2007). Position of the first residue is indicated (left), and (poly)protein size is shown in parentheses.

within a monophyletic clade, which supports their common origin (Fig. 7B). The P2-2 domains show phylogenetical relatedness to CP of Virgaviridae, and of wheat stripe mosaic virus (WhSMV), a putative benyvirus (Fig. 7B). Besides plant viruses, the TMV-like CP fold is found in algae and invertebrate RNA viruses but has negligible homology with cellular proteins (Nasir and Caetano-Anollés 2015, Shi et al. 2016, Vlok, Gibbs and Suttle 2019).

Helper component proteinase (HC-pro)

HC-pro is a multifunctional leader proteinase with roles in virus transmission, polyprotein processing, and suppression of antiviral RNA silencing (Valli et al. 2018). The HC-pro RNA silencing suppressor activity is indispensable for potyvirus infection (Garcia-Ruiz et al. 2015), and, based on current data, it can be safely considered a Potyvirus core component. Within a family-wide perspective, the reported genomic variation beyond the genus Potyvirus supports the HC-pro classification as a non-core module. Several ipomoviruses naturally lack HC-pro and its sequence is absent in ~3% of potyvirid genomes (Figs 3A and 8A). Experimental evidence using a clone of wheat streak mosaic virus (WSMV; Tritimovirus) with complete HC-pro deletion shows the protein is dispensable for virus replication and movement (Stenger, French and Gildow 2005). Tritimoviruses as well as ipomoviruses enlist Type-B P1 as the viral silencing suppressor (Valli, Dujovny and García 2008, Mbanzibwa et al. 2009, Giner et al. 2010, Young et al. 2012). HC-pro is thus dispensable in potyvirids that encode proteins evolved to take over key functions originally described for homologs of model potyviruses. Adaptive HC-pro functional loss and dependency evolution were reported for onion yellow dwarf

virus, which encodes a defective HC-pro *trans*-complemented by a co-infecting potyvirus (Jayasinghe *et al.* 2021). Bymovirus RNA1 lacks leader proteases, whereas RNA2 encodes the HC-Pro homolog P2-1 (Adams, Antoniw and Beaudoin 2005), two functionally and phylogenically divergent HC-pro copies are present in arepaviruses (Qin *et al.* 2020) (Fig. 8A and B).

HC-pro has a papain-like cysteine protease domain that autocatalytically hydrolyzes its C terminus (Guo, Lin and Ye 2011), and shows significant sequence divergence within the family that can be possibly rooted close to the bevemovirus ortholog and bymovirus P2-1 (Fig. 8B). HC-pro shows homology to the nsP2 main proteinase of alphaviruses, as well as leader proteinases of closteroviruses, picornaviruses and arteriviruses (Gorbalenya, Koonin and Lai 1991, Mann and Sanfaçon 2019). Homology identification between HC-pro and *Cryphonectria* hypovirus 1 (CHV1) p29 was instrumental to postulate the evolutionary relationship between Potyviridae and Hypoviridae, a family of fungal RNA viruses (Koonin et al. 1991) (Figs 1B and 8C). Papain-like cysteine proteases are common in cellular organisms, and main components of plant immunity (Misas-Villamil, van der Hoorn and Doehlemann 2016).

Celery latent virus—an outlier

Celavirus is a single-member genus with celery latent virus (CeLV) as the largest and most divergent of recognized potyvirids (Gibbs *et al.* 2020). CeLV polyprotein initiates with a signal peptide that could translocate reporter proteins to the endoplasmic reticulum (Rose *et al.* 2019), no other *N*-terminal signal peptides are known in potyvirids. Sensitive HMM-profile scans failed to identify P1, HC-pro, or other potyvirid non-core modules. Inspection



II. Furovirus, Pomovirus II. Hordeivirus, Pecluvirus, Goravirus IV. Tobamovirus

Figure 7. Monophyletic origin of the pseudo TMV-like CP domains of *Bymovirus.* **(A)** Genomic diagrams of the potyvirid barley yellow mosaic virus (BaYMV; genus *Bymovirus*) and soil-borne cereal mosaic virus (SBCMV; *Furovirus*), a Virgaviridae vectored by soil-borne plasmodiophorids. Relevant domains are colored; TMV-like CP, tobacco mosaic virus-like coat protein domain. BaYMV RNA2 encodes P2-1, with HC-pro homology, and P2-2, with a conserved 'pseudo' TMV-like CP domain. **(B)** TMV-like CP phylogeny of bymoviruses and reference RNA viruses. Protein sequences were aligned (Figure S3), and phylogeny was inferred; number beside branches indicates the bootstrap support value; scale bar = 1. Bymovirus accessions are in red—barley mild mosaic virus (BaMMV), rice necrosis mosaic virus (RNMV), wheat spindle streak mosaic virus (WSSMV) wheat yellow mosaic virus (WYMV). Wheat stripe mosaic virus (WSSMV) is a putative benyvirus (Valente *et al.* 2019).

of the CeLV leader nonetheless reveals the presence of a putative papain-like protease domain with sequence similarity to HC-pro and bymovirus P2-1, as well as p29 of the hypovirus CHV1 (Fig. 8C). The identified catalytic residues and cleavage site are conserved in *Striga* potyvirus B (QVG60634.1; Fig. 8C), a virus phylogenetically related to CeLV and recently reported as *Striga*-associated poty-like virus 2 (Choi *et al.* 2022).

Mechanisms of non-core module evolution

Our pan-family, quantitative survey of the Potyviridae proteomes defines the abundance of non-core modules and highlights discrete distribution patterns along the evolutionary tree of the family (Fig. 9A). High mutation rates, recombination, gene duplication and *de novo* emergence as well as extensive gene loss and gain, and host-niche adaptation drive virus evolution. Which are the main molecular mechanisms behind non-core proteome expansion in the family's evolutionary radiation?

Recombination is common in RNA viruses and an important component of potyvirid speciation (Sztuba-Solińska et al. 2011). Its significance in potyvirid non-core module evolution can be clearly exemplified by AlkB identification in the potyvirus ENMV and the brambyvirus BlVY (Fig. 9B), possibly linked to independent acquisition events occurred in mixed infections with unrelated plant viruses.

Gene duplication is a major source of phenotypic novelty in cellular organism (Innan and Kondrashov 2010). It is however rare in RNA viruses, with the Closteroviridae coat protein duplication as a notable exception in plant viruses (Simon-Loriere and Holmes 2013). Tandem P1 or HC-pro copies in ipomoviruses and arepaviruses, respectively, were related to duplication events (Valli, López-Moya and García 2007, Qin et al. 2020). Empirical results nonetheless show that redundant sequences are rapidly purged from potyvirid genomes despite the potentially beneficial effect of the encoded proteins. Artificial insertion of a second HCpro copy in the genome of TEV was deleterious and rapidly lost (Willemsen et al. 2016). Experimental evolution of a PPV clone encoding its own Type-A P1 and a second ortholog from a phylogenically distant potyvirus led to an array of progeny viruses with enhanced fitness that were characterized by an almost or complete duplication loss (Rodamilans, Casillas and García 2021). Together the results indicate that both sequence identity and functional redundance constrain gene duplication in potyvirids. Further supported by the polyphyletic origin of duplicated copies (e.g. see Ipomovirus-encoded P1s labeled by II and V in Fig. 3), it can be concluded that gene duplication events detected in potyvirids are likely by-products of interspecific, ortholog recombination (Fig. 9B).

Neofunctionalization and functional specialization in potyvirids can be inferred from biochemical and biological characterization of P1 lineages (Fig. 9B). P1 was identified as a host adaptation determinant based on gene swapping and infection assays, and on genome-wide analysis of nucleotide variation (Salvador et al. 2008, Maliogka et al. 2012, Shan et al. 2015, Nigam et al. 2019), at a protein level, it shows conserved structural disorder (Pasin, Simón-Mateo and García 2014). Structurally flexible segments in viral proteins increase mutation tolerance and adaptability through acquisition of new linear motifs or protein domains (Gitlin et al. 2014, Charon et al. 2018, Mishra et al. 2020). Strong evolvability and adaptation capacities of P1 are corroborated by family-wide identification of heterogenous motifs and domains within the P1 N termini, as well as the de novo emergence of PISPO through overprinting. Subfunctionalization allows the division of functions in duplicated genes (Innan and Kondrashov 2010). A zinc finger motif is conserved in all Type-B proteins but absent in most of P1s (Fig. 4); subfunctionalization of a Type-B-like ancestor could have participated in Type-A specialization.

Recently proposed scenarios place the bisegmented bymoviruses at the evolutionary diversification root of potyvirid genera, which are suggested to have originated through genomic segment fusion (Qin et al. 2020). Celaviruses have monopartite genomes (Rose et al. 2019, Choi et al. 2022), and their ancestral status compared to bymoviruses as supported by RdRp phylogeny makes direction of the multipartitism transition uncertain. Experimental examples are known of transitions from an originally non-segmented virus to a bisegmented one (Lucía-Sanz and Manrubia 2017). Supported by identification of a conserved pseudo TMV-like CP in bymoviruses, an intriguing possibility to explain the bipartidism emergence is the recruitment by a monopartite ancestor of a new genomic segment from co-infecting tobamoviruses to access to a vector transmission mode unprecedented within the family; transfer of the HC-pro homolog P2-1 could have been required to stabilize this de novo association and the gained multipartite state.



Figure 8. HC-pro and papain-like protease domains in Potyviridae. (A) HC-pro homologs across genera of Potyviridae. Absolute numbers (Counts) and counts per species (%) are shown (Table S2). Unassigned includes CRCSV, SpMoV, and LWBD. (B) HC-pro phylogeny. Protease domain sequences were aligned (Figure S4), and phylogeny was inferred; number beside branches indicates the bootstrap support value; scale bar = 1. Roman numerals indicate genera; branch II includes Bymovirus P2-1; Arepavirus members encode an HC-pro tandem and branches of the first and second encoded copies are labeled with III(1) and III(2), respectively; LWBD, CRCSV, SpMoV are orphans. (C) Putative HC-pro-like domain in *Celavirus*. Alignment of papain-like protease domains of celery latent virus (CeLV; *Celavirus*) and Striga potyvirus B (QVG60634.1), with turnip mosaic virus (TuMV) HC-pro, barley mild mosaic virus (BaMMV) P2-1, and Cryphonectria hypovirus 1 (CHV1) p29; polyprotein accession numbers and positions are indicated; diamonds, catalytic residues; inverted triangle, known or putative cleavage sites.

Immune evasion through leader and non-core modules of potyvirids and other RNA viruses

In microbial systems, large taxonomic variability can be summarized by sets of redundant, polyphyletic functions (Louca *et al.* 2018). Functional analyses of cellular pangenomes suggest that gain of non-core genes influences adaptation of plant microbes to ecological niches (Box 1). Notwithstanding the low level of structural and biochemical conservation: Do Potyviridae non-core modules share biological function(s) and a common evolutionary driver? We present evidence supporting a main biological role of these non-core modules in counteracting host defensive reactions and thus host adaptation.

Potyvirid leader modules and RNA silencing evasion

Plant RNA viruses have evolved a variety of strategies to modulate disease severity and escape cellular antiviral responses (Paudel



Figure 9. Potyviridae non-core proteome diversity and evolution drivers. (A) Abundance of non-core proteome components in genera of Potyviridae and recognized orphan species (CRCSV, SpMoV, LWBD); species numbers per genus are in parentheses; QVG60634.1, Striga potyvirus B; question marks indicate presence of putative homologs. Right, family phylogeny; RdRp domains were identified, protein sequences were aligned (Figure S5), and phylogeny was inferred; branches with bootstrap support \geq 95 are in red, scale bar = 1. (B) Mechanisms and examples of non-core component evolution in Potyviridae. Non-core module inventory and virus complete names are given in Table S2.

and Sanfaçon 2018, Li and Wang 2019, Križnik, Baebler and Gruden 2020). Being the first translation products, viral leader cistrons are considered important virulence and pathogenicity factors that can coordinate the early infection stages.

Potyvirid leaders are enriched in non-core modules (Fig. 2), and experimental evidence supports their roles in immune evasion and symptom development (Figs 10 and 11). RNA si-

lencing is a major antiviral mechanism of plants. Potyvirus HC-pro is among the best characterized silencing suppressors, with multiple roles that include direct sequestration of small RNA molecules and inhibition of RNA silencing factors (Valli *et al.* 2018). In addition to HC-pro, other potyvirid leader proteins have been implicated in evasion of antiviral RNA silencing (Fig. 10A).



Figure 10. Immune evasion roles of non-core modules of Potyviridae. (A) Diagram of Potyviridae infection and replication stages, as well as main antiviral pathways in which non-core modules participate. Question marks indicate roles with unknown mechanisms or unavailable *in vivo* validation data; HR, hypersensitive response; NLR, NOD-like receptor; NPR1, NONEXPRESSOR OF PATHOGENESIS-RELATED GENES 1; SA, salicylic acid; ROS, reactive oxygen species; JA, jasmonic acid; ABA, abscisic acid; ET, ethylene. (B) Immune evasion by virus post-translational negative autoregulation. Left, P1 autoinhibition allows PPV virulence attenuation and evasion of defense responses; right, removal of the P1 autoinhibitory domain leads to an accelerated PPV amplification that activates hormone responses, which restrict virus accumulation in the long term (Pasin *et al.* 2020). Plots depict levels of virus accumulation (blue) and defense response activation (red).

Type-A P1 does not suppress RNA silencing in transient expression assays, and enhances potyviral infection in plants defective in RNA silencing (Young *et al.* 2012, Pasin, Simón-Mateo and García 2014), which suggests that the protein has additional roles independent of silencing suppression. P1 *cis*-expression strengthens the HC-pro activity, and improved translation in heterologous systems was implicated to this effect (Tena Fernández *et al.* 2013). P1 evolution for the mere enhancement of HC-pro expression appears unlikely, since optimization of the nucleotide Kozak context would be a more economic strategy.

Robust RNA silencing suppressor activity was nonetheless reported for several Type-B proteins common in Ipomovirus, Poacevirus, Tritimovirus, Roymovirus and Brambyvirus members. These viruses encode HC-pro with no detectable suppressor activity or silencing suppressor motifs, or lack the cistron altogether, as seen in some ipomoviruses. Type-B P1 binds short RNA molecules; this ability correlates with its silencing suppression activity (Valli, Dujovny and García 2008, Kenesi et al. 2017, Gupta and Tatineni 2019a). Other data show that RNA silencing suppressor activity of Type-B proteins is conferred by GW motifs that guide recognition and inhibition of the antiviral silencing component ARGONAUTE 1 (AGO1) (Giner et al. 2010, Kenesi et al. 2017). Motif requirements for silencing suppression activity has been investigated in Type-B homologs of Ipomovirus, Poacevirus, and Tritimovirus (Giner et al. 2010, Gupta and Tatineni 2019a,b, Chen et al. 2020). Type-B P1 proteolytic activity is not needed for silencing suppression of the ipomovirus SPMMV and the poacevirus Triticum mosaic virus (TriMV) (Giner et al. 2010, Gupta and Tatineni 2019a). The P1 cistron was expressed alone, however, and it is unclear if proteolysis is needed during infection to release mature, active silencing suppressors from polyproteins.

GW motifs involved Type-B protein activity are also present in the potyviral PISPO, and the SPFMV P1N-PISPO fusion acts as a silencing suppressor that functionally replaces HC-pro (Mingot *et al.* 2016, Untiveros *et al.* 2016). GW motifs are present in HC-pro; although not involved in silencing suppression, they are needed to recruit AGO1 for pro-viral functions (Pollari *et al.* 2020), which further highlights the leader cistron multifunctionality in host adaptation.

Negative autoregulation of potyvirid infection for immune evasion

Diseases result from failures of cellular homeostasis (Kotas and Medzhitov 2015). Negative feedback and incoherent feedforward loops are major autocontrol mechanisms that allow biological systems to adapt to changing environment and perturbations without homeostasis loss, disease or autoimmunity. They regulate natural and engineered cellular systems, as well as phase transitions and adaptation to resource changes of bacteriophages (Pitsili, Phukan and Coll 2020, Brady *et al.* 2021, Frei and Khammash 2021, Yao *et al.* 2021). The importance of negative autoregulation in plant virus infection just starts to be appreciated.

Mechanisms have been reported in plant viruses that avoid cellular toxicity or excessive inhibition of antiviral pathways, which can trigger host damage and pathogen fitness loss (Paudel and Sanfaçon 2018, Križnik, Baebler and Gruden 2020). Promotion of RNA silencing spread was reported for tobamovirus movement protein and sobemovirus P1 (phylogenically unrelated to potyvirid P1), restriction of the silencing suppressor activity of cucumber mosaic virus 2b, geminiviral β C1 and polerovirus P0 was proposed to minimize host homeostasis perturbation (Vogler *et al.* 2008, Lacombe *et al.* 2010, Zhang *et al.* 2017, Ismayil *et al.* 2020, Watt *et al.* 2020, Clavel *et al.* 2021, Shukla *et al.* 2021).

HC-pro is a symptom determinant; its uncontrolled expression severely affects plant physiology, growth, fertility, and can trigger hypersensitive response or lethal necrosis (Pacheco et al. 2012, Valli et al. 2018). These findings suggest that tight control of proteins with strong silencing suppressor activity is desirable for optimal viral fitness, but how can it be achieved by an RNA virus that lacks transcriptional regulations? Recent data on P1 highlight a post-translational negative autoregulation that provides an evolutionary answer to the virus dilemma of counteracting defenses of the host without killing it. P1 can antagonize HC-pro, since the P1-HC-pro fusion lacks RNA silencing suppressor activity and could not sustain viral infection in hosts with unopposed antiviral immunity (Pasin, Simón-Mateo and García 2014). P1 is itself under autoinhibitory control. Several proteases display autoinhibitory domains or are synthesized as precursors that undergo structural rearrangements to activate (Hedstrom 2002, Gohara and Di Cera 2011, Trudeau et al. 2013). Nterminal deletions of PPV P1 identified a gain-of-function phenotype consistent with an autoinhibitory mechanism in which the N terminus negatively regulates P1 proteolysis, and selfcleavage results from autoinhibition relief by plant co-factor(s) (Pasin, Simón-Mateo and García 2014, Shan et al. 2018). A recent study model proposes the autoinhibited P1 self-cleavage as an immune evasion mechanism that regulates PPV replication through controlled release of the functional silencing suppressor HC-pro (Pasin et al. 2020). Self-controlled P1 processing kinetics would thus balance the strength of RNA silencing suppression with magnitude of phytohormone-mediated defense activation to mitigate resource burden and promote long-term viral fitness (Fig. 10B).

Additional immune evasion roles of potyvirid non-core modules

RNA silencing and other RNA metabolic pathways contribute to plant defense against potyvirids (Li and Wang 2019, Xu *et al.* 2020). They are further interconnected with autoimmunity, hormonal, and autophagic responses to provide robust plant immunity and tolerance to viruses (Cui *et al.* 2020, Pasin *et al.* 2020, Pitzalis *et al.* 2020, Shukla *et al.* 2021).

HC-pro interacts physically with RNA turnover components and inhibits EXORIBONUCLEASE 4 (XRN4) to counteract antiviral RNA decay (Fig. 10A) (Li and Wang 2018, De *et al.* 2020). Clover yellow vein virus (ClYVV) P1 was involved in overcoming the recessive resistance conferred by eukaryotic translation initiation factor 4E in pea (Nakahara *et al.* 2010). Selective translation enhancement of viral genomes by TEV P1 has been reported, which might contribute to suppressing expression of host immune factors (Martínez and Daròs 2014). An evasion strategy of alphaviruses relies on disruption of stress granule formation by G3BP targeting mediated by peptide motifs that resemble the IxFG motif conserved in P1 N termini (Pasin, Simón-Mateo and García 2014, Panas et al. 2015, Reuper and Krenz 2021); P1 roles in stress granule processes are unknown.

Methylation impacts small RNA stability and loading in silencing complexes, and it is modulated by several silencing suppressors (Ji and Chen 2012, Csorba, Kontra and Burgyán 2015). HC-pro alters small RNA methylation through HUA ENHANCER1 methyltransferase interference and local disruption of the methionine cycle (Ji and Chen 2012, Ivanov et al. 2016, Del Toro et al. 2021). Roles of AlkB and its RNA demethylase activity in potyvirid infection are less clear (Fig. 10A). It has been suggested that regulation of RNA methylation during infection contributes to viral immune evasion by fine-tuning viral replication rates or by post-transcriptional control of host gene expression (van den Born et al. 2008, Zhang, Qian and Jia 2021). N⁶-methyladenosine amount modulation was recently proposed as a new plant antiviral mechanism that hinders long-distance viral movement (Martínez-Pérez et al. 2021), and recent data indicate it could be effective against potyvirids as supported by reported changes in N⁶-methyladenosine levels upon bymovirus infection (Zhang et al. 2021).

HAM1 and its ITPase activity were recently shown to be CBSV necrosis determinants (Tomlinson *et al.* 2019). Although the mechanistic details were not studied, levels of inosine triphosphate (an ITPase substrate) were shown to regulate key factors potentially involved in antiviral immunity such as the viral RdRp catalytic speed and possibly viral replication rates, as well as plant stress response activation (Dulin *et al.* 2015, Kazibwe *et al.* 2020).

Recent results indicate possible P1 roles in coordinating plant homeostasis during mixed infections, since the protein impaired activity of the crinivirus silencing suppressor P25 (Domingo-Calap et al. 2021).

Expansion and immune evasion roles of RNA virus leaders

Diversification and immune evasion roles have been described for leaders or 5' genomic cistrons of phylogenetically divergent groups of RNA viruses of plants, fungi and animals.

Plant RNA viruses of the Sobemoviridae family, genera Enamovirus and Polerovirus (family Luteoviridae), as well as Waikavirus and Fabavirus (family Secoviridae) belong to the picorna-like supergroup (Wolf et al. 2018). Cistrons encoded by their 5' genomic portions show RNA silencing suppressor activity (Csorba, Kontra and Burgyán 2015, Sõmera, Sarmiento and Truve 2015, Stewart et al. 2017, Carpino et al. 2020). Leader proteinases of Closteroviridae (phylum Kitrinoviricota) affect pathogen virulence, superinfection exclusion, and promote viral amplification, possibly by viral replicase activation or subversion of host antiviral defenses (Dolja, Kreuze and Valkonen 2006, Atallah et al. 2016, Kang et al. 2018). Similar to potyvirids, proliferation of closterovirus leader proteases is reported. A single, a tandem, or three copies of leader proteinases are found, respectively, in genomes of beet yellows virus, citrus tristeza virus, and actinidia virus 1, among others (Fig. 11).

Fungal RNA viruses of *Hypoviridae* recruit leader cistrons to counteract antiviral immunity. RNA silencing suppressor activity was reported for the CHV1 leader protease p29, and p24 of *Cryphonectria* hypovirus 4 (CHV4; Fig. 11) (Segers *et al.* 2006, Aulia *et al.* 2021).

Among animal viruses and similar to potyvirids, picornaviruses show expansion of genomic layouts with highly divergent leaders (Fig. 11) (Gorbalenya and Lauber 2010, Zell 2018). Their leader proteins have a low level of structural and biochemical conservation, but share common biological functions in immune evasion





(Agol and Gmyl 2010). Leader proteinase (L^{pro}) of foot-and-mouth disease virus (FMDV; Aphthovirus) suppresses host cellular translation and antiviral responses by direct proteolysis of host translation factors and other RNA-binding proteins, signaling components, and conjugated ubiquitins (Saiz and Martinez-Salas 2021). Cardiovirus L, which is not a protease, antagonizes immune responses by suppressing interferon production, and can be functionally replaced by FMDV L^{pro} (Freundt, Drappier and Michiels 2018, Visser et al. 2020). Theiler's murine encephalomyelitis virus (TMEV; Cardiovirus) encodes the accessory L*, which directly targets RNase L ankyrin domains for interferon pathway inhibition and virus persistence promotion (Drappier et al. 2018). Murine but not human RNase L was found to be inhibited by L* (Drappier et al. 2018); this species-specific activity brings to mind the host-dependent activation of potyvirid Type-A P1 proteolysis (see above). The small protein UP was recently identified in the 5' region of diverse enteroviruses; UP modulates virus infection and tropism, and was suggested to participate in autophagy subversion for virus particle release (Lulla et al. 2019).

Leader size and domain organization vary considerably among members of the order Nidouirales. Arteriviruses are important veterinary disease agents; nsp1 is the first and most variable protein encoded. It is a leader proteinase, and up to three active copies are found in *Deltaarterivirus* (Vatter *et al.* 2014, Gulyaeva *et al.* 2017). Nsp1 proliferation resembles those of potyvirids encoding tandems of P1 or HC-pro (Fig. 2). Arterivirus nsp1 and its copies nsp1 α , nsp1 β , and nsp1 γ counteract host immune defenses through interferon pathway suppression (Han and Yoo 2014, Lunney *et al.* 2016). Nsp1 of betacoronaviruses is released from polyprotein N termini to rapidly repress translation of cellular transcript and expression of innate immunity factors by 40S ribosomal subunit association (Nakagawa and Makino 2021). Immune evasion roles are conserved in nsp1 of alphacoronaviruses (Shen *et al.* 2020).

Pestivirus N^{pro} (phylum Kitrinoviricota) is an accessory leader proteinase that acts as an interferon pathway antagonist to prevent cell apoptosis (Tautz, Tews and Meyers 2015, Jo et al. 2019).

Roles in immune evasion thus appear to be a functional link that connects Potyviridae non-core modules to each other, as well as leader cistrons of potyvirids with those of multiple RNA viruses (Fig. 11).

Biotech appeal of non-core modules Infectious clones—established tools for potyvirid biological characterization and biotechnological advances

The accessory nature identified in non-core modules warrants the use of suitable experimental systems for their biological role characterization. Full-length infectious clones are universal, indispensable tools for virus biology research and the development of experimental systems for investigating diseases (Pasin, Menzel and Daròs 2019, Kannan et al. 2020). They have been generated for members of Potyvirus (Domier et al. 1989), Tritimovirus (Choi et al. 1999), Macluravirus (Kondo and Fujita 2012), Poacevirus (Tatineni et al. 2015), Ipomovirus (Pasin et al. 2017), Celavirus (Rose et al. 2019), Arepavirus (Qin et al. 2020), as well as for bipartite viruses of Bymovirus (You and Shirako 2010, Ohki, Sasaya and Maoka 2019). Homology-based cloning methods are revolutionizing the potyvirid infectious clone construction, since they are efficient and require limited viral sequence information (Desbiez et al. 2012, Zhao et al. 2020). T-DNA vectors with stabilizing features



Figure 12. Potyvirid vectors for sequence delivery and expression in plants. (A) Diagram of a potyvirid genome and its insertion sites suitable for gene expression; GOI, gene of interest; site labels are as follows: I, upstream of P1; II, P1 N terminus; III, P1/HC-pro junction; IV, P3 N terminus; V, CI N terminus; VI, NIb N terminus; VII, NIb/CP junction (Rajamäki et al. 2005, Chen et al. 2007, Majer, Navarro and Daròs 2015); VIII indicates the CP N terminus, which has been used for heterologous peptide expression (Sánchez and Ponz 2018). Sites involved in natural gene gain events are in cyan. (B) Transient, potyvirid-mediated manipulation of crop traits. Right, a potyvirid vector is used to confer plant dwarfism through over-expression of a gibberellin catabolic enzyme gene (GA20x8) inserted at the P1/HC-pro junction (Torti et al. 2021); left, growth of an untreated plant with unaltered gibberellin levels.

have been used for one-step assembly of potyvirid clones suitable for *Agrobacterium*-mediated delivery (Pasin *et al.* 2017, 2018). A recently developed synthetic genomics framework with plant virome capacity could streamline characterization and engineering of plant viruses with no biological material need (Pasin 2021).

Non-core module characterization to guide plant expression vector development

Virus infectious clones can be engineered and optimized as expression vectors for plant biotechnology and synthetic biology (Fig. 12) (Pasin, Menzel and Daròs 2019, Khakhar and Voytas 2021). Vectors based on potyvirids have been applied for disparate uses, ranging from production of heterologous peptides in plants, to flowering induction, gene silencing, metabolic engineering, CRISPR/Cas-targeted plant genome editing, and reprogramming of crops and their organelles (Lin et al. 2007, Llorente et al. 2020, Martí et al. 2020, Torti et al. 2021, Tuo et al. 2021, Uranga et al. 2021, Xie et al. 2021). Knowledge of Type-A P1 and its proteolytic activity has been instrumental in generating the first potyviral vectors (Fig. 12A). The bacterial β -glucuronidase (GUS) gene was inserted between TEV P1 and HC-pro, and the heterologous protein was released by polyprotein proteolysis mediated by P1 alone or in combination with NIa-pro (Dolja, McBride and Carrington 1992, Carrington et al. 1993). The same approach was used successfully in potyvirids encoding Type-B P1. GUS or fluorescent proteins were expressed using viral vectors derived from WSMV (Tritimovirus) (Choi et al. 2002, Tatineni et al. 2011), and TriMV

(*Poacevirus*) (Tatineni *et al.* 2015). The 2A 'self-cleaving' peptides of FMDV or *Thosea asigna* virus were applied to engineer NIa-pro independent processing of potyvirid polyproteins (Tatineni *et al.* 2011, Pasin, Simón-Mateo and García 2014).

The ClYVV P1/HC-pro junction was engineered for coexpression of multiple heterologous proteins that were released by P1 and NIa-pro proteolysis (Masuta *et al.* 2000). More recently, ClYVV was used for plant overexpression of a gibberellin catabolic enzyme inserted between P1 and HC-pro (Fig. 12B). Infections of pea and broad bean plants with the recombinant ClYVV conferred dwarfism, an agronomically important trait (Torti *et al.* 2021). Traditional plant breeding is time- and cost-consuming, and innovative strategies are needed for accelerated and tailored crop trait manipulation (Steinwand and Ronald 2020, French *et al.* 2021). Transient, viral-mediated manipulation of plant size and other agronomic performance traits holds promise to become a new standard for fast, flexible crop reprogramming.

The NIb/CP polyprotein junction is an insertion site used for heterologous gene expression that mimics the natural HAM1 location in potyvirids (Fig. 2). Simultaneous insertions at the P1/HCpro and NIb/CP junctions allowed production of two recombinant proteins from a single potyviral vector (Beauchemin, Bougie and Laliberté 2005). Consistent with the AlkB location found in potyvirids (Fig. 2), a new insertion site suitable for heterologous gene expression was identified within the P1 N terminus (Fig. 12A). It was used alone or in combination with inserts at the P1/HC-pro and NIb/CP junctions for production of up to three recombinant proteins (Rajamäki et al. 2005, Kelloniemi, Mäkinen and Valkonen 2008). Heterologous protein expression has been reported by gene insertion upstream of TEV P1 (Fig. 12A). This strategy allowed correct targeting of heterologous proteins to subcellular compartments (Majer, Navarro and Daròs 2015), and was used for metabolite production by potyvirus-mediated enzyme delivery to chloroplasts (Martí et al. 2020).

In addition to protein overexpression, heterologous sequences inserted within P1 or at the P1/HC-pro junction can trigger silencing of plant homologs (Gammelgård, Mohan and Valkonen 2007, Xie *et al.* 2021). Potyvirid vectors have been used for virus-induced gene silencing, as well as for simultaneous plant gene silencing and heterologous protein production (Gammelgård, Mohan and Valkonen 2007, Tuo *et al.* 2021, Xie *et al.* 2021).

Non-core modules—untapped synthetic biology resources

Given their stringent specificity and orthogonality, potyvirid proteinases have been engineered for commercial purposes as well as for synthetic biology applications to control cellular functions (Chung and Lin 2020, Dyer and Weiss 2021). These proteins have been integrated into synthetic signaling pathways with designs that included induction of degron-dependent protein depletion, autoinhibition release of transcription regulators, and enzyme reconstitution through dimerization inhibition or activation (Fernandez-Rodriguez and Voigt 2016, Gao *et al.* 2018, Fink *et al.* 2019). Use of potyvirid leader proteinases in synthetic genetic circuitries has not yet been reported. Given its activation requirements and strict cis-cleavage activity, Type-A P1 could nonetheless be an appealing choice for biodesigns with high host specificity or biocontainment levels.

Synthetic, tight control over protein activity can be achieved by destabilizing tags, oligomerization domains, inhibitory modules, or subcellar sequestration signals (Alberstein, Guo and Kortemme 2021, Chen and Elowitz 2021). Type-A P1 was shown to undergone

rapid degradation in plants and to inhibit activity of downstream fusion partners, such as HC-Pro or GUS (Verchot and Carrington 1995, Martínez and Daròs 2014, Pasin, Simón-Mateo and García 2014, Shan *et al.* 2015), and could be repurposed for conditional, fine-tuned activation of recombinant proteins.

HC-pro and other RNA silencing suppressors from plant viruses are used routinely to enhance protein yields of plant transient expression systems (Csorba, Kontra and Burgyán 2015, Sainsbury 2020); Type-B proteins and P1N-PISPO could be also useful in similar applications. In-depth characterization of potyvirid AlkB and ITPase activities may also lead to novel tools for epigenetic or metabolic engineering applications.

Virus-directed continuous evolution has been used to obtain biomolecules with improved or new functions in prokaryotic and mammalian systems (Morrison, Podracky and Liu 2020), but suitable methods are lacking for plants. Experimental studies aiming to evaluated the evolutionary fate of sequences inserted in potyvirid genomes have revealed constraints linked to the evolutionary time, as well as genome position and specific insert features (Willemsen *et al.* 2016, 2017, Willemsen and Zwart 2019). A potyviral reverse genetic system was nonetheless engineered for forced evolution of P1 proteins (Rodamilans, Casillas and García 2021).

Leader proteinases—overlooked targets for antiviral strategies

Human viruses are targeted by proteinase inhibitor therapies to a clinically useful level (Agbowuro et al. 2018); yet the use of similar antiviral strategies for plant virus control is lagging. Use of protease inhibitors for potyvirus control has shown limited success so far (Gutierrez-Campos et al. 1999), but new promising antiviral strategies have been reported. A plant protein involved in bacterial immunity was successfully repurposed to specifically sense NIa-pro and trigger antiviral cell death (Kim et al. 2016). This synthetic antiviral system has been implemented in soybean, and further optimized for enhanced control of the NIa-pro-induced cell necrosis (Helm et al. 2019, Pottinger et al. 2020). Potyvirid leader proteinases are attractive antiviral targets, since P1 and HC-pro defects preclude infectivity (Kasschau and Carrington 1995, Verchot and Carrington 1995, Pasin, Simón-Mateo and García 2014, Shan et al. 2015). A zucchini yellow mosaic virus isolate with reduced HC-pro silencing suppressor activity has been registered since 2007 for the US market as a cross-protection agent of cucurbits (U.S. Environmental Protection Agency 2007). RNA silencing transgenic approaches that target P1 or HC-pro confer potyvirid resistance in crops (Di Nicola-Negri et al. 2005).

Investigation of non-core module roles in plant-potyvirid interactions recently allowed identification of new host factors and signaling pathways that could be exploited in antiviral strategies. High abscisic acid (ABA) levels were found to accumulate during infection of a PPV mutant having a truncated P1 (Fig. 10B); the finding prompted evaluation of ABA effects on infection. Defects of the cap-binding complex components ABA HYPERSENSI-TIVE1/CAP BINDING PROTEIN 80 (ABH1/CBP80) and CAP BIND-ING PROTEIN 20 (CBP20) are known to confer ABA hypersensitivity and were shown to significantly delay PPV infection (Pasin et al. 2020). Cap-binding complex contribution in antiviral defense was reported in other organisms, including insects and mammals (Gebhardt et al. 2019, Blagrove and Barribeau 2021). ABA treatments promote resistance to PPV, and possibly to other potyvirids (Alazem, Widyasari and Kim 2019, Zhang et al. 2019, Pasin et al. 2020, Chiu et al. 2021). Rapid catabolism, photolability, and chemical instability make ABA unsuited for agricultural purposes.

Availability of synthetic ABA receptor agonists with high stability and binding affinities nonetheless paves the way for crop antiviral strategies based on chemical manipulation of ABA signaling (Hewage *et al.* 2020).

Research outlooks and conclusions

The phylum Pisuriviricota includes extremely diversified RNA viruses whose radiation was proposed to be concomitant with key eukaryogenesis events (Koonin *et al.* 2008). Potyviridae is currently the largest family of Riboviria (Fig. 1A), yet thousands of novel RNA viruses await accommodation in recognized taxa (Callanan *et al.* 2020, Edgar *et al.* 2022).

Gene gain and loss, specialization, and de novo emergence have promoted the diversification of leader layouts of Potyviridae (Figs 2 and 9), as well as of divergent RNA viruses of plants and animals, e.g. closteroviruses, picornaviruses and arteriviruses (Dolja, Kreuze and Valkonen 2006, Valli, López-Moya and García 2007, Agol and Gmyl 2010, Gorbalenya and Lauber 2010, Gulyaeva et al. 2017, Zell 2018). Functional expansion of a polyprotein core through domain gain is hypothesized to have taken part in the evolutionary transition from plastroviruses to modern potyvirids (Lauber et al. 2019). We point out that evolution of potyvirid noncore domains is diverse and can potentially be traced to multiple or single acquisition events (see AlkB in Potyvirus and Brambyvirus, or the pseudo TMV-like CP in Bymovirus, respectively; Figs 5 and 7), recombination and retention of functionally divergent homologs (P1 tandem in Ipomovirus; Fig. 3), as well as emergence of a new, overlapping protein module through overprinting (PISPO in Potyvirus). In-depth database search and sequence analyses uncovered a putative HC-pro-like domain within Celavirus (Fig. 8C), as well as the presence of the ITPase fold (HAM1, a former oddity of a narrow group of potyvirids) in taxonomically divergent RNA and DNA viruses (Fig. 6).

Identification of factors that interact with potyvirid non-core proteins and elucidation of host perturbations linked to their functional alteration are indeed major research priorities for dissecting their niche adaptation roles. Complete kinetic models were described for RNA viruses that share with potyvirids similar genome replication and protein expression strategies (Zitzmann et al. 2020, Lopacinski et al. 2021). Mathematical models could provide a quantitative understanding of the complex dynamics that regulate potyvirid replication as well as host immune responses and viral counterstrategies (Pasin et al. 2020). Accessory genes shape cellular pangenome diversity and are enriched in plant-microbe interaction determinants (Box 1), and models have been developed to describe pangenome gene content variation (Domingo-Sananes and McInerney 2021). Can empirical data from plant-potyvirid systems contribute to theoretical frameworks for understanding cellular pangenome evolution and ecological niche adaptation?

Functional redundancy of non-core modules has allowed the establishment of relationships between potyviral HC-pro and the non-canonical silencing suppressors P1N-PISPO and Type-B P1. HC-pro counteracts RNA decay antiviral defenses and associates with RNA turnover components for infection enhancement (Li and Wang 2018, De *et al.* 2020). Are these and additional HC-pro activities performed by other non-core modules? P1, HC-pro, and HAM1 are involved in symptom development (Valli *et al.* 2018, Tomlinson *et al.* 2019, Pasin *et al.* 2020). Is there any mechanistic connection between these otherwise structurally unrelated modules? Members of Nidovirales, the largest known RNA viruses, have evolved proofreading replication for maintaining in-

tegrity of genomes that can reach ~40 kb (Robson et al. 2020). Potyvirids are among the largest plant viruses and have unusually low mutation rates estimated to be in the range 10^{-5} – 10^{-6} mutations/site/generation (Sanjuán et al. 2010, Tromas and Elena 2010). Are functions carried out by AlkB, HAM1, or other modules conditioning the potyvirid evolution rates? Answers to these questions will assist in the better understanding contribution of the noncore proteome expansion in the Potyviridae evolutionary radiation and RNA virus evolution.

Finally, driven by advances in high-throughput sequencing technologies and easy access to underexplored geographical and ecological areas (Villamor *et al.* 2019, Maclot *et al.* 2020, Sommers *et al.* 2021), discovery of new potyvirids and poty-like ancestors with unusual genomic organization, atypical protein modules, and niche-optimized traits is likely to be further expanded in the near future (Lauber *et al.* 2019, Wolf *et al.* 2020). Extending efforts for potyvirid discovery and proteome functional characterization would improve understanding of non-core module roles in host adaptation evolution to eventually guide design of novel antiviral strategies and synthetic biology solutions.

Box 1. Non-core genes in plant-microbe interactions

Host niche adaptation is a major driving force of virus evolution (Simmonds, Aiewsakun and Katzourakis 2019), and functional characterization of the Potyviridae non-core modules supports their roles in symptom development and antiviral immunity evasion (main text). Pangenomes are increasingly used to represent known structural variants of cellular taxa, wherein the adaptive nature of accessory and rare genes is subject to debate (Domingo-Sananes and McInerney 2021, Coelho et al. 2022). Recent data nonetheless highlight the critical contribution of gene content variation in the evolution of plantmicrobe dynamics and the genetic potential of holobionts (Badet and Croll 2020, Zilber-Rosenberg and Rosenberg 2021). Plant pathogenicity and adaptation factors are components of accessory genomes, lineage-specific replicons or chromosomes of bacterial and fungal species (Ma et al. 2010, Levy et al. 2017, Laflamme et al. 2020, Langner et al. 2021, Chou et al. 2022). Among plants, non-core genes of Arabidopsis thaliana, rice, rapeseed, cabbage, sunflower, and wheat are known actuators of the host-pathogen warfare (Zhao et al. 2018, Hübner et al. 2019, Van de Weyer et al. 2019, Bayer et al. 2020, Upadhyaya et al. 2021).

Supplementary data

Supplementary data are available at FEMSRE online.

Author contribution

FP conceived the work, performed data analyses, wrote the manuscript and prepared the figures; J-AD, and IET collaborated in the manuscript preparation. All authors revised and approved the final version.

Acknowledgements

The authors are grateful to AE Gorbalenya and AJ Gibbs for their comments, and to C Mark for editorial assistance. Computational resources from Galaxy are acknowledged.

Funding

FP is the recipient of a 'Juan de la Cierva Incorporación' contract [grant IJC2019-039970-I] from Ministerio de Ciencia e Innovación (Spain). J-AD is funded by Ministerio de Ciencia e Innovación (Spain), and co-financed by European Regional Development Fund [grant PID2020-114691RB-I00]. IET is funded from crop commodity boards, the berry and ornamental industry, the United States Department of Agriculture and the National Science Foundation (USA).

Conflicts of interest. None declared.

References

- Adams MJ, Antoniw JF, Beaudoin F. Overview and analysis of the polyprotein cleavage sites in the family Potyviridae. Mol Plant Pathol 2005;6:471–87.
- Agbowuro AA, Huston WM, Gamble AB et al. Proteases and protease inhibitors in infectious diseases. Med Res Rev 2018;**38**:1295–331.
- Agol VI, Gmyl AP. Viral security proteins: counteracting host defences. Nat Rev Microbiol 2010;**8**:867–78.
- Alazem M, Widyasari K, Kim K-H. An avirulent strain of soybean mosaic virus reverses the defensive effect of abscisic acid in a susceptible soybean cultivar. Viruses 2019;**11**:E87.
- Alberstein RG, Guo AB, Kortemme T. Design principles of protein switches. Curr Opin Struct Biol 2022;**72**:71–8.
- Alicai T, Ndunguru J, Sseruwagi P *et al.* Cassava brown streak virus has a rapidly evolving genome: implications for virus speciation, variability, diagnosis and host resistance. Sci Rep 2016;**6**:36164.
- Atallah OO, Kang S-H, El-Mohtar CA et al. A 5'-proximal region of the Citrus tristeza virus genome encoding two leader proteases is involved in virus superinfection exclusion. Virology 2016;**489**: 108–15.
- Aulia A, Hyodo K, Hisano S *et al.* Identification of an RNA silencing suppressor encoded by a symptomless fungal hypovirus, Cryphonectria hypovirus 4. *Biology* 2021;**10**:100.
- Badet T, Croll D. The rise and fall of genes: origins and functions of plant pathogen pangenomes. Curr Opin Plant Biol 2020;56:65–73.
- Bayer PE, Golicz AA, Scheben A et al. Plant pan-genomes are the new reference. Nat Plants 2020;**6**:914–20.
- Beauchemin C, Bougie V, Laliberté J-F. Simultaneous production of two foreign proteins from a potyvirus-based vector. Virus Res 2005;112:1–8.
- Bin Jang H, Bolduc B, Zablocki O et al. Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. Nat Biotechnol 2019;**37**:632–9.
- Blagrove M, Barribeau SM. Immune priming can prevent WNV establishment in *Culex quinquefasciatus* mosquitoes: evidence for immune priming based reversal of WNV-mediated immune suppression. *bioRxiv* 2021. DOI: 10.1101/2021.05.05.442826.
- Brady A, Felipe-Ruiz A, Gallego Del Sol F et al. Molecular basis of lysislysogeny decisions in Gram-positive phages. Annu Rev Microbiol 2021;75:563–81.
- Bratlie MS, Drabløs F. Bioinformatic mapping of AlkB homology domains in viruses. BMC *Genomics* 2005;**6**:1.
- Callanan J, Stockdale SR, Shkoporov A *et al*. Expansion of known ssRNA phage genomes: from tens to over a thousand. *Sci Adv* 2020;**6**:eaay5981.
- Carpino C, Ferriol Safont I, Elvira-González L et al. RNA2-encoded VP37 protein of *Broad bean wilt virus* 1 is a determinant of pathogenicity, host susceptibility, and a suppressor of posttranscriptional gene silencing. Mol Plant Pathol 2020;**21**:1421–35.

- Carrington JC, Haldeman R, Dolja VV *et al.* Internal cleavage and *trans*-proteolytic activities of the VPg-proteinase (NIa) of tobacco etch potyvirus in vivo. *J Virol* 1993;**67**:6995–7000.
- Cassandri M, Smirnov A, Novelli F et al. Zinc-finger proteins in health and disease. Cell Death Discovery 2017;**3**:17071.
- Charon J, Barra A, Walter J *et al.* First experimental assessment of protein intrinsic disorder involvement in an RNA virus natural adaptive process. *Mol Biol Evol* 2018;**35**:38–49.
- Chen C-C, Chen T-C, Raja JAJ et al. Effectiveness and stability of heterologous proteins expressed in plants by *Turnip mosaic virus* vector at five different insertion sites. *Virus Res* 2007;**130**:210–27.
- Chen J-S, Liang S-S, Sun S-R et al. Diverse conserved domains and a positively selected site in the sugarcane streak mosaic virus P1 protein are essential for RNA silencing suppression and protein stability. Plant Pathol 2020;**69**:1390–400.
- Chen Z, Elowitz MB. Programmable protein circuit design. Cell 2021;**184**:2284–301.
- Chiu Y-H, Hung Y-L, Wang H-P et al. Investigation of P1/HC-Promediated ABA/calcium signaling responses via gene silencing through high- and low-throughput RNA-seq approaches. Viruses 2021;13:2349.
- Choi D, Shin C, Shirasu K et al. Two novel poty-like viruses identified from the transcriptome data of purple witchweed (Striga hermonthica). Acta Virol 2022;**65**:365–72.
- Choi I-R, French R, Hein GL et al. Fully biologically active in vitro transcripts of the eriophyid mite-transmitted wheat streak mosaic tritimovirus. *Phytopathology* 1999;**89**:1182–5.
- Choi I-R, Horken KM, Stenger DC et al. Mapping of the P1 proteinase cleavage site in the polyprotein of Wheat streak mosaic virus (genus Tritimovirus). J Gen Virol 2002;**83**:443–50.
- Chou L, Lin Y-C, Haryono M *et al*. Modular evolution of secretion systems and virulence plasmids in a bacterial species complex. *BMC Biol* 2022;**20**:16.
- Chung HK, Lin MZ. On the cutting edge: protease-based methods for sensing and controlling cell biology. Nat Methods 2020;17:885–96.
- Clark CA, Davis JA, Abad JA et al. Sweetpotato viruses: 15 years of progress on understanding and managing complex diseases. Plant Disease 2012;**96**:168–85.
- Clavel M, Lechner E, Incarbone M et al. Atypical molecular features of RNA silencing against the phloem-restricted polerovirus TuYV. *Nucleic Acids Res* 2021;**49**:11274–93.
- Coelho LP, Alves R, Del Río ÁR et al. Towards the biogeography of prokaryotic genes. Nature 2022;**601**:252–6.
- Corley M, Burns MC, Yeo GW. How RNA-binding proteins interact with RNA: molecules and mechanisms. *Mol Cell* 2020;**78**:9–29.
- Csorba T, Kontra L, Burgyán J. Viral silencing suppressors: tools forged to fine-tune host-pathogen coexistence. Virology 2015;479-480:85–103.
- Cui C, Wang J-J, Zhao J-H et al. A Brassica miRNA regulates plant growth and immunity through distinct modes of action. *Mol Plant* 2020;**13**:231–45.
- De S, Pollari M, Varjosalo M *et al.* Association of host protein VARICOSE with HCPro within a multiprotein complex is crucial for RNA silencing suppression, translation, encapsidation and systemic spread of potato virus A infection. *PLoS Pathog* 2020;**16**:e1008956.
- Deeg CM, Chow C-ET, Suttle CA. The kinetoplastid-infecting Bodo saltans virus (BsV), a window into the most abundant giant viruses in the sea. *eLife* 2018;**7**:e33014.
- Del Toro F, Sun H, Robinson C et al. In planta vs viral expression of HCPro affects its binding of non-plant 21-22 nt small RNAs, but not its preference for 5'-terminal adenines, or its effects

on small RNA methylation. New Phytol 2022;**233**:2266–81. DOI: 10.1111/nph.17935.

- Desbiez C, Chandeysson C, Lecoq H et al. A simple, rapid and efficient way to obtain infectious clones of potyviruses. J Virol Methods 2012;**183**:94–7.
- Desbiez C, Schoeny A, Maisonneuve B *et al.* Molecular and biological characterization of two potyviruses infecting lettuce in southeastern France. *Plant Pathol* 2017;**66**:970–9.
- Desbiez C, Verdin E, Tepfer M et al. Characterization of a new cucurbit-infecting ipomovirus from Sudan. Arch Virol 2016;**161**:2913–5.
- Dessens JT, Meyer M. Identification of structural similarities between putative transmission proteins of *Polymyxa* and *Spongospora* transmitted bymoviruses and furoviruses. Virus Genes 1996;**12**:95–9.
- Di Nicola-Negri E, Brunetti A, Tavazza M et al. Hairpin RNA-mediated silencing of Plum pox virus P1 and HC-Pro genes for efficient and predictable resistance to the virus. *Transgenic Res* 2005;**14**:989–94.
- Dolja VV, Kreuze JF, Valkonen JPT. Comparative and functional genomics of closteroviruses. Virus Res 2006;**117**:38–51.
- Dolja VV, Krupovic M, Koonin EV. Deep roots and splendid boughs of the global plant virome. *Annu Rev Phytopathol* 2020;**58**:23–53.
- Dolja VV, McBride HJ, Carrington JC. Tagging of plant potyvirus replication and movement by insertion of β -glucuronidase into the viral polyprotein. Proc Natl Acad Sci 1992;**89**:10208–12.
- Dombrovsky A, Reingold V, Antignus Y. *Ipomovirus* an atypical genus in the family Potyviridae transmitted by whiteflies. *Pest Manage Sci* 2014;**70**:1553–67.
- Domier LL, Franklin KM, Hunt AG et al. Infectious in vitro transcripts from cloned cDNA of a potyvirus, tobacco vein mottling virus. Proc Natl Acad Sci 1989;**86**:3509–13.
- Domingo-Calap ML, Chase O, Estapé M et al. The P1 protein of Watermelon mosaic virus compromises the activity as RNA silencing suppressor of the P25 protein of Cucurbit yellow stunting disorder virus. Front in Microbiol 2021;**12**:645530.
- Domingo-Sananes MR, McInerney JO. Mechanisms that shape microbial pangenomes. Trends Microbiol 2021;29:493–503.
- Drappier M, Jha BK, Stone S et al. A novel mechanism of RNase L inhibition: Theiler's virus L* protein prevents 2-5A from binding to RNase L. PLoS Pathog 2018;**14**:e1006989.
- Dulin D, Vilfan ID, Berghuis BA *et al.* Elongation-competent pauses govern the fidelity of a viral RNA-dependent RNA polymerase. *Cell Rep* 2015;**10**:983–92.
- Dyer RP, Weiss GA. Making the cut with protease engineering. Cell Chem Biol 2021: S2451-9456(21)00515-8.
- Edgar RC, Taylor J, Lin V et al. Petabase-scale sequence alignment catalyses viral discovery. *Nature* **602**, 2022;142–7. DOI: 10.1038/s41586-021-04332-2.
- Fedeles BI, Singh V, Delaney JC et al. The AlkB family of $Fe(II)/\alpha$ ketoglutarate-dependent dioxygenases: repairing nucleic acid alkylation damage and beyond. J Biol Chem 2015;**290**:20734–42.
- Fernandez-Rodriguez J, Voigt CA. Post-translational control of genetic circuits using Potyvirus proteases. Nucleic Acids Res 2016;44:6493–502.
- Fink T, Lonzarić J, Praznik A et al. Design of fast proteolysis-based signaling and logic circuits in mammalian cells. Nat Chem Biol 2019;15:115–22.
- Frei T, Khammash M. Adaptive circuits in synthetic biology. Curr Opin in Syst Biol 2021;**28**:100399.
- French E, Kaplan I, Iyer-Pascuzzi A et al. Emerging strategies for precision microbiome management in diverse agroecosystems. Nat Plants 2021;7:256–67.

- Freundt EC, Drappier M, Michiels T. Innate immune detection of cardioviruses and viral disruption of interferon signaling. Front Microbiol 2018;9:2448.
- Gammelgård E, Mohan M, Valkonen JPT. Potyvirus-induced gene silencing: the dynamic process of systemic silencing and silencing suppression. J Gen Virol 2007;88:2337–46.
- Gao XJ, Chong LS, Kim MS et al. Programmable protein circuits in living cells. Science 2018;**361**:1252–8.
- Gao Y, Cui Y, Fox T et al. Structures and operating principles of the replisome. Science 2019;**363**:eaav7003.
- Garcia-Ruiz H, Carbonell A, Hoyer JS et al. Roles and programming of Arabidopsis ARGONAUTE proteins during Turnip mosaic virus infection. PLoS Pathog 2015;11:e1004755.
- Gebhardt A, Bergant V, Schnepf D et al. The alternative cap-binding complex is required for antiviral defense in vivo. PLoS Pathog 2019;15:e1008155.
- Gibbs AJ, Hajizadeh M, Ohshima K *et al*. The potyviruses: an evolutionary synthesis is emerging. Viruses 2020;**12**:132.
- Giner A, Lakatos L, García-Chapa M *et al*. Viral protein inhibits RISC activity by Argonaute binding through conserved WG/GW motifs. *PLoS Pathog* 2010;**6**:e1000996.
- Gitlin L, Hagai T, LaBarbera A *et al.* Rapid evolution of virus sequences in intrinsically disordered protein regions. *PLoS Pathog* 2014;**10**:e1004529.
- Gohara DW, Di Cera E. Allostery in trypsin-like proteases suggests new therapeutic strategies. *Trends Biotechnol* 2011;**29**:577–85.
- Gorbalenya AE, Koonin EV, Lai MM. Putative papain-related thiol proteases of positive-strand RNA viruses. Identification of rubiand aphthovirus proteases and delineation of a novel conserved domain associated with proteases of rubi-, alpha- and coronaviruses. FEBS Lett 1991;**288**:201–5.
- Gorbalenya AE, Lauber C. Origin and evolution of the Picornaviridae proteome. In: Ehrenfeld E Domingo E Roos RP (eds.), *The Picornaviruses*. Washington, DC, USA: ASM Press, 2010, 253–70.
- Gulyaeva A, Dunowska M, Hoogendoorn E et al. Domain organization and evolution of the highly divergent 5' coding region of genomes of arteriviruses, including the novel possum nidovirus. J Virol 2017;91:e02096–16.
- Guo B, Lin J, Ye K. Structure of the autocatalytic cysteine protease domain of potyvirus helper-component proteinase. J Biol Chem 2011;286:21937–43.
- Gupta AK, Tatineni S. RNA silencing suppression mechanisms of Triticum mosaic virus P1: dsRNA binding property and mapping functional motifs. Virus Res 2019a;269:197640.
- Gupta AK, Tatineni S. Wheat streak mosaic virus P1 binds to dsRNAs without size and sequence specificity and a GW motif is crucial for suppression of RNA silencing. Viruses 2019b;**11**:472.
- Gutierrez-Campos R, Torres-Acosta JA, Saucedo-Arias LJ et al. The use of cysteine proteinase inhibitors to engineer resistance against potyviruses in transgenic tobacco plants. Nat Biotechnol 1999;17:1223–6.
- Han M, Yoo D. Modulation of innate immune signaling by nonstructural protein 1 (nsp1) in the family Arteriviridae. Virus Res 2014;**194**:100–9.
- Hedstrom L. Serine protease mechanism and specificity. Chem Rev 2002;102:4501–24.
- Helm M, Qi M, Sarkar S et al. Engineering a decoy substrate in soybean to enable recognition of the soybean mosaic virus NIa protease. Mol Plant Microbe Interact 2019;**32**:760–9.
- Hewage KAH, Yang J-F, Wang D *et al*. Chemical manipulation of abscisic acid signaling: a new approach to abiotic and biotic stress management in agriculture. *Adv Sci* 2020;**7**:2001265.

- Holmes EC, Goldstein SA, Rasmussen AL et al. The origins of SARS-CoV-2: a critical review. Cell 2021;**184**:4848–56.
- Hübner S, Bercovich N, Todesco M et al. Sunflower pan-genome analysis shows that hybridization altered gene content and disease resistance. Nat Plants 2019;**5**:54–62.
- Innan H, Kondrashov F. The evolution of gene duplications: classifying and distinguishing between models. *Nat Rev Genet* 2010;**11**: 97–108.
- International Committee on Taxonomy of Viruses 2020. ICTV master species list 2019.v1 (MSL #35). https://talk.ictvonline.org/files/m aster-species-lists/m/msl/9601 (June 2021, date last accessed).
- Ismayil A, Yang M, Haxim Y et al. Cotton leaf curl Multan virus βC1 protein induces autophagy by disrupting the interaction of autophagy-related protein 3 with glyceraldehyde-3-phosphate dehydrogenases. Plant Cell 2020;**32**:1124–35.
- Ivanov KI, Eskelin K, Bašić M et al. Molecular insights into the function of the viral RNA silencing suppressor HCPro. Plant J 2016; 85:30–45.
- Jayasinghe WH, Kim H, Sasaki J et al. Aphid transmissibility of onion yellow dwarf virus isolates with an N-terminal truncated HC-Pro is aided by leek yellow stripe virus. J Gen Plant Pathol 2021;**87**: 178–83.
- Jeudy S, Rigou S, Alempic J-M et al. The DNA methylation landscape of giant viruses. Nat Commun 2020;**11**:2657.
- Ji L, Chen X. Regulation of small RNA stability: methylation and beyond. Cell Res 2012;**22**:624–36.
- Jiang C, Kan J, Ordon F et al. Bymovirus-induced yellow mosaic diseases in barley and wheat: viruses, genetic resistances and functional aspects. Theor Appl Genet 2020;133:1623–40.
- Jo WK, van Elk C, van de Bildt M *et al*. An evolutionary divergent pestivirus lacking the N^{pro} gene systemically infects a whale species. *Emerg Microbes Infect* 2019;**8**:1383–92.
- Jones RAC, Naidu RA. Global dimensions of plant virus diseases: current status and future perspectives. Annu Rev Virol 2019;6: 387–409.
- Kang S-H, Atallah OO, Sun Y-D et al. Functional diversification upon leader protease domain duplication in the citrus tristeza virus genome: role of RNA sequences and the encoded proteins. Virology 2018;514:192–202.
- Kannan M, Zainal Z, Ismail I et al. Application of reverse genetics in functional genomics of potyvirus. Viruses 2020;**12**:803.
- Kasschau KD, Carrington JC. Requirement for HC-Pro processing during genome amplification of tobacco etch potyvirus. Virology 1995;209:268–73.
- Kazibwe Z, Soto-Burgos J, MacIntosh GC et al. TOR mediates the autophagy response to altered nucleotide homeostasis in an RNase mutant. J Exp Bot 2020;71:6907–20.
- Kelloniemi J, Mäkinen K, Valkonen JPT. Three heterologous proteins simultaneously expressed from a chimeric potyvirus: infectivity, stability and the correlation of genome and virion lengths. Virus Res 2008;**135**:282–91.
- Kenesi E, Carbonell A, Lózsa R et al. A viral suppressor of RNA silencing inhibits ARGONAUTE 1 function by precluding target RNA binding to pre-assembled RISC. Nucleic Acids Res 2017;45: 7736–50.
- Khakhar A, Voytas DF. RNA viral vectors for accelerating plant synthetic biology. Front Plant Sci 2021;**12**:668580.
- Kiljunen S, Hakala K, Pinta E et al. Yersiniophage φR1-37 is a tailed bacteriophage having a 270 kb DNA genome with thymidine replaced by deoxyuridine. Microbiology 2005;151:4093–102.
- Kim SH, Qi D, Ashfield T *et al.* Using decoys to expand the recognition specificity of a plant disease resistance protein. *Science* 2016;**351**:684–7.

- Knierim D, Menzel W, Winter S. Analysis of the complete genome sequence of euphorbia ringspot virus, an atypical member of the genus Potyvirus. Arch Virol 2017;**162**:291–3.
- Kondo T, Fujita T. Complete nucleotide sequence and construction of an infectious clone of Chinese yam necrotic mosaic virus suggest that macluraviruses have the smallest genome among members of the family *Potyviridae*. Arch Virol 2012;**157**:2299–307.
- Koonin EV, Choi GH, Nuss DL et al. Evidence for common ancestry of a chestnut blight hypovirulence-associated double-stranded RNA and a group of positive-strand RNA plant viruses. Proc Natl Acad Sci 1991;88:10647–51.
- Koonin EV, Dolja VV, Krupovic M et al. Global organization and proposed megataxonomy of the virus world. Microbiol Mol Biol Rev 2020;**84**:e00061–19.
- Koonin EV, Wolf YI, Nagasaki K et al. The Big Bang of picornalike virus evolution antedates the radiation of eukaryotic supergroups. Nat Rev Microbiol 2008;6:925–39.
- Kotas ME, Medzhitov R. Homeostasis, inflammation, and disease susceptibility. Cell 2015;**160**:816–27.
- Križnik M, Baebler Š, Gruden K. Roles of small RNAs in the establishment of tolerant interaction between plants and viruses. Curr Opin Virol 2020;42:25–31.
- Lacombe S, Bangratz M, Vignols F *et al*. The rice yellow mottle virus P1 protein exhibits dual functions to suppress and activate gene silencing. *Plant J* 2010;**61**:371–82.
- Laflamme B, Dillon MM, Martel A *et al*. The pan-genome effectortriggered immunity landscape of a host-pathogen interaction. Sci*ence* 2020;**367**:763–8.
- Langner T, Harant A, Gomez-Luciano LB et al. Genomic rearrangements generate hypervariable mini-chromosomes in hostspecific isolates of the blast fungus. PLoS Genet 2021;**17**:e1009386.
- Lauber C, Seifert M, Bartenschlager R et al. Discovery of highly divergent lineages of plant-associated astro-like viruses sheds light on the emergence of potyviruses. Virus Res 2019;**260**:38–48.
- Le Lay C, Shi M, Buček A *et al*. Unmapped RNA virus diversity in termites and their symbionts. *Viruses* 2020;**12**:1145.
- Lefeuvre P, Martin DP, Elena SF et al. Evolution and ecology of plant viruses. Nat Rev Microbiol 2019;**17**:632–44.
- Leiva AM, Jimenez J, Sandoval H *et al*. Complete genome sequence of a novel secovirid infecting cassava in the Americas. *Arch Virol* 2022;**167**:665–8.
- Levy A, Salas Gonzalez I, Mittelviefhaus M et al. Genomic features of bacterial adaptation to plants. Nat Genet 2018;**50**:138–50.
- Li F, Wang A. RNA decay is an antiviral defense in plants that is counteracted by viral RNA silencing suppressors. *PLoS Pathog* 2018;**14**:e1007228.
- Li F, Wang A. RNA-targeted antiviral immunity: more than just RNA silencing. *Trends Microbiol* 2019;**27**:792–805.
- Liang G, Bushman FD. The human virome: assembly, composition and host interactions. Nat Rev Microbiol 2021;**19**,514–27.
- Lin M-K, Belanger H, Lee Y-J *et al.* FLOWERING LOCUS T protein may act as the long-distance florigenic signal in the cucurbits. *Plant Cell* 2007;**19**:1488–506.
- Llorente B, Torres-Montilla S, Morelli L *et al*. Synthetic conversion of leaf chloroplasts into carotenoid-rich plastids reveals mechanistic basis of natural chromoplast development. *Proc Natl Acad Sci* 2020;**117**:21796–803.
- Lopacinski AB, Sweatt AJ, Smolko CM et al. Modeling the complete kinetics of coxsackievirus B3 reveals human determinants of hostcell feedback. Cell Syst 2021;**12**:304–23.
- Louca S, Polz MF, Mazel F et al. Function and functional redundancy in microbial systems. Nat Ecol Evol 2018;**2**:936–43.

Lucía-Sanz A, Manrubia S. Multipartite viruses: adaptive trick or evolutionary treat? NPJ Syst Biol Appl 2017;**3**:34.

- Lulla V, Dinan AM, Hosmillo M et al. An upstream protein-coding region in enteroviruses modulates virus infection in gut epithelial cells. Nat Microbiol 2019;**4**:280–92.
- Lunney JK, Fang Y, Ladinig A et al. Porcine reproductive and respiratory syndrome virus (PRRSV): pathogenesis and interaction with the immune system. Annu Rev Anim Biosci 2016;4: 129–54.
- Ma L-J, van der Does HC, Borkovich KA et al. Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 2010;**464**:367–73.
- Maclot F, Candresse T, Filloux D *et al*. Illuminating an ecological blackbox: using high throughput sequencing to characterize the plant virome across scales. *Front Microbiol* 2020;**11**:578064.
- Majer E, Navarro J-A, Daròs J-A. A potyvirus vector efficiently targets recombinant proteins to chloroplasts, mitochondria and nuclei in plant cells when expressed at the amino terminus of the polyprotein. Biotechnol J 2015;**10**:1792–802.
- Maliogka VI, Salvador B, Carbonell A *et al*. Virus variants with differences in the P1 protein coexist in a *Plum pox virus* population and display particular host-dependent pathogenicity features. *Mol Plant Pathol* 2012;**13**:877–86.
- Mann K, Sanfaçon H. Expanding repertoire of plant positive-strand RNA virus proteases. Viruses 2019;**11**:66.
- Martí M, Diretto G, Aragonés V et al. Efficient production of saffron crocins and picrocrocin in Nicotiana benthamiana using a virusdriven system. Metab Eng 2020;**61**:238–50.
- Martínez F, Daròs J-A. *Tobacco etch virus* protein P1 traffics to the nucleolus and associates with the host 60S ribosomal subunits during infection. J Virol 2014;**88**:10725–37.
- Martínez-Pérez M, Aparicio F, López-Gresa MP et al. Arabidopsis m⁶A demethylase activity modulates viral infection of a plant virus and the m⁶A abundance in its genomic RNAs. Proc Natl Acad Sci 2017;**114**:10755–60.
- Martínez-Pérez M, Gómez-Mena C, Alvarado-Marchena L et al. The m⁶A RNA demethylase ALKBH9B plays a critical role for vascular movement of alfalfa mosaic virus in Arabidopsis. Front Microbiol 2021;**12**:745576.
- Masuta C, Yamana T, Tacahashi Y *et al*. Development of clover yellow vein virus as an efficient, stable gene-expression system for legume species. *Plant J* 2000;**23**:539–46.
- Mbanzibwa DR, Tian Y, Mukasa SB et al. Cassava brown streak virus (Potyviridae) encodes a putative Maf/HAM1 pyrophosphatase implicated in reduction of mutations and a P1 proteinase that suppresses RNA silencing but contains no HC-Pro. J Virol 2009;**83**:6934–40.
- Mingot A, Valli A, Rodamilans B et al. The P1N-PISPO trans-frame gene of sweet potato feathery mottle potyvirus is produced during virus infection and functions as an RNA silencing suppressor. *J* Virol 2016;**90**:3543–57.
- Misas-Villamil JC, van der Hoorn RAL, Doehlemann G. Papainlike cysteine proteases as hubs in plant immunity. *New Phytol* 2016;**212**:902–7.
- Mishra PM, Verma NC, Rao C *et al*. Intrinsically disordered proteins of viruses: involvement in the mechanism of cell regulation and pathogenesis. *Prog Mol Biol Transl Sci* 2020;**174**:1–78.
- Morrison MS, Podracky CJ, Liu DR. The developing toolkit of continuous directed evolution. Nat Chem Biol 2020;**16**:610–9.
- Nakagawa K, Makino S. Mechanisms of coronavirus nsp1-mediated control of host and viral gene expression. *Cells* 2021;**10**:300.
- Nakahara KS, Shimada R, Choi S-H et al. Involvement of the P1 cistron in overcoming eIF4E-mediated recessive resistance against Clover

yellow vein virus in pea. Mol Plant Microbe Interact 2010;**23**: 1460–9.

- Nasir A, Caetano-Anollés G. A phylogenomic data-driven exploration of viral origins and evolution. *Sci Adv* 2015;**1**:e1500527.
- Nigam D, LaTourrette K, Souza PFN et al. Genome-wide variation in potyviruses. Front Plant Sci 2019;**10**:1439.
- Ohki T, Sasaya T, Maoka T. Cylindrical inclusion protein of wheat yellow mosaic virus is involved in differential infection of wheat cultivars. Phytopathol 2019;**109**:1475–80.
- Pacheco R, García-Marcos A, Manzano A et al. Comparative analysis of transcriptomic and hormonal responses to compatible and incompatible plant-virus interactions that lead to cell death. Mol Plant Microbe Interact 2012;25:709–23.
- Panas MD, Schulte T, Thaa B et al. Viral and cellular proteins containing FGDF motifs bind G3BP to block stress granule formation. PLoS Pathog 2015;**11**:e1004659.
- Pasin F, Bedoya LC, Bernabé-Orts JM et al. Multiple T-DNA delivery to plants using novel mini binary vectors with compatible replication origins. ACS Synthetic Biology 2017;6:1962–8.
- Pasin F, Menzel W, Daròs J-A. Harnessed viruses in the age of metagenomics and synthetic biology: an update on infectious clone assembly and biotechnologies of plant viruses. *Plant Biotechnol J* 2019;**17**:1010–26.
- Pasin F, Shan H, García B *et al*. Abscisic acid connects phytohormone signaling with RNA metabolic pathways and promotes an antiviral response that is evaded by a self-controlled RNA virus. *Plant Commun* 2020;**1**:100099.
- Pasin F, Simón-Mateo C, García JA. The hypervariable aminoterminus of P1 protease modulates potyviral replication and host defense responses. PLoS Pathog 2014;**10**:e1003985.
- Pasin F, Tseng X-A, Bedoya LC et al. Streamlined generation of plant virus infectious clones using the pLX mini binary vectors. J Virol Methods 2018;262:48–55.
- Pasin F. Oligonucleotide abundance biases aid design of a type IIS synthetic genomics framework with plant virome capacity. *Biotechnol J* 2021;**16**:2000354.
- Paudel DB, Sanfaçon H. Exploring the diversity of mechanisms associated with plant tolerance to virus infection. Front Plant Sci 2018;9:1575.
- Pitsili E, Phukan UJ, Coll NS. Cell death in plant immunity. Cold Spring Harb Perspect Biol 2020;**12**:a036483.
- Pitzalis N, Amari K, Graindorge S *et al*. Turnip mosaic virus in oilseed rape activates networks of sRNA-mediated interactions between viral and host genomes. *Commun* Biol 2020;**3**:702.
- Pollari M, De S, Wang A *et al*. The potyviral silencing suppressor HCPro recruits and employs host ARGONAUTE1 in pro-viral functions. PLoS Pathog 2020;**16**:e1008965.
- Pottinger SE, Bak A, Margets A et al. Optimizing the PBS1 decoy system to confer resistance to potyvirus infection in *Arabidopsis* and soybean. Mol Plant Microbe Interact 2020;**33**:932–44.
- Qin L, Shen W, Tang Z *et al*. A newly identified virus in the family *Potyviridae* encodes two leader cysteine proteases in tandem that evolved contrasting RNA silencing suppression functions. *J* Virol 2020;**95**:e01414–20.
- Rajamäki ML, Kelloniemi J, Alminaite A et al. A novel insertion site inside the potyvirus P1 cistron allows expression of heterologous proteins and suggests some P1 functions. Virology 2005;342: 88–101.
- Reuper H, Krenz B. Comparison of two Turnip mosaic virus P1 proteins in their ability to co-localize with the Arabidopsis thaliana G3BP-2 protein. Virus Genes 2021;**57**:233–7.
- Revers F, García JA. Molecular biology of potyviruses. Adv Virus Res 2015;92:101–99.

- Robson F, Khan KS, Le TK *et al.* Coronavirus RNA proofreading: molecular basis and therapeutic targeting. *Mol Cell* 2020;**79**: 710–27.
- Rodamilans B, Casillas A, García JA. P1 of Sweet potato feathery mottle virus shows strong adaptation capacity, replacing P1-HCPro in a chimeric Plum pox virus. J Virol 2021;**95**:e0015021.
- Rodamilans B, Shan H, Pasin F et al. Plant viral proteases: beyond the role of peptide cutters. Front Plant Sci 2018;**9**:666.
- Rodamilans B, Valli A, García JA. Mechanistic divergence between P1 proteases of the family *Potyviridae*. J Gen Virol 2013;**94**:1407–14.
- Rohožková J, Navrátil M. P1 peptidase a mysterious protein of family Potyviridae. J Biosci 2011;**36**:189–200.
- Rose H, Döring I, Vetten H-J et al. Complete genome sequence and construction of an infectious full-length cDNA clone of celery latent virus – an unusual member of a putative new genus within the Potyviridae. J Gen Virol 2019;**100**:308–20.
- Sainsbury F. Innovation in plant-based transient protein expression for infectious disease prevention and preparedness. *Curr Opin Biotechnol* 2020;**61**:110–5.
- Saiz M, Martinez-Salas E. Uncovering targets of the Leader protease: linking RNA-mediated pathways and antiviral defense. Wiley Interdiscip Rev RNA 2021:e1645.
- Salvador B, Saénz P, Yangüez E et al. Host-specific effect of P1 exchange between two potyviruses. Mol Plant Pathol 2008;9:147–55.
- Sánchez F, Ponz F. Presenting peptides at the surface of potyviruses in planta. In: Wege C, Lomonossoff GP (eds.), Virus-Derived Nanoparticles for Advanced Technologies: Methods and Protocols. New York, NY: Springer, 2018, 471–85.
- Sanjuán R, Nebot MR, Chirico N et al. Viral mutation rates. J Virol 2010;**84**:9733–48.
- Savary S, Willocquet L, Pethybridge SJ et al. The global burden of pathogens and pests on major food crops. Nat Ecol Evol 2019;**3**:430–9.
- Savchenko A, Proudfoot M, Skarina T et al. Molecular basis of the antimutagenic activity of the house-cleaning inosine triphosphate pyrophosphatase RdgB from Escherichia coli. J Mol Biol 2007;374:1091–103.
- Segers GC, van Wezel R, Zhang X *et al*. Hypovirus papain-like protease p29 suppresses RNA silencing in the natural fungal host and in a heterologous plant system. *Eukaryotic Cell* 2006;**5**:896–904.
- Shan H, Pasin F, Tzanetakis IE *et al.* Truncation of a P1 leader proteinase facilitates potyvirus replication in a non-permissive host. *Mol Plant Pathol* 2018;**19**:1504–10.
- Shan H, Pasin F, Valli A *et al*. The Potyviridae P1a leader protease contributes to host range specificity. Virology 2015;**476**:264–70.
- Shen Z, Yang Y, Yang S *et al*. Structural and biological basis of alphacoronavirus nsp1 associated with host proliferation and immune evasion. Viruses 2020;**12**:812.
- Shi M, Lin X-D, Tian J-H et al. Redefining the invertebrate RNA virosphere. Nature 2016;**540**:539–43.
- Shukla A, Hoffmann G, Kushwaha NK et al. Salicylic acid and the viral virulence factor 2b regulate the divergent roles of autophagy during cucumber mosaic virus infection. Autophagy 2021:1–13.
- Simmonds P, Aiewsakun P, Katzourakis A. Prisoners of war host adaptation and its constraints on virus evolution. Nat Rev Microbiol 2019;**17**:321–8.
- Simon-Loriere E, Holmes EC. Gene duplication is infrequent in the recent evolutionary history of RNA viruses. Mol Biol Evol 2013;**30**:1263–9.
- Simone PD, Pavlov YI, Borgstahl GEO. ITPA (inosine triphosphate pyrophosphatase): from surveillance of nucleotide pools to human disease and pharmacogenetics. *Mutat Res Rev Mutat Res* 2013;**753**:131–46.

- Sõmera M, Sarmiento C, Truve E. Overview on sobemoviruses and a proposal for the creation of the family Sobemoviridae. Viruses 2015;7:3076–115.
- Sommers P, Chatterjee A, Varsani A et al. Integrating viral metagenomics into an ecological framework. Annu Rev Virol 2021;8: 133–58.
- Steinwand MA, Ronald PC. Crop biotechnology and the future of food. Nat Food 2020;1:273–83.
- Stenger DC, French R, Gildow FE. Complete deletion of Wheat streak mosaic virus HC-Pro: a null mutant is viable for systemic infection. J Virol 2005;79:12077–80.
- Stewart LR, Jarugula S, Zhao Y et al. Identification of a maize chlorotic dwarf virus silencing suppressor protein. Virology 2017;504:88–95.
- Sun T-W, Ku C. Unraveling gene content variation across eukaryotic giant viruses based on network analyses and host associations. *Virus Evol* 2021;**7**:veab081.
- Susaimuthu J, Tzanetakis IE, Gergerich RC et al. A member of a new genus in the Potyviridae infects Rubus. Virus Res 2008;**131**:145–51.
- Sztuba-Solińska J, Urbanowicz A, Figlerowicz M et al. RNA-RNA recombination in plant virus replication and evolution. Annu Rev Phytopathol 2011;**49**:415–43.
- Tamada T, Kondo H. Biological and genetic diversity of plasmodiophorid-transmitted viruses and their vectors. J Gen Plant Pathol 2013;79:307–20.
- Tatineni S, McMechan AJ, Bartels M et al. In vitro transcripts of wild-type and fluorescent protein-tagged Triticum mosaic virus (family Potyviridae) are biologically active in wheat. Phytopathol 2015;105:1496–505.
- Tatineni S, McMechan AJ, Hein GL et al. Efficient and stable expression of GFP through Wheat streak mosaic virus-based vectors in cereal hosts using a range of cleavage sites: Formation of dense fluorescent aggregates for sensitive virus tracking. Virology 2011;410:268–81.
- Tautz N, Tews BA, Meyers G. The molecular biology of pestiviruses. Adv Virus Res 2015;**93**:47–160.
- Tena Fernández F, González I, Doblas P *et al*. The influence of cisacting P1 protein and translational elements on the expression of *Potato virus* Y helper-component proteinase (HCPro) in heterologous systems and its suppression of silencing activity. *Mol Plant Pathol* 2013;**14**:530–41.
- Tomlinson KR, Pablo-Rodriguez JL, Bunawan H et al. Cassava brown streak virus Ham1 protein hydrolyses mutagenic nucleotides and is a necrosis determinant. Mol Plant Pathol 2019;**20**:1080–92.
- Torti S, Schlesier R, Thümmler A et al. Transient reprogramming of crop plants for agronomic performance. Nat Plants 2021;**7**:159–71.
- Tromas N, Elena SF. The rate and spectrum of spontaneous mutations in a plant RNA virus. *Genetics* 2010;**185**:983–9.
- Trudeau T, Nassar R, Cumberworth A et al. Structure and intrinsic disorder in protein autoinhibition. *Structure* 2013;**21**:332–41.
- Tuo D, Yan P, Zhao G et al. An efficient papaya leaf distortion mosaic potyvirus vector for virus-induced gene silencing in papaya. Hortic Res 2021;**8**:144.
- U.S. Environmental Protection Agency Zucchini yellow mosaic virus-weak strain - chemical details. 2007. https://iaspub.epa.g ov/apex/pesticides/f?p=CHEMICALSEARCH:3::::1,3,31,7,12,25: P3_XCHEMICAL_ID:4462 (June 2021, date last accessed).
- Untiveros M, Olspert A, Artola K *et al*. A novel sweet potato potyvirus open reading frame (ORF) is expressed via polymerase slippage and suppresses RNA silencing. *Mol Plant Pathol* 2016;**17**: 1111–23.
- Upadhyaya NM, Mago R, Panwar V *et al*. Genomics accelerated isolation of a new stem rust avirulence gene-wheat resistance gene pair. Nat Plants 2021;**7**:1220–8.

- Uranga M, Vazquez-Vilar M, Orzáez D *et al*. CRISPR-Cas12a genome editing at the whole-plant level using two compatible RNA virus vectors. CRISPR J 2021;**4**:761–9.
- Valente JB, Pereira FS, Stempkowski LA et al. A novel putative member of the family *Benyviridae* is associated with soilborne wheat mosaic disease in Brazil. Plant Pathol 2019;**68**:588–600.
- Valli A, Dujovny G, García JA. Protease activity, self interaction, and small interfering RNA binding of the silencing suppressor P1b from *Cucumber vein yellowing ipomovirus*. J Virol 2008;**82**:974–86.
- Valli A, López-Moya JJ, García JA. Recombination and gene duplication in the evolutionary diversification of P1 proteins in the family *Potyviridae*. J Gen Virol 2007;88:1016–28.
- Valli AA, Gallo A, Rodamilans B et al. The HCPro from the Potyviridae family: an enviable multitasking Helper Component that every virus would like to have. Mol Plant Pathol 2018;**19**:744–63.
- Van de Weyer A-L, Monteiro F, Furzer OJ et al. A species-wide inventory of NLR genes and alleles in Arabidopsis thaliana. Cell 2019;**178**:1260–72.
- van den Born E, Omelchenko MV, Bekkelund A et al. Viral AlkB proteins repair RNA damage by oxidative demethylation. Nucleic Acids Res 2008;**36**:5451–61.
- Vatter HA, Di H, Donaldson EF *et al*. Functional analyses of the three simian hemorrhagic fever virus nonstructural protein 1 papainlike proteases. *J* Virol 2014;**88**:9129–40.
- Verchot J, Carrington JC. Debilitation of plant potyvirus infectivity by P1 proteinase-inactivating mutations and restoration by secondsite modifications. J Virol 1995;**69**:1582–90.
- Villamor DEV, Ho T, Al Rwahnih M et al. High throughput sequencing for plant virus detection and discovery. Phytopathol 2019;**109**: 716–25.
- Visser LJ, Aloise C, Swatek KN *et al.* Dissecting distinct proteolytic activities of FMDV L^{pro} implicates cleavage and degradation of RLR signaling proteins, not its deISGylase/DUB activity, in type I interferon suppression. PLoS Pathog 2020;**16**:e1008702.
- Vlok M, Gibbs AJ, Suttle CA. Metagenomes of a freshwater charavirus from British Columbia provide a window into ancient lineages of viruses. Viruses 2019;11,299.
- Vogler H, Kwon M-O, Dang V et al. Tobacco mosaic virus movement protein enhances the spread of RNA silencing. PLoS Pathog 2008;**4**:e1000038.
- Wang G, Zheng C. Zinc finger proteins in the host-virus interplay: multifaceted functions based on their nucleic acid-binding property. FEMS Microbiol Rev 2021;**45**:fuaa059.
- Watt LG, Crawshaw S, Rhee S-J *et al.* The cucumber mosaic virus 1a protein regulates interactions between the 2b protein and ARG-ONAUTE 1 while maintaining the silencing suppressor activity of the 2b protein. PLoS Pathog 2020;**16**:e1009125.
- Willemsen A, Zwart MP, Ambrós S et al. 2b or not 2b: experimental evolution of functional exogenous sequences in a plant RNA virus. *Genome Biol Evol* 2017;**9**:297–310.
- Willemsen A, Zwart MP, Higueras P et al. Predicting the stability of homologous gene duplications in a plant RNA virus. *Gen Biol Evol* 2016;**8**:3065–82.
- Willemsen A, Zwart MP. On the stability of sequences inserted into viral genomes. Virus Evol 2019;**5**:vez045.
- Wolf YI, Kazlauskas D, Iranzo J *et al*. Origins and evolution of the global RNA virome. MBio 2018;**9**:e02329–18.
- Wolf YI, Silas S, Wang Y et al. Doubling of the known set of RNA viruses by metagenomic analysis of an aquatic virome. Nat Microbiol 2020;**5**:1262–70.

- Xie W, Marty DM, Xu J et al. Simultaneous gene expression and multigene silencing in Zea mays using maize dwarf mosaic virus. BMC Plant Biol 2021;**21**:208.
- Xu M, Mazur MJ, Tao X et al. Cellular RNA hubs: friends and foes of plant viruses. Mol Plant Microbe Interact 2020;**33**:40–54.
- Yang X, Li Y, Wang A. Research advances in potyviruses: from the laboratory bench to the field. Annu Rev Phytopathol 2021;59: 1–29.
- Yao T, Coleman S, Nguyen TVP et al. Bacteriophage selfcounting in the presence of viral replication. Proc Natl Acad Sci 2021;118:e2104163118.
- Yoshikawa G, Askora A, Blanc-Mathieu R et al. Xanthomonas citri jumbo phage XacN1 exhibits a wide host range and high complement of tRNA genes. Sci Rep 2018;**8**:4486.
- You Y, Shirako Y. Bymovirus reverse genetics: requirements for RNA2-encoded proteins in systemic infection. Mol Plant Pathol 2010;11:383–94.
- Young BA, Stenger DC, Qu F et al. Tritimovirus P1 functions as a suppressor of RNA silencing and an enhancer of disease symptoms. Virus Res 2012;**163**:672–7.
- Yu B, Edstrom WC, Benach J et al. Crystal structures of catalytic complexes of the oxidative DNA/RNA repair enzyme AlkB. Nature 2006;439:879–84.
- Zamora M, Méndez-López E, Agirrezabala X et al. Potyvirus virion structure shows conserved protein fold and RNA binding site in ssRNA viruses. Sci Adv 2017;3:eaao2182.
- Zell R. Picomaviridae the ever-growing virus family. Arch Virol 2018;**163**:299–317.
- Zhang T, Liu P, Zhong K *et al.* Wheat yellow mosaic virus NIb interacting with host light induced protein (LIP) facilitates its infection through perturbing the abscisic acid pathway in wheat. *Biology* 2019;**8**:80.
- Zhang T, Wang Z, Hu H et al. Transcriptome-wide N⁶methyladenosine (m⁶A) profiling of susceptible and resistant wheat varieties reveals the involvement of variety-specific m⁶A modification involved in virus-host interaction pathways. Front Microbiol 2021;**12**:656302.
- Zhang W, Qian Y, Jia G. The detection and functions of RNA modification m6A based on m6A writers and erasers. J Biol Chem 2021;297:100973.
- Zhang X-P, Liu D-S, Yan T et al. Cucumber mosaic virus coat protein modulates the accumulation of 2b protein and antiviral silencing that causes symptom recovery in planta. PLoS Pathog 2017;**13**:e1006522.
- Zhao M, García B, Gallo A et al. Home-made enzymatic premix and Illumina sequencing allow for one-step Gibson assembly and verification of virus infectious clones. Phytopathol Res 2020; 2:36.
- Zhao Q, Feng Q, Lu H et al. Pan-genome analysis highlights the extent of genomic variation in cultivated and wild rice. Nat Genet 2018;**50**:278–84.
- Zilber-Rosenberg I, Rosenberg E. Microbial-driven genetic variation in holobionts. FEMS Microbiol Rev 2021;**45**:fuab022.
- Zimmerman AE, Howard-Varona C, Needham DM et al. Metabolic and biogeochemical consequences of viral infection in aquatic ecosystems. Nat Rev Microbiol 2020;**18**:21–34.
- Zitzmann C, Schmid B, Ruggieri A *et al.* A coupled mathematical model of the intracellular replication of dengue virus and the host cell immune response to infection. *Front Microbiol* 2020;**11**:725.