

PROCEEDINGS

Open Access

Down-regulation of promoter methylation level of *CD4* gene after MDV infection in MD-susceptible chicken line

Juan Luo^{1†}, Ying Yu^{1,3†}, Huanmin Zhang², Fei Tian¹, Shuang Chang², Hans H Cheng², Jiuzhou Song^{1*}

From International Symposium on Animal Genomics for Animal Health (AGAH 2010)
Paris, France. 31 May – 2 June 2010

Abstract

Background: Marek's disease virus (MDV) is an oncovirus that induces lymphoid tumors in susceptible chickens, and may affect the epigenetic stability of the *CD4* gene. The purpose of this study was to find the effect of MDV infection on DNA methylation status of the *CD4* gene differed between MD-resistant (L6₃) and -susceptible (L7₂) chicken lines.

Methods: Chickens from each line were divided into two groups with one group infected by MDV and the other group as uninfected controls. Then, promoter DNA methylation levels of the *CD4* gene were measured by Pyrosequencing; and gene expression analysis was performed by quantitative PCR.

Results: Promoter methylation of the *CD4* gene was found to be down-regulated in L7₂ chickens only after MDV infection. The methylation down-regulation of the *CD4* promoter is negatively correlated with up-regulation of *CD4* gene expression in the L7₂ spleen at 21 dpi.

Conclusions: The methylation fluctuation and mRNA expression change of *CD4* gene induced by MDV infection suggested a unique epigenetic mechanism existed in MD-susceptible chickens.

Background

CD4 encodes a glycoprotein, located on the surface of T helper (Th) cells and regulatory T cells. Through interaction with MHC class II molecules, CD4 directs the lineage development of Th cells in immune organs and activates the CD4⁺ T cell maturation process [1]. Thus, the transcriptional level of *CD4* is directly related to T cell development [2]. In mice, *CD4* transcription is controlled by several *cis*-acting elements including enhancers, silencers and DNA methylation [3,4]. However, the epigenetic regulation of *CD4* gene in chicken and its relationship with any virus infection are still unclear...

Marek's disease (MD), a T cell lymphoma of chickens caused by the Marek's disease virus (MDV), is

characterized by mononuclear cell-infiltration in various organs including peripheral nerves, skin, muscle, and visceral organs [5], and is a worldwide problem for the poultry industry. A complex MDV life cycle was found in susceptible chickens during MD progression, which includes an early cytolitic phase (2-7 days post infection, dpi), latent phase (7-10 dpi), late cytolitic phase (from 18 dpi) and transformation phase (28 dpi and onwards) [6].

Epigenetics is the study of alterations that result in inherited changes in phenotypes despite the lack of DNA sequence polymorphisms and include DNA methylation, histone modification and chromatin remodeling [7]. It is described as the interaction between genes and environmental factors. Aberrant CpG methylation levels of the gene promoter region contribute to oncogenesis [8]. Viruses are one of the environmental agents that can cause alterations of DNA methylation level in host genes [9].

* Correspondence: songj88@umd.edu

† Contributed equally

¹Animal & Avian Sciences Department, University of Maryland, College Park, Maryland, 20740, USA

Full list of author information is available at the end of the article

The focus of this study was to better understand the expression control of *CD4* by ascertaining the epigenetic status in the *CD4* promoter and the *CD4* expression in relation to MDV infection. Two inbred chicken lines, MD-resistant or -susceptible with the same MHC (major histocompatibility complex) haplotypes, from Avian Disease and Oncology laboratory (ADOL) were used [5]. We, therefore, measured the promoter methylation and transcription of the *CD4* gene before and after MDV infection of both lines. We found methylation alterations in the *CD4* promoter region after MDV infection differ between these two lines.

Methods

Animals, virus infection experiments and sample collection

USDA, Avian Disease and Oncology Laboratory (ADOL) chicken lines 6 (L6₃) and lines 7 (L7₂) chickens, which are MD-resistant and MD-susceptible, respectively, were obtained. For each line, the chickens were divided into two groups with 30 chickens infected by MDV and 30 uninfected controls. A very virulent plus strain of MDV (648A passage 40, VV+) was injected intra-abdominally on the fifth day after hatching with 500 plaque-forming units (PFU). Spleen samples were collected at 5 dpi, 10 dpi and 21 dpi, put in RNAlater (Qiagen, USA) immediately, and then stored at -80°C. All procedures followed the standard animal ethics and user guidelines.

DNA extraction, bisulfite treatment and pyrosequencing

DNA was extracted from 20-30 mg spleen by NucleoSpin® Tissue Kits (Macherey-Nagel, Germany). 500 ng DNA was treated with sodium bisulfite and purified by EZ DNA Methylation-Gold Kit™ (ZYMO Research, USA). Primers for pyrosequencing were designed by PSQ Assay Design software (Biotage, Sweden) (Table 1). For cost reduction, a universal primer (5'-GGGACACCGCTGATCGTTTA-3') was used in the PCR assays [10]. DNA methylation level analysis was performed with Pyro Q-CpG system (PyroMark ID, Biotage, Sweden) as previously described [10,11].

RNA extraction and quantitative real-time RT-PCR

RNA from 30-50mg spleen was extracted using the RNAeasy Mini Kit (Qiagen, USA). Reverse transcription was carried out in 20 µl with 1 µg of total RNA by using SuperScript™ III Reverse Transcriptase (Invitrogen, USA) and oligo (dT)₁₂₋₁₈ primers (Invitrogen, USA). Primers (Table 1) for quantitative real-time RT-PCR were designed by Primer3 online primer designer system (<http://frodo.wi.mit.edu/>). qPCR was performed on the iCycler iQ PCR system (Bio-Rad, USA) in a final volume of 20 µl using QuantiTect SYBR Green PCR Kit (Qiagen, USA) with the following procedure: denatured at 95°C for 15 min, followed by 40 cycles at 95 °C for 30 s, 60 °C for 30 s, 72 °C for 30 s, then extended at 72 °C for 10 min. Each reaction was replicated twice. The housekeeping gene *GAPDH* (glyceraldehyde-3-phosphate dehydrogenase) was used to normalize the assays.

Statistical analysis

Promoter methylation levels and gene expression before and after MDV infection were compared by Student's *t* test. An exact *F* test was performed to distinguish different methylation patterns [10]. Correlation between *CD4* DNA methylation and expression was tested by Pearson's correlation coefficient.

Results


CD4 promoter methylation analysis before and after MDV infection

To determine the promoter methylation level of the *CD4* gene, a DNA sequence containing the CpG islands from the *CD4* gene promoter region (sequence shown in Table 1) was downloaded from UCSC (<http://genome.ucsc.edu>) and the methylation level was determined by pyrosequencing. The CpGs in the promoter of *CD4* exhibits a high (>70%) methylation level in both L6₃ and L7₂ chickens before MDV infection. During MD progression, no significant methylation changes of *CD4* promoter were detected in L6₃ chickens at 5, 10 and 21 dpi or in L7₂ chickens at 5 and 10 dpi (*P*>0.05, Figure 1, and Figure 1A and 1B); however, the significant down-regulation of *CD4* promoter methylation level was observed at 21 dpi in L7₂ chickens

Table 1 Primers used in Pyrosequencing and quantitative PCR

Genes	Primers	Sequence	Purpose
<i>CD4</i>	F	5'- TTGAGATTATAYGTATTTGGAAGA -3'	Pyrosequencing
	R	5'- GGGACACCGCTGATCGTTTA ACCTTTATATCTCCTCTCCA -3'	
	Sequencing Assay	5'- AGTATTATTGAGAGAAGIT -3'	
		5'- YGTAGATTGTAGTAGAGTTGGATYG GTAGTAAGATYGTGTTGAYGTTTT -3'	
<i>GAPDH</i>	F	5'-GAGGGTAGTGAAGGCTGCTG-3'	quantitative PCR
	R	5'-ACCAGGAAACAAGCTTGACG-3'	
<i>CD4</i>	F	5'- TGTCAACGCCGGATGTATAA-3'	quantitative PCR
	R	5'- CTTGCCAATGGCTCCTCTC-3'	

Y stands for C/T. Bold Y in the assay sequence is the CpG sites analyzed.



($P < 0.05$, Figure 1C). The result from the exact F test revealed that the $CD4$ promoter methylation pattern in L7₂ infected samples at 21 dpi was significantly different from any other groups (Figure 2).

CD4 gene expression at 21 dpi


To ascertain if the $CD4$ gene transcription level is influenced by its promoter methylation changes at 21 dpi, we conducted quantitative PCR. We found a significantly higher expression of $CD4$ gene in L7₂ infected samples compared with noninfected control samples ($P < 0.05$) (Figure 3), whereas no significant up or down-regulation of $CD4$ expression was detected in L6₃ chickens after MDV infection ($P > 0.05$). Hereinafter, further correlation analysis showed that methylation level of all the detected CpG sites existed a negatively relationship with $CD4$ gene expression in L7₂ chicken at 21dpi (Figure 4).

Discussion



The $CD4$ gene and its regulatory sequences are conserved [12]. In human and mouse, multiple protein or

transcription factor binding sites, including the Myb binding site, Elf-1 binding site, and Ikaros binding site, were found in the promoter region of $CD4$, which is involved in the on/off switching of $CD4$ gene expression [4]. These regulatory sites were also found in the chicken $CD4$ promoter with potential functions in its expression [12]. It is well known that epigenetic factors such as DNA methylation and histone modifications play important roles in transcriptional regulation in mammals [7]. For example, the methylation change in at least one CpG site of $CD4$ gene in mouse is related to $CD4^+$ T cell differentiation [3]. In this study, we thus examined the methylation status in the promoter region of $CD4$ gene in chickens related to MDV infection.

MDV is an oncovirus using $CD4^+$ T cell as a target for latent infection and transformation, which may have interactions with the $CD4$ gene at the epigenetic level [13]. In our previously study, two mutations (CG→TG) were identified in the DNMT3b gene between L63 and L72 chickens [10], which implied that the DNA methylation machinery may be different in the two lines in response to MDV infection. In this study, the



methylation levels on the promoter region of the *CD4* gene were fluctuated over different time points of MDV infection in MD-susceptible chickens, especially during the late cytolitic phase. The quantitative PCR results confirmed that *CD4* expression in *L72* chicken during the late stages of MDV infection was upregulated while the *CD4* promoter methylation was down-regulation. Since the expression of *CD4* is essential for *CD4*⁺ T cell development and activation, it may suggest that there are different epigenetic machineries of activation of *CD4*⁺ T cells by MDV infection through regulation of *CD4* methylation levels between MD-resistant and susceptible chicken lines. From previous studies, it was found that the number of infected *CD4*⁺ T cells were similar



during the early phase (cytolitic phase) of MDV infection between MD-resistant and –susceptible chicken lines, but was increased during cytolitic phase in MD-susceptible chicken line and decreased in MD-resistant chicken line [14]. Additionally, in MD-resistant chicken line, *CD4*⁺ T cell is latently infected, but cannot be transformed, whereas in MD-susceptible chicken lines the infected *CD4*⁺ T cell can be transformed after the latent phase [5,15]. Taken together, the methylation change of *CD4* gene gives us an important clue that epigenetic alteration could associate with MD etiology. Therefore, future efforts will disclose the epigenetic landscapes, including genome-wide DNA methylation and histone modifications, in immune organs and specific cell types, such as the *CD4*⁺ T cell, which will supply rich information to explore the epigenetic machinery related to chemical and physiological mechanisms of MD resistance or susceptibility.

Conclusions

In conclusion, the methylation fluctuation and mRNA expression of *CD4* gene induced by MDV infection suggested a unique epigenetic mechanism existed in MD-susceptible chickens.

Acknowledgements

The study was grant-supported by NIFA 2008-35204-04660. This article has been published as part of *BMC Proceedings* Volume 5 Supplement 4, 2011: Proceedings of the International Symposium on Animal Genomics for Animal Health (AGAH 2010). The full contents of the supplement are available online at <http://www.biomedcentral.com/1753-6561/5?issue=S4>.

Author details

¹Animal & Avian Sciences Department, University of Maryland, College Park, Maryland, 20740, USA. ²USDA, ARS, Avian Disease and Oncology Laboratory, East Lansing, MI 48823, USA. ³College of Animal Sciences, China Agricultural University, Haidian, Beijing, 100193, P.R. China.

Authors' contributions

JL performed the experiments and prepare the manuscript. YY and FT performed the experiments. HMZ and SC designed and performed the viral challenge experiment. JZS designed and wrote the paper.

Competing interests

The authors declare that they have no competing interests.

Published: 3 June 2011

References

1. Huang Z, Xie H, Ioannidis V, Held W, Clevers H, Sadim MS, Sun Z: **Transcriptional regulation of CD4 gene expression by T cell factor-1/ beta-catenin pathway.** *J Immunol* 2006, **176**(8):4880-4887.
2. Paillard F, Sterkers G, Vaquero C: **Transcriptional and post-transcriptional regulation of TcR, CD4 and CD8 gene expression during activation of normal human T lymphocytes.** *Embo J* 1990, **9**(6):1867-1872.
3. Tutt Landolfi MM, Scollay R, Parnes JR: **Specific demethylation of the CD4 gene during CD4 T lymphocyte differentiation.** *Mol Immunol* 1997, **34**(1):53-61.
4. Ellmeier W, Sawada S, Littman DR: **The regulation of CD4 and CD8 coreceptor gene expression during T cell development.** *Annu Rev Immunol* 1999, **17**:523-524.
5. Davison F, Nair V: **Marek's Disease: An Evolving Problem.** Oxford: Elsevier Academic Press; 2004.
6. Calnek BW: **Pathogenesis of Marek's disease virus infection.** *Curr Top Microbiol Immunol* 2001, **255**:25-55.
7. Allis CD, Jenuwein T, Reinberg D, Caparros M-L: **Epigenetics.** Cold Spring Harbor Laboratory Press; 2006.
8. Herman JG, Baylin SB: **Gene silencing in cancer in association with promoter hypermethylation.** *N Engl J Med* 2003, **349**(21):2042-2054.
9. Amara K, Trimeche M, Ziadi S, Laatiri A, Hachana M, Sriha B, Mokni M, Korbi S: **Presence of simian virus 40 DNA sequences in diffuse large B-cell lymphomas in Tunisia correlates with aberrant promoter hypermethylation of multiple tumor suppressor genes.** *Int J Cancer* 2007, **121**(12):2693-2702.
10. Yu Y, Zhang H, Tian F, Zhang W, Fang H, Song J: **An integrated epigenetic and genetic analysis of DNA methyltransferase genes (DNMTs) in tumor resistant and susceptible chicken lines.** *PLoS ONE* 2008, **3**(7):e2672.
11. Colella S, Shen L, Baggerly KA, Issa JP, Krahe R: **Sensitive and quantitative universal Pyrosequencing methylation analysis of CpG sites.** *Biotechniques* 2003, **35**(1):146-150.
12. Koskinen R, Salomonsen J, Tregaskes CA, Young JR, Goodchild M, Bumstead N, Vainio O: **The chicken CD4 gene has remained conserved in evolution.** *Immunogenetics* 2002, **54**(7):520-525.
13. Rouse BT, Wells RJ, Warner NL: **Proportion of T and B lymphocytes in lesions of Marek's disease: theoretical implications for pathogenesis.** *J Immunol* 1973, **110**(2):534-539.
14. Burgess SC, Basaran BH, Davison TF: **Resistance to Marek's disease herpesvirus-induced lymphoma is multiphasic and dependent on host genotype.** *Vet Pathol* 2001, **38**(2):129-142.
15. Burgess SC, Davison TF: **Identification of the neoplastically transformed cells in Marek's disease herpesvirus-induced lymphomas: recognition by the monoclonal antibody AV37.** *J Virol* 2002, **76**(14):7276-7292.

doi:10.1186/1753-6561-5-S4-S7

Cite this article as: Luo *et al.*: Down-regulation of promoter methylation level of CD4 gene after MDV infection in MD-susceptible chicken line. *BMC Proceedings* 2011 **5**(Suppl 4):S7.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

