

Original article

Exertional heat illness risk factors and physiological responses of youth football players

Susan W. Yeargin^{a,*}, John J. Dickinson^{a,b}, Dawn M. Emerson^{a,c}, Jessica Koller^{a,d},
Toni M. Torres-McGehee^a, Zachary Y. Kerr^e

^a Department of Exercise Science, University of South Carolina, Columbia, SC 29208, USA

^b Palmetto Health/USC Orthopedic Center, Keenan High School, Columbia, SC 29203, USA

^c Department of Health, Sport, and Exercise Sciences, University of Kansas, Lawrence, KS 66045, USA

^d Surgi-Care Inc., Boston, MA 02115, USA

^e Department of Exercise and Sport Science, University of North Carolina, Chapel Hill, NC 27599-8700, USA

Received 8 June 2018; revised 7 September 2018; accepted 9 October 2018

Available online 7 March 2019

2095-2546/© 2021 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Objective: To determine which intrinsic and extrinsic exertional heat illness (EHI) risk factors exist in youth American football players and observe perceptual and physiological responses of players during events (games and practices).

Methods: Cross-sectional cohort study observing 63 youth football players, varying in position. Independent variables were league (weight-restricted (WR, $n = 27$) and age-restricted (AR, $n = 36$)) and event type. Dependent variables were anthropometrics, work-to-rest ratio, and wet bulb globe temperature. Descriptive variables included preparticipation examination and uniform configuration. A subset of 16 players participated in physiological variables (heart rate and gastrointestinal temperature). Data collection occurred on 7 AR and 8 WR nonconsecutive practices and the first 3 games of the season.

Results: Mean values for anthropometric variables were higher ($p < 0.05$) in the AR league than the WR league. Work time ($\chi^2(1,111) = 4.232$; $p = 0.039$) and rest time ($\chi^2(1,111) = 43.41$; $p < 0.001$) were significantly greater for games, but ratios were significantly higher for practices ($\chi^2(1,111) = 40.62$; $p < 0.001$). The majority of events (77%) observed were in black and red flag wet bulb globe temperature risk categories. A total of 57% of the players had a preparticipation examination, and up to 82% of events observed were in full uniforms. Individual gastrointestinal temperature and heart rate responses ranged widely and no players reached critical thresholds.

Conclusion: Extrinsic (disproportionate work ratios, environmental conditions) and intrinsic (higher body mass index) EHI risk factors exist in youth football. Certain risk factors may be influenced by event and league type. National youth football organizations need to create thorough guidelines that address EHI risk factors for local leagues to adopt.

Keywords: Adolescents; Anthropometrics; Core temperature; Heat exchange; Pediatric; Wet bulb globe temperature

1. Introduction

In the past decade, the diagnosis of exertional heat illness (EHI) has increased in athletes, as well as the number of confirmed deaths from EHI.^{1–3} Youth and adolescent (<19 years of age) male athletes participating in American football compose the majority of EHI cases that present to emergency rooms.⁴ Recent research reveals that youth football has the highest EHI rate compared with high school and college

teams.⁵ Epidemiological research indicates that youth football players sustain EHI at a rate of 1.82 per 10,000 athlete exposures.⁵ Emergency department data reported approximately 1000 EHI cases sustained per year in football players younger than 19 years of age in a sample from the National Electronic Injury Surveillance System.⁴ EHI rates are highest in the pre-season regardless of competitive level,^{5–7} but differ between practices and games depending on the level.⁵

EHI has numerous risk factors, including lack of heat acclimatization, increased body mass index (BMI), illness, and inappropriate work-to-rest ratios.^{8,9} More specifically, football is an intense sport and this high exercise intensity is the largest driver

Peer review under responsibility of Shanghai University of Sport.

* Corresponding author.

E-mail address: syeargin@mailbox.sc.edu (S.W. Yeargin).

<https://doi.org/10.1016/j.jshs.2019.03.002>

Cite this article: Yeargin SW, Dickinson JJ, Emerson DM, Koller J, Torres-McGehee TM, Kerr ZY. Exertional heat illness risk factors and physiological responses of youth football players. *J Sport Health Sci* 2021;10:91–8.

of increased core body temperature.^{10,11} Football players have additional risk factors, such as their equipment and playing in hot environmental conditions (wet bulb globe temperature (WBGT)) during preseason.^{12–16} Although there has been immense research on EHI risk factors in adults^{4,9,12,13,17–20} and high school athletes,^{5,6,21–24} research is lacking for youth football players.^{5,14} Youth football players may have supplementary thermoregulatory predisposing factors^{25–28} and team dynamics. Teams have small rosters, leading to athletes often playing both offense and defense,²⁹ which may result in uneven work-to-rest ratios. Youth football leagues vary across the country by organization structure, team demographics, and guidelines/rules. For example, some leagues create teams based on age only, whereas others are based on age plus weight. Leagues can be part of a national organization or independent and may or may not have guidelines on starting dates, practice length, or playing time. These factors in turn can significantly impact environmental conditions players are exposed to,¹⁴ as well as athlete exposures²⁹ and possibly the types of players that participate in each league.

To date, no existing literature has examined the risk factors associated with EHI in youth football players beside environmental conditions.¹⁴ Describing which factors are prevalent in youth players and how their bodies respond to football events (practices and games) may improve prevention strategies and help to develop guidelines for safer participation. Therefore, the primary purpose of our study was to determine which intrinsic (anthropometrics, previous history) and extrinsic (WBGT, work-to-rest ratio) EHI risk factors exist in youth football players. The secondary purpose was to describe the perceptual responses (heat illness symptoms), maximum physiological responses (gastrointestinal temperature (T_{GI}) and heart rate (HR)) that youth football players experienced in a warm environment.

2. Methods

2.1. Design

We used a cross-sectional research design to observe youth football teams during 1 season. Data were collected from 2 types of youth football leagues (weight restricted (WR) and age restricted (AR)) during 2 types of events (games and practices). Dependent variables included subjects' anthropometrics (age, height, weight, BMI category, and body surface area (BSA)), each team's work and rest times, environmental conditions measured by WBGT, and perceptual heat illness symptoms. Physiological variables including HR and T_{GI} were acquired from a subgroup of participants. Additional risk factor history (preparticipation examination, previous EHI, sickle cell trait status, medications, sleep habits, signs of illness, and uniform configuration), preseason and regular season start dates, and event times were collected for descriptive purposes.

2.2. Participants

Youth football players ($n = 63$) from local recreational leagues in the southeastern region of the United States participated in this study. The youth players in these leagues were 13 years of age or younger and were not participating on an

interscholastic team. The convenience sample was recruited during the team parent meetings for each league before the season began. There were twenty-seven of 80 participants from the WR league (34%) and thirty-six of 100 from the AR league (36%) who volunteered to participate. The WR league included participants from 4 teams of specific weight and age categories. Each player within this league format had to be within the weight range before the season began to be eligible for that division. Additionally, at each game, participants had to weigh in with full pads and meet requirements to play that day. The AR league had 11 teams with participants from 8 teams separated only by age ranges with no weight specifications. There were no exclusions to participate in the study. However, for those interested in participating in the T_{GI} portion of the study, exclusions (i.e., less than 80 lbs) were followed per manufacturer instructions.³⁰ Of the 63 youth football players, 16 participants (8 from each league) met the criteria and participated. The University of South Carolina's Institutional Review Board approved the protocol before recruitment. An informational team meeting was held before the preseason began for each league. This meeting included describing the study and the risks involved and answering questions from the parents and athletes. Those interested in participating signed consent and assent forms.

2.3. Measurements and instrumentation

2.3.1. Risk factor history

League websites were examined to determine if preparticipation examinations were mandated. Preseason start dates, regular season start dates, practice times, and game times were also examined. The results were confirmed with the league's director. A baseline survey asked participants to self-report any past EHI history, general sleep habits, and known sickle cell trait history. Players also completed a short survey at each observed game and practice asking about sleep (time to bed and time of waking) and general illness symptoms (flu, diarrhea, vomiting, fever, or malaise) within the last 24 h and 48 h, respectively. Researchers recorded uniform configurations on activity logs for each practice and game (1 (helmet only), 2 (helmet and shoulder pads), 3 (full uniform)).

2.3.2. Anthropometrics

Age was self-reported in whole years. Participants were weighed in shorts only on the first day of practice as a baseline measure using a digital scale (TBF-300A; Tanita, Arlington Heights, IL, USA), and height was measured using a portable stadiometer (ShorrBoard Weight and Measure, Olney, MD, USA). These variables were used to calculate BMI and then categorized (underweight, healthy, overweight, and obese) by Centers for Disease Control and Prevention delimiters.^{31,32} BSA was calculated with the Mosteller formula.³³

2.3.3. Work-to-rest ratio

In relation to EHI, work represents any activity or drill that could result in metabolic heat generation, and rest represents the opportunity to optimize heat loss. Researchers recorded the time of day each team began and ended an activity or drill

(i.e., work) on activity logs. Any time a team had a break (i.e., rest), the beginning and ending times were recorded. Activity logs were filled out at all events ($n = 11$). The team's work and rest times were averaged together to represent the leagues' experience as a collective identity. A work-to-rest ratio was calculated using the number of minutes the teams engaged in work divided by the number of minutes of rest.

2.3.4. Environmental conditions

A portable WBGT device (4600 Heat Stress Meter; Kestrel Meters, Birmingham, MI, USA) was used to measure environmental conditions. The average and maximum WBGT values were analyzed to determine the typical and highest heat stresses experienced by players.²⁴ The average WBGT was calculated using all measurements obtained (1 measurement every 15 min) during all events. The maximum WBGT was derived by extracting the highest WBGT measurement recorded for that event and then averaged. Average results were compared with the American College of Sports Medicine recommendation chart to determine frequencies in each flag risk category.²⁰ The green risk category was any event played at less than 24.0°C and the yellow category was any event played at 24.0°C–25.9°C. The red category events were played at 26.0°C–29.0°C and the black flag risk categories were any events played at more than 29.0°C.²⁰

2.3.5. Perceptual and physiological variables

A total of 14 selected items from an Environmental Symptoms Questionnaire (ESQ)^{24,34,35} evaluated common EHI signs and symptoms for each event. Ingestible T_{GI} sensors (HQ Inc., Palmetto, FL, USA) were used to measure T_{GI} during selected practices ($n = 3$) and games ($n = 3$).³⁰ The youth ingested the sensors in the morning (approximately 6 h before the start of the event) to ensure the sensor reached the intestines. HR chest straps (Polar Electro Inc T31, Lake Success, NY, USA) were used to monitor HR simultaneously with T_{GI} . The maximum T_{GI} and HR were derived by extracting the highest measurement recorded for each player during each event and then averaged.

2.4. Procedures

Data collection occurred on 7 AR or 8 WR nonconsecutive practices and the first 3 games of the season for both leagues. The WR had 2 practices on Day 8. Anthropometric data and risk factor history were collected before the first day of practice. Sleep habits and general illness symptoms were collected before every practice and game. WBGT was recorded throughout each practice or game in approximately 15-min intervals. During practices and games, researchers noted the uniform configuration and logged work and rest time intervals for each team. Immediately after each practice and game, subjects completed the ESQ.

A subset of football players ($n = 16$) agreed to participate and completed data collection. At 3 predetermined practices and 3 games, the T_{GI} and HR were measured. Parents were given specific instructions to have their youth ingested the

sensor during the morning hours (i.e., with breakfast) to ensure proper location within the intestines. On the day of the practice or game, a baseline T_{GI} was measured to confirm the sensor had reached the intestines (i.e., to ensure a nonerroneous measurement). Participants were instructed to practice and play normally. During practices, T_{GI} and HR were checked at approximately 15- to 20-min intervals throughout the 2-h practice. During games, the T_{GI} and HR were recorded when the athlete returned to the sideline and at timeouts.

2.5. Statistical analyses

Descriptive statistics were calculated for all dependent and descriptive variables. The average and maximum WBGT were used to describe environmental conditions experienced by leagues.²⁴ Independent t tests compared means between leagues for anthropometrics variables. Independent t tests were also used to determine differences between leagues and events for WBGT and between events for ESQ scores. The Kruskal–Wallis test was used to compare overall work time, rest time, and ratios for each event type. Pearson's r correlation coefficients were used to determine if relationships existed for HR and T_{GI} with work-to-rest ratio, WBGT, and BMI. Maximum physiological variables from all 16 players at each football event were used to represent thermal strain.²⁴ No statistical comparisons were made for physiological variables because the data sets were incomplete. Significance was set *a priori* at $p = 0.05$. Statistical Analysis Software (Version 4.3; SAS Institute, Cary, NC, USA) was used for all analyses.

3. Results

3.1. Risk factor history

Preparticipation examination mandates, start dates, and times for practices and games differed depending on the league (Table 1). Of the 63 players participating in the study, thirty-four completed a baseline questionnaire. About one-quarter (26%) reported experiencing symptoms in the heat consistent with an EHI (i.e., struggling, faint, dizzy, nauseous, and falling down) in a previous season, but zero reported an actual previous EHI episode. When the participants were asked if they were taking a medication, 29% indicated yes. There were no clarifications of medication type provided by the players. One player (3%) was sickle cell trait positive as confirmed by a parent. Players reported an average of 8.7 ± 1.3 h of sleep on the baseline questionnaire and 9.4 ± 0.3 h of sleep on the prepractice/pregame questionnaire. Players reported symptoms consistent with a general illness within 48 h of a practice or game 9% of the time (range: 0–22%). Players in 1 league we observed in the study wore shorts and T-shirts for their first 2 practices, added helmets for the third practice, and then wore full uniforms for the remaining practices and games (72.7%). In the other league we observed, players wore helmets only for the first 2 days and then proceeded to full uniform for the remaining practices and games (82%).

Table 1
Descriptive values for anthropometrics and risk factors by league.

Variable	Aggregate (<i>n</i> = 63)	Weight-restricted league (<i>n</i> = 27)	Age-restricted league (<i>n</i> = 36)	<i>p</i> ^a
Anthropometrics				
Age (year)	11 ± 1 (9–13)	10 ± 1 (9–12)	11 ± 1 (9–13)	0.004
Height (cm)	148 ± 9 (127–166)	144 ± 9 (127–163)	152 ± 8 (133–167)	0.001
Weight (kg)	44.7 ± 15.2 (26.2–95.7)	37.6 ± 8.0 (26.2–56.3)	50.1 ± 17.2 (27.7–95.7)	<0.001
BMI (kg/m ²)	20.0 ± 5.3 (14.8–36.3)	18.0 ± 3.3 (14.8–26.9)	21.5 ± 6.1 (14.9–36.3)	0.006
BSA (m ²)	1.4 ± 0.3 (1.0–2.1)	1.2 ± 0.1 (1.0–1.5)	1.4 ± 0.3 (1.0–2.1)	<0.001
Risk and event factors				
PPE		Mandated	Not required	
Preseason start date		August 2	August 26	
Practice start times		6:00 p.m.	6:00 p.m.	
Regular season start date		August 31	September 16	
Game times		9:00 a.m.–2:00 p.m.	6:00 p.m.–8:00 p.m.	
Flag risk category^b				
Green and yellow flag	5 (22.7)	1 (8.3)	4 (40.0)	
Red and black flag	17 (77.3)	11 (91.7)	6 (60.0)	

Note: Values are mean ± SD (minimum–maximum) or number (%) unless otherwise indicated.

^a *p* values originate from independent *t* tests comparison means in weight-restricted and age-restricted leagues.

^b The flag risk category refers to the practices/games were observed. Flag risk categories per ACSM recommendations.²⁰

Abbreviations: ACSM = American College of Sports Medicine; BMI = body mass index; BSA = body surface area; PPE = preparticipation examination.

3.2. Anthropometrics

The overall and league demographics are provided in Table 1. Mean values for anthropometrics variables were higher (*p* < 0.05) in the AR league than in the WR league. The BMI category percentages in the WR league were 3.7% underweight (*n* = 1), 66.7% healthy (*n* = 18), 18.5% overweight (*n* = 5), and 11.1% obese (*n* = 3). For the AR league, the percentages were 2.8% underweight (*n* = 1), 61.1% healthy (*n* = 22), 8.3% overweight (*n* = 3), and 27.8% obese (*n* = 10).

3.3. Work-to-rest ratio

The average work and rest time are provided with ratios in Table 2 by league. Work time was significantly greater for games than practices (98 ± 18 min vs. 89 ± 25 min; χ^2 (1,111) = 4.232, *p* = 0.039), as were rest time (29 ± 15 min vs. 9 ± 6 min; χ^2 (1,111) = 43.41, *p* < 0.001). However, work-to-rest ratios were significantly higher for practices ((13 ± 7):1 vs. (4 ± 1):1; χ^2 (1,111) = 40.62, *p* < 0.001).

3.4. Environmental conditions

The maximum and average WBGT conditions are presented in Fig. 1 by league and events. Of the practices and games observed

(*n* = 11) in each league, 9% (*n* = 2) were held in black flag conditions, 68% (*n* = 15) were held in red flag conditions, 18% (*n* = 4) were held in yellow flag conditions, and 5% (*n* = 1) were held in green flag conditions, according to the American College of Sports Medicine WBGT risk table.²⁰ The average (*t* = 0.468, *p* = 0.641) and maximum (*t* = 0.662, *p* = 0.534) practice WBGT compared with game WBGT were not significantly different. However, there was a significant difference (*t* = 4.402, *p* < 0.001) for the WR league's average WBGT compared with the AR league in games and all events. Upon further analysis with pairwise comparisons, the WR league's average and maximum WBGT was significantly higher (*t* = 11.158, *p* < 0.001) than the AR league's WBGT during games. There were no other significant comparisons (*p* > 0.05).

3.5. Perceptual and physiological responses

The average practice and game ESQ score were 8 ± 7 (0–42) and 7 ± 8 (0–30) (mean ± SD (minimum–maximum)), respectively. There were no significant differences between events (*t* = 1.121, *p* = 0.263). During games, individual *T*_{GI} and HR responses ranged from 37.5°C to 39.3°C and 80 bpm to 224 bpm, respectively. Practices ranged from 36.8°C to 39.2°C and 75 bpm to 200 bpm, respectively, for the same variables. Maximum measurements for physiological variables are presented in Fig. 2. There

Table 2
Work time, rest time, and work-to-rest ratios for all youth football teams, by league and event type (mean ± SD).

Variable	Aggregate (<i>n</i> = 12 teams)			Weight-restricted league (<i>n</i> = 4 teams)			Age-restricted league (<i>n</i> = 8 teams)		
	Work (min)	Rest (min)	Ratio	Work (min)	Rest (min)	Ratio	Work (min)	Rest (min)	Ratio
Practice	89 ± 25	9 ± 6	(13 ± 7):1**	108 ± 26	13 ± 6	(11 ± 6):1	75 ± 13	7 ± 4	(14 ± 8):1
Game	98 ± 18*	29 ± 15**	(4 ± 1):1	103 ± 23	43 ± 16	(3 ± 1):1	95 ± 15	20 ± 3	(5 ± 1):1

Note: Practice data were collected from 8 sessions; game data were collected from 3 games.

* *p* < 0.05, ** *p* < 0.001, compared with game determined by the Kruskal–Wallis test.




Fig. 1. Maximum WBGT (A) and average WBGT (B) between games ($n = 3$) and practices ($n = 8$) for each league. Red flag (dotted line), black flag (solid line) American College of Sports Medicine recommendations.²⁰ The maximum WBGT was derived by extracting the highest WBGT measurement recorded for each event and then averaged; the average WBGT was calculated using all collected measurements for each event. * $p < 0.05$, compared with AR league determined by independent t tests. Agg = aggregate; AR = age-restricted; WBGT = wet bulb globe temperature; WR = weight-restricted.

were no significant correlations between physiological variables and work-to-rest ratio, WBGT, or BMI (all $p > 0.05$).

4. Discussion

4.1. Risk factor history

Preparticipation examinations are recommended for youth who participate in sports,³⁶ yet very few recreational youth leagues require them. One league within our study required preparticipation examinations and the other did not. Preparticipation examinations allow pediatricians to assess the youth's medical and family history to determine if any conditions may predispose him or her to unnecessary risk during sports.³⁶ Almost one-third of our participants indicated that they were currently taking medications. Certain medications have the potential to change body temperature,⁹ decreasing the heat storage capacity of the youth player.

A previous history of an EHI and a lack of sleep are individual risk factors that have been examined in adult populations.^{8,20} Even though a previous history of EHI was not directly assessed in our study, participating youth did not seem to understand what constituted an EHI episode but reported symptoms alluding to previous occurrences (i.e., struggling, faint, dizzy, nauseous, and falling down). Survey research has indicated a similar lack of knowledge among high school




Fig. 2. Maximum T_{GI} (A) and maximum HR (B) between games ($n = 3$) and practices ($n = 3$) for each league. T_{GI} and HR were derived by extracting the highest measurement recorded for each player during each event and then averaged. Agg = aggregate; AR = age-restricted; HR = heart rate; T_{GI} = gastrointestinal temperature; WR = weight-restricted.

football players.²⁴ As previously reported,³⁷ sleep habits in our subjects were good throughout the study observation period. This EHI risk factor was not evident in our players and instead seems to manifest itself as youth enter high school.³⁸

Uniforms are a considerable risk factor because they increase physiological strain in hot conditions during full and partial configurations.¹⁵ The youth teams involved in our study wore partial uniforms during the first week of practices before full pad configurations were added at the beginning of their second week. However, both leagues approached integration differently, and full uniform configurations were never removed after incorporation despite environmental conditions in red flag conditions (26°C – 29°C).³⁹ Youth take longer to acclimatize to the heat.⁴⁰ Not allowing players to fully make adaptations before adding increased stressors (uniforms) increases EHI risk.

4.2. Anthropometrics

Anthropometric differences existed between leagues, revealing that players in the AR league were taller, heavier, and had a higher BMI and larger BSA than players in the WR league. League formation influences the types of players who participate, thereby impacting risk factors within a league. Larger athletes may be forced to participate in a different league or drop out owing to the difficulty of “making weight” for games.

Nevertheless, we found some portion of players (approximately 30%) in each league who were either overweight or obese according to criteria set forth by the Centers for Disease Control and Prevention. Our proportion of players was lower than in previous research,⁴¹ possibly owing to a smaller sample size. Overweight and obese categories are a risk factor that both leagues' organizations should consider when developing safety guidelines. Players with a high BMI are at risk for an EHI because of greater metabolic heat production during football.⁴²

It is also important to remember that, although they are not inferior, prepubertal athletes thermoregulate differently than adults.²⁸ Evaporative sweat is the principal means of heat dispersal during exercise in hot climatic conditions for adults. However, prepubertal boys have a lower overall sweat rate, lower sweat output per gland, and decreased sensitivity of sweat gland output in response to a given ambient temperature.⁴⁰ With evaporative heat loss minimized, prepubertal boys instead rely heavily on conductive and convective heat loss mechanisms. During thermoregulation, greater BSA as compared with mass results in a greater exchange of heat.^{28,43} Yet in hot temperatures this relationship is detrimental because the higher gradient from ambient temperature to skin reduces heat loss.²⁸ Maturational thermoregulatory differences in youth players lead to a slightly longer acclimatization process.⁴⁰

4.3. Work and rest variables

Leagues had a greater work-to-rest ratio during practice compared with games. Work length, without adequate time to rest and particularly in high-intensity situations, is a primary risk factor of exertional heat stroke and exertional sickling.^{8,9,20} Adequate rest time should be a priority for this population during practices and games to encourage heat dissipation and provide fair playing time. An informal study of the National Football League reported an average 10:1 ratio during games,⁴⁴ whereas a high school scrimmage had a 6:1 ratio.⁴⁵ The National Football League ratio is much greater than our game data owing to inherent differences, yet the high school ratio was similar to ours. Only 1 other study has examined practice exercise and rest intervals in a football team, which was at the collegiate level. Even though it was not a primary outcome measure, a 6:1 ratio could be calculated from the data given in the study.⁴⁶ Our practice ratio was, surprisingly, much higher for youth football players. The difference could be explained by a more accurate assessment of intervals by global positioning system units as compared with our observational timing of the whole team, yet our data call for attention to be drawn to the youth football population. Coaching and medical staff should plan to make work-to-rest ratios during practice closer to what the youth athlete will experience during games. Appropriate ratios are especially important on days with hot environmental conditions to help decrease EHI risk in this population.^{14,25}

4.4. Environmental conditions

Leagues in our study participated in events with stressful environmental conditions, as is typical in the southeast. Both

leagues participated in games or practice during 1 black flag condition²⁰ (WBGT of $>29^{\circ}\text{C}$). These temperatures could primarily be attributed to the start date of practices. Practices began in early August for WR leagues, similar to colleges and high schools; this time is the hottest of the year across the nation. Research clearly indicates that the majority of EHI occurs in the summer months when the temperatures are at the highest,^{12,47} particularly the first one-half of August.¹⁴ However, owing to starting 25 days later in August, players in AR leagues experienced milder maximum WBGT conditions. Some of the more alarming temperatures were during WR games in which a maximum WBGT of 34°C was recorded. The difference in temperatures that players in the WR and AR leagues experienced was due the time the games or practices were scheduled. The WR leagues played their games were between 9:00 a.m. and 2:00 p.m., the hottest part of the day.¹⁴ AR leagues played their games or practiced during evening hours (6:00 p.m. to 8:00 p.m.). The timing of football practices and games may play a significant role in EHI rates⁵ and therefore should be considered when scheduling events.

4.5. Perceptual and physiological responses

Perceptual response to thermal strain (as measured by ESQ scores) did not differ between games and practices. Yet the score ranges for practice and games were both wide, with some players reporting scores upwards of 42 (of a maximum of 70). These scores indicate a higher number of heat-related symptoms experienced, with greater severity. Players routinely experienced symptoms such as headaches, dizziness, nausea, and feeling hot. Other symptoms reported were thirst, tiredness, and trouble concentrating. This study was the first time an ESQ had been used in the youth football population, but the average and range of scores were similar to those of high school football players during preseason practices.²⁴

T_{GI} also varied widely depending on event type and the time point within the event. Of those players who participated in this portion of the study, 56% ($n=9$) reached a T_{GI} of 39°C or higher at least once during the observation period, but no players reached a T_{GI} indicative of exertional heat stroke (40.5°C).⁹ Previous research has examined natural fluctuations in physiological variables during football practices,^{16,19,24,46} but not during game situations or in the youth football population. Our results can provide a descriptive foundation of responses for future research. There were no correlations between the physiological variables and environmental conditions or anthropometrics. Typically, there is a relationship with anthropometrics with T_{GI} , with higher T_{GI} in larger individuals,^{13,16,46} but our sample size was small and relatively homogeneous owing to temperature sensor criteria, which may explain the lack of correlations.

4.6. Limitations

Work and rest times were monitored for the team as a whole and, therefore, cannot accurately represent what an individual player may have experienced. Physiological variables were measured only in a subset of players and events and,

Table 3
Recommendations based on dependent variables.

Variable	Recommendation
Preparticipation examinations	Leagues should mandate examinations by a pediatrician to evaluate exertional heat illness risk factors.
Uniform integration	Gradual integration of uniforms during the heat acclimatization process, and guidelines for removal based on environmental conditions, should be consistent across organizations.
Anthropometrics	Adequate rest breaks should be provided to all youth players when intense exercise is a goal, but particularly in leagues with higher percentages of overweight and obese youth. Local youth teams may consider the creation of extra shade locations, the use of fans, and other means of cooling to promote conductive and convective heat loss.
Work-to-rest ratio	Work-to-rest ratio modifications based on environmental conditions exist and should be supported by national leagues for local teams to adopt to ensure that youth participate in a safety-centric atmosphere.
Environmental conditions	Teams should practice in the evening hours. Morning hours can be used for games, but multiple fields should be used to decrease the number of hours players are exposed to heat.
General	Youth leagues should begin to provide medical coverage, such as a certified athletic trainer to help mitigate exertional heat illness risk factors.

therefore, do not provide a complete picture of the thermal strain in this population. Practice times and lengths were determined by the league or the coach and, therefore, were not consistent across events. One league cannot fully represent a league type; therefore, future research should include more leagues of each type (AR and WR) to confirm the differences we found and provide the capability to expand comparisons to more variables. Recommendations (Table 3) can be made based on the dependent variables from the current study and previous literature. However, future research should also examine a greater number of players over the first 2 weeks of practices to provide better descriptions of thermal stress during the preseason in this population.

5. Conclusion

Extrinsic and intrinsic EHI risk factors do exist in youth football leagues. Some leagues do not mandate a preparticipation physical, which removes the pediatrician's ability to screen for sport-specific injury risk. Even at this young age, a previous history of EHI, thermoregulatory-affecting medications, and the presence of general illnesses during football are possible. Once full uniforms were integrated, they were not removed for environmental conditions. WR and AR leagues influence the anthropometrics of players who participate in those leagues, with overweight and obese players still being prevalent in both types. During practices, ratios indicate that players work more with little rest, which is of concern when leagues practice in hot conditions. Rest break guidelines similar to those adopted by organizations in older competitive leagues should be used by youth football coaches. Some player's experienced T_{GI} temperatures of greater than 39°C, but all remained below the EHI threshold for both practices and games. National leagues need to take steps to ensure their handbooks include accurate heat safety information and comprehensive guidelines for local leagues to adopt.

Acknowledgments

We would like to acknowledge the hard work and significant time commitment of all the research assistants who

helped during data collection. We would also like to thank the players and their parents for their willingness to participate in the study and patience during data collection. Last, we recognize Ross Hayden for his help with statistical analysis. This work was supported in part by the University of South Carolina's College of Education Internal Grant Program. The study was also supported in part by the Master's Grant program of the Datalys Center for Sports Injury Prevention and Research. Funding sources did not influence the study design, data collection, analysis, or writing of the manuscript.

Authors' contributions

SWY and JJD developed the research questions, study design, planned and executed data collection, aided in the statistical analysis, interpreted the analysis, and drafted the manuscript; DME, JK, and TMTM helped with study design, executed data collection, interpreted the analysis, and helped to draft the manuscript; ZYK conducted the statistical analysis, interpreted the analysis, and helped to draft the manuscript. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

References

1. Mueller FO, Colgate B. *Annual survey of football injury research: 1931–2011*. Chapel Hill, NC: University of North Carolina; 2012.
2. Boden BP, Breit I, Beachler JA, Williams A, Mueller FO. Fatalities in high school and college football players. *Am J Sports Med* 2013;**41**:1108–16.
3. Maron BJ, Doerer JJ, Haas TS, Tierney DM, Mueller FO. Sudden deaths in young competitive athletes: analysis of 1866 deaths in the United States, 1980–2006. *Circulation* 2009;**119**:1085–92.
4. Nelson NG, Collins CL, Comstock RD, McKenzie LB. Exertional heat-related injuries treated in emergency departments in the U.S., 1997–2006. *Am J Prev Med* 2011;**40**:54–60.
5. Yeargin SW, Kerr ZY, Casa DJ, et al. Epidemiology of exertional heat illnesses in youth, high school, and college football. *Med Sci Sports Exerc* 2016;**48**:1523–9.

6. Kerr ZY, Casa DJ, Marshall SW, Comstock RD. Epidemiology of exertional heat illness among U.S. high school athletes. *Am J Prev Med* 2013;**44**:8–14.
7. Dick R, Ferrara MS, Agel J, et al. Descriptive epidemiology of collegiate men's football injuries: National Collegiate Athletic Association Injury Surveillance System, 1988–1989 through 2003–2004. *J Athl Train* 2007;**42**:221–33.
8. Rav-Acha M, Hadad E, Epstein Y, Heled Y, Moran DS. Fatal exertional heat stroke: a case series. *Am J Med Sci* 2004;**328**:84–7.
9. Casa DJ, DeMartini JK, Bergeron MF, et al. National athletic trainers' association position statement: exertional heat illnesses. *J Athl Train* 2015;**50**:986–1000.
10. Kenny GP, Niedre PC. The effect of exercise intensity on the post-exercise esophageal temperature response. *Eur J Appl Physiol* 2002;**86**:342–6.
11. Mora-Rodriguez R, Del Coso J, Estevez E. Thermoregulatory responses to constant versus variable-intensity exercise in the heat. *Med Sci Sports Exerc* 2008;**40**:1945–52.
12. Cooper ER, Ferrara MS, Broglio SP. Exertional heat illness and environmental conditions during a single football season in the southeast. *J Athl Train* 2006;**41**:332–6.
13. Godek SF, Bartolozzi AR, Burkholder R, Sugarman E, Dorshimer G. Core temperature and percentage of dehydration in professional football linemen and backs during preseason practices. *J Athl Train* 2006;**41**:8–14.
14. Yeargin SW, Cahoon E, Hosokawa Y, Mensch JM, Dompier TP, Kerr ZY. Environmental conditions and seasonal variables in American youth football leagues. *Clin Pediatr (Phila)* 2017;**56**:1209–18.
15. Armstrong LE, Johnson EC, Casa DJ, et al. The American football uniform: uncompensable heat stress and hyperthermic exhaustion. *J Athl Train* 2010;**45**:117–27.
16. Fowkes Godek S, Godek JJ, Bartolozzi AR. Thermal responses in football and cross-country athletes during their respective practices in a hot environment. *J Athl Train* 2004;**39**:235–40.
17. Carter 3rd R, Chevront SN, Williams JO, et al. Epidemiology of hospitalizations and deaths from heat illness in soldiers. *Med Sci Sports Exerc* 2005;**37**:1338–44.
18. DeMartini JK, Casa DJ, Belval LN, et al. Environmental conditions and the occurrence of exertional heat illnesses and exertional heat stroke at the Falmouth Road Race. *J Athl Train* 2014;**49**:478–85.
19. Yeargin SW, Casa DJ, Armstrong LE, et al. Heat acclimatization and hydration status of American football players during initial summer workouts. *J Strength Cond Res* 2006;**20**:463–70.
20. American College of Sports Medicine, Armstrong LE, Casa DJ, Millard-Stafford M, Moran DS, Pyne SW, Roberts WO. American College of Sports Medicine position stand. Exertional heat illness during training and competition. *Med Sci Sports Exerc* 2007;**39**:556–72.
21. Huffman EA, Yard EE, Fields SK, Collins CL, Comstock RD. Epidemiology of rare injuries and conditions among United States high school athletes during the 2005–2006 and 2006–2007 school years. *J Athl Train* 2008;**43**:624–30.
22. Kerr ZY, Marshall SW, Comstock RD, Casa DJ. Exertional heat stroke management strategies in United States high school football. *Am J Sports Med* 2014;**42**:70–7.
23. Yard EE, Gilchrist J, Haileyesus T, et al. Heat illness among high school athletes—United States, 2005–2009. *J Safety Res* 2010;**41**:471–4.
24. Yeargin SW, Casa DJ, Judelson DA, et al. Thermoregulatory responses and hydration practices in heat-acclimatized adolescents during preseason high school football. *J Athl Train* 2010;**45**:136–46.
25. Bergeron MF. Youth sports in the heat: recovery and scheduling considerations for tournament play. *Sports Med* 2009;**39**:513–22.
26. Bergeron MF, Laird MD, Marinik EL, Brenner JS, Waller JL. Repeated-bout exercise in the heat in young athletes: physiological strain and perceptual responses. *J Appl Physiol (1985)* 2009;**106**:476–85.
27. McDermott BP, Casa DJ, Yeargin SW, Ganio MS, Lopez RM, Mooradian EA. Hydration status, sweat rates, and rehydration education of youth football campers. *J Sport Rehabil* 2009;**18**:535–52.
28. Rowland T. Thermoregulation during exercise in the heat in children: old concepts revisited. *J Appl Physiol (1985)* 2008;**105**:718–24.
29. Kerr ZY, Yeargin SW, Djoko A, Dalton SL, Niceley MM, Dompier TP. Examining play counts and measurements of injury incidence in youth football. *J Athl Train* 2017;**52**:955–65.
30. Casa DJ, Becker SM, Ganio MS, et al. Validity of devices that assess body temperature during outdoor exercise in the heat. *J Athl Train* 2007;**42**:333–42.
31. Fonseca H, Silva AM, Matos MG, et al. Validity of BMI based on self-reported weight and height in adolescents. *Acta Paediatr* 2010;**99**:83–8.
32. Freedman DS, Sherry B. The validity of BMI as an indicator of body fatness and risk among children. *Pediatrics* 2009;**124**(Suppl 1):S23–34.
33. El Edelmi R, Lindemalm S, Eksborg S. Estimation of body surface area in various childhood ages—validation of the Mosteller formula. *Acta Paediatr* 2012;**101**:540–4.
34. Sampson JB, Kobrick JL. The environmental symptoms questionnaire: revisions and new filed data. *Aviat Space Environ Med* 1980;**51**:872–7.
35. Yamamoto LM, Casa DJ, Stearns RL, et al. Validation of a modified Environmental Symptoms Questionnaire for exercise in the heat. *Med Sci Sports Exerc* 2008;**40**:S190. doi:10.1249/01.mss.0000322285.63050.08.
36. LaBotz M, Bernhardt D. Preparticipation physical evaluation. *Open Access J Sports Med* 2013;**4**:151–60.
37. Galland BC, Taylor BJ, Elder DE, Herbison P. Normal sleep patterns in infants and children: a systematic review of observational studies. *Sleep Med Rev* 2012;**16**:213–22.
38. Eaton DK, McKnight-Eily LR, Lowry R, Perry GS, Presley-Cantrell L, Croft JB. Prevalence of insufficient, borderline, and optimal hours of sleep among high school students—United States, 2007. *J Adolesc Health* 2010;**46**:399–401.
39. Casa DJ, Csillan D. Inter-association task force for preseason secondary school athletics. Preseason heat-acclimatization guidelines for secondary school athletics. *J Athl Train* 2009;**44**:332–3.
40. Anderson SJ, Griesemer BA, Johnson MD, Martin TJ, Small E. Climatic heat stress and the exercising child and adolescent. *Pediatrics* 2000;**106**:158–9.
41. Malina RM, Morano PJ, Barron M, et al. Overweight and obesity among youth participants in American football. *J Pediatr* 2007;**151**:378–82.
42. Chung NK, Pin CH. Obesity and the occurrence of heat disorders. *Mil Med* 1996;**161**:739–42.
43. Tsuzuki-Hayakawa K, Tochiwara Y, Ohnaka T. Thermoregulation during heat exposure of young children compared to their mothers. *Eur J Appl Physiol Occup Physiol* 1995;**72**:12–7.
44. Biderman D. 11 minutes of action. *Wall Street Journal* January 15, 2010.
45. Gleason BH, Sams M, Salley JT, Pustina A, Stone MH. GPS analysis of a high school football scrimmage. *J Strength Cond Res* 2016;**31**:2183–8.
46. DeMartini-Nolan JK, Martschinske JL, Casa DJ, et al. Examining the influence of exercise intensity and hydration on gastrointestinal temperature in collegiate football players. *J Strength Cond Res* 2018;**32**:2888–96.
47. Grundstein AJ, Ramseyer C, Zhao F, et al. A retrospective analysis of American football hyperthermia deaths in the United States. *Int J Biometeorol* 2012;**56**:11–20.