

RESEARCH ARTICLE

Open Access

# Impact of gene variants on sex-specific regulation of human Scavenger receptor class B type 1 (SR-BI) expression in liver and association with lipid levels in a population-based study

Ornit Chiba-Falek<sup>1</sup>, Marshall Nichols<sup>1</sup>, Sunil Suchindran<sup>1</sup>, John Guyton<sup>2</sup>, Geoffrey S Ginsburg<sup>1</sup>, Elizabeth Barrett-Connor<sup>3</sup>, Jeanette J McCarthy<sup>1\*</sup>

## Abstract

**Background:** Several studies have noted that genetic variants of *SCARB1*, a lipoprotein receptor involved in reverse cholesterol transport, are associated with serum lipid levels in a sex-dependent fashion. However, the mechanism underlying this gene by sex interaction has not been explored.

**Methods:** We utilized both epidemiological and molecular methods to study how estrogen and gene variants interact to influence *SCARB1* expression and lipid levels. Interaction between 35 *SCARB1* haplotype-tagged polymorphisms and endogenous estradiol levels was assessed in 498 postmenopausal Caucasian women from the population-based Rancho Bernardo Study. We further examined associated variants with overall and *SCARB1* splice variant (SR-BI and SR-BII) expression in 91 human liver tissues using quantitative real-time PCR.

**Results:** Several variants on a haplotype block spanning intron 11 to intron 12 of *SCARB1* showed significant gene by estradiol interaction affecting serum lipid levels, the strongest for rs838895 with HDL-cholesterol ( $p = 9.2 \times 10^{-4}$ ) and triglycerides ( $p = 1.3 \times 10^{-3}$ ) and the triglyceride:HDL cholesterol ratio ( $p = 2.7 \times 10^{-4}$ ). These same variants were associated with expression of the SR-BI isoform in a sex-specific fashion, with the strongest association found among liver tissue from 52 young women <45 years old ( $p = 0.002$ ).

**Conclusions:** Estrogen and *SCARB1* genotype may act synergistically to regulate expression of *SCARB1* isoforms and impact serum levels of HDL cholesterol and triglycerides. This work highlights the importance of considering sex-dependent effects of gene variants on serum lipid levels.

## Background

The scavenger receptor class B type 1 (SR-BI) is a plasma membrane protein that binds high density lipoprotein (HDL) with high affinity and mediates selective uptake of cholesterol esters by the liver[1,2]. Besides its role as a functional HDL receptor involved in reverse cholesterol transport, SR-BI also participates in the metabolism of Apolipoprotein B-containing lipoproteins, including low density lipoprotein (LDL)[3,4] and very low density lipoprotein (VLDL)[5,6]. In addition, studies have implicated SR-BI as a key co-receptor mediating

infection with the hepatitis C virus[7], where chronic infection is characterized by marked lipid changes reflecting viral dependence on host lipid metabolism for replication and assembly[8]. The full length gene encoding SR-BI (gene symbol *SCARB1*) is comprised of 13 exons that are alternatively spliced to produce two major transcripts: the full length SR-BI and the splice variant SR-BII, in which exon 12 is skipped. SR-BI and SR-BII splice forms are conserved in both the mouse and rat genomes, have different tissue distributions and may influence cellular cholesterol trafficking and homeostasis in different ways[9]. SR-BII is reported to be a minor splice variant in human liver and has been shown to be less efficient at reverse cholesterol transport[10].

\* Correspondence: jeanette.mccarthy@duke.edu

<sup>1</sup>Institute for Genome Sciences and Policy, Duke University Medical Center, Durham, NC 27710, USA

In humans, with the exception of genome-wide association studies, targeted investigations into the association between *SCARB1* single nucleotide polymorphisms (SNPs) and serum lipids have been limited to a handful of polymorphisms in either exon 1, intron 5 or exon 8. These studies have found polymorphisms of *SCARB1* associated with serum levels of triglycerides [11-14], HDL cholesterol [11-18], LDL cholesterol [11,12,17] and VLDL [12]. Interestingly, many of these studies show different effects of the polymorphisms in men and women, suggesting a possible mediating role of sex hormones [11-15,17,18].

Estrogen is known to have a profound impact on serum lipid levels, resulting in a decrease in LDL cholesterol and triglycerides, affording younger women relative protection from coronary artery disease [19]. Estrogen is also a potent regulator of *SCARB1*, influencing the relative expression of SR-BI and SR-BII isoforms [20]. In rodent models, treatment with high doses of estrogen suppressed liver expression of full length SR-BI isoform, but increased expression of the splice variant SR-BII [21,26]. Similarly, treatment of human HepG2 liver cell lines with estradiol has been shown to result in a down-regulation of SR-BI and up-regulation of the SR-BII splice form [24,27].

We previously described how the association between *SCARB1* gene variants and HDL cholesterol levels varied by estrogen therapy status among postmenopausal women from the community-based Rancho Bernardo Study [28]. In the current study, we took a comprehensive SNP tagging approach to identify *SCARB1* genetic variation influencing lipid levels in an estrogen dependent manner. We examined interaction between *SCARB1* genotypes and endogenous estradiol levels among postmenopausal women, where levels of estradiol are lower than premenopausal women but not subject to monthly fluctuations. We further explored the functional consequences of *SCARB1* genetic variation on alternative splicing of *SCARB1* in a large series of human liver tissues. We report here how both endogenous estrogen and *SCARB1* genotype synergistically influence serum levels of HDL cholesterol and triglycerides in human populations and how this effect may be mediated through expression of the SR-BI isoform.

## Methods

### Rancho Bernardo Study

The Rancho Bernardo Study is a population-based cohort of white men and women, established during 1972-1974 in the suburban community of Rancho Bernardo, California [29]. The present study utilizes medical history, current medication use and health behaviors assessed from standardized questionnaires, anthropometric measures, and laboratory measures collected on

women between 1984 and 1987 during a follow-up visit focusing on diabetes and cardiovascular disease (visit 4). At the time of this visit, the average age of subjects was 70 years and the vast majority of women were postmenopausal. Therefore, we restricted our analyses to postmenopausal women. Subjects who participated in this visit provided morning blood samples obtained after an overnight fast and used to measure lipid and lipoprotein levels. Fasting serum cholesterol, triglyceride and HDL cholesterol levels were measured in a Center for Disease Control-Certified Lipid Research Clinic laboratory. Total cholesterol and triglyceride levels were measured by enzymatic techniques using an ABA-200 biochromatic analyzer (Abbott Laboratories, Irving, TX). HDL cholesterol was measured after precipitation of the other lipoproteins with heparin and manganese chloride. LDL cholesterol was estimated using the Friedewald formula [30]. Current use of estrogen therapy was validated by examination of pills or prescriptions brought to the clinic for that purpose and was classified as current versus non-current use. Endogenous estradiol in non-hormone using women was measured between 1992 and 1993 by radioimmunoassay after solvent extraction and column-chromatography [31] using first-thawed fasting morning specimens from the 1984-1987 venipuncture. Measurements were performed in the endocrinology research laboratory of S. S. C. Yen (Department of Reproductive Medicine, University of California-San Diego, La Jolla, CA). Serum estradiol is stable for >10 years when kept frozen at -70C (personal communication from S. Yen). The sensitivity and the intra- and inter-assay coefficients of variation, respectively, were 11 pmol/L, 5.9% and 7.1%. Because of a skewed distribution, estradiol measures were grouped into tertiles with the following cut-offs: low (7-17 pmol/L); medium (18-24 pmol/L); high (25-58 pmol/L). The 52 women with estradiol levels below the lower limit of detection (representing about 10% of the cohort) were included in the low estradiol group. Five women with estradiol levels above normal for postmenopausal women (>217 pmol/L) were excluded.

There were a total of 899 post-menopausal women who participated in visit 4 and provided a blood sample for DNA. There were 83 *SCARB1* SNPs with minor allele frequency >0.05 in Caucasians in the HapMap database. The Tagger program in Haploview software version 3.2 [32] was used to select 35 haplotype tagged SNPs in the *SCARB1* gene having minor allele frequencies of >0.05 and capturing all alleles with correlations ( $r^2$ ) >0.80 among Caucasians (Table 1). Of the 899 women with blood samples available, 793 were successfully genotyped for *SCARB1* SNPs. We further excluded six women missing primary lipid data; an additional nine women on lipid lowering therapy; and 280 women

**Table 1 SCARB1 SNPs evaluated in the Rancho Bernardo Study.**

	Location	rs number	Minor allele	Allele frequency	Hardy-Weinberg p value
Tag SNPs					
	3'	rs838881	G	0.31	0.07
	3'	rs838883	T	0.07	0.26
	IVS12	rs701106	A	0.16	0.41
	IVS12	rs7977729	C	0.29	0.60
	IVS11	rs838893	T	0.31	0.41
	IVS9	rs9919713	T	0.04	0.40
	IVS9	rs1031605	T	0.20	0.12
	IVS7	rs989892*	G	0.47	0.46
	IVS7	rs838905	C	0.04	0.31
	IVS6	rs838900	G	0.08	0.90
	IVS2	rs4765615	T	0.47	0.40
	IVS2	rs745529	T	0.37	0.006
	IVS1	rs11057830	A	0.15	0.45
	IVS1	rs10846745	G	0.44	0.85
	IVS1	rs7135223	A	0.40	0.12
	IVS1	rs12581963	A	0.08	0.69
	IVS1	rs12580803	G	0.17	0.59
	IVS1	rs10773107	A	0.48	0.50
	IVS1	rs10744182	C	0.42	0.84
	IVS1	rs4765180	T	0.43	0.97
	IVS1	rs7954519	G	0.20	0.44
	IVS1	rs4765621	T	0.33	0.12
	IVS1	rs12229555	C	0.21	0.86
	IVS1	rs10773109	G	0.45	0.90
	IVS1	rs12582221	A	0.44	0.48
	IVS1	rs12370382	T	0.39	0.84
	IVS1	rs3924313	A	0.33	0.51
	IVS1	rs11057851	A	0.11	0.53
	IVS1	rs11057852	T	0.05	0.76
	IVS1	rs10773111	T	0.42	0.38
	IVS1	rs4765181	A	0.39	0.99
	IVS1	rs11615630	T	0.40	1.00
	5'	rs4379922	C	0.37	1.00
	5'	rs10846760	T	0.36	1.00
Additional SNPs					
	IVS11	rs838892	A	0.31	0.47
	IVS11	rs838895	G	0.32	0.82
	IVS11	rs838896	C	0.34	0.52

\*Proxy SNP for exon 8 SNP evaluated in previous studies (rs5888)

currently taking estrogen therapy on whom estradiol measures were not measured, leaving a final sample of 498 women.

#### Liver Samples

Post-mortum liver tissues from 93 otherwise healthy Caucasian adults were purchased from the National Institute of Child Health and Human Development

Brain and Tissue Bank for Developmental Disorders (University of Maryland, Baltimore, MD). All had a post mortem interval (PMI) <24 hours. A demographic description is included in Table 2. The human hepatoma (HepG2) cell line was obtained from American Type Culture Collection (Manassas, VA).

#### RNA extraction and cDNA synthesis from liver tissue

Total RNA was extracted from liver sample (100 mg) using TRIzol reagent (Invitrogen, Carlsbad, CA) followed by purification with RNeasy kit using an on-column DNase treatment (Qiagen, Inc., Valencia, CA) following the manufacturer's protocol. RNA concentration was determined spectrophotometrically at 260 nm, while the quality of the purification was determined by 260 nm/280 nm ratio. Additionally, quality of sample and lack of significant degradation products was confirmed on an Agilent Bioanalyzer. Next, cDNA was synthesized using MultiScribe RT enzyme (Applied Biosystems, Foster City, CA) under the following conditions: 10 min at 25°C and 120 min at 37°C.

#### Real time PCR

Real-time PCR was used to quantify human *SCARB1* mRNA levels (SR-BI isoform, SR-BII isoform and overall *SCARB1*) using commercially available TaqMan assays. Briefly, each liver sample was assayed by quantitative real-time PCR using the ABI 7900 to determine the level of the target message relative to mRNA encoding the human peptidylprolyl isomerase A (*PPIA*). This housekeeping gene has been shown to be among the most stable in studies of mouse liver, making it ideal for normalization in liver gene expression studies[33]. Each cDNA (50 ng) was amplified in duplicate using TaqMan Gene Expression PCR master mix reagent (Applied Biosystems, Foster City, CA) under the following conditions: 2 min at 50°C, 10 min at 95°C, 40 cycles: 15 sec at 95°C, and 1 min at 60°C. The different target cDNAs were amplified using the following ABI MGB probe and primer set assays (Applied Biosystems, Foster City, CA): ID HS00969822\_m1 for overall *SCARB1*, targeting the junction of *SCARB1* exon 2-3 which is conserved across all isoforms; HS00969819\_m1 for SR-BI (includes exon 12), targeting the junction of exons 11-12; and a custom assay designed for SR-BII (excludes exon 12), with forward primer 5'TCCCTGTCATCTGCCAAATCC-3', reverse primer 5'-GGCTGGCTCACGGTGT-3' and probe 5'-CCTCAGGACCTTGGCTCC-3' targeting the junction of exons 11-13. All three mRNA assays were normalized to a *PPIA* mRNA control (ABI MGB probe and primer set assay ID Hs99999904\_m1).

The data were analyzed using the  $\Delta\Delta C_t$  method, with a threshold set in the linear range of amplification. The cycle number at which any particular sample crossed that threshold ( $C_t$ ) was then used to determine fold difference. Fold difference was calculated

**Table 2 Demographic characteristics and *SCARB1* SNP minor allele carrier frequencies in 93 post-mortem liver samples, obtained from otherwise healthy Caucasian adults.**

Characteristic	Males (all) (n = 41)	Females (all) (n = 52)	Females < 45 years old (n = 43)
Age (years)	40.9 ± 12.1	32.2 ± 13.8	27.8 ± 9.4
PMI (hours)	14.1 ± 5.9	17.2 ± 7.3	17.0 ± 7.5
rs7977729 (C)	0.41	0.50	0.49
rs838891 (G)	0.56	0.54	0.53
rs838892 (A)	0.56	0.53	0.52
rs838893 (T)	0.56	0.52	0.51
rs838895 (G)	0.54	0.54	0.53
rs838896 (C)	0.51	0.58	0.58
rs10846744 (G)	0.39	0.25	0.28

PMI = post mortem interval. Age and PMI are presented as means ± standard deviation.

as  $2^{-\Delta\Delta Ct}$ ;  $\Delta Ct = [Ct(target) - Ct(PPIA)]$ .  $\Delta\Delta Ct = [\Delta Ct(sample)] - [\Delta Ct(calibrator)]$ , where target is *SCARB1*, SR-BI or SR-BII. The calibrator was a single, randomly selected control liver sample used in each plate for normalization within and across runs and used in creation of standard curves. Each sample was run in duplicate in two independent plates, over all 4 repeats. The  $\Delta\Delta Ct$  results obtained with the four repeats were average to determine the fold expression level used in the association analysis.

We controlled for DNA contamination by running three randomly selected, RNA control samples that were not converted to cDNA and no-cDNA/RNA sample in each plate. No observable amplification was detected. In addition for assay validation we generated standard curves for each target and reference assay, using different amounts of human liver total RNA (1-1000 ng). In addition, the slope of the relative efficiency plot for each target and internal control were determined to validate the assays, i.e. *SCARB1*, SR-BI or SR-BII and *PPIA*. The slope in the relative efficiency plot for each target and *PPIA* were determined, and showed a standard value (<0.1) required for the validation of the relative quantitative method. (Additional file 1, Figure S1).

#### **SCARB1 genotyping**

Genomic DNA was isolated using conventional protocol by Qiagen kits (Qiagen, Inc., Valencia, CA). *SCARB1* genotypes were assayed in genomic DNA isolated from frozen whole blood samples from subjects in the Rancho Bernardo Study and from the liver tissues and HepG2 cell line using either the Sequenom iPLEX™ multiplex mass spectrometry genotyping system (Sequenom, Inc, La Jolla, CA) or using the 5' nuclease assay with allele specific Taq-Man probes[34]. The percent missing genotypes and Hardy-Weinberg Equilibrium were evaluated using Haploview software[32] as a means of quality control.

#### **Statistical analysis**

SAS statistical software, Version 9.1 (SAS Institutes, Cary, NC) was used for all statistical analyses. The

ggplot2 package[35] of the R language was also used for plots[36]. The association between each *SCARB1* SNP and levels of total cholesterol, LDL cholesterol, triglycerides, HDL cholesterol and the triglyceride:HDL cholesterol ratio in the Rancho Bernardo Study was assessed with Analysis of Variance (ANOVA). Triglycerides, HDL cholesterol, LDL cholesterol, and the TG/HDL cholesterol ratio were natural log (ln) transformed to improve normality of the residuals. SNP genotypes were coded in an additive model, except for SNPs where the minor allele frequency was below 5%, in which case heterozygous and homozygous variant genotypes were pooled to test a dominant model. All models controlled for age (continuous) and current estrogen therapy status. Interactions between SNPs and estradiol levels (based on tertile cutoffs) were determined by introduction of a cross-product term in the model.

Differences in overall *SCARB1*, SR-BI and SR-BII expression in liver samples were examined by sex and by presence/absence of *SCARB1* allelic variants using standard linear regression analysis, controlling for post-mortum interval and age. Genotypes were coded in a dominant model (homozygous variant and heterozygote genotypes pooled) because of small numbers. Genotype by sex interaction was determined by inclusion of a cross product term in the model. Expression levels were natural log (ln) transformed for analysis. The proportion of SR-BI and SR-BII out of total *SCARB1* was determined by calculating the  $\Delta\Delta Ct$  ratios (i.e. SR-BI or SR-BII/overall *SCARB1*).

This study was approved by the Institutional Review Boards at Duke University.

## **Results**

### **Genotype associations with lipid traits in women from the Rancho Bernardo Study**

Minor allele frequencies for the *SCARB1* SNPs typed in this study ranged from 0.04 to 0.44. The average call rate was 97% and one SNP had nominally significant

( $p < 0.006$ ) deviation from Hardy-Weinberg equilibrium (Table 1). Characteristics of women from the Rancho Bernardo Study are shown in Table 3. We modeled the interaction between the *SCARB1* SNPs and estradiol levels on serum total cholesterol, triglycerides, HDL cholesterol, LDL cholesterol and the triglyceride:HDL cholesterol ratio among women not currently taking estrogen therapy. Two SNPs on the same haplotype block showed significant evidence of genotype by estradiol interaction ( $p < 0.05$ ). The tag SNP rs838893 showed the strongest evidence of interaction for HDL cholesterol ( $p = 0.006$ ), triglycerides ( $p = 0.001$ ) and the triglyceride:HDL cholesterol ratio ( $p = 0.0005$ ). This common variant (minor allele frequency 0.31) lies on a haplotype block spanning intron 11 to intron 12 of *SCARB1*. We examined additional SNPs on this block, all of which showed a similar pattern of significant associations (Figure 1). Figure 2 illustrates how at low levels of estradiol, the variant allele, G', of the most significant of these SNPs, rs838895, was associated with lower levels of triglycerides and high levels of HDL cholesterol, but at increasing levels of estradiol, the opposite was found. Importantly, this SNP showed no association with estradiol levels ( $p = 0.40$ ). Results for another significant SNP, rs838896, are also shown in Figure 2 for comparison with results from the livers, presented below. Similar trends were seen for all assayed SNPs on that haplotype block. No significant interaction was found between any of the *SCARB1* tagging SNPs and current use of estrogen therapy for HDL-C ( $p = 0.34$ ) or triglycerides ( $p = 0.53$ ).

#### Expression levels and presence of alternative splice variants in liver tissue


Given the existence of alternative splice forms of *SCARB1* involving inclusion/exclusion of exon 12, we hypothesized that a SNP located within the associated haplotype block may modulate splicing efficiency. We quantified the amounts of the full-length SR-BI isoform (including exon 12), the alternative SR-BII isoform (exon 12 skipping) and overall *SCARB1* mRNA (all splice forms) in human liver tissues from 93 healthy individuals. SR-BI showed higher expression levels than SR-BII, consistent with other studies, but accounted for less than half of the overall *SCARB1* expression: on average SR-BI accounted for  $39 \pm 19\%$  and SR-BII  $3.5 \pm 1.8\%$  of overall *SCARB1* expression.


#### The effect of sex and genotype on alternative splicing of SCARB1 in human liver tissues

In order to determine the functional consequences of *SCARB1* SNPs, we carried out genetic association analysis of the associated SNPs and expression of *SCARB1*. We genotyped seven SNPs, including six within the associated haplotype block, which spans intron 11 to intron 12 of *SCARB1*, and one outside the region in 93

commercially available liver samples. All SNPs were in Hardy-Weinberg Equilibrium and showed linkage disequilibrium patterns consistent with those seen in the Rancho Bernardo Study (not shown). One outlier sample was removed after performing linear regression diagnostics and the reference liver sample was also not included, leaving a sample size of 91 (57% female). Because of the apparent estrogen dependency of *SCARB1* variants on lipid levels, we examined primary sex effects, primary genotype effects and interactive effects between sex and *SCARB1* genotype on mRNA expression.

Previous studies found that estrogen treatment of human HepG2 cells results in a decrease in SR-BI and increase of SR-BII[24,27]. First, as a proxy for estrogen effects, we examined male-female differences in liver expression of SR-BI, SR-BII and overall *SCARB1*. We found significantly lower levels of overall *SCARB1*, SR-BI isoform and SR-BII isoform in female versus male livers (Figure 3). Next, we examined the effect of each of the seven SNPs on expression in the combined liver samples from men and women, but no significant associations were found. We then repeated this analysis, assessing SNP by sex interaction for all SNPs. We found modest evidence of interaction between sex and *SCARB1* SNP rs838896 influencing expression of the SR-BI isoform ( $p = 0.09$ ), which became stronger when restricting analysis to subjects  $\leq 45$  years old ( $p = 0.04$ ), leading us to carry out sex-stratified analyses. As shown in Figure 4, presence (+) of the rs838896 minor allele, C, was associated with significantly lower levels of SR-BI in liver tissue from women ( $p = 0.01$ ) but not in men ( $p = 0.65$ ). Restricting the analysis to liver tissue from women  $< 45$  years of age, the genotype association became stronger: in younger women, presence of at least one copy of the rs838896C allele conferred a 38% lower level of SR-BI ( $p = 0.002$ ), a non-significant 26% lower level of SR-BII ( $p = 0.16$ ) and a 28% lower level of overall *SCARB1* ( $p = 0.02$ ), compared to young women not carrying this allele. Similar, but less significant associations were seen for other SNPs on the haplotype block (rs7977729, rs838891, rs838892, rs838893, rs838895, and rs838896). It is noteworthy that rs838895, which showed the strongest association with lipids in the Rancho Bernardo Study, did not reach significance in the liver gene expression analysis (e.g. in females  $< 45$ , the  $p$  values were 0.07, 0.13 and 0.54 for association with overall *SCARB1*, SR-BI and SR-BII expression). This may be due to differences in linkage disequilibrium between rs838895 and rs838896 in the two studies:  $r^2 = 0.80$  in Rancho Bernardo and  $r^2 = 0.62$  in the livers. No associations were found for the one SNP outside of the block (rs10846744). We then compared relative levels of each splice form (ratio to overall *SCARB1* expression) in





young women and found no significant differences by genotype: Carriers of the rs838896 minor allele (rs838896+) had on average  $41 \pm 4\%$  SR-BI and  $3.9 \pm 0.4\%$  SR-BII while non-carriers (rs838896-) had average levels of  $47 \pm 5\%$  and  $3.6 \pm 0.4\%$ , respectively.

**Discussion**

The results of our study in a community-based cohort of post-menopausal women demonstrate that polymorphisms of SCARB1 are associated with HDL cholesterol and triglyceride levels in an endogenous estrogen-

dependent fashion. Specifically, variant alleles of SCARB1 in a block encompassing intron 11 to intron 12 were associated with increased triglycerides and decreased HDL cholesterol in the presence of high levels of endogenous estradiol, but the opposite was seen at lower levels of estradiol. Importantly, these effects were observed among post-menopausal women who already have low levels of estradiol, suggesting that small changes in endogenous estradiol can impact the association of SCARB1 genotype on serum lipid levels.

**Table 3 Baseline characteristics of postmenopausal Caucasian women from the Rancho Bernardo Study in the overall cohort and by low, moderate or high estradiol (E2) tertile in the subset with estradiol measures available.**

Characteristic (mean ± SD)	All women	Low E2	Mod E2	High E2
Sample size	(784)	(180)	(140)	(173)
Age (years)	69.1 ± 8.6	71.1 ± 8.2	70.8 ± 8.4	68.5 ± 9.2
Waist circumference (cm)	78.0 ± 9.1	75.8 ± 8.0	78.2 ± 8.6	80.9 ± 9.6
BMI (kg/m <sup>2</sup> )	24.1 ± 3.6	23.0 ± 3.0	23.9 ± 2.9	25.5 ± 4.0
Total cholesterol (mmol/L)	5.9 ± 1.0	5.9 ± 1.1	6.1 ± 1.0	6.0 ± 0.94
HDL cholesterol (mmol/L)	1.8 ± 0.5	1.8 ± 0.45	1.8 ± 0.47	1.6 ± 0.43
LDL cholesterol (mmol/L)	3.5 ± 1.0	3.6 ± 1.0	3.7 ± 1.0	3.8 ± 0.9
Triglycerides (mmol/L)*	1.1 ± 0.8	1.0 ± 0.7	1.1 ± 0.8	1.2 ± 1.0
Estradiol levels (pmol/L)*	-	11.0 ± 7.3	18.4 ± 3.7	29.4 ± 11


\* Median and interquartile range

In contrast, we did not find evidence of interaction between these same genotypes and current use of oral estrogen therapy, as one might expect, nor did we find evidence of genotype by endogenous estradiol interaction for two SNPs reported to interact with oral estrogen therapy in our previous study of HDL cholesterol (data not shown). These discrepancies may be related to the marked perturbation of hepatic metabolism produced by oral estrogen therapy, including increased plasma levels of VLDL, and HDL, and reduced plasma LDL[37,38]. Oral estrogen is administered in high doses due to poor systemic bioavailability averaging 5%[39] and thus hepatocyte exposure to oral estrogen may be magnified, exceeding the physiological range. Another difference between oral and endogenous estrogen is that the former has typically involved equine estrogens (particularly during the period of sample collection for this study) whereas estradiol is the most active endogenous estrogen. Their effects on gene expression relevant to lipid metabolism may be quite different. Another


possibility is that the SNPs that interacted with estrogen therapy in our previous study could reflect other differences between users and non-users of estrogen therapy, the former being younger and having lower fasting glucose[28], as well as having lower testosterone (unpublished data).

We further demonstrate in the current study that in liver tissues from young, adult females (but not males) the *SCARB1* rs838896 minor allele from the identified haplotype block is associated with lower overall expression, driven in large part by the SR-BI isoform. Hence, increased expression of SR-BI may underlie the protective lipid profiles observed in women with both high levels of estradiol and homozygous for the rs838896 wildtype allele as observed in the Rancho Bernardo Study, and could account for underlying sex differences in *SCARB1* genetic associations with lipids described in previous studies[11-15,17,18].

Previous studies in rodent models and human HepG2 cell lines indicate that estrogen down-regulates SR-BI and up-regulates SR-BII isoforms[24,27]. Thus, one would expect that women (specifically pre-menopausal women with higher levels of estrogen) would have lower levels of SR-BI and higher levels of SR-BII than men. Our findings in a series of human liver tissues confirm that females have significantly lower levels of hepatic SR-BI expression compared to males, but SR-BII levels were not higher; in fact levels of SR-BII were lower in females, especially among carriers of the *SCARB1* rs838896 minor allele (the same genotype of HepG2 cell lines, data not shown). Although experimental results in human liver tissue may not equate with those in rodent liver or cultured human HepG2 cell lines, these discrepant findings warrant further investigation. In the study by Graf[27], antibodies specific to the 3' end of the *SCARB1* protein were used to distinguish SR-BI from SR-BII in human HepG2 cell lines treated with 17 estradiol. In the study by Zhang[24], treatment of HepG2 cell lines under the same conditions as Graf *et al.* did







**Figure 4 mRNA expression in human liver tissue by SCARB1 genotype and sex.** Results shown are average  $\Delta\Delta C_t$  values with standard error bars for A.) Full length SR-BI isoform; B.) SR-BII and C.) overall SCARB1, all relative to PPIA control. Rs838896+ refers to carrier of the G allele (GG and GC); Rs838896- refers to genotype CC.

not produce any changes in SR-BI or SR-BII. Only after subsequent introduction of estrogen receptor and treatment with synthetic equine estrogen did they recapitulate the increase in SR-BII expression noted by Graf *et al.* However, SR-BII was not measured directly, but was determined by calculating the difference between overall *SCARB1* and SR-BI levels. SR-BII, resulting from skipping of exon 12, is the most studied alternatively splice form, but may not be the only splice form. Additional splice forms have been described in genomic databases such as Entrez Gene Aceview[40], including one form that differs dramatically in the 3' end of the gene, missing both exons 12 and 13 and retaining a large segment of intron 11 as coding. We confirmed the expression of this splice form in human brain and liver tissues (data not shown). Thus, differences in our findings from others may relate to the method of detection of SR-BII expression and the possible existence of additional isoforms of *SCARB1* not accounted for in any published analyses to date.

Recently, Zhang and colleagues proposed that estrogen regulation of alternative splicing of *SCARB1* in the rat occurs via regulatory splicing factors that interact with regulatory sequences in intron 11 of *SCARB1*[24,41]. The human *SCARB1* polymorphisms most strongly associated with estrogen-dependent splicing and lipid traits in our study are located on a haplotype block spanning intron 11 to intron 12. The coincidental location of the putative regulatory region identify by Zhang and the presence of SNPs controlling SR-BI isoform expression in our study lead us to suspect that the polymorphism underlying our observed association impacts the estrogen-dependent regulation of *SCARB1* expression, perhaps of specific isoforms. Our data do not support an effect of *SCARB1* SNPs on relative amounts of SR-BI or SR-BII in liver, but the possibility that other isoforms are impacted by these gene variants should be explored. Whether the rs838896 variant is the causal variant or in linkage disequilibrium with the causal variant remains to be determined in follow up functional studies using cell culture and other systems.

Both HDL cholesterol and triglyceride levels are sexually dimorphic traits: levels of HDL cholesterol are 20-25% higher in women than men and these differences persist over time; triglyceride levels, on the other hand, start out slightly lower in women compared to men in young adulthood, but around the time of menopause, triglyceride levels rise dramatically in women[42]. We propose that this rise in triglycerides may be attributable in part to estrogen regulation of *SCARB1*, and that the protection afforded by estrogen in younger women may be limited to women with a certain *SCARB1* genotype. It is interesting to note that *SCARB1* is also a co-receptor for the Hepatitis C virus (HCV)

and that increased expression of both SR-BI and SR-BII isoforms increase infectivity *in vitro*[43]. Additionally, women generally have better outcomes of HCV infection, perhaps related to estrogen [44], leading us to speculate on the role of estrogen-mediated expression of *SCARB1* in outcomes of Hepatitis C infection.

Our study of the phenotypic consequences of *SCARB1* variation was carried out in a community-based sample of post-menopausal Caucasian women who were not selected based on any disease phenotype. The samples and data were collected before the widespread use of lipid lowering therapy, thus reducing confounding by treatment. Yet, there are some caveats of our study, including that estradiol assays were measured only once. Although single measurements of most hormones reliably characterize average levels over a two to three year period, estradiol levels are less reproducible[45,46]. However, estradiol was measured in morning fasting samples, and among postmenopausal women estradiol does not show significant diurnal variation[47]. Moreover, it should be noted that this was an elderly population of women, with relatively low levels of estradiol; we might expect estrogen effects to be more pronounced in a younger population. As with most large scale SNP association studies, ours is also prone to false positive associations, the result of multiple testing. Replication of the observed associations in other study populations is warranted.

## Conclusions

In summary, our work has detected sex-specific expression of *SCARB1* isoforms *in vivo* in human liver tissue and identified polymorphisms in intron 11 of the gene that modify this effect. We further found these same genetic variants to influence triglyceride and HDL cholesterol levels in an endogenous estrogen-dependent manner in a human population study. Thus, estrogen regulation of *SCARB1* may contribute to the observed sexual dimorphism of lipid levels and highlights the importance of considering sex-dependent effects of gene variants on serum lipid levels.

**Additional file 1: Figure S1. Validation curves of the  $\Delta\Delta$  real-time gene expression assays.** Shown are curves for overall *SCARB1*, SR-BI and SR-BII versus endogenous control (PPIA). Relative efficiency plots were formed by plotting the log input amount (ng of total RNA) versus the  $\Delta\text{Ct}$  ( $\text{Ct } SCARB1 - \text{Ct } PPIA$ ), for example. The slopes are  $<0.1$  which indicates the validation of the  $\Delta\Delta\text{Ct}$  calculation relative to the reference controls in the range between 1-100 ng RNA.  
Click here for file  
[ <http://www.biomedcentral.com/content/supplementary/1471-2350-11-9-S1.TIFF> ]

## Acknowledgements

We would like to thank Sonal Patel and Colton Linnertz for their technical assistance in the laboratory. This study was supported in part by grant RO1

HL085191-01 from the National Heart, Lung and Blood Institute (P.I. McCarthy), grant P60 MD00220, grant DK31801 from the National Institute of Diabetes and Digestive and Kidney Diseases (P.I. Barrett-Connor) and grant AG07181 (P.I. Barrett-Connor) from the National Institute on Aging. Liver tissues were purchased from the National Institute of Child Health and Human Development Brain and Tissue Bank for Developmental Disorders under contracts N01-HD-4-3368 and N01-HD-4-3383.

#### Author details

<sup>1</sup>Institute for Genome Sciences and Policy, Duke University Medical Center, Durham, NC 27710, USA. <sup>2</sup>Division of Endocrinology, Duke University Medical Center, Durham, NC 27710, USA. <sup>3</sup>Department of Family and Preventive Medicine, University of California at San Diego, San Diego, CA 92121, USA.

#### Authors' contributions

OCF contributed to the conception and design, oversaw the molecular genetic studies, and contributed to drafting of the manuscript; MN carried out the molecular genetic analysis; SS carried out the genetic association analysis; JG, GG and EBC participated in interpretation of data and manuscript revisions; JJM conceived and designed the study, drafted the manuscript and oversaw all genetic analyses. All authors have read and approved the final manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 13 July 2009

Accepted: 19 January 2010 Published: 19 January 2010

#### References

- Acton S, Rigotti A, Landschulz KT, Xu S, Hobbs HH, Krieger M: **Identification of scavenger receptor SR-BI as a high density lipoprotein receptor.** *Science* 1996, **271**(5248):518-520.
- Murao K, Terpstra V, Green SR, Kondratenko N, Steinberg D, Quehenberger O: **Characterization of CLA-1, a human homologue of rodent scavenger receptor BI, as a receptor for high density lipoprotein and apoptotic thymocytes.** *J Biol Chem* 1997, **272**(28):17551-17557.
- Trigatti BL, Rigotti A, Braun A: **Cellular and physiological roles of SR-BI, a lipoprotein receptor which mediates selective lipid uptake.** *Biochim Biophys Acta* 2000, **1529**(1-3):276-286.
- Brodeur MR, Luangrath V, Bourret G, Falstra L, Brissette L: **Physiological importance of SR-BI in the in vivo metabolism of human HDL and LDL in male and female mice.** *J Lipid Res* 2005, **46**(4):687-696.
- Van Eck M, Hoekstra M, Out R, Bos IS, Kruijt JK, Hildebrand RB, Van Berkel TJ: **Scavenger receptor BI facilitates the metabolism of VLDL lipoproteins in vivo.** *J Lipid Res* 2008, **49**(1):136-146.
- Hu L, Hoogt van der CC, Espirito Santo SM, Out R, Kypreos KE, van Vijmen BJ, Van Berkel TJ, Romijn JA, Havekes LM, van Dijk KW, et al: **The hepatic uptake of VLDL in lrp-ldlr-/vldlr- mice is regulated by LPL activity and involves proteoglycans and SR-BI.** *J Lipid Res* 2008, **49**(7):1553-1561.
- Dreux M, Dao Thi VL, Fresquet J, Guerin M, Julia Z, Verney G, Durantel D, Zoulim F, Lavillette D, Cosset FL, et al: **Receptor complementation and mutagenesis reveal SR-BI as an essential HCV entry factor and functionally imply its intra- and extra-cellular domains.** *PLoS Pathog* 2009, **5**(2):e1000310.
- Burlone ME, Budkowska A: **Hepatitis C virus cell entry: role of lipoproteins and cellular receptors.** *J Gen Virol* 2009, **90**(Pt 5):1055-1070.
- Eckhardt ER, Cai L, Sun B, Webb NR, Westhuyzen van der DR: **High density lipoprotein uptake by scavenger receptor SR-BII.** *J Biol Chem* 2004, **279**(14):14372-14381.
- Webb NR, Connell PM, Graf GA, Smart EJ, de Villiers WJ, de Beer FC, Westhuyzen van der DR: **SR-BII, an isoform of the scavenger receptor BI containing an alternate cytoplasmic tail, mediates lipid transfer between high density lipoprotein and cells.** *J Biol Chem* 1998, **273**(24):15241-15248.
- Acton S, Osgood D, Donoghue M, Corella D, Pocovi M, Cenarro A, Mozas P, Keilty J, Squazzo S, Woolf EA, et al: **Association of polymorphisms at the SR-BI gene locus with plasma lipid levels and body mass index in a white population.** *Arterioscler Thromb Vasc Biol* 1999, **19**(7):1734-1743.
- Tai ES, Adiconis X, Ordovas JM, Carmena-Ramon R, Real J, Corella D, Ascaso J, Carmena R: **Polymorphisms at the SRBI locus are associated with lipoprotein levels in subjects with heterozygous familial hypercholesterolemia.** *Clin Genet* 2003, **63**(1):53-58.
- McCarthy JJ, Lehner T, Reeves C, Moliterno DJ, Newby LK, Rogers WJ, Topol EJ: **Association of genetic variants in the HDL receptor, SR-B1, with abnormal lipids in women with coronary artery disease.** *J Med Genet* 2003, **40**(6):453-458.
- McCarthy JJ, Lewitzky S, Reeves C, Permutt A, Glaser B, Groop LC, Lehner T, Meyer JM: **Polymorphisms of the HDL receptor gene associated with HDL cholesterol levels in diabetic kindred from three populations.** *Hum Hered* 2003, **55**(4):163-170.
- Hong SH, Kim YR, Yoon YM, Min WK, Chun SI, Kim JQ: **Association between Haell polymorphism of scavenger receptor class B type I gene and plasma HDL-cholesterol concentration.** *Ann Clin Biochem* 2002, **39**(Pt 5):478-481.
- Hsu LA, Ko YL, Wu S, Teng MS, Peng TY, Chen CF, Lee YS: **Association between a novel 11-base pair deletion mutation in the promoter region of the scavenger receptor class B type I gene and plasma HDL cholesterol levels in Taiwanese Chinese.** *Arterioscler Thromb Vasc Biol* 2003, **23**(10):1869-1874.
- Morabia A, Ross BM, Costanza MC, Cayanis E, Flaherty MS, Alvin GB, Das K, James R, Yang AS, Evagrafov O, et al: **Population-based study of SR-BI genetic variation and lipid profile.** *Atherosclerosis* 2004, **175**(1):159-168.
- Roberts CG, Shen H, Mitchell BD, Damcott CM, Shuldiner AR, Rodriguez A: **Variants in scavenger receptor class B type I gene are associated with HDL cholesterol levels in younger women.** *Hum Hered* 2007, **64**(2):107-113.
- Bush TL, Fried LP, Barrett-Connor E: **Cholesterol, lipoproteins, and coronary heart disease in women.** *Clin Chem* 1988, **34**(8B):B60-70.
- Lopez D, McLean MP: **Estrogen regulation of the scavenger receptor class B gene: Anti-atherogenic or steroidogenic, is there a priority?.** *Mol Cell Endocrinol* 2006, **247**(1-2):22-33.
- Landschulz KT, Pathak RK, Rigotti A, Krieger M, Hobbs HH: **Regulation of scavenger receptor, class B, type I, a high density lipoprotein receptor, in liver and steroidogenic tissues of the rat.** *J Clin Invest* 1996, **98**(4):984-995.
- Fluiter K, Westhuyzen van der DR, van Berkel TJ: **In vivo regulation of scavenger receptor BI and the selective uptake of high density lipoprotein cholesteryl esters in rat liver parenchymal and Kupffer cells.** *J Biol Chem* 1998, **273**(14):8434-8438.
- Stangl H, Graf GA, Yu L, Cao G, Wyne K: **Effect of estrogen on scavenger receptor BI expression in the rat.** *J Endocrinol* 2002, **175**(3):663-672.
- Zhang X, Moor AN, Merkle KA, Liu Q, McLean MP: **Regulation of alternative splicing of liver scavenger receptor class B gene by estrogen and the involved regulatory splicing factors.** *Endocrinology* 2007, **148**(11):5295-5304.
- Fluiter K, Sattler W, De Beer MC, Connell PM, Westhuyzen van der DR, van Berkel TJ: **Scavenger receptor BI mediates the selective uptake of oxidized cholesterol esters by rat liver.** *J Biol Chem* 1999, **274**(13):8893-8899.
- Serougne C, Feurgard C, Hajri T, Champarnaud G, Ferezou J, Mathe D, Lutton C: **Catabolism of HDL1 cholesteryl ester in the rat. Effect of ethinyl estradiol treatment.** *C R Acad Sci III* 1999, **322**(7):591-596.
- Graf GA, Roswell KL, Smart EJ: **17beta-Estradiol promotes the up-regulation of SR-BII in HepG2 cells and in rat livers.** *J Lipid Res* 2001, **42**(9):1444-1449.
- Richard E, von Muhlen D, Barrett-Connor E, Alcaraz J, Davis R, McCarthy JJ: **Modification of the effects of estrogen therapy on HDL cholesterol levels by polymorphisms of the HDL-C receptor, SR-BI: the Rancho Bernardo Study.** *Atherosclerosis* 2005, **180**(2):255-262.
- Criqui MH, Barrett-Connor E, Austin M: **Differences between respondents and non-respondents in a population-based cardiovascular disease study.** *Am J Epidemiol* 1978, **108**(5):367-372.
- Friedewald WT, Levy RI, Fredrickson DS: **Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge.** *Clin Chem* 1972, **18**(6):499-502.
- Anderson DC, Hopper BR, Lasley BL, Yen SS: **A simple method for the assay of eight steroids in small volumes of plasma.** *Steroids* 1976, **28**(2):179-196.
- Barrett JC, Fry B, Maller J, Daly MJ: **Haploview: analysis and visualization of LD and haplotype maps.** *Bioinformatics* 2005, **21**(2):263-265.

33. Tatsumi K, Ohashi K, Taminishi S, Okano T, Yoshioka A, Shima M: **Reference gene selection for real-time RT-PCR in regenerating mouse livers.** *Biochem Biophys Res Commun* 2008, **374**(1):106-110.
34. Livak KJ: **Allelic discrimination using fluorogenic probes and the 5' nuclease assay.** *Genet Anal* 1999, **14**(5-6):143-149.
35. Wickham H: **R package ggplot2: An implementation of the Grammar of Graphics.** *version 0.8.1. edn* 2008.
36. Team RDC: **R: A language and environment for statistical computing.** Vienna, Austria: R Foundation for Statistical Computing 2008.
37. Shifren JL, Rifai N, Desindes S, McIlwain M, Doros G, Mazer NA: **A comparison of the short-term effects of oral conjugated equine estrogens versus transdermal estradiol on C-reactive protein, other serum markers of inflammation, and other hepatic proteins in naturally menopausal women.** *J Clin Endocrinol Metab* 2008, **93**(5):1702-1710.
38. Walsh BW, Schiff I, Rosner B, Greenberg L, Ravnikar V, Sacks FM: **Effects of postmenopausal estrogen replacement on the concentrations and metabolism of plasma lipoproteins.** *N Engl J Med* 1991, **325**(17):1196-1204.
39. Fotherby K: **Bioavailability of orally administered sex steroids used in oral contraception and hormone replacement therapy.** *Contraception* 1996, **54**(2):59-69.
40. Thierry-Mieg D, Thierry-Mieg J: **AceView: a comprehensive cDNA-supported gene and transcripts annotation.** *Genome Biol* 2006, **7**(Suppl 1):S12-11-14.
41. Zhang X, Merkler KA, McLean MP: **Characterization of regulatory intronic and exonic sequences involved in alternative splicing of scavenger receptor class B gene.** *Biochem Biophys Res Commun* 2008, **372**(1):173-178.
42. Carroll MD, Lacher DA, Sorlie PD, Cleeman JI, Gordon DJ, Wolz M, Grundy SM, Johnson CL: **Trends in serum lipids and lipoproteins of adults, 1960-2002.** *JAMA* 2005, **294**(14):1773-1781.
43. Grove J, Huby T, Stamataki Z, Vanwolleghem T, Meuleman P, Farquhar M, Schwarz A, Moreau M, Owen JS, Leroux-Roels G, et al: **Scavenger receptor BI and BII expression levels modulate hepatitis C virus infectivity.** *J Virol* 2007, **81**(7):3162-3169.
44. Codes L, Matos L, Parana R: **Chronic hepatitis C and fibrosis: evidences for possible estrogen benefits.** *Braz J Infect Dis* 2007, **11**(3):371-374.
45. Hankinson SE, Manson JE, Spiegelman D, Willett WC, Longcope C, Speizer FE: **Reproducibility of plasma hormone levels in postmenopausal women over a 2-3-year period.** *Cancer Epidemiol Biomarkers Prev* 1995, **4**(6):649-654.
46. Muti P, Trevisan M, Micheli A, Krogh V, Bolelli G, Sciajno R, Berrino F: **Reliability of serum hormones in premenopausal and postmenopausal women over a one-year period.** *Cancer Epidemiol Biomarkers Prev* 1996, **5**(11):917-922.
47. Mortola JF, Laughlin GA, Yen SS: **A circadian rhythm of serum follicle-stimulating hormone in women.** *J Clin Endocrinol Metab* 1992, **75**(3):861-864.

#### Pre-publication history

The pre-publication history for this paper can be accessed here:<http://www.biomedcentral.com/1471-2350/11/9/prepub>

doi:10.1186/1471-2350-11-9

**Cite this article as:** Chiba-Falek et al.: Impact of gene variants on sex-specific regulation of human Scavenger receptor class B type 1 (SR-BI) expression in liver and association with lipid levels in a population-based study. *BMC Medical Genetics* 2010 **11**:9.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

