

Review Article

Gene transcription and chromatin regulation in hypoxia

 Michael Batie and  Sonia Rocha

Department of Biochemistry, Institute of Integrative Biology, University of Liverpool, Biosciences Building, Crown Street, Liverpool L69 7ZB, U.K.

Correspondence: Michael Batie (m.batie@liverpool.ac.uk) and Sonia Rocha (s.rocha@liverpool.ac.uk)



Oxygen sensing is an essential feature of metazoan biology and reductions in oxygen availability (hypoxia) have both physiological and pathophysiological implications. Co-ordinated mechanisms have evolved for sensing and responding to hypoxia, which involve diverse biological outputs, with the main aim of restoring oxygen homeostasis. This includes a dynamic gene transcriptional response, the central drivers of which are the hypoxia-inducible factor (HIF) family of transcription factors. HIFs are regulated in an oxygen-dependent manner and while their role in hypoxia is well established, it is apparent that other key players are required for gene expression control in hypoxia. In this review, we highlight the current understanding of the known and potential molecular mechanisms underpinning gene transcriptional responses to hypoxia in mammals, with a focus on oxygen-dependent effects on chromatin structure.

Introduction

Molecular oxygen is a co-factor in various biochemical reactions and is essential for many aerobic organisms in their maintenance of intracellular ATP levels [1]. Responses to low oxygen (hypoxia) are highly conserved and vital to function and survival [2]. Dysregulated hypoxia response systems can lead to developmental issues as well as diseases [3], a notable example of which is renal clear cell carcinoma [4]. Hypoxia can occur in the context of a whole organism, for example during embryonic development where it is an important physiological cue, or when exposed to high altitude environments [3]. Hypoxia can also occur at the cellular level, when oxygen supply is reduced and/or metabolic demand is increased, for example in tumour hypoxia [5]. Importantly, normal physiological oxygen levels and the oxygen concentration at which hypoxia responses are activated are tissue and developmental stage specific.

The core aim of the hypoxia response is to restore oxygen homeostasis. To achieve this, a co-ordinated response involving changes in the regulation of gene expression, and protein stability and function are triggered, which impinges on many aspects of energy-related processes [6]. Given the importance of hypoxia to both normal and disease biology, understanding the mechanisms involved in oxygen sensing and adaptations is an attractive challenge to researchers.

A major breakthrough in our understanding of the hypoxia response came through the discovery of the von Hippel–Lindau tumour suppressor (VHL)-prolyl hydroxylase (PHD)-hypoxia inducible factor (HIF) pathway [2,7]. The HIF transcription factors are the primary mediators of hypoxia-induced gene transcriptional responses but do not act alone. There are additional factors that control HIF transcriptional activities and non-HIF dependent gene regulation in response to low oxygen, including regulators of the chromatin environment [8–11].

Hypoxia inducible factor (HIF)

The HIF transcription factor family consists of oxygen labile α subunits, of which there are three homologues in vertebrates, HIF-1 α , HIF-2 α and HIF-3 α , and a β subunit, HIF-1 β . HIFs are the master regulator of the transcriptional response to hypoxia and bind to hypoxia response elements (5-(A/G)CGTG-3), typically functioning as transactivators [12,13]. To date, there are over 100

Received: 17 March 2020
Revised: 2 April 2020
Accepted: 6 April 2020

Version of Record published:
5 May 2020

validated, direct HIF target genes, which are involved in biological processes (Figure 1). Canonical regulation of HIF- α subunits is via proline hydroxylation in their oxygen-dependent degradation domain by PHDs, which are 2-oxoglutarate (2-OG) dependent dioxygenases, requiring oxygen, iron and 2-OG for their catalytic activity [2]. Hydroxylated HIF- α provides a high affinity-binding site for the VHL E3 ubiquitin ligase complex. This leads to proteasomal degradation of HIF- α . PHDs can act as molecular oxygen sensors and, in hypoxia PHDs ability to hydroxylate HIF- α is impaired, enabling stable HIF- α subunits to translocate to the nucleus and dimerise typically with HIF-1 β . Another HIF hydroxylase, factor inhibiting HIF1 (FIH), hydroxylates an asparagine residue in the C-terminal transactivation domain (CTAD) of HIF-1 α and HIF-2 α , inhibiting interaction with the co-activator CREB-binding protein (CBP)/p300 [14,15]. Similar to PHDs, FIH is a 2-OG dependent dioxygenase, and inhibition of FIH activity in hypoxia alleviates blockage of CTAD co-activator interactions [16]. Although controversial, there is an expanding list of non-HIF putative hydroxylation targets of PHDs and FIH, providing additional mechanisms by which they can influence the hypoxia response [17–19]. There are additional post-translation modifications of HIF subunits that regulate their stability and function [11], and control of HIF subunit transcription is also pertinent to HIF activity and responses to low oxygen [20].

There is a high degree of tissue-specific HIF transcriptional responses and the numbers of estimated HIF regulated genes can vary massively depending on the study and cell types analysed. Whilst in some cases, technological differences in approaches used may account for this, evidence points towards altered isoform expression and activity, and chromatin accessibility and local chromatin environment, including RNA pol II availability around HREs, as being a major determinant of cell-type specificity [21–27]. Chromatin regulation is likely also important in controlling HIF independent gene transcriptional responses to hypoxia and efforts are ongoing to determine the role of chromatin in hypoxia.

Chromatin and hypoxia

Reduced oxygen can alter chromatin post-translational modifications, both directly via inhibition of oxygen-dependent enzymes and indirectly through additional mechanisms such as increased expression and/or HIF recruitment to chromatin [8,10,11,28–33].

Eukaryotic DNA is stored in the nucleus in the form of chromatin, consisting of DNA wrapped around histone octamers to form nucleosomes, which are linked together by linker DNA and histones and packaged into higher-order structures [34]. DNA replication, DNA repair and gene transcription are dependent upon the




Figure 1. HIF pathways.

Some of the biological processes regulated by hypoxia-inducible factor (HIF) target genes.

chromatin environment. Furthermore, chromatin can act as an accessibility barrier for gene transcription [35,36]. A closed chromatin conformation with nucleosomes tightly packaged can block the accessibility of transcriptional regulators and machinery to target genomic loci, silencing gene transcription. Alternatively, a more open conformation at gene regulatory regions is permissive to the binding of transcriptional regulators and machinery.

Mechanisms utilised by cells to modulate chromatin structure include, chromatin remodelling by ATP dependent chromatin remodellers [37], incorporation of histone variants [38], action of non-coding RNAs [39], DNA methylation [40] and histone post-translational modifications [41]. These mechanisms of chromatin modulation combine to dictate gene transcriptional output and cell fate decisions.

The impact of low-oxygen signalling on the chromatin environment, focusing on chromatin post-translational modification and chromatin organisation will be discussed in this review. Roles chromatin remodeller complexes, non-coding RNAs and incorporation of histone variants in hypoxia are also important factors but will not be detailed here.

Histone methylation

Histone methylation is a reversible modification to lysine residues of histone N-terminal tails. Some histone methylation modifications are associated with active gene transcription such as H3K4me3 and H3K36me3 whereas as others, such as H3K9me3 and H3K27me3 are associated with gene silencing [41]. The major family of histone demethylases, which remove methyl groups from histones lysine residues are the Jumonji-C (JmjC) proteins [42]. There are 32 JmjC proteins which have been shown to demethylate histones in humans and are classified into subgroups based on sequence homology and histone substrates (lysine specific demethylase, KDM2–8) [8]. JmjC proteins, like the HIF hydroxylases, PHDs and FIH, are 2-OG dependent dioxygenases. Several groups have reported hypoxia causes increases in total levels of various histone methylations; some have attributed this to inhibition of JmjC protein activity [29,30,33,43–49]. Recently, Batie et al. [29] and Chakraborty et al. [30] provide evidence that oxygen sensing by JmjC proteins in hypoxia results in increased levels of certain histone methylations, which co-ordinates the transcriptional response to hypoxia (Figure 2). It is clear from both *in vitro* and cell culture studies that different JmjC proteins have different oxygen sensitivities and this combined with their relative expression will contribute to their activity in oxygen-deprived environments. Many JmjC proteins are hypoxia induced, with some being HIF targets [8,50]. KDM4C and KDM3A can also function as co-activators of hypoxia-inducible gene expression in hypoxia via H3K9me3 demethylation, facilitating gene transcriptional activation [51,52] (Figure 2). Increased expression of JmjC proteins may in part act as a feedback mechanism to help retain activity, as is seen with PHD2 and PHD3.

SET domain bifurcated 1 (SETDB1) and euchromatic histone lysine methyltransferase 2 (EHMT2) are H3K9 targeting histone methyltransferases (HMTs) that have been found to mediate gene repression in hypoxia via H3K9 methylation (Figure 2), with implications hypoxia-induced p53-dependent apoptosis and tumour growth, respectively [53,54]. Increased H3K9me3 in hypoxia is also required for A-T mutated (ATM) activation via epigenetic silencing of ATM-specific phosphatases, which facilitates DNA replication in a low-oxygen environment [47].

Direct oxygen sensing may also affect DNA methylation. DNA methylation is associated with gene silencing and ten-eleven translocation (TET) enzymes, which are involved in DNA demethylation, are also 2-OG dependent dioxygenases. It has recently been shown that TET enzyme inhibition in hypoxia increases global levels of DNA methylation, thus oxygen sensing via TETs in hypoxia may also contribute to transcriptional regulation [31]. Hypoxia-induced DNA hypermethylation has also been attributed to HIF-1 α and transcriptional up-regulation of methionine adenosyltransferase 2A (MAT2A) [55].

Histone acetylation

Histone acetylation is associated with transcriptional activation [41]. It involves the transfer of an acetyl group from acetyl CoA to certain lysine residues on side chains within the positively charged N-terminal tail region of histones. This is a reversible histone modification, with acetyl group addition catalysed by histone acetyl transferases (HATs) and removed catalysed by histone deacetylases (HDACs). Acetylation neutralises the positive charge at histone tails and increases local chromatin accessibility. Whereas removal of acetyl groups is typically associated with increased chromatin compaction and localised transcriptional repression. Histone acetylation can also function as a binding site for chromatin-associated proteins containing a bromodomain, such as chromatin remodellers [37].




Figure 2. Histone methylation mediated regulation of gene transcription in hypoxia.

Hypoxia-induced changes in histone methylation, their effect on gene expression and the enzymes directing these changes. Lysine-specific demethylase (KDM), SET domain bifurcated 1 (SETDB1), euchromatic histone lysine methyltransferase 2 (EHMT2).

HATs and HDACs play a complex and dynamic role in the regulation of gene transcription in hypoxia via a multitude of mechanisms [9]. Focusing on direct regulation of gene expression through histone acetylation/deacetylation functions, lysine acetyltransferase 5 (KAT5) (also known as TIP60) and CBP/p300 are known HIF co-activators which interact with HIF, acetylate histones at HIF bound loci and are required for transcriptional activation at a subsets of HIF target genes [11,56]. Whereas HDAC1 interacts with HIF and reduces gene transcription at HIF targets via histone deacetylation [57]. SIN3 transcription regulator family member A (SIN3A) forms co-repressor complexes with HDACs and Tiana et al. [58] found that SIN3A binding is enriched both at hypoxia induced and repressed genes in human endothelial cells. SIN3A was required for down-regulation of

75% of the hypoxia-repressed genes and for the up-regulation of 47% of hypoxia-induced genes. In this study, SIN3A binding was unaffected by hypoxia, but H3K27Ac levels correlated with gene transcription, suggesting that HDAC activity at SIN3A complexes in hypoxia is a key factor in how SIN3As effects gene transcription. Detailed analysis of local chromatin accessibility in response to altered oxygen availability is lacking, as is the contribution of HATs and HDACs through histone acetylation/deacetylation. Johnson et al. [44] found that promoter nucleosome occupancy is altered in hypoxia at small group of hypoxia-responsive genes, with levels of nucleosome occupancy inversely correlating with hypoxia-induced gene expression. More recently, Suzuki et al. [59] identified hypoxia-inducible, nucleosome-free regions within the promoters of some hypoxia-inducible genes. These changes were HIF dependent and reversible upon reoxygenation. Interestingly, nucleosome reassembly through reoxygenation required SIN3A but not HDAC classes I and II activity.

Although limited, there are reports of hypoxia effecting total levels of acetylated histones. Decreased H3K9Ac [43,44] and H3K27Ac [60] and increased H3K14Ac [44] has been shown in cell culture models of hypoxia. Mechanistic insight into these changes and the extent to which they are driven by altered HAT/HDAC enzymatic activity or differential chromatin recruitment remains unclear.

3D chromatin organisation

Eukaryotic genomes inside the nucleus require a complex three-dimensional organisation. This organisation includes chromatin loops and topological associating chromatin domains, which modulate gene transcription [61]. Sequencing approaches employed by various groups have increased our understanding of HIF mediated transcriptional regulation on genome-wide scale, providing some insights into chromatin arrangement. Through the use of RNA-seq, ChIP-seq, DNase-seq, GRO-seq and Capture-C, researchers find that HIF functions mainly by binding to permissive HRE containing regulatory elements with pre-bound and paused RNAPol II [21–27,33,62,63]. HIF binding and coactivation subsequently triggers the release of paused RNA pol II at the majority of targets. It is now appreciated that HIFs bind to promoter distal, as well as, proximal regulatory elements, and that promoter enhancer interactions through chromatin looping is important in the HIF response. Work from Mole and Ratcliffe laboratories, using Capture-C and ChIP-seq, mapped chromatin–chromatin interactions at a subset of HIF target promoters [27]. This study found that HIF promoter distal binding occurs at pre-established and primed, promoter enhancer loops in VHL reconstituted 786-O cells and MCF7 cells. Using DNase-seq, Choudhry et al. [26] demonstrated that the average accessibility signal at proximal promoters of the top 100 hypoxia-inducible genes determined by RNAs-seq was unchanged in MCF-7 cells treated to 1% oxygen for 24 h [26].

Despite these insights, there is limited information regarding the effects of hypoxia on chromatin organisation in hypoxia, particularly in an unbiased manner. Using single-molecule localisation microscopy (SMLM) and *in situ* DNA digestion coupled with fluorescent microscopy, Kirmes et al. [64] observed a rapid change in chromatin architecture and elevated chromatin compaction in cardiomyocytes deprived of oxygen and nutrients. Furthermore, total transcription rates were reduced and chromatin and transcription changes were rapidly reversed upon restoration of oxygen and nutrients. Mechanistically changes in chromatin architecture were linked to reduced intracellular ATP and elevated polyamines. We can speculate that this chromatin conformational change in ischaemic conditions is strategic for the conservation of cellular energy. This is also occurring for the process of translation in severe hypoxia and ischaemia, with observed reduced translational rates in cells [65]. Reduced sensitivity to Mononuclease digestion, indicative of increased heterochromatin and increased global chromatin compaction, and increased levels of heterochromatin protein 1 binding protein 3 (HP1BP3), a heterochromatin structural protein, was determined in A431 cells exposed to severe hypoxia [66]. Therefore, HP1BP3 may be involved in hypoxia-induced chromatin compaction. Given that some chromatin-modifying proteins utilise oxygen and metabolites, chromatin may directly and dynamically sense changes in oxygen and metabolites during hypoxia, resulting in changes to chromatin organisation.

Summary

There are still many questions concerning how different cell and tissue responses to low-oxygen environments are determined. There is growing evidence for a dynamic role of chromatin in sensing and responding to hypoxia to facilitate transcriptional changes, both via and independent of HIF. This includes changes in histone and DNA modification and localised chromatin compaction. Further elucidating the complex, co-ordinated

mechanisms of chromatin regulation in low oxygen will be essential to better understanding hypoxia in normal physiological responses and disease.

Perspectives

- Sensing and responding to changes in oxygen availability is paramount for organismal and cellular survival. Changes in gene expression are at the forefront of cellular responses to hypoxia, with the transcription factor family hypoxia inducible factor (HIF) as a master regulator.
- Very recently, chromatin epigenetic marks have been shown to be essential for gene expression changes in hypoxia as well as cellular fate.
- Over the coming years, the use of advanced techniques such as chromatin conformation capture, assay for transposase-accessible chromatin using sequencing (ATAC-seq), microscopy-based chromosome conformation capture (Hi-M) and single molecule localisation microscopy (SMLM) should be employed to characterise changes in chromatin compaction and 3D arrangement in response to changes in oxygen availability.

Competing Interests

The authors declare that there are no competing interests associated with the manuscript.

Funding

This work was supported by Cancer Research U.K. [C99667/A12918] (S.R.), the Wellcome Trust [206293/Z/17/Z] (S.R.) the University of Liverpool (S.R. and M.B.).

Open Access

Open access for this article was enabled by the participation of the University of Liverpool in an all-inclusive *Read and Publish* pilot with Portland Press and the Biochemical Society under a transformative agreement with JISC.

Author Contributions

M.B. and S. R. researched the literature and wrote the manuscript.

Abbreviations

2-OG, 2-oxoglutarate; ATAC-seq, assay for transposase-accessible chromatin using sequencing; ATM, A-T mutated; CBP, CREB-binding protein; CTAD, C-terminal transactivation domain; EHMT2, euchromatic histone lysine methyltransferase 2; FIH, factor inhibiting HIF1; HATs, histone acetyl transferases; HDACs, histone deacetylases; HIF, hypoxia-inducible factor; Hi-M, microscopy-based chromosome conformation capture; HMTs, histone methyltransferases; HP1BP3, heterochromatin protein 1 binding protein 3; JmjC, Jumonji-C; KAT5, lysine acetyltransferase 5; KDM, lysine-specific demethylase; MAT2A, methionine adenosyltransferase 2A; PHD, prolyl hydroxylase; SETDB1, SET domain bifurcated 1; SIN3A, SIN3 transcription regulator family member A; SMLM, single-molecule localisation microscopy; TET, ten-eleven translocation; VHL, von Hippel–Lindau tumor suppressor.

References

- 1 Lee, P., Chandel, N.S. and Simon, M.C. (2020) Cellular adaptation to hypoxia through hypoxia inducible factors and beyond. *Nat. Rev. Mol. Cell Biol.* **21**, 268–283 <https://doi.org/10.1038/s41580-020-0227-y>
- 2 Kaelin, Jr., W.G. and Ratcliffe, P.J. (2008) Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol. Cell* **30**, 393–402 <https://doi.org/10.1016/j.molcel.2008.04.009>
- 3 Giaccia, A.J., Simon, M.C. and Johnson, R. (2004) The biology of hypoxia: the role of oxygen sensing in development, normal function, and disease. *Genes Dev.* **18**, 2183–2194 <https://doi.org/10.1101/gad.1243304>
- 4 Shen, C. and Kaelin, Jr., W.G. (2013) The VHL/HIF axis in clear cell renal carcinoma. *Semin. Cancer Biol.* **23**, 18–25 <https://doi.org/10.1016/j.semcancer.2012.06.001>

- 5 Xie, H. and Simon, M.C. (2017) Oxygen availability and metabolic reprogramming in cancer. *J. Biol. Chem.* **292**, 16825–16832 <https://doi.org/10.1074/jbc.R117.799973>
- 6 Rocha, S. (2007) Gene regulation under low oxygen: holding your breath for transcription. *Trends Biochem. Sci.* **32**, 389–397 <https://doi.org/10.1016/j.tibs.2007.06.005>
- 7 Semenza, G.L. (2001) HIF-1, O₂, and the 3 PHDs: how animal cells signal hypoxia to the nucleus. *Cell* **107**, 1–3 [https://doi.org/10.1016/S0092-8674\(01\)00518-9](https://doi.org/10.1016/S0092-8674(01)00518-9)
- 8 Shmakova, A., Batie, M., Druker, J. and Rocha, S. (2014) Chromatin and oxygen sensing in the context of JmJc histone demethylases. *Biochem. J.* **462**, 385–395 <https://doi.org/10.1042/BJ20140754>
- 9 Batie, M., Del Peso, L. and Rocha, S. (2018) Hypoxia and chromatin: a focus on transcriptional repression mechanisms. *Biomedicines* **6**, 47 <https://doi.org/10.3390/biomedicines6020047>
- 10 Hancock, R.L., Dunne, K., Walport, L.J., Flashman, E. and Kawamura, A. (2015) Epigenetic regulation by histone demethylases in hypoxia. *Epigenomics* **7**, 791–811 <https://doi.org/10.2217/epi.15.24>
- 11 Dengler, V.L., Galbraith, M. and Espinosa, J.M. (2014) Transcriptional regulation by hypoxia inducible factors. *Crit. Rev. Biochem. Mol. Biol.* **49**, 1–15 <https://doi.org/10.3109/10409238.2013.838205>
- 12 Semenza, G.L. (2014) Oxygen sensing, hypoxia-inducible factors, and disease pathophysiology. *Annu. Rev. Pathol.* **9**, 47–71 <https://doi.org/10.1146/annurev-pathol-012513-104720>
- 13 Ortiz-Barahona, A., Villar, D., Pescador, N., Amigo, J. and del Peso, L. (2010) Genome-wide identification of hypoxia-inducible factor binding sites and target genes by a probabilistic model integrating transcription-profiling data and in silico binding site prediction. *Nucleic Acids Res.* **38**, 2332–2345 <https://doi.org/10.1093/nar/gkp1205>
- 14 Mahon, P.C., Hirota, K. and Semenza, G.L. (2001) FIH-1: a novel protein that interacts with HIF-1 α and VHL to mediate repression of HIF-1 transcriptional activity. *Genes Dev.* **15**, 2675–2686 <https://doi.org/10.1101/gad.924501>
- 15 Lando, D., Peet, D.J., Gorman, J.J., Whelan, D.A., Whitelaw, M.L. and Bruck, R.K. (2002) FIH-1 is an asparaginyl hydroxylase enzyme that regulates the transcriptional activity of hypoxia-inducible factor. *Genes Dev.* **16**, 1466–1471 <https://doi.org/10.1101/gad.991402>
- 16 Schofield, C.J. and Ratcliffe, P.J. (2004) Oxygen sensing by HIF hydroxylases. *Nat. Rev. Mol. Cell Biol.* **5**, 343–354 <https://doi.org/10.1038/nrm1366>
- 17 Strowitzki, M.J., Cummins, E.P. and Taylor, C.T. (2019) Protein hydroxylation by hypoxia-inducible factor (HIF) hydroxylases: unique or ubiquitous? *Cells* **8**, 384 <https://doi.org/10.3390/cells8050384>
- 18 Bersten, D.C. and Peet, D.J. (2019) When is a target not a target? *eLife* **8**, e50585 <https://doi.org/10.7554/eLife.50585>
- 19 Lee, F.S. (2019) Substrates of PHD. *Cell Metab.* **30**, 626–627 <https://doi.org/10.1016/j.cmet.2019.08.008>
- 20 Kenneth, N.S. and Rocha, S. (2008) Regulation of gene expression by hypoxia. *Biochem. J.* **414**, 19–29 <https://doi.org/10.1042/BJ20081055>
- 21 Xia, X. and Kung, A.L. (2009) Preferential binding of HIF-1 to transcriptionally active loci determines cell-type specific response to hypoxia. *Genome Biol.* **10**, R113 <https://doi.org/10.1186/gb-2009-10-10-r113>
- 22 Schodel, J., Mole, D.R. and Ratcliffe, P.J. (2013) Pan-genomic binding of hypoxia-inducible transcription factors. *Biol. Chem.* **394**, 507–517 <https://doi.org/10.1515/hsz-2012-0351>
- 23 Mole, D.R., Blancher, C., Copley, R.R., Pollard, P.J., Gleadle, J.M., Ragoussis, J. et al. (2009) Genome-wide association of hypoxia-inducible factor (HIF)-1 α and HIF-2 α DNA binding with expression profiling of hypoxia-inducible transcripts. *J. Biol. Chem.* **284**, 16767–16775 <https://doi.org/10.1074/jbc.M901790200>
- 24 Schodel, J., Oikonomopoulos, S., Ragoussis, J., Pugh, C.W., Ratcliffe, P.J. and Mole, D.R. (2011) High-resolution genome-wide mapping of HIF-binding sites by ChIP-seq. *Blood* **117**, e207–e217 <https://doi.org/10.1182/blood-2010-10-314427>
- 25 Galbraith, M.D., Allen, M.A., Bensard, C.L., Wang, X., Schwinn, M.K., Qin, B., Long, H.W. et al. (2013) HIF1A employs CDK8-mediator to stimulate RNAPII elongation in response to hypoxia. *Cell* **153**, 1327–1339 <https://doi.org/10.1016/j.cell.2013.04.048>
- 26 Choudhry, H., Schodel, J., Oikonomopoulos, S., Camps, C., Grampp, S., Harris, A.L. et al. (2014) Extensive regulation of the non-coding transcriptome by hypoxia: role of HIF in releasing paused RNAPII. *EMBO Rep.* **15**, 70–76 <https://doi.org/10.1002/embr.201337642>
- 27 Platt, J.L., Salama, R., Smythies, J., Choudhry, H., Davies, J.O., Hughes, J.R. et al. (2016) Capture-C reveals preformed chromatin interactions between HIF-binding sites and distant promoters. *EMBO Rep.* **17**, 1410–1421 <https://doi.org/10.15252/embr.201642198>
- 28 Batie, M. and Rocha, S. (2019) JmJc histone demethylases act as chromatin oxygen sensors. *Mol. Cell. Oncol.* **6**, 1608501 <https://doi.org/10.1080/23723556.2019.1608501>
- 29 Batie, M., Frost, J., Frost, M., Wilson, J.W., Schofield, P. and Rocha, S. (2019) Hypoxia induces rapid changes to histone methylation and reprograms chromatin. *Science* **363**, 1222–1226 <https://doi.org/10.1126/science.aau5870>
- 30 Chakraborty, A.A., Laukka, T., Myllykoski, M., Ringel, A.E., Booker, M.A., Tolstorukov, M.Y. et al. (2019) Histone demethylase KDM6A directly senses oxygen to control chromatin and cell fate. *Science* **363**, 1217–1222 <https://doi.org/10.1126/science.aaw1026>
- 31 Thienpont, B., Steinbacher, J., Zhao, H., D'Anna, F., Kuchnio, A., Ploumaki, A. et al. (2016) Tumour hypoxia causes DNA hypermethylation by reducing TET activity. *Nature* **537**, 63–68 <https://doi.org/10.1038/nature19081>
- 32 Camuzi, D., de Amorim, I.S.S., Ribeiro Pinto, L.F., Oliveira Trivilin, L., Mencialha, A.L. and Soares Lima, S.C. (2019) Regulation is in the air: the relationship between hypoxia and epigenetics in cancer. *Cells* **8**, 300 <https://doi.org/10.3390/cells8040300>
- 33 Xia, X., Lemieux, M.E., Li, W., Carroll, J.S., Brown, M., Liu, X.S. et al. (2009) Integrative analysis of HIF binding and transactivation reveals its role in maintaining histone methylation homeostasis. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 4260–4265 <https://doi.org/10.1073/pnas.0810067106>
- 34 Luger, K., Dechassa, M.L. and Tremethick, D.J. (2012) New insights into nucleosome and chromatin structure: an ordered state or a disordered affair? *Nat. Rev. Mol. Cell Biol.* **13**, 436–447 <https://doi.org/10.1038/nrm3382>
- 35 Owen-Hughes, T. and Gkikopoulos, T. (2012) Making sense of transcribing chromatin. *Curr. Opin. Cell Biol.* **24**, 296–304 <https://doi.org/10.1016/j.ceb.2012.02.003>
- 36 Li, B., Carey, M. and Workman, J.L. (2007) The role of chromatin during transcription. *Cell* **128**, 707–719 <https://doi.org/10.1016/j.cell.2007.01.015>
- 37 Narlikar, G.J., Sundaramoorthy, R. and Owen-Hughes, T. (2013) Mechanisms and functions of ATP-dependent chromatin-remodeling enzymes. *Cell* **154**, 490–503 <https://doi.org/10.1016/j.cell.2013.07.011>
- 38 Weber, C.M. and Henikoff, S. (2014) Histone variants: dynamic punctuation in transcription. *Genes Dev.* **28**, 672–682 <https://doi.org/10.1101/gad.238873.114>

- 39 Mattick, J.S. and Makunin, I.V. (2006) Non-coding RNA. *Hum. Mol. Genet.* **15**, R17–R29 <https://doi.org/10.1093/hmg/ddi046>
- 40 Greenberg, M.V.C. and Bourc'his, D. (2019) The diverse roles of DNA methylation in mammalian development and disease. *Nat. Rev. Mol. Cell Biol.* **20**, 590–607 <https://doi.org/10.1038/s41580-019-0159-6>
- 41 Kouzarides, T. (2007) Chromatin modifications and their function. *Cell* **128**, 693–705 <https://doi.org/10.1016/j.cell.2007.02.005>
- 42 Kooistra, S.M. and Helin, K. (2012) Molecular mechanisms and potential functions of histone demethylases. *Nat. Rev. Mol. Cell Biol.* **13**, 297–311 <https://doi.org/10.1038/nrm3327>
- 43 Chen, H., Yan, Y., Davidson, T.L., Shinkai, Y. and Costa, M. (2006) Hypoxic stress induces dimethylated histone H3 lysine 9 through histone methyltransferase G9a in mammalian cells. *Cancer Res* **66**, 9009–9016 <https://doi.org/10.1158/0008-5472.CAN-06-0101>
- 44 Johnson, A.B., Denko, N. and Barton, M.C. (2008) Hypoxia induces a novel signature of chromatin modifications and global repression of transcription. *Mutat. Res.* **640**, 174–179 <https://doi.org/10.1016/j.mrfmmm.2008.01.001>
- 45 Zhou, X., Sun, H., Chen, H., Zavadil, J., Kluz, T., Arita, A. et al. (2010) Hypoxia induces trimethylated H3 lysine 4 by inhibition of JARID1A demethylase. *Cancer Res.* **70**, 4214–4221 <https://doi.org/10.1158/0008-5472.CAN-09-2942>
- 46 Tausendschon, M., Dehne, N. and Brune, B. (2011) Hypoxia causes epigenetic gene regulation in macrophages by attenuating Jumonji histone demethylase activity. *Cytokine* **53**, 256–262 <https://doi.org/10.1016/j.cyto.2010.11.002>
- 47 Olcina, M.M., Foskolou, I.P., Anbalagan, S., Senra, J.M., Pires, I.M., Jiang, Y. et al. (2013) Replication stress and chromatin context link ATM activation to a role in DNA replication. *Mol. Cell* **52**, 758–766 <https://doi.org/10.1016/j.molcel.2013.10.019>
- 48 Prickaerts, P., Adriaens, M.E., Beucken, T.V., Koch, E., Dubois, L., Dahlmans, V.E. et al. (2016) Hypoxia increases genome-wide bivalent epigenetic marking by specific gain of H3K27me3. *Epigenet. Chromatin* **9**, 46 <https://doi.org/10.1186/s13072-016-0086-0>
- 49 Dobrynin, G., McAllister, T.E., Leszczynska, K.B., Ramachandran, S., Krieg, A.J., Kawamura, A. et al. (2017) KDM4A regulates HIF-1 levels through H3K9me3. *Sci. Rep.* **7**, 11094 <https://doi.org/10.1038/s41598-017-11658-3>
- 50 Batie, M., Druker, J., D'Ignazio, L. and Rocha, S. (2017) KDM2 family members are regulated by HIF-1 in hypoxia. *Cells* **6**, 8 <https://doi.org/10.3390/cells6010008>
- 51 Luo, W., Chang, R., Zhong, J., Pandey, A. and Semenza, G.L. (2012) Histone demethylase JMJD2C is a coactivator for hypoxia-inducible factor 1 that is required for breast cancer progression. *Proc. Natl. Acad. Sci. U.S.A.* **109**, E3367–E3376 <https://doi.org/10.1073/pnas.1217394109>
- 52 Krieg, A.J., Rankin, E.B., Chan, D., Razorenova, O., Fernandez, S. and Giaccia, A.J. (2010) Regulation of the histone demethylase JMJD1A by hypoxia-inducible factor 1 α enhances hypoxic gene expression and tumor growth. *Mol. Cell. Biol.* **30**, 344–353 <https://doi.org/10.1128/MCB.00444-09>
- 53 Olcina, M.M., Leszczynska, K.B., Senra, J.M., Isa, N.F., Harada, H. and Hammond, E.M. (2016) H3K9me3 facilitates hypoxia-induced p53-dependent apoptosis through repression of APAK. *Oncogene* **35**, 793–799 <https://doi.org/10.1038/onc.2015.134>
- 54 Casciello, F., Al-Ejeh, F., Kelly, G., Brennan, D.J., Ngjoui, S.F., Young, A. et al. (2017) G9a drives hypoxia-mediated gene repression for breast cancer cell survival and tumorigenesis. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 7077–7082 <https://doi.org/10.1073/pnas.1618706114>
- 55 Liu, Q., Liu, L., Zhao, Y., Zhang, J., Wang, D., Chen, J. et al. (2011) Hypoxia induces genomic DNA demethylation through the activation of HIF-1 α and transcriptional upregulation of MAT2A in hepatoma cells. *Mol. Cancer Ther.* **10**, 1113–1123 <https://doi.org/10.1158/1535-7163.MCT-10-1010>
- 56 Perez-Perri, J.I., Dengler, V.L., Audetat, K.A., Pandey, A., Bonner, E.A., Urh, M. et al. (2016) The TIP60 complex is a conserved coactivator of HIF1A. *Cell Rep.* **16**, 37–47 <https://doi.org/10.1016/j.celrep.2016.05.082>
- 57 Lee, K.J., Lee, K.Y. and Lee, Y.M. (2010) Downregulation of a tumor suppressor RECK by hypoxia through recruitment of HDAC1 and HIF-1 α to reverse HRE site in the promoter. *Biochim. Biophys. Acta* **1803**, 608–616 <https://doi.org/10.1016/j.bbamcr.2010.01.004>
- 58 Tiana, M., Acosta-Iborra, B., Puente-Santamaria, L., Hernansanz-Agustin, P., Worsley-Hunt, R., Masson, N. et al. (2018) The SIN3A histone deacetylase complex is required for a complete transcriptional response to hypoxia. *Nucleic Acids Res.* **46**, 120–133 <https://doi.org/10.1093/nar/gkx951>
- 59 Suzuki, N., Vojnovic, N., Lee, K.L., Yang, H., Gradin, K. and Poellinger, L. (2018) HIF-dependent and reversible nucleosome disassembly in hypoxia-inducible gene promoters. *Exp. Cell Res.* **366**, 181–191 <https://doi.org/10.1016/j.yexcr.2018.03.020>
- 60 Gao, X., Lin, S.H., Ren, F., Li, J.T., Chen, J.J., Yao, C.B. et al. (2016) Acetate functions as an epigenetic metabolite to promote lipid synthesis under hypoxia. *Nat. Commun.* **7**, 11960 <https://doi.org/10.1038/ncomms11960>
- 61 Zheng, H. and Xie, W. (2019) The role of 3D genome organization in development and cell differentiation. *Nat. Rev. Mol. Cell Biol.* **20**, 535–550 <https://doi.org/10.1038/s41580-019-0132-4>
- 62 Schodel, J., Bardella, C., Sciesielski, L.K., Brown, J.M., Pugh, C.W., Buckle, V. et al. (2012) Common genetic variants at the 11q13.3 renal cancer susceptibility locus influence binding of HIF to an enhancer of cyclin D1 expression. *Nat. Genet.* **44**, 420–425. S1–2 <https://doi.org/10.1038/ng.2204>
- 63 Salama, R., Masson, N., Simpson, P., Sciesielski, L.K., Sun, M., Tian, Y.M. et al. (2015) Heterogeneous effects of direct hypoxia pathway activation in kidney cancer. *PLoS One* **10**, e0134645 <https://doi.org/10.1371/journal.pone.0134645>
- 64 Kirmes, I., Szczurek, A., Prakash, K., Charapitsa, I., Heiser, C., Musheev, M. et al. (2015) A transient ischemic environment induces reversible compaction of chromatin. *Genome Biol.* **16**, 246 <https://doi.org/10.1186/s13059-015-0802-2>
- 65 Chee, N.T., Lohse, I. and Brothers, S.P. (2019) mRNA-to-protein translation in hypoxia. *Mol. Cancer* **18**, 49 <https://doi.org/10.1186/s12943-019-0968-4>
- 66 Dutta, B., Yan, R., Lim, S.K., Tam, J.P. and Sze, S.K. (2014) Quantitative profiling of chromatome dynamics reveals a novel role for HP1BP3 in hypoxia-induced oncogenesis. *Mol. Cell. Proteom.* **13**, 3236–3249 <https://doi.org/10.1074/mcp.M114.038232>