

Can Aptameric Ligands Specific to Plasma Coagulation Factor VII Bind the Recombinant Form with High Affinity: Affinity Measurement by Fluorescence Method

Maryam Tabarzad ^{1*}, Marzieh Jafari ², and Nastaran Nafissi-varcheh ³

1. Protein Technology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Pharmacology and Toxicology, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3. Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Abstract

Background: Among diverse protein purification systems, affinity chromatography is the most attractive one in the purification process of coagulation factors. Coagulation factor VII is a plasma serine protease that has a significant role in natural human hemostasis and its recombinant form such as AryoSevenTM, has been applied in clinical treatment of bleeding disorders. Immunoaffinity chromatography is the purification method of choice that is currently applied in the development of coagulation factor VIIa products. Aptamers as nucleic acid based affinity ligands are more promising than monoclonal antibodies. In addition, DNA aptamers are more acceptable than RNA ones in this regard.

Methods: In this study, two of the aptameric DNA oligonucleotides that showed acceptable affinities for purification of coagulation factor VIIa from plasma, were selected to evaluate their affinity against Aryoseven. A serial dilution of fluorescence labeled aptamers was incubated against the concentration of 1 nM from Aryoseven. Then, a fluorescence index was calculated according to the fluorescence intensity data measured from test and control samples. The dissociation constant of aptamers was calculated according to the fluorescence index using Prism5 software.

Results: Results showed that the binding affinity of the 44 nucleotide aptamer was more than 81 nucleotide aptamer sequence. As a result, this aptamer could be optimized in order to develop aptamer based affinity chromatography process for this form of recombinant coagulation factor VIIa.

Discussion: Aptamers with shorter length of sequence could show higher affinity in target binding, as they could adapt more easily to suitable conformation according to target interaction. However, it should be considered that the selectivity of affinity ligands is also important for target purification and analytical applications.

* Corresponding author:
Maryam Tabarzad, Ph.D.,
Protein Technology Research
Center, Shahid Beheshti
University of Medical Sciences,
Tehran, Iran
Tel: +98 21 88648124
Fax: +98 21 88647456
E-mail:
m_tabarzad@sbmu.ac.ir
Received: 8 May 2016
Accepted: 22 Jun 2016

Avicenna J Med Biotech 2017; 9(2): 109-112

Keywords: Affinity, Aptamer, Factor VIIa, Fluorescence

Introduction

Aptamers are small ligands with high affinity and selectivity to their targets, which have been developed by *in vitro* selection process against a wide range of target molecules ¹⁻³. Aptamers are appropriate ligands for affinity separation and purification of biomolecules using affinity chromatography ⁴. Aptamers as oligosorbents have been applied in the purification of various targets ⁵⁻⁹.

Coagulation factors are the first interesting protein targets in aptamer development ¹⁰. One of the valuable members, involved in extrinsic pathway, is coagulation factor VII and related biopharmaceuticals have been administered as a replacement therapy in hemophilia

and non-hemophilia bleeding disorders ¹¹⁻¹³. Coagulation factor VII specific RNA aptamers have been selected by independent scientists to inhibit coagulation process ^{14,15}.

DNA aptamers are more appropriate for analytical applications such as protein detection and purification, because of more stability than RNA aptamers. There is one patent report of DNA aptamers specific to coagulation factor VIIa developed for affinity purification of this protein from plasma samples ¹⁶.

Aryoseven[®] is a recombinant form of coagulation factor VIIa and a biosimilar of Novoseven[®], which is manufactured in Iran. Development of an efficient af-

finity chromatography technique for protein purification would result in a decrease in time and cost of product manufacturing¹⁷. Aptamers as affinity ligands are valuable choices for affinity chromatography^{4,8,18}.

In this study, binding affinities of two aptameric sequences against Aryoseven[®] were determined by a fluorescent method to evaluate their potentials for the development of protein purification process.

Materials and Methods

Chemicals and oligonucleotide sequences

Two DNA sequence of 44 and 81 nucleotides were supplied by chemical synthesis. The FAM labeled sequences of Apt81: 5'GGGAGATAGCCACGACC TATGCAGCCAGCCGCAGTGTAAAGTGAATGCAG ACATGGTCTAAGTGTCCAGGCTGTGCGAAAGC 3' and Apt 44:5'CCG CACACCACGCGCATAATC CCGCGCACACGACTTGAAGTAGC3' were purchased from SinaClon (Iran). Recombinant coagulation factor VIIa (Aryose-ven[®]) was gifted from Aryogene company (Iran). All chemicals of molecular biology grade were supplied from Sigam-Aldrich (USA).

Binding affinity measurement

The oligonucleotides were dissolved in binding buffer (Tris 40 mM, NaCl 117 mM, CaCl₂ 5 mM, MgCl₂ 5 mM and pH=7.4) to the final concentration of 100 μM. The solutions were thermally treated before analysis (5 min at 95 °C, 10 min on ice, 10 min in room temperature). At each run, a serial dilution of aptamers was prepared from 100 pM to 10 μM in two set and duplicate tests. The protein solution was added to the final concentration of 1 nM. Controls were aptamer free and protein free samples. After incubation at 4 °C for 1 hr, fluorescence intensities of the samples were measured at 492 nm excitation and 521 nm emission wavelength using Cytation5 Multi-Mode Reader (BioTek, USA). Fluorescent index was calculated according to the equation 1.

A aptamer, A protein and A test are the absorption of aptamer control, protein control and test samples, respectively, at the 521 nm that is the emission wavelength of fluorescence label.

$$\text{Fluorescence Index} = \frac{[(A \text{ aptamer} + A \text{ protein}) - A \text{ test}]}{A \text{ aptamer}}$$

Statistics and dissociation constant determination

Mean±SE of duplicated tests were calculated by IBM-SPSS21. The results from three independent runs during different days were analyzed. The non-linear regression was performed using Prism5 to calculate the Kd according to the binding-saturation model and one-site total sub-deviation.

Results

Binding affinity characterization

Fluorescence methods for aptamer-target binding affinity determination are simpler and faster than the




Figure 1. Predicted secondary structures of aptamers; A: Apt44 and B: Apt81 predicted structure by fold method. The boxes include the non-structured part

other ones⁵. In this study, changes in fluorescence intensity were measured as the consequence of aptamer binding to the target protein. Raw data from fluorescence measurement showed significant variation between duplicated runs. Therefore, mean of measurement data for each sample was calculated and then applied to a formula (Equation 1) in order to derive validated data. Similar method was evaluated in aptamer related study¹⁹. Kd±SE of Apt81 and Apt44 were calculated as 938±149 and 167.47±18 nM, respectively.

Accordingly, lower value of the calculated Kd related to the smaller aptamer and it was revealed that adaptive changes in folding structures of aptamers could happen more easily than long sequences such as truncated aptameric sequences, as widely confirmed before^{7,20-22}. Some other studies showed that truncated aptameric sequences would have similar Kd as a full length aptamer^{8,23}. It should also be considered that small length aptamers are more useful in synthesis and application⁶.

Prediction of aptamer secondary structure

Binding affinity of aptamers is related to their 3D structure. Accordingly, secondary structure of the 44nts and 81nts aptamers was predicted by RNAStructure[®] online server, setting the parameters for DNA entry. Predicted structures using fold method (structure with minimum free energy) were presented in figure 1. As presented, a larger part of the Apt44 nucleotide sequence was predicted as the non-structured part that could be adapted to a proper binding conformation during protein interaction. Consequently, this short aptameric ligand had more flexible structures for binding.

Discussion

The binding affinity between nucleic acid aptamers and the target molecules could be measured by different methods. Fluorescence methods including measurement the fluorescent intensity using labeled aptamers are more feasible one. Fluorescent intensity of the labeled aptamer might change after binding to the target protein and measurement of these changes help to

determine the binding affinity and kinetic. In this study, changes in fluorescence intensity were measured as the consequence of aptamer binding to the target protein, recombinant form of coagulation factor VIIa. The smaller calculated K_d of smaller aptamer revealed that adaptive changes in folding structures of aptamers happen in short sequences more easily than long sequences. Presence of non-structured positions that could be adapted to a proper binding conformation during protein binding led to flexible structures for binding, but, it would be undesirable in the development of an aptamer based affinity chromatography process if this flexibility resulted in non-specific binding to unrelated target structures. Therefore, cross binding studies should be performed to indicate that the aptamer has significant binding affinity to the background impurities or not.

Conclusion

The study showed that plasma coagulation factor VIIa specific aptamers could bind to the recombinant form of protein with a considerable affinity, because of the structural similarities between same proteins of different sources. In addition, short length aptamers have more flexibility and higher affinity in target binding. However, during the process of designing the purification or analysis based on aptamers, it should be considered that more adaptive structure of small length aptamers could result in less selectivity of binding.

Acknowledgement

The author reports no conflicts of interest. The authors acknowledge the financial supports granted by Protein Technology Research Center of Shahid Beheshti University of Medical Sciences (Iran) and gratefully acknowledge the Aryogen Company for supplying the Aryoseven®.

References

- Wandtke T, Woźniak J, Kopiński P. Aptamers in diagnostics and treatment of viral infections. *Viruses* 2015; 7(2):751-780.
- Li F, Zhang H, Wang Z, Newbigging AM, Reid MS, Li XF, et al. Aptamers facilitating amplified detection of biomolecules. *Anal Chem* 2015;87(1):274-292.
- Ku TH, Zhang T, Luo H, Yen TM, Chen PW, Han Y, et al. Nucleic acid aptamers: an emerging tool for biotechnology and biomedical sensing. *Sensors (Basel)* 2015;15(7):16281-16313.
- Zhao Q, Wu M, Chris Le X, Li XF. Applications of aptamer affinity chromatography. *Trends Analyt Chem* 2012; 41:46-57.
- Jing M, Bowser MT. Methods for measuring aptamer-protein equilibria: a review. *Anal Chim Acta* 2011;686(1-2):9-18.
- Wang G, Zhu Y, Chen L, Zhang X. Photoinduced electron transfer (PET) based label-free aptasensor for platelet-derived growth factor-BB and its logic gate application. *Biosens Bioelectron* 2015;63:552-557.
- Deng Q, German I, Buchanan D, Kennedy RT. Retention and separation of adenosine and analogues by affinity chromatography with an aptamer stationary phase. *Anal Chem* 2001;73(22):5415-5421.
- Lim HK, Kim IH, Nam HY, Shin S, Hah SS. Aptamer-based alternatives to the conventional immobilized metal affinity chromatography for purification of His-tagged proteins. *Anal Lett* 2013;46(3):407-415.
- Brothier F, Pichon V. Miniaturized DNA aptamer-based monolithic sorbent for selective extraction of a target analyte coupled on-line to nanoLC. *Anal Bioanal Chem* 2014;406(30):7875-7886.
- Bock LC, Griffin LC, Latham JA, Vermaas EH, Toole JJ. Selection of single-stranded DNA molecules that bind and inhibit human thrombin. *Nature* 1992;355(6360): 564-566.
- Tiede A, Amano K, Ma A, Arkhammar P, El Fegoun SB, Rosholm A, et al. The use of recombinant activated factor VII in patients with acquired haemophilia. *Blood Rev* 2015;29 Suppl 1:S19-25.
- Lin Y, Stanworth S, Birchall J, Doree C, Hyde C. Recombinant factor VIIa for the prevention and treatment of bleeding in patients without haemophilia. *Cochrane Database Syst Rev* 2011;(2):CD005011.
- Neufeld EJ, Négrier C, Arkhammar P, Benchikh el Fegoun S, Simonsen MD, Rosholm A, et al. Safety update on the use of recombinant activated factor VII in approved indications. *Blood Rev* 2015;29:S34-S41.
- Rusconi CP, Yeh A, Lysterly HK, Lawson JH, Sullenger BA. Blocking the initiation of coagulation by RNA aptamers to factor VIIa. *Thromb Haemost* 2000;84(5):841-848.
- Layzer JM, Sullenger BA. Simultaneous generation of aptamers to multiple gamma-carboxyglutamic acid proteins from a focused aptamer library using DeSELEX and convergent selection. *Oligonucleotides* 2007;17(1):1-11.
- Perret G, Nogre M. Means for purifying a protein of blood plasma and methods for implementing same. U.S. Patent Application No. 13/201,690;2012.
- Hage DS, Anguizola JA, Bi C, Li R, Matsuda R, Papanastavros E, et al. Pharmaceutical and biomedical applications of affinity chromatography: recent trends and developments. *J Pharm Biomed Anal* 2012;69:93-105.
- Romig TS, Bell C, Drolet DW. Aptamer affinity chromatography: combinatorial chemistry applied to protein purification. *J Chromatogr B Biomed Sci Appl* 1999;731(2):275-284.
- Ramezani M, Danesh NM, Lavaee P, Abnous Kh, Taghdisi SM. A selective and sensitive fluorescent aptasensor for detection of kanamycin based on catalytic recycling activity of exonuclease III and gold nanoparticles. *Sens Actuators B Chem*. 2016;222:1-7.
- Dassie JP, Liu XY, Thomas GS, Whitaker RM, Thiel KW, Stockdale KR, et al. Systemic administration of optimized aptamer-siRNA chimeras promotes regression

Fluorescence Affinity Measurement of FVIIa Aptamers

- of PSMA-expressing tumors. *Nat Biotechnol* 2009;27(9):839-849.
21. Stoltenburg R, Schubert T, Strehlitz B. In vitro selection and interaction studies of a DNA aptamer targeting protein A. *PLoS One* 2015;10(7):e0134403.
 22. Biesecker G, Dihel L, Enney K, Bendele R. Derivation of RNA aptamer inhibitors of human complement C5. *Immunopharmacology* 1999;42(1-3):219-230.
 23. Kiani Z, Shafiei M, Ebrahimi A. Selection of DNA aptamers for digoxin and assaying of their binding profile and inhibitory effects. *Res Pharm Sci* 2012;7(5):S490.