

CORRECTION

Correction: Serum *Wisteria floribunda* agglutinin-positive Mac-2-binding protein levels predict the presence of fibrotic nonalcoholic steatohepatitis (NASH) and NASH cirrhosis

Naim Alkhouri, Casey Johnson, Leon Adams, Sachiko Kitajima, Chikayuki Tsuruno, Tracey L. Colpitts, Kazuki Hatcho, Eric Lawitz, Rocio Lopez, Ariel E. Feldstein

The seventh author's name is spelled incorrectly. The correct name is: Ariel E. Feldstein

Reference

1. Alkhouri N, Johnson C, Adams L, Kitajima S, Tsuruno C, Colpitts TL, et al. (2018) Serum *Wisteria floribunda* agglutinin-positive Mac-2-binding protein levels predict the presence of fibrotic nonalcoholic steatohepatitis (NASH) and NASH cirrhosis. PLoS ONE 13(8): e0202226. <https://doi.org/10.1371/journal.pone.0202226> PMID: 30161179



OPEN ACCESS

Citation: Alkhouri N, Johnson C, Adams L, Kitajima S, Tsuruno C, Colpitts TL, et al. (2018) Correction: Serum *Wisteria floribunda* agglutinin-positive Mac-2-binding protein levels predict the presence of fibrotic nonalcoholic steatohepatitis (NASH) and NASH cirrhosis. PLoS ONE 13(10): e0205541. <https://doi.org/10.1371/journal.pone.0205541>

Published: October 4, 2018

Copyright: © 2018 Alkhouri et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.