

RESEARCH ARTICLE

High-Frequency Patterns in the Abundance of Benthic Species near a Cold-Seep – An Internet Operated Vehicle Application

Damianos Chatzievangelou^{1*}, Carolina Doya², Laurenz Thomsen¹, Autun Purser³, Jacopo Aguzzi²

1 Jacobs University Bremen, Bremen, Germany, **2** Instituto de Ciencias del Mar (ICM-CSIC), Barcelona, Spain, **3** Alfred-Wegener-Institut (AWI), Bremerhaven, Germany

* d.hatzievangelou@jacobs-university.de



OPEN ACCESS

Citation: Chatzievangelou D, Doya C, Thomsen L, Purser A, Aguzzi J (2016) High-Frequency Patterns in the Abundance of Benthic Species near a Cold-Seep – An Internet Operated Vehicle Application. PLoS ONE 11(10): e0163808. doi:10.1371/journal.pone.0163808

Editor: Cristiano Bertolucci, University of Ferrara, ITALY

Received: April 8, 2016

Accepted: September 14, 2016

Published: October 12, 2016

Copyright: © 2016 Chatzievangelou et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant numerical data are provided within the paper and its Supporting Information files. All video footage is available online at the Ocean Networks Canada database by selecting the adequate temporal window at the following URL: <http://dmas.uvic.ca/DataSearch?location=BACWL.V1&deviceCategory=VIDEOCAM>.

Funding: This work was supported by Robotic Exploration of Extreme Environments (ROBEX) project by Helmholtz Alliance (HA-304), <http://www.helmholtz.de/> (DC and AP); RITFIM project

Abstract

Three benthic megafaunal species (i.e. sablefish *Anoplopoma fimbria*; pacific hagfish *Eptatretus stoutii* and a group of juvenile crabs) were tested for diel behavioral patterns at the methane hydrates site of Barkley Canyon (890 m depth), off Vancouver Island (BC, Canada). Fluctuations of animal counts in linear video-transects conducted with the Internet Operated Deep-Sea Crawler “Wally” in June, July and December of 2013, were used as proxy of population activity rhythms. Count time series and environmental parameters were analyzed under the hypothesis that the environmental conditioning of activity rhythms depends on the life habits of particular species (i.e. movement type and trophic level). Non-linear least squares modeling of biological time series revealed significant diel periods for sablefish in summer and for hagfish and crabs in December. Combined cross-correlation and redundancy (RDA) analyses showed strong relationships among environmental fluctuations and detected megafauna. In particular, sablefish presence during summer months was related to flow magnitude, while the activity of pacific hagfish and juvenile crabs in December correlated with change in chemical parameters (i.e. chlorophyll and oxygen concentrations, respectively). Waveform analyses of animal counts and environmental variables confirmed the phase delay during the 24 h cycle. The timing of detection of sablefish occurred under low flow velocities, a possible behavioral adaptation to the general hypoxic conditions. The proposed effect of chlorophyll concentrations on hagfish counts highlights the potential role of phytodetritus as an alternative food source for this opportunistic feeder. The juvenile crabs seemed to display a cryptic behavior, possibly to avoid predation, though this was suppressed when oxygen levels were at a minimum. Our results highlight the potential advantages such mobile observation platforms offer in multiparametric deep-sea monitoring in terms of both spatial and temporal resolution and add to the vastly understudied field of diel rhythms of deep-sea megafauna.

(ref. CTM2010-16274) of the Spanish national RTD program (CD). Development and deployment of the crawler was co-funded by Ocean Networks Canada and Neptune Canada, <http://www.oceannetworks.ca/>. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Ocean Networks Canada and Neptune Canada (www.oceannetworks.ca/) partially funded the development and deployment of the crawler. The authors declare that this does not alter our adherence to PLOS ONE policies on sharing data and materials.

1. Introduction

The study of gas hydrates systems has gained progressive attention in both the geochemical and biological research fields, with the systems clearly showing the lack of homogeneity in the deep sea [1]. The potential exploitation of the hydrocarbon sources available at these sites generated further interest from an economic point of view, in relation to the quantity and quality of utilizable material, as well as their potential impact on the structural stability of the seabed [2–3]. Cold-seeps are considered biodiversity hotspots capable of sustaining a greater biomass than comparable areas of adjacent continental slopes, due to their high spatio-temporal habitat heterogeneity [4–6]. These systems may provide the principal energy pathways through bacterial chemosynthetic production [7–9]. Gas seepage, their characterizing element, can be affected by periodic bottom currents at scales of few hours (i.e. of semi-diurnal nature; [3]). Within this context, the importance of high-frequency, long-term and multidisciplinary ecological monitoring of such environments is evident.

Nowadays, the rapid technological advances in seabed exploration and monitoring techniques allow scientists have access to real-time biological and environmental data from various marine sites within the world ocean [10–13]. As described in [3, 14], an Internet Operated Deep-Sea Crawler (Wally) has been deployed at the gas hydrates site in Barkley Canyon (NE Pacific, Canada) and in operation since September of 2010, as a part of the NEPTUNE Canada Cabled Observatory network (www.oceannetworks.ca/). The uniqueness of this monitoring platform lies in the combination of mobility with the high-frequency acquisition of multiparametric data throughout long-term deployment, whilst remaining in communication with land-based researchers 24/7 via the Internet. Until now all these features could be found distributed among the previously existing mobile and fixed benthic technologies (i.e. ROVs, AUVs, and landers *versus* cabled observatory nodes), rather than on one mobile platform.

Remote, high-frequency and long-lasting image monitoring of the continental margin and abyssal communities is nowadays increasingly required in order to link measurable biodiversity to species turnover as a product of activity rhythms [15–17]. Animal activity patterns may come as a response to a persistent temporal structure within an environment (i.e. geophysical cycles in light intensity and internal tides), resulting in 4 behaviorally different phenotypes, as identified in laboratory controlled conditions: diurnal, nocturnal, crepuscular and arrhythmic animal behaviors [18–21]. Species use time as an ecological resource while they are competing with others of a partially overlapping habitat niche, forming the dynamics of predator-prey relationships and finally in behavioral adaptations that come as a response to their own physiological limitations [22–24]. In benthic communities, the “zeitgeber” (the environmental cue forcing animals to keep their internal clock synchronized) can be either sunlight or periodic hydrodynamism (e.g. current velocity and direction, internal tides) [25–27]. Intraspecific plasticity of activity patterns is often observed [28], as local variability of the aforementioned variables may lead to distinct populations facing different conditions. On continental margins, animal response to such triggers is commonly expressed as diel displacement, either vertically, horizontally or in and out of the substrate (i.e. pelagic migrations, nektobenthic or endobenthic movements) [15]. Apart from regulating the dynamics of benthic communities, vertical migrations (benthopelagic coupling) can affect the redistribution of nutrients and organic matter in the ocean [29–30]. Additionally, periodical change in local abundances of benthic species can affect their detection in a certain space and time, leading to potential bias in traditional scientific sampling by trawling and affecting commercial fishing activities [31–32].

In this study, crawler imaging and environmental data were used to identify any diel patterns within the visual counts of the most abundant megabenthos at the Barkley Canyon gas hydrates site, and whether they exhibited a relationship with concomitant environmental

forcing. We hypothesized that the activity of megafauna with different ecological traits in relation to motility and trophic level, is modulated by different environmental variables.

2. Material and Methods

2.1 Study Site and Data Acquisition

The study was conducted at the gas hydrates site of NEPTUNE Cabled Observatory network (www.oceannetworks.ca), located on a small (1 Km²) plateau in Barkley Canyon (Fig 1; 48° 18' 46" N, 126° 03' 57" W), at approx. 890 m depth. Authorization for conducting research was provided by the Transport Canada (www.tc.gc.ca/), after Fisheries and Oceans Canada (<http://www.dfo-mpo.gc.ca/>) assessed that the cable installation would not have a negative impact on fish habitat.

Oceanographic conditions in Barkley Canyon at these depths are known to differ between seasons, with distinct water flow, temperature and chlorophyll flux patterns previously reported for summer and winter months [3, 14, 33]. Following [33], the annual cycle was divided into two main seasons (summer and winter). The months analyzed in this current study are June, July and December of 2013, since they were deemed as a sampling period representative of the oceanographic conditions and community structure of the location and also provided material that matched the study's criteria (see Section 2.2).




Fig 1. Map of the study site. The position of the gas hydrates site in Barkley Canyon, off Vancouver Island, and the path followed by Wally when the linear transects were performed (i.e. between waypoints 16–10). Canyon map modified from [34]. Transect representation modified from [14].

doi:10.1371/journal.pone.0163808.g001

Linear, constant back and forth imaging census transects of 20 m length (i.e. between waypoints 16 and 10; Fig 1) were carried out with Wally every 4 hours during 5 days, in the first week of each calendar month. All transects were video-recorded with a Panasonic web camera (BB-HCM580; 720 x 480 pixel) mounted in a glass globe designed to resist high pressure levels up to 6000 m depth [14]. Driving and recording were stopped by the researcher operating the crawler every time a sediment resuspension event was impeding visibility (i.e. < 70% of the Region of Interest visible). All video footage is archived and can be viewed online in the Ocean Networks Canada database (dmas.uvic.ca/).

Water flow and temperature were measured with an ADCP 2 MHz (Nortek Aquadopp Profiler AQD 9917) fixed on the crawler. Oxygen concentration measurements were taken using a Sea-Bird SBE 63 Dissolved Oxygen Sensor (Device Code 630109) deployed at the hydrates site. Density data were provided by the CTD Sea-Bird SeaCAT SBE16plus V2 7027 deployed at the POD4 site (mid-canyon east, at 896 m depth). Finally, the sensor used to measure chlorophyll levels was an ECO-FL-NTU (RT) 2973 (WET Laboratories), deployed also at POD4.

2.2 Data processing

Visual count variations can be considered as a direct proxy of populational behavioral rhythms, with animal rate of displacement controlling their chance to be spotted in the camera's field of view during each sampling window [26–27, 35].

Stationary observations of the seafloor and video sequences with low visibility were discarded and the remaining video duration was further used. Pictured animals were classified to the lowest possible taxonomic level following [36] and then counted. Since there was no information available on the exact surface covered in each transect, animal counts data were summed for each pair of transects (i.e. back and forth between waypoints) and then normalized to number of individuals per 10 min. This normalization was based on the assumption that the differences in surface covered during the video transects were minimal and the underlying error was constant, leading to no effect on the final outcome. As a result, the biological time series consisted of a total of 92 observations taken at a 4 h frequency, each of these being the normalized sum of 2 transects. In total, over 21 h of footage were considered to be of acceptable quality for the faunal analysis.

For the statistical analyses, species time series containing 0 (zero) counts in more than 1/3 of the observations were discarded from further analyses to prevent misleading results, as it would be extremely hard to detect periodicities with statistical significance and low bias in short, zero-inflated time series [15]. As a result, the three most abundant and regularly spotted species that were subsequently studied were the sablefish *Anoplopoma fimbria* (can also be found by the name black cod) in June and July, the pacific hagfish *Eptatretus stoutii* in December and a group of juvenile crabs in the same month, which the authors believe are grooved tanner crabs, *Chionoecetes tanneri* (Fig 2). The small size of specimens and the regularly turbid conditions at the site did not permit a 100% precise identification, so hereafter they will be referred to as a juvenile functional group and treated as such. A complete species list, along with a table containing the abundances of all the observed species is provided in [37].

Environmental data were downloaded from the Ocean Networks Canada database (dmas.uvic.ca/) and averaged into 1 h bins for high resolution and flexibility in data treatment. This step was performed with MATLAB R2014b [38]. Current velocity data that were initially provided in the form of 2 components-vectors (North-South and East-West) were transformed to flow magnitude and flow direction with the use of the package “circular” [39] of the R statistical language [40]. These two variables were later treated with the R package “plotrix” [41] to construct polar plots, which allowed the visual assessment of water flow variability. All




Fig 2. Photos of the studied species. The three species studied, as recorded by Wally in transects at the hydrates site; a) *A. fimbria* b) *E. stoutii* and c) juvenile crab. The same scale bar applies to all 3 plates.

doi:10.1371/journal.pone.0163808.g002

environmental data were subsequently averaged by 4 h using the R package “zoo” [42] and used in cross-correlations and RDA analyses with visual count time series (see Section 2.4). Hourly environmental data are provided in [S1 Table](#). Visual counts and environmental data at 4 h frequency are provided in [S2 Table](#).

2.3 Periodogram analysis and modeling

We carried out an implemented periodogram analysis in order to match the peculiar features of the crawler’s video-censusing frequency and duration. To tackle this methodological challenge, we customized a statistical approach based on non-linear least squares models. Accordingly, faunal data were plotted against time (i.e. normalized abundances as recorded every 4 h) and non-linear (sinusoidal) curves were fitted to each time series with R statistical language [40], as a commonly accepted alternative to traditional time series analyses for the detection of biological rhythms [43]. In this way, the best approximations of the underlying periods in visual counts were obtained. All models follow [Eq 1.1](#), with a linear ascending trend added in

the case of sablefish (Eq 1.2).

$$\alpha_1 \cos(2\pi t/\alpha_4) + \alpha_2 \sin(2\pi t/\alpha_4) + \alpha_3 \quad (1.1)$$

$$\alpha_1 \cos(2\pi t/\alpha_4) + \alpha_2 \sin(2\pi t/\alpha_4) + \alpha_3 + \alpha_5 t \quad (1.2)$$

Where t is an integer [1:n] representing a time step of 4 h and α_4 is the corresponding period.

We screened for significant periodicities in oceanographic parameters, which were later coupled to the visual counts of the different megafaunal species and used as a proxy of environmental modulation on animal behavior. Lomb-Scargle periodogram analyses were applied on hourly datasets using the “lomb” R package [44]. In output plots, the periodicity is indicated as the peak of this statistic’s power that crossed the significant threshold set for $\alpha = 0.05$.

2.4 Cross-correlations and RDAs

Cross-correlation analyses were performed in R statistical language [40] between biological and environmental (i.e. normalized abundances as recorded every 4 h and 4 h averages respectively) time series, in order to reveal the significant ($\alpha = 0.05$) response of animals to the fluctuations of oceanographic parameters. As this is not a two-way process (i.e. animal abundances do not affect the oceanographic parameters tested), negative lags are not presented. The power of the statistic is presented at normalized scale (i.e. from -1 to 1). Since cross-correlations were used as a complementary method to interpret periodicity in megafaunal abundances, oceanographic parameters that did not display periodic signal were not tested.

Furthermore, the overall effects of environmental parameters on animal abundances were tested with Redundancy Analysis (RDA) performed in the “vegan” R package [45], for each of the three months studied. Based on the results of the cross-correlations (see 3.2), the biological data input for the RDAs was lagged for 12 h. Environmental data were centered and scaled (i.e. use of z-scores of the 4 h averages as, input instead of raw values) prior to analyses. The output is presented as biplots, an advanced version of scatterplots that depicts ordination (here expressed as Euclidean distance) of both observations and variables.

2.5 Waveforms

The 4 h sampling frequency for biological data allowed the creation of 6 bins per day, with each bin containing the averaged corresponding value across the 5 sampled days of each month. The obtained waveform curves for each species were plotted together with a horizontal threshold line, the Midline Estimated Statistic Of Rhythm (MESOR; see [46]). Values above this line represented the peak, hence describing the temporal amplitude of biological data. The MESOR was computed by re-averaging all curve values. Waveform curves were also calculated for the hourly environmental data and added onto the output plots when a significant correlation between the abundance of a species and an environmental variable was indicated.

3. Results

3.1 Assessment of periodicities

The sinusoidal models of biological data (Fig 3) revealed significant diel periodicities in the number of animal counts, close to the 24 h period. The periods varied from 21.62 h for the sablefish in June, to 25 h for juvenile crabs in December. The shortest periodicity could be a product of the scattered sampling frequency (i.e. 4 h) and of the reduced length of the time series, and should be interpreted together with the results of the cross-correlation analysis (see




Fig 3. Non-linear least squares models of biological time series. Sinusoidal curves (red) fitted to the animal counts series and their corresponded equations with calculated coefficients; a) and b) *A. fimbria* in June and July respectively, c) *E. stoutii* in December and finally, d) juvenile crabs in December. Black points represent animal abundances for each pair of transects (i.e. back and forth, 4 h frequency). Note the difference of scale among species in the number of counts, as well as the ascending trend in sablefish counts (see Section 4.1).

doi:10.1371/journal.pone.0163808.g003

below). A graphic representation of the model residuals (i.e. plots of residuals vs fitted values and histograms of residuals) is provided in S1 Fig.

The periodogram analysis output for hourly environmental data (Fig 4) showed the presence of significant rhythmicity in flow magnitude and the concentrations of chlorophyll and oxygen. In all 4 cases the dominant period was of diel character (i.e. $P = 24.79$ h, 25.63 h, 24.60 h and finally, 24.12 h for plates a) to d) respectively), with shorter periods also present in flow data, indicating distinct tidal signals.




Fig 4. Periodograms of environmental parameters. Lomb-Scargle periodograms for: a) and b) flow magnitude in June and July respectively, c) chlorophyll concentration in December and finally, d) oxygen concentration in December. The horizontal dashed lines indicate significance for $\alpha = 0.05$. The p values correspond to the peak in each periodogram (i.e. red, vertical dashed lines).

doi:10.1371/journal.pone.0163808.g004




Fig 5. Cross-correlations between animal counts and environmental parameters. Analyses used to assess the presence of temporal lag in the response of animal behavior to environmental fluctuations; a) and b) *A. fimbria* and flow magnitude in June and July respectively, c) *E. stoutii* and chlorophyll concentration in December and finally, d) juvenile crabs and oxygen concentration in December. “Cross-Correlation Function, (CCF)” is the measure of the power of the statistic. The dashed blue lines represent a CCF significance for $\alpha = 0.05$ (critical values: $\pm 1.96/\sqrt{n} = \text{approx. } \pm 0.36$ in a, and ± 0.35 in b, c and d, where n is the sample size). The dashed red lines represent a CCF significance for $\alpha = 0.1$ (accordingly, critical values: $\pm 1.65/\sqrt{n} = \text{approx. } \pm 0.3$ in all four plates).

doi:10.1371/journal.pone.0163808.g005

3.2 Response to environmental fluctuations

Significant correlations between biological and 4 h averaged environmental time series are presented in Fig 5. In the first three cases, sablefish and hagfish responded positively to environmental variations (i.e. power of ~ 0.6 against flow magnitude and chlorophyll concentration, respectively) after 12 h for a level of significance $\alpha = 0.05$, while juvenile crabs had a similar response (i.e. power ~ 0.4 to changes in oxygen concentration) for $\alpha = 0.1$.

The RDA biplots (Fig 6) between environmental parameters at t and biological data at $t + 12$ h provided a general view of the lagged effect of environmental variations on the species studied. Regarding the species that displayed periodicities in their abundances, sablefish numbers appeared to be strongly related to flow magnitude. In a more detailed view, this species scores were higher for the RDA1 axis than for the RDA2 axis (i.e. 1.56 vs ~ 0 in June and 1.31 vs -0.53 in July). In both months, the stronger vector on the RDA1 was flow magnitude (i.e. scores 0.77 in June and 0.73 in July). On the other hand, the pacific hagfish and juvenile crab abundances in December were influenced by chlorophyll and oxygen concentrations, respectively. In this month, the hagfish score was higher for the RDA1 axis than for the RDA2 axis (i.e. -1.51 vs 0.23), while the opposite was observed for the crabs (i.e. 0.46 vs 0.76). The dominant vectors in each axis were chlorophyll for RDA1 (i.e. score -0.75) and oxygen for RDA2 (i.e. score 0.53). For detailed RDA scores for species and environmental parameters see S3 Table.

As displayed visually in the output of waveform analysis (Fig 7), sablefish counts during the summer months peaked (i.e. mean values above the MESOR) right before midday, at a time of low currents. In contrast, hagfish were more abundant at crepuscular hours (dawn), following a peak of chlorophyll during the night. Finally, juvenile crab abundance reached a maximum in the early night hours, when oxygen concentrations were close to their minimum. The peaks in the waveforms of these two species come with a phase difference of 12 h over the same study period (i.e. hagfish and crabs in December of 2013). The waveform curves for visual count time series presented high standard deviation at their respective peak points. Nevertheless, the curve shape remains the same when the waveforms are calculated excluding the maximum values for each time bin (not shown), which reduces the possibility of biased/misleading results.




Fig 6. Redundancy Analysis (RDA) of the hydrates benthic community. RDA biplots for each one of the three months of the study period; a) June, b) July and c) December. The direction and length of the blue vectors depicting environmental variables represent positive or negative correlation with the ordination and proportionality of correlation respectively. Their biplot scores are depicted on the blue axes (normalized range -1 to 1). The scores for sites and species are weighted for scaling reasons and depicted on the black axes.

doi:10.1371/journal.pone.0163808.g006

Table 1 gives a summary of the characteristics of species rhythmic activity and the environmental parameters which correlate with their abundances and life habits (i.e. swimming in the water column for sablefish and close relationship to the seabed for hagfish and juvenile crabs).

4. Discussion

The major objective of the current study was to test for the presence of diel patterns in visual counts of the most abundant and regularly recorded species in video transects performed with Wally at the methane hydrates site of Barkley Canyon. We established a relationship between




Fig 7. Waveform analyses of concomitant biological and environmental data. Waveforms of animal counts (black, 4 h sampling frequency) as the average (\pm sd) of the 5 days sampling period and of environmental parameters (red, hourly); a) and b) *A. fimbria* in June and July respectively, c) *E. stoutii* in December and finally, d) juvenile crabs in December. Note the difference in scale among different species. The dashed horizontal lines correspond to the MESOR. Shaded parts represent the approximate night duration at the time of the study. X axis values correspond to local time (PST).

doi:10.1371/journal.pone.0163808.g007

animal activity rhythms and the environmental parameters that could substitute light intensity as the main “zeitgeber” at such depths, according to different species ecological habits. In the aphotic deep-sea, diel rhythms are a product of cycles mainly triggered by oceanographic processes, varying locally and regionally. As a result, our knowledge on such patterns remains limited, and generalizations should be treated cautiously. Anyway, present results fall within the expected scenarios on the expression of diel activity at such depths, with animals entering and leaving the field of view as they move on a daily basis in the water column or on the seafloor [15, 17, 28].

All targeted species showed a similar, near 24 h periodicity (see Table 1), as indicated directly by the least squares models calculated (see Fig 3) and indirectly by the cross-correlations (see Fig 5) between sablefish and flow magnitude, hagfish and chlorophyll and finally, juvenile crabs and oxygen concentrations respectively, also showing diel fluctuations (see Fig 4). The combined outcome of these analyses revealed a 12 h lagged response of the animals to such environmental parameters, depending on their ecology. Species inhabiting the benthic boundary layer with different locomotion strategies can react differently to changes in hydrodynamic conditions, with animals in closer relation to the seafloor being less susceptible to flow variations than swimmers [26, 47]. Accordingly, sablefish swimmers previously reported performing nektobenthic movements within the canyon (see Section 4.1), were highly affected by diel variation of flow magnitude. On the other hand, the two species that are more closely associated to the seabed (i.e. the endobenthic burrower hagfish and the epibenthic crabs; see Sections 4.2 and 4.3 respectively) were less affected by currents but responded primarily to diel changes in the concentrations of chlorophyll and oxygen. This was also confirmed by the RDA analyses (see Fig 6), where flow magnitude and chlorophyll and oxygen concentrations were the variables most closely associated with the respective sablefish, hagfish and crab abundances after a 12 h lag. Since no reverse flow conditions or major stochastic events occurred during the study period (see S2 Fig), our results should be considered an accurate representation of the community under normal oceanographic conditions.

4.1 Activity rhythms of sablefish

Sablefish activity in both summer months was connected to flow magnitude, with a peak right before midday (i.e. 11:00 local time PST; see Fig 7). This coincided with a time of low flow, which reached maximum velocity at night. A similar pattern was found at a nearby site [27], with most individuals recorded during times of low currents, as previously reported for the

Table 1. Species life habits (i.e. SW for swimmer, WL for walker, NKB for nektobenthic, ENB for endobenthic and EPB for epibenthic) in relation to their mode of displacement, the summary of their periodic activity (i.e. D for diurnal, C for crepuscular and N for nocturnal) and the most important environmental variables that regulate it.

Species	Period—Coefficient a_4 <i>p</i> value	Phase	Movement Type	Env. Variable
<i>A. fimbria</i>	21.62	<0.001	SW—NKB	Flow magnitude
	23.53	<0.001		D
<i>E. stoutii</i>	25.24	<0.001	SW—ENB	Chlorophyll
Crabs	25	<0.001	WL—EPB	Oxygen

doi:10.1371/journal.pone.0163808.t001

ecologically similar Atlantic cod *Gadus morhua* [48]. Periodic changes of the current regime in deep-sea environments can be detected by fish *via* their sensitive, well-developed lateral line systems, enabling them to keep their activity patterns synchronized [25]. Another study at similar depths in Barkley canyon reported the relationship between sablefish activity and tidal phenomena, this time expressed as water density [35]. In accordance, we believe that the ascending trend in sablefish counts in June and July of 2013 is a reflection of a lower frequency tidal signal (i.e. lunar tides), since the 5 days sampling periods for both months were entirely integrated in the spring tide phase (i.e. towards new moon).

Sablefish found at these depths are most commonly adult predators with demersal life habits [27, 49], in contrast to younger pelagic individuals [50–51]. As active swimmers preying on smaller benthic fish and invertebrates [50, 52], adult sablefish could be expected to react positively to increases of current velocity that lead to odor transport [53–54]. Sablefish have characteristically low-affinity hemoglobin, a characteristic which favors their active swimming, predatory life habits, but may result in reduced oxygen uptake within hypoxic waters [55]. Oxygen consumption in fish is directly related to the metabolic cost of locomotion [56], and the timing of sablefish activity within the hypoxic waters of Barkley Canyon during low flow conditions could be interpreted as a compromising strategy between search for prey and energy saving due to physiological limitations.

A diel displacement of the sablefish population within Barkley Canyon has been proposed, possibly in search of prey or to avoid predation by pelagic mammals, based on the phase difference between peaks in counts recorded from different fixed cameras [27]. Peaks in fish detection at consecutive temporal frames for nearby sampling sites can be used as an indication of populational movement [16, 57]. Nevertheless, the trajectory of the individuals in our study could not be followed with Wally. Thus, whether sablefish population moves from the hydrates site further up towards the canyon axis and back, on a daily basis, cannot be determined with 100% certainty. Further, coordinated research on that matter with the use of additional means (e.g. acoustic techniques), is needed to enlighten if the same population or distinct sub-populations are recorded at Barkley Canyon sites.

4.2 Activity rhythms of hagfish

Hagfish were found to be more associated with chemical variables such as chlorophyll concentration, rather than with flow. Their lifestyle is predominantly endobenthic, with animals spending most of their time resting coiled on the seabed (see Fig 2) or in self-built burrows [58–59]. Hagfish can opportunistically feed on several trophic levels [60], combining preying on small invertebrates and scavenging carcasses with the ability to take up dissolved organic matter *via* their skin [61–62]. While a high percentage of the organic matter within their diet can also originate from chemoautotrophs, many individuals' isotopic signatures correspond to macroalgal sources (*Eptatretus sp.*; [60, 63–65]). No predation or scavenging activity was observed in the video recordings analyzed in this study. The robot did not film the nearby hydrate mound, where chemosynthetic bacteria formed white mats with seasonally varying surface coverage [3, 14], that may supply resuspended food to the surrounding environment. Chlorophyll rich organic matter settling to the seafloor could provide an additional food source and regulate the animal burrow emergence and activity rhythms on and above it.

Regarding their phase during the day-night cycle, hagfish have been reported as mainly nocturnal species both in laboratory conditions and *in situ* at various depths, from shallow waters down to depths of 300 m [32, 59, 66]. In contrast with these findings, two chronobiological studies conducted recently at Barkley Canyon reported arrhythmic activity for the hagfish [27, 35]. In our results, hagfish emerged mainly at crepuscular hours, at dawn, in a display of

intraspecific plasticity of activity patterns [28]. Visual counts for hagfish in this study are higher than in the aforementioned studies, if the sampling period duration is taken into account, thus facilitating the detection of behavioral patterns. Hagfish are relatively sedentary animals [67] and the mobile nature of Wally enables the coverage of a greater area, providing a methodological advantage over fixed platforms. Although the underdeveloped eyes of species of the family Myxinidae can detect photoperiodic changes [68]), this inconsistency of Barkley Canyon case studies allows us to infer that at these depths environmental signals other than sunlight, such as the periodic chlorophyll influxes acting as an alternative food source, play an important role in synchronizing their circadian rhythms.

4.3 Activity rhythms of juvenile crabs

Our results suggest that the temporal pattern of juvenile crabs during the video transects is a combination of avoiding predation and physiological regulation by oxygen concentration. In the absence of sablefish from the study site during December of 2013, hagfish emerged as the major potential predator, known to prey on small benthic invertebrates, along with detritivory and a more opportunistic scavenging feeding behavior [69–71]. The waveforms of hagfish and crabs during that time have a semi-diurnal phase difference, inferring a possible avoidance behavior of the latter towards a potential predator. Minimizing predation risk through synchronization of diel activity is a common strategy in benthic decapod crustaceans [15–16]. In addition, this group appeared mostly during early night transects, when oxygen concentration was minimum. Benthic decapods of similar size have been observed to suppress their cryptic behavior under the effect of sharply induced hypoxia at a scale of hours (i.e. *Pilumnus spinifer*, *Pisidia longimana*; [72–73]), as well as in seasonally hypoxic conditions (*Munida quadrispina* small size class; [74]), which could explain their appearance in the field of view during minimum oxygen levels.

4.4 Methodology remarks

Extracting statistically meaningful results from relatively small faunal sample sizes was the principal methodological challenge we faced. Curve fitting based on non-linear least squares methods was used as an efficient alternative to the Lomb-Scargle periodogram analysis, since the latter can be sensitive to outliers at reduced sample sizes [75]. In contrast, although the non-linear least squares model may not provide a perfect fit from time to time (see Fig 3), the estimated wave period coefficients (i.e. a_4) are indeed statistically significant (see Table 1).

The sampling frequency of 4 h was deemed adequate for a chronobiological study, as it complied with the Nyquist criterion for sampling. Indeed, the temporal windows of activity for any type of rhythm (diurnal, nocturnal, crepuscular) were not missed. Even though fewer samples were generated in comparison to fixed cabled observatory cameras with higher imaging rate, greater spatial coverage was achieved due to the mobility of the crawler, resulting in observed animal abundances comparable to previous studies. Nevertheless, a new generation of crawlers able to perform autonomous transects will provide even better temporal resolution and bigger datasets over longer period of time; a further tool to be used to gain a better understanding of deep-sea environments. The range of the crawler lights (~50 m) and the transect duration (< 20 min) restricted any possible attraction bias to fast swimming organisms. Sablefish that entered the field of view should be close enough, and they were not observed following the crawler during transects. Therefore, we are confident that results were not affected by attraction or double-counting. On a similar note, linear video-recording transects performed by the crawler simulate the visual census methods used by divers in shallower waters and can offer a solution to the ever-increasing need for volunteer citizen scientists [13, 76].

5. Conclusions

This study highlights the potential of the use of Internet Operated deep-sea Crawlers for generating high-quality biological and environmental datasets. Diel patterns were detected in the activity of the sablefish *A. fimbria*, the pacific hagfish *E. stoutii* and a group of juvenile crabs. These patterns were conditioned by different environmental parameters, depending on the ecology of each species. Flow magnitude was the variable mostly associated with swimmer sablefish, with the preference for mild flow conditions being a possible adaptation to the hypoxic waters of Barkley Canyon. On the other hand, the activity of species closely related to the seafloor was regulated by chemical properties such as chlorophyll and oxygen levels. Predator—prey relationships were also indicated in the abundance patterns, with the crabs possibly avoiding the hagfish and suppressing this avoidance behavior at times of minimum oxygen concentration.

Supporting Information

S1 Fig. Plots of residuals vs fitted values and histograms of residuals for the models presented in Fig 3. The plates correspond to a) and e) *A. fimbria* in June, b) and f) *A. fimbria* in July, c) and g) *E. stoutii* in December and finally, d) and h) juvenile crabs in December. (TIF)

S2 Fig. Polar plots of flow conditions, depicting the magnitude (in m/s) and direction of water flow during the study period; a) June, b) July, and c) December. Each vector is an hourly averaged value. (TIF)

S1 Table. Hourly environmental data corresponding to the study period. (XLSX)

S2 Table. Visual counts of studied species, together with concomitant environmental data averaged at 4 h frequency. (XLSX)

S3 Table. Species and environmental variables scores for each RDA axis, per month. (XLSX)

Acknowledgments

This work was conducted within the framework of Ocean Networks Canada and NEPTUNE Canada. The authors would like to thank Joachim Vogt (Jacobs University Bremen) for his valuable consulting during the preparation of the manuscript, as well as the shipboard teams and staff scientists of Ocean Networks Canada and OceanLab Bremen. Steven Mihály and Akash Sastri from Ocean Networks Canada provided sensor data from POD4 (i.e. CTD and fluorometer data, respectively).

Author Contributions

Conceptualization: DC JA.

Data curation: CD DC.

Formal analysis: DC.

Funding acquisition: LT.

Investigation: CD.

Methodology: DC JA.

Project administration: LT.

Resources: LT.

Software: DC.

Supervision: LT JA.

Visualization: DC.

Writing – original draft: DC.

Writing – review & editing: CD LT AP JA.

References

1. Jørgensen BB, Boetius A. Feast and famine—microbial life in the deep-sea bed. *Nat Rev Micro*. 2007; 5: 770–781.
2. Chapman R, Pohlman J, Coffin R, Chanton J, Lapham L. Thermogenic gas hydrates in the northern Cascadia margin. *Eos Trans AGU*. 2004; 85: 361–365. doi: [10.1029/2004eo380001](https://doi.org/10.1029/2004eo380001)
3. Thomsen L, Barnes C, Best M, Chapman R, Pirenne B, Thomson R, et al. Ocean circulation promotes methane release from gas hydrate outcrops at the NEPTUNE Canada Barkley Canyon node. *Geophys Res Lett*. 2012; 39: L16605. doi: [10.1029/2012gl052462](https://doi.org/10.1029/2012gl052462)
4. Sibuet M, Olu K. Biogeography, biodiversity and fluid dependence of deep-sea cold-seep communities at active and passive margins. *Deep-Sea Res II*. 1998; 45: 517–567. doi: [10.1016/s0967-0645\(97\)00074-x](https://doi.org/10.1016/s0967-0645(97)00074-x)
5. Cordes EE, Cunha MR, Galéron J, Mora C, Roy KO-L, Sibuet M, et al. The influence of geological, geochemical, and biogenic habitat heterogeneity on seep biodiversity. *Mar Ecol*. 2010; 31: 51–65. doi: [10.1111/j.1439-0485.2009.00334.x](https://doi.org/10.1111/j.1439-0485.2009.00334.x)
6. Boetius A, Wenzhöfer F. Seafloor oxygen consumption fuelled by methane from cold seeps. *Nat Geosci*. 2013; 6: 725–734. doi: [10.1038/ngeo1926](https://doi.org/10.1038/ngeo1926)
7. Levin LA, James DW, Martin CM, Rathburn AE, Harris LH, Michener RH. Do methane seeps support distinct macrofaunal assemblages? Observations on community structure and nutrition from the northern California slope and shelf. *Mar Ecol Prog Ser*. 2000; 208: 21–39. doi: [10.3354/meps208021](https://doi.org/10.3354/meps208021)
8. Levin LA, Mendoza GF, Gonzalez JP, Thurber AR, Cordes EE. Diversity of bathyal macrofauna on the northeastern Pacific margin: the influence of methane seeps and oxygen minimum zones. *Mar Ecol*. 2010; 31: 94–110. doi: [10.1111/j.1439-0485.2009.00335.x](https://doi.org/10.1111/j.1439-0485.2009.00335.x)
9. Sahling H, Galkin SV, Salyuk A, Greinert J, Foerstel H, Piepenburg D, et al. Depth-related structure and ecological significance of cold-seep communities—a case study from the Sea of Okhotsk. *Deep-Sea Res I*. 2003; 50: 1391–1409. doi: [10.1016/j.dsr.2003.08.004](https://doi.org/10.1016/j.dsr.2003.08.004)
10. Glover AG, Gooday AJ, Bailey DM, Billett DSM, Chevaldonné P, Colaço A, et al. Temporal Change in Deep-Sea Benthic Ecosystems: A Review of the Evidence From Recent Time-Series Studies. *Adv Mar Biol*. 2010; 58: 1–95. doi: [10.1016/B978-0-12-381015-1.00001-0](https://doi.org/10.1016/B978-0-12-381015-1.00001-0) PMID: 20959156
11. Bahamon N, Aguzzi J, Bernardello R, Ahumada-Sempoal MA, Puigdefabregas J, Cateura J, et al. The new pelagic Operational Observatory of the Catalan Sea (OOCs) for the multisensor coordinated measurement of atmospheric and oceanographic conditions. *Sensors*. 2011; 11: 11251–11272. doi: [10.3390/s111211251](https://doi.org/10.3390/s111211251) PMID: 22247664
12. Ruhl HA, André M, Beranzoli L, Çağatay MN, Colaço A, Cannat M, et al. Societal need for improved understanding of climate change, anthropogenic impacts, and geo-hazard warning drive development of ocean observatories in European Seas *Prog Oceanogr*. 2011; 91: 1–33. doi: [10.1016/j.pocean.2011.05.001](https://doi.org/10.1016/j.pocean.2011.05.001)
13. Aguzzi J, Doya C, Tecchio S, De Leo FC, Azzurro E, Costa C, et al. Coastal observatories for monitoring of fish behaviour and their responses to environmental changes. *Rev Fish Biol Fisheries*. 2015; 25: 463–483. doi: [10.1007/s11160-015-9387-9](https://doi.org/10.1007/s11160-015-9387-9)
14. Purser A, Thomsen L, Barnes C, Best M, Chapman R, Hofbauer M, et al. Temporal and spatial benthic data collection via an internet operated Deep Sea Crawler. *Methods in Oceanography*. 2013; 5: 1–18. doi: [10.1016/j.mio.2013.07.001](https://doi.org/10.1016/j.mio.2013.07.001)

15. Aguzzi J, Company JB. Chronobiology of deep-water decapod crustaceans on continental margins. *Adv Mar Biol.* 2010; 58: 155–225. doi: [10.1016/B978-0-12-381015-1.00003-4](https://doi.org/10.1016/B978-0-12-381015-1.00003-4) PMID: [20959158](https://pubmed.ncbi.nlm.nih.gov/20959158/)
16. Aguzzi J, Company JB, Costa C, Menesatti P, Garcia JA, Bahamon N, et al. Activity rhythms in the deep-sea: a chronobiological approach. *Frontiers in Bioscience.* 2011; 16: 131–150. doi: [10.2741/3680](https://doi.org/10.2741/3680) PMID: [21196163](https://pubmed.ncbi.nlm.nih.gov/21196163/)
17. Aguzzi J, Company JB, Costa C, Matabos M, Azzurro E, Mànuel A, et al. Challenges to the assessment of benthic populations and biodiversity as a result of rhythmic behaviour: Video solutions from cabled observatories. *Oceanogr Mar Biol.* 2012; 50: 235–286. doi: [10.1201/b12157-6](https://doi.org/10.1201/b12157-6)
18. Sbragaglia V, Aguzzi J, García JA, Chiesa JJ, Angelini C, Sardà F. Dusk but not dawn burrow emergence rhythms of *Nephrops norvegicus* (Crustacea: Decapoda). *Scientia Marina.* 2013; 77: 641–647. doi: [10.3989/scimar.03902.28c](https://doi.org/10.3989/scimar.03902.28c)
19. Sbragaglia V, García JA, Chiesa JJ, Aguzzi J. Effect of simulated tidal currents on the burrow emergence rhythms of the Norway lobster (*Nephrops norvegicus*). *Mar Biol.* 2015; 162: 2007–2016. doi: [10.1007/s00227-015-2726-5](https://doi.org/10.1007/s00227-015-2726-5)
20. Sbragaglia V, Lamanna F, Mat AM, Rottlant G, Joly V, Ketmaier V, et al. Identification, characterization, and diel pattern of expression of canonical clock genes in *Nephrops norvegicus* (Crustacea: Decapoda) eyestalk. *PLoS ONE.* 2015; 10: e0141893. doi: [10.1371/journal.pone.0141893](https://doi.org/10.1371/journal.pone.0141893) PMID: [26524198](https://pubmed.ncbi.nlm.nih.gov/26524198/)
21. Bosiger YJ, McCormic MI. Temporal links in daily activity patterns between coral reef predators and their prey. *PLoS ONE.* 2014; 9: e111723. doi: [10.1371/journal.pone.0111723](https://doi.org/10.1371/journal.pone.0111723) PMID: [25354096](https://pubmed.ncbi.nlm.nih.gov/25354096/)
22. Kronfeld-Schor N, Dayan T. Partitioning of time as an ecological resource. *Annu Rev Ecol Evol Syst.* 2003; 34: 153–181. doi: [10.1146/annurev.ecolsys.34.011802.132435](https://doi.org/10.1146/annurev.ecolsys.34.011802.132435)
23. Navarro J, Votier SC, Aguzzi J, Chiesa JJ, Forero MG, Phillips RA. Ecological segregation in space, time and trophic niche of sympatric planktivorous petrels. *PLoS ONE.* 2013; 8: e62897. doi: [10.1371/journal.pone.0062897](https://doi.org/10.1371/journal.pone.0062897) PMID: [23646155](https://pubmed.ncbi.nlm.nih.gov/23646155/)
24. Aiken CM, Navarrete SA. Coexistence of competitors in marine metacommunities: environmental variability, edge effects, and the dispersal niche. *Ecology.* 2014; 95: 2289–2302. doi: [10.1890/13-0472.1](https://doi.org/10.1890/13-0472.1) PMID: [25230479](https://pubmed.ncbi.nlm.nih.gov/25230479/)
25. Wagner HJ, Kemp K, Mattheus U, Priede IG. Rhythms at the bottom of the deep sea: Cyclic current flow changes and melatonin patterns in two species of demersal fish. *Deep-Sea Res I.* 2007; 54: 1944–1956. doi: [10.1016/j.dsr.2007.08.005](https://doi.org/10.1016/j.dsr.2007.08.005)
26. Aguzzi J, Costa C, Furushima Y, Chiesa JJ, Company JB, Menesatti P, et al. Behavioral rhythms of hydrocarbon seep fauna in relation to internal tides. *Mar Ecol Prog Ser.* 2010; 418: 47–56. doi: [10.3354/meps08835](https://doi.org/10.3354/meps08835)
27. Doya C, Aguzzi J, Pardo M, Matabos M, Company JB, Costa C, et al. Diel behavioral rhythms in sablefish (*Anoplopoma fimbria*) and other benthic species as recorded by the Deep-sea cabled observatories in Barkley canyon (NEPTUNE-Canada). *J Mar Syst.* 2014; 130: 69–78. doi: [10.1016/j.jmarsys.2013.04.003](https://doi.org/10.1016/j.jmarsys.2013.04.003)
28. Colmenero A, Aguzzi J, Lombarte A, Bozzano A. Sensory constraints in temporal segregation in two species of anglerfish, *Lophius budegassa* and *L. piscatorius*. *Mar Ecol Prog Ser.* 2010; 416: 255–265. doi: [10.3354/meps08766](https://doi.org/10.3354/meps08766)
29. Schmidt K, Atkinson A, Steigenberger S, Fielding S, Lindsay MCM, Pond DW, et al. Seabed foraging by Antarctic krill: Implications for stock assessment, benthic-pelagic coupling, and the vertical transfer of iron. *Limnol Oceanogr.* 2011; 56: 1411–1428. doi: [10.4319/lo.2011.56.4.1411](https://doi.org/10.4319/lo.2011.56.4.1411)
30. Aguzzi J, Sbragaglia V, Tecchio S, Navarro J, Company JB. Rhythmic behaviour of marine benthopelagic species and the synchronous dynamics of benthic communities. *Deep-Sea Res I.* 2015; 95: 1–11. doi: [10.1016/j.dsr.2014.10.003](https://doi.org/10.1016/j.dsr.2014.10.003)
31. Bahamon N, Sardà F, Aguzzi J. Fuzzy diel patterns in catchability of deep-water species on the continental margin. *J Mar Sci.* 2009; 66: 2211–2218. doi: [10.1093/icesjms/fsp190](https://doi.org/10.1093/icesjms/fsp190)
32. Hart TD, Clemons JER, Wakefield WW, Heppell SS. Day and night abundance, distribution, and activity patterns of demersal fishes on Heceta Bank, Oregon. *Fish Bull.* 2010; 108: 466–477.
33. Juniper SK, Matabos M, Mihály S, Ajayamohan RS, Gervais F, Bui AOV. A year in Barkley Canyon: A time-series observatory study of mid-slope benthos and habitat dynamics using the NEPTUNE Canada network. *Deep-Sea Res II.* 2013; 92: 114–123. doi: [10.1016/j.dsr2.2013.03.038](https://doi.org/10.1016/j.dsr2.2013.03.038)
34. Lundsten EM, Anderson K, Paull CK, Caress DW, Thomas HJ, Riedel M. Morphology of Neptune Node Sites, Barkley Canyon, Cascadia Margin. Abstract OS31A-0977 presented at 2014 Fall Meeting, AGU, San Francisco CA.

35. Matabos M, Bui AOV, Mihály S, Aguzzi J, Juniper SK, Ajayamohan RS. High-frequency study of epibenthic megafaunal community dynamics in Barkley Canyon: A multi-disciplinary approach using the NEPTUNE Canada network. *J Mar Syst.* 2014; 130: 56–68. doi: [10.1016/j.jmarsys.2013.05.002](https://doi.org/10.1016/j.jmarsys.2013.05.002)
36. Gervais F, Juniper SK, Matabos M, Spicer A. *Marine Life Field Guide*, First edition. NEPTUNE-Canada publications; 2012.
37. Doya C, Chatzievangelou D, Bahamon N, Purser A, De Leo F, Juniper SK, et al. Yearly epibenthic megafaunal community dynamics at a cold seep (Barkley Canyon) using a tracked Sea Crawler. In Prep.
38. MATLAB R2014b, The Mathworks Inc, Natick, Massachusetts, USA.
39. Agostinelli C, Lund U. R package 'circular': Circular Statistics (version 0.4–7). 2013. <https://r-forge.r-project.org/projects/circular/>.
40. R Development Core Team. *R: A Language and Environment for Statistical Computing*. 2008.
41. Lemon J. Plotrix: a package in the red light district of R. *R-News*. 2006; 6: 8–12.
42. Zeileis A, Grothendieck G. zoo: S3 Infrastructure for Regular and Irregular Time Series. *J Stat Soft.* 2005; 14: 1–27. doi: [10.18637/jss.v014.i06](https://doi.org/10.18637/jss.v014.i06)
43. Refinetti R. Non-stationary time series and the robustness of circadian rhythms. *J Theor Biol.* 2004; 227: 571–581. doi: [10.1016/j.jtbi.2003.11.032](https://doi.org/10.1016/j.jtbi.2003.11.032) PMID: [15038991](https://pubmed.ncbi.nlm.nih.gov/15038991/)
44. Ruf T. The Lomb-Scargle Periodogram in Biological Rhythm Research: Analysis of Incomplete and Unequally Spaced Time-Series. *Biol Rhythm Res.* 1999; 30: 178–201. doi: [10.1076/brhm.30.2.178.1422](https://doi.org/10.1076/brhm.30.2.178.1422)
45. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, et al. *vegan: Community Ecology Package*. 2006. <https://CRAN.R-project.org/package=vegan>.
46. Aguzzi J, Bullock NM, Tosini G. Spontaneous internal desynchronization of locomotor activity and body temperature rhythms from plasma melatonin rhythm in rats exposed to constant dim light. *J Circad Rhythms.* 2006; 4: 6. doi: [10.1186/1740-3391-4-6](https://doi.org/10.1186/1740-3391-4-6) PMID: [16594995](https://pubmed.ncbi.nlm.nih.gov/16594995/)
47. Drew CA, Eggleston DB. Currents, landscape structure, and recruitment success along a passive-active dispersal gradient. *Landscape Ecology* 2006; 21: 917–931. doi: [10.1007/s10980-005-5568-6](https://doi.org/10.1007/s10980-005-5568-6)
48. Løkkeborg S, Bjordal Å, Fernö A. Responses of cod (*Gadus morhua*) and haddock (*Melanogrammus aeglefinus*) to baited hooks in the natural environment. *Can J Fish Aquat Sci.* 1989; 46: 1478–1483. doi: [10.1139/f89-189](https://doi.org/10.1139/f89-189)
49. King JR, McFarlane GA, Beamish RJ. Incorporating the dynamics of marine systems into the stock assessment and management of sablefish. *Prog Oceanogr.* 2001; 49: 619–639. doi: [10.1016/s0079-6611\(01\)00044-1](https://doi.org/10.1016/s0079-6611(01)00044-1)
50. Laidig TE, Adams PB, Samiere WM. Feeding habits of sablefish, *Anoplopoma fimbria*, off the coast of Oregon and California. *NOAA Tech Rep NMFS.* 1997; 130: 65–79.
51. Kimura DK, Shimada AM, Shaw FR. Stock structure and movement of tagged sablefish, *Anoplopoma fimbria*, in offshore northeast Pacific waters and the effects of El Niño-Southern Oscillation on migration and growth. *Fishery Bulletin* 1998; 96: 462–481.
52. Orlov AM. Ecological characteristics of the feeding of some Pacific predatory fish of South-East Kamchatka and northern Kuril Islands. *Rus J Aquat Ecol.* 1997; 6: 59–74.
53. Løkkeborg S, Olla BL, Pearson WH, Davies MW. Behavioural responses of sablefish, *Anoplopoma fimbria*, to bait odour. *J Fish Biol.* 1995; 46: 142–155. doi: [10.1111/j.1095-8649.1995.tb05953.x](https://doi.org/10.1111/j.1095-8649.1995.tb05953.x)
54. Aguzzi J, Jamieson AJ, Fujii T, Sbragaglia V, Costa C, Menesatti P, et al. Shifting behaviour of deep-sea buccinid gastropods at natural and simulated food falls. *Mar Ecol Prog Ser.* 2012; 458: 247–253. doi: [10.3354/meps09758](https://doi.org/10.3354/meps09758)
55. Rummer JL, Roshan-Moniri M, Balfry SK, Brauner CJ. Use it or lose it? Sablefish, *Anoplopoma fimbria*, a species representing a fifth teleostean group where the bNHE associated with the red blood cell adrenergic stress response has been secondarily lost. *J Exp Biol.* 2010; 213: 1503–1512. doi: [10.1242/jeb.038844](https://doi.org/10.1242/jeb.038844) PMID: [20400635](https://pubmed.ncbi.nlm.nih.gov/20400635/)
56. Liao JC. A review of fish swimming mechanics and behaviour in altered flows. *Phil Trans R Soc B.* 2007; 362: 1973–1993. doi: [10.1098/rstb.2007.2082](https://doi.org/10.1098/rstb.2007.2082) PMID: [17472925](https://pubmed.ncbi.nlm.nih.gov/17472925/)
57. Bozzano A, Sardà F, Ríos J. Vertical distribution and feeding patterns of the juvenile European hake, *Merluccius merluccius*, in the NW Mediterranean. *Fish Res.* 2005; 73: 29–36. doi: [10.1016/j.fishres.2005.01.006](https://doi.org/10.1016/j.fishres.2005.01.006)
58. Barss WH. Pacific hagfish, *Eptatretus stouti*, and black hagfish, *E. deani*: the Oregon fishery and port sampling observations, 1988–92. *Mar Fish Rev.* 1993; 55: 19–30.

59. Davies S, Griffiths A, Reimchen TE. Pacific hagfish, *Eptatretus stoutii*, Spotted rattfish, *Hydrolagus coliei*, and scavenger activity on tethered carrion in subtidal benthic communities off Western Vancouver Island. *Canadian Field-Naturalist* 2006; 120: 363:366.
60. McLeod RJ, Wing SR. Hagfish in the New Zealand fjords are supported by chemoautotrophy of forest carbon. *Ecology*. 2007; 88: 809–816. doi: [10.1890/06-1342](https://doi.org/10.1890/06-1342) PMID: [17536697](https://pubmed.ncbi.nlm.nih.gov/17536697/)
61. Stephens GC. Dissolved organic matter as a potential source of nutrition for marine organisms. *Am Zoologist*. 1968; 8: 95–106. doi: [10.1093/icb/8.1.95](https://doi.org/10.1093/icb/8.1.95)
62. Bucking C, Glover CN, Wood CM. Digestion under duress: Nutrient acquisition and metabolism during hypoxia in the pacific hagfish. *Physiol Biochem Zool*. 2011; 84: 607–617. doi: [10.1086/662630](https://doi.org/10.1086/662630) PMID: [22030853](https://pubmed.ncbi.nlm.nih.gov/22030853/)
63. MacAvoy SE, Carney RS, Fisher CR, Macko SA. Use of chemosynthetic biomass by large, mobile, benthic predators in the Gulf of Mexico. *Mar Ecol Prog Ser*. 2002; 225: 65–78. doi: [10.3354/meps225065](https://doi.org/10.3354/meps225065)
64. MacAvoy SE, Macko SA, Carney RS. Links between chemosynthetic production and mobile predators on the Louisiana continental slope: Stable carbon isotopes of specific fatty acids. *Chem Geol*. 2003; 201: 229–237. doi: [10.1016/s0009-2541\(03\)00204-3](https://doi.org/10.1016/s0009-2541(03)00204-3)
65. MacAvoy SE, Fisher CR, Carney RS, Macko SA. Nutritional associations among fauna at hydrocarbon seep communities in the Gulf of Mexico. *Mar Ecol Prog Ser*. 2005; 292: 51–60. doi: [10.3354/meps292051](https://doi.org/10.3354/meps292051)
66. Cox GK, Sandblom E, Richards JG, Farrell AP. Anoxic survival of the pacific hagfish (*Eptatretus stoutii*). *J Comp Physiol B*. 2011; 181: 361–371. doi: [10.1007/s00360-010-0532-4](https://doi.org/10.1007/s00360-010-0532-4) PMID: [21085970](https://pubmed.ncbi.nlm.nih.gov/21085970/)
67. Miyashita T, Palmer AR. Handed behavior in hagfish—an ancient vertebrate lineage—and a survey of lateralized behaviors in other invertebrate chordates and elongate vertebrates. *Biol Bull*. 2014; 226: 111–120. PMID: [24797093](https://pubmed.ncbi.nlm.nih.gov/24797093/)
68. Fernholm B. Diurnal variations in the behaviour of the Hagfish *Eptatretus burgeri*. *Mar. Biol*. 1974; 27: 351–356. doi: [10.1007/bf00394371](https://doi.org/10.1007/bf00394371)
69. Martini FH. The ecology of hagfishes. In: Jørgensen J.J., Lomholt J.P., Weber R.E., Malte H., (Eds.), *The Biology of Hagfishes*. Springer Netherlands. 1998; pp. 57–77.
70. Zintzen V, Roberts CD, Anderson MJ, Steward AL, Struthers CD, Harvey ES. Hagfish predatory behaviour and slime defense mechanism. *Scientific Reports*. 2011; 1: 131. doi: [10.1038/srep00131](https://doi.org/10.1038/srep00131) PMID: [22355648](https://pubmed.ncbi.nlm.nih.gov/22355648/)
71. Zintzen V, Rogers KM, Roberts CD, Steward AL, Anderson MJ. Hagfish feeding habits along a gradient inferred from stable isotopes. *Mar Ecol Prog Ser*. 2013; 485: 223–234. doi: [10.3354/meps10341](https://doi.org/10.3354/meps10341)
72. Haselmair A, Stachowitsch A, Zuschin M, Riedel B. Behaviour and mortality of benthic crustaceans in response to experimentally induced hypoxia and anoxia *in situ*. *Mar Ecol Prog Ser*. 2010; 414: 195–208. doi: [10.3354/meps08657](https://doi.org/10.3354/meps08657)
73. Riedel B, Pados T, Pretterebner K, Schiemer L, Steckbauer A, Haselmair A, et al. Effect of hypoxia and anoxia on invertebrate behaviour: ecological perspectives from species to community level. *Bio-geosciences* 2014; 11: 1491–1518. doi: [10.5194/bg-11-1491-2014](https://doi.org/10.5194/bg-11-1491-2014)
74. Doya C, Aguzzi J, Chatzievangelou D, Costa C, Company JB, Tunnicliffe V. The seasonal use of small-scale space by benthic species in a transiently hypoxic area. *J Mar Syst*. 2016; 154: 280–290. doi: [10.1016/j.jmarsys.2015.09.005](https://doi.org/10.1016/j.jmarsys.2015.09.005)
75. Schimmel M. Emphasizing difficulties in the detection of rhythms with Lomb-Scargle periodograms. *Biol Rhythm Res*. 2001; 32: 341–345. doi: [10.1076/brhm.32.3.341.1340](https://doi.org/10.1076/brhm.32.3.341.1340) PMID: [11665689](https://pubmed.ncbi.nlm.nih.gov/11665689/)
76. Azzurro E, Aguzzi J, Maynou F, Chiesa JJ, Savini D. Diel rhythms in shallow Mediterranean rocky-reef fishes: a chronobiological approach with the help of trained volunteers. *J Mar Biol Ass UK*. 2013; 93: 461–470. doi: [10.1017/s0025315412001166](https://doi.org/10.1017/s0025315412001166)