




OPEN

High level of fatty liver index predicts new onset of diabetes mellitus during a 10-year period in healthy subjects

Yukimura Higashiura^{1,4}, Masato Furuhashi^{1,4}, Marenao Tanaka^{1,4}, Satoko Takahashi¹, Masayuki Koyama^{1,2}, Hirofumi Ohnishi^{1,2}, Keita Numata³, Takashi Hisasue³, Nagisa Hanawa³, Norihito Moniwa¹, Kazufumi Tsuchihashi¹ & Tetsuji Miura¹

Fatty liver index (FLI), a predictor of nonalcoholic fatty liver disease, has been reported to be associated with several metabolic disorders. This study aimed to evaluate the relationship between FLI and new onset of diabetes mellitus (DM). We investigated the association of FLI with new onset of DM during a 10-year period in subjects who received annual health examinations ($n = 28,990$). After exclusion of subjects with DM at baseline and those with missing data, a total of 12,290 subjects (male/female: 7925/4365) who received health examinations were recruited. FLI was significantly higher in males than in females. During the 10-year period, DM was developed in 533 males (6.7%) and 128 females (2.9%). Multivariable Cox proportional hazard models with a restricted cubic spline showed that the risk of new onset of DM increased with a higher FLI at baseline in both sexes after adjustment of age, fasting plasma glucose, habits of alcohol drinking and current smoking, family history of DM and diagnosis of hypertension and dyslipidemia at baseline. When the subjects were divided into subgroups according to tertiles of FLI level at baseline (T1–T3) in the absence and presence of impaired fasting glucose (IFG), hazard ratios after adjustment of the confounders gradually increased from T1 to T3 and from the absence to presence of IFG in both male and female subjects. In conclusion, a high level of FLI predicts new onset of DM in a general population of both male and female individuals.

Diabetes mellitus (DM) is one of major medical concerns in metabolic diseases¹. Because of changes in lifestyle including habits of eating and exercise, the number of patients with DM is continuously increasing worldwide². Since patients with DM have compromised healthy longevity due to multiple complications including diabetic nephropathy and atherosclerotic cardiovascular disease^{3,4}, prevention of DM is a critical issue. Therefore, it is crucial to find out subjects at high risk for development of DM for performing appropriate intervention such as exercise encouragement and dietary advice at an early stage.

Nonalcoholic fatty liver disease (NAFLD), a chronic liver disease, has been highlighted as a lifestyle-related disease^{5,6}, and the prevalence of NAFLD has been increasing worldwide, leading to a prominent cause of liver-related prognosis^{7,8}. It was shown in a cohort study that subjects with NAFLD were at a higher risk for the development of DM than were those without NAFLD⁹. Meta-analyses also showed that NAFLD diagnosed by altered serum liver enzymes, radiological findings or histological evidence increases the risk of type 2 DM^{10,11}.

For diagnosis of NAFLD, liver biopsy as an invasive procedure is required¹², but several noninvasive procedures in adequate concordance with histological findings have recently been established using imaging tools and several biochemical markers including fatty liver index (FLI)¹³. FLI calculated by using waist circumference (WC), body mass index (BMI), and levels of triglycerides and γ -glutamyl transferase (γ GTP)¹⁴ has been reported to be highly concordant with the histological criteria for NAFLD^{15–17}. It has recently been reported that NAFLD diagnosed by FLI is a good predictor for incidence of type 2 DM^{18–25} (Supplementary Table S1).

¹Department of Cardiovascular, Renal and Metabolic Medicine, Sapporo Medical University School of Medicine, S-1, W-16, Chuo-ku, Sapporo 060-8543, Japan. ²Department of Public Health, Sapporo Medical University School of Medicine, Sapporo, Japan. ³Department of Health Checkup and Promotion, Keijinkai Maruyama Clinic, Sapporo, Japan. ⁴These authors contributed equally: Yukimura Higashiura, Masato Furuhashi and Marenao Tanaka. ✉email: furuhashi@sapmed.ac.jp




Figure 1. Flow chart of the selected study participants. Among 28,990 subjects enrolled in 2006, a total of 12,290 subjects (male/female: 7925/4365) were finally recruited for analyses in the present study. *BMI* body mass index, *DM* diabetes mellitus, *FPG* fasting plasma glucose, *γGTP* γ -glutamyl transferase, *HbA1c* hemoglobin A1c, *WC* waist circumference.

However, the relationship of FLI with new onset of DM has not yet been investigated in a large cohort with a sufficiently long observational period and/or as a continuous variable for FLI. Therefore, in the present study, we investigated the effect of FLI level at baseline on new onset of DM during a 10-year period in a large number of subjects divided by sex.

Results

Characteristics of the study subjects. A flow chart of the study participants is shown in Fig. 1. The characteristics of the enrolled and excluded subjects are shown in Supplementary Table S2. The excluded subjects were significantly younger and more metabolic healthy than the enrolled subjects. Demographic parameters and metabolic profiles of the recruited subjects are shown in Table 1. Male subjects had significantly larger BMI and WC, higher systolic and diastolic blood pressures, higher levels of albumin, uric acid, fasting plasma glucose (FPG), hemoglobin A1c (HbA1c) and triglycerides and lower levels of estimated glomerular filtration rate (eGFR) and high-density lipoprotein cholesterol than did female subjects. FLI was significantly higher in male subjects than in female subjects. The frequencies of habits of smoking and alcohol drinking were lower in female subjects than in male subjects.

Basal characteristics of male and female subjects divided into the three subgroups according to tertiles of FLI levels at baseline are shown in Tables 2 and 3, respectively. There were significant differences in levels of FPG and HbA1c, prevalence of alcohol drinking habit, comorbidity of hypertension and dyslipidemia, and family history of DM between the three groups of FLI in both male and female subjects. Levels of FPG and HbA1c in the T3 group of FLI tended to be higher than those in the T1 group of FLI in both male and female subjects.

Cumulative incidence of new onset of DM during the follow-up period. Among the 12,290 subjects (male/female: 7925/4365), 533 male subjects (6.7%) and 128 female subjects (2.9%) developed new onset of DM during a 10-year period. The mean follow-up period was 9.5 years (range: 1–10 years), and follow-up summation was 82,709 (male/female: 53,320/29,389) person-years. The cumulative incidence of new onset of DM was 6.1% (male/female: 7.7%/3.1%).

Prediction of new onset of DM by levels of FLI and FPG. Receiver operating characteristic (ROC) curve analyses for predicting new onset of DM showed that the area under curves (AUCs) of FLI at baseline in males (Fig. 2A) and females (Fig. 2B) were 0.71 and 0.71, respectively. The cutoff points of FLI at baseline in males and females were 56.6 and 23.7, respectively. On the other hand, the AUCs of FPG at baseline for predicting new onset of DM in ROC analyses in males (Fig. 2C) and females (Fig. 2D) were 0.82 and 0.80, respectively.

	Total	Male	Female	P
	n = 12,290	n = 7925	n = 4365	
Age (years)	48 ± 10	48 ± 10	48 ± 10	<0.001
Body mass index	23.2 ± 3.4	24.0 ± 3.1	21.7 ± 3.3	<0.001
Waist circumference (cm)	83.4 ± 9.2	85.9 ± 8.3	79.9 ± 9.1	<0.001
Systolic blood pressure (mmHg)	117 ± 16	120 ± 16	111 ± 16	<0.001
Diastolic blood pressure (mmHg)	74 ± 11	77 ± 11	70 ± 10	<0.001
Smoking habit	4231 (34.4)	3431 (43.3)	800 (18.3)	<0.001
Alcohol drinking habit	5634 (45.8)	4492 (56.7)	1142 (26.1)	<0.001
Family history				
Diabetes mellitus	2115 (17.2)	1220 (15.4)	895 (20.5)	<0.001
Comorbidity				
Hypertension	2068 (16.8)	1582 (20.0)	486 (11.1)	<0.001
Dyslipidemia	2720 (22.1)	1674 (21.1)	1046 (24.0)	<0.001
Biochemical data				
Albumin (g/dL)	4.4 ± 0.2	4.4 ± 0.2	4.3 ± 0.2	<0.001
eGFR (mL/min/1.73 m ²)	84.5 ± 14.6	83.3 ± 14.0	86.7 ± 15.4	<0.001
Uric acid (mg/dL)	5.5 ± 1.4	6.1 ± 1.2	4.4 ± 0.9	<0.001
FPG (mg/dL)	90 ± 9.3	92 ± 9.3	86 ± 8.4	<0.001
Hemoglobin A1c (%)	5.2 ± 0.4	5.2 ± 0.4	5.2 ± 0.4	<0.001
AST (U/L)	23 (18–26)	25 (19–28)	20 (16–22)	<0.001
ALT (U/L)	26 (15–31)	31 (18–36)	18 (12–20)	<0.001
γGTP (U/L)	50 (19–56)	63 (26–72)	26 (14–26)	<0.001
Total cholesterol (mg/dL)	204 ± 34	205 ± 34	204 ± 34	0.05
LDL cholesterol (mg/dL)	122 ± 31	124 ± 31	119 ± 31	<0.001
HDL cholesterol (mg/dL)	61 ± 16	56 ± 14	69 ± 15	<0.001
Non-HDL cholesterol (mg/dL)	144 ± 35	149 ± 35	135 ± 34	<0.001
Triglycerides (mg/dL)	113 (63–136)	131 (77–158)	78 (50–93)	<0.001
FLI	30.1 (7.9–48.2)	38.6 (16.5–58.2)	14.5 (3.5–17.5)	<0.001

Table 1. Characteristics of the recruited subjects. Variables are expressed as number (%), means ± SD or medians (interquartile ranges). *AST* aspartate aminotransferase, *ALT* alanine aminotransferase, *eGFR* estimated glomerular filtration rate, *FLI* fatty liver index, *FPG* fasting plasma glucose, *γGTP* γ-glutamyl transferase, *HDL* high-density lipoprotein, *LDL* low-density lipoprotein.

The cutoff points of FPG at baseline in male and female subjects were 100 mg/dL and 93 mg/dL, respectively. The AUCs of FPG tended to be higher than those of FLI in both male and female subjects.

Risk of FLI at baseline for new onset of DM during a 10-year follow-up period. Multivariable Cox proportional hazard models with a restricted cubic spline showed that the hazard ratio (HR) of DM development increased with a higher FLI at baseline in both males (Fig. 3A) and females (Fig. 3B) after adjustment of age, FPG, habits of smoking and alcohol drinking, family history of DM and diagnosis of hypertension and dyslipidemia at baseline.

When the 1st tertile (T1) group of FLI was used as the reference, multivariable Cox proportional hazard model analysis after adjustment of age, FPG, habits of smoking and alcohol drinking, family history of DM and diagnosis of hypertension and dyslipidemia showed that HRs in the T2 and T3 groups were significantly higher than HR in the T1 group of FLI in male subjects (Table 4). In female subjects, the adjusted HR in the T3 group of FLI was significantly higher than that in the T1 group of FLI in female subjects (Table 4).

When further divided by the absence and presence of impaired fasting glucose (IFG), HRs after adjustment of covariates gradually increased from T1 to T3 and from the absence to presence of IFG in both male and female subjects (Table 5). Even in the absence of IFG, HRs in the T3 group were significantly higher than those in the T1 group in both male and female subjects.

Discussion

The present study showed that a high level of FLI was significantly associated with the risk of DM development during a 10-year period in both male and female subjects. It has been reported that FLI, originally developed for diagnosis of NAFLD, is associated with several metabolic diseases²⁶. Furthermore, FLI has been proposed to be a marker for incidence of type 2 DM (Supplementary Table S1)^{18–25}. Earlier studies showed an association between a high FLI level (≥60–70) as diagnosis of NAFLD and incidence of DM in logistic regression analyses, but the cumulative incidence of DM was not analyzed^{18–20}. Several studies also showed a high FLI level (FLI ≥ 60) was associated with the cumulative incidence of DM in Cox proportional regression analyses using a relatively small

	T1 (0.9–21.8)	T2 (21.9–49.7)	T3 (49.8–99.7)	P
	n = 2650	n = 2633	n = 2642	
Age (years)	47 ± 11	49 ± 9	48 ± 9	<0.001
Body mass index	21.4 ± 1.9	23.9 ± 1.9	26.6 ± 3.0	<0.001
Waist circumference (cm)	78.7 ± 5.3	86.1 ± 4.9	93.1 ± 7.0	<0.001
Systolic blood pressure (mmHg)	115 ± 15	120 ± 15	125 ± 15	<0.001
Diastolic blood pressure (mmHg)	73 ± 10	77 ± 10	81 ± 10	<0.001
Smoking habit	1139 (44.3)	1121 (43.5)	1219 (47.5)	0.007
Alcohol drinking habit	1011 (37.6)	1189 (44.2)	1271 (47.4)	<0.001
Family history				
Diabetes mellitus	372 (14.0)	405 (15.3)	443 (16.8)	0.023
Comorbidity				
Hypertension	152 (5.7)	299 (11.1)	447 (16.7)	<0.001
Dyslipidemia	39 (1.5)	113 (4.2)	176 (6.6)	<0.001
Biochemical data				
Albumin (g/dL)	4.4 ± 0.2	4.4 ± 0.2	4.5 ± 0.2	<0.001
eGFR (mL/min/1.73 m ²)	85.1 ± 14.0	82.3 ± 13.9	82.6 ± 14.1	<0.001
Uric acid (mg/dL)	5.7 ± 1.1	6.1 ± 1.2	6.6 ± 1.2	<0.001
FPG (mg/dL)	89 ± 9	92 ± 9	95 ± 12	<0.001
Hemoglobin A1c (%)	5.1 ± 0.4	5.3 ± 0.4	5.4 ± 0.4	<0.001
AST (U/L)	21 (17–23)	24 (19–26)	31 (22–34)	<0.001
ALT (U/L)	20 (15–24)	28 (19–33)	45 (27–53)	<0.001
γGTP (U/L)	30 (20–34)	53 (30–61)	108 (51–125)	<0.001
Total cholesterol (mg/dL)	194 ± 30	206 ± 32	216 ± 35	<0.001
LDL cholesterol (mg/dL)	116 ± 28	126 ± 30	126 ± 33	<0.001
HDL cholesterol (mg/dL)	62 ± 15	56 ± 14	51 ± 12	<0.001
Non-HDL cholesterol (mg/dL)	132 ± 30	150 ± 32	165 ± 35	<0.001
Triglycerides (mg/dL)	78 (58–94)	120 (87–144)	197 (125–230)	<0.001

Table 2. Characteristics of male subjects divided by tertiles of FLI at baseline (n = 7925). Variables are expressed as number (%), means ± SD or medians (interquartile ranges). *AST* aspartate aminotransferase, *ALT* alanine aminotransferase, *eGFR* estimated glomerular filtration rate, *FLI* fatty liver index, *FPG* fasting plasma glucose, *γGTP* γ-glutamyl transferase, *HDL* high-density lipoprotein, *LDL* low-density lipoprotein.

number of subjects (n = 1142–1922)^{21,22,24}. Furthermore, in only one study using 389 subjects with prediabetes defined as FPG level of 110–125 mg/dL, the association of DM development with level of FLI as a continuous variable was examined by Cox proportional regression analysis²³. The present study showed that the risk of DM development continuously increased with a higher FLI at baseline in both male and female subjects in multivariable Cox proportional hazard models with a restricted cubic spline using a large number of subjects (n = 12,290, male/female: 7935/4365).

Since there is a sex difference in components of FLI calculation, including BMI, WC, triglycerides and γ-GTP, being higher in male subjects than in female subjects^{27–30}, it is necessary to analyze the FLI value divided by sex. In fact, there was a significant sex difference in FLI level in the present study as well as in a previous study¹⁶. Definitions of FLI ≥ 60 as NAFLD and FLI < 30 as non-NAFLD have been used in several studies^{20–24}. However, it has been reported that cutoff levels of FLI for diagnosis of NAFLD seem to be lower in Asians than in Europeans: FLI ≥ 30 in China¹⁷ and FLI ≥ 60 in Italy¹⁴, in which subjects were not divided by sex in the both studies^{14,17}. Although there have been no studies about the validation of FLI in other races including African Americans and Hispanics, it has been reported that the prevalence of NAFLD is higher in Hispanics than in non-Hispanic whites and African Americans³¹. These findings suggest that there might be racial and sex differences in cutoff levels of FLI for diagnosis of NAFLD. In the present study, the optimal cutoff value of FLI to predict incidence of DM was higher in male subjects than in female subjects (FLI: 56.6 vs. 23.7) (Fig. 2A,B). Therefore, instead of the conventional division of FLI (FLI ≥ 60 and < 30), we analyzed HRs in subgroups according to tertiles of FLI level at baseline in both sexes in the present study (Table 4).

The level of FPG at baseline was found to be a strong predictor of new onset of DM in the present study. The ability of FLI at baseline to predict new onset of DM was comparable to that of FPG in both male and female subjects. When further divided by the absence and presence of IFG in tertiles of FLI level, HRs after adjustment of confounders gradually increased from T1 to T3 and from the absence to presence of IFG in both male and female subjects (Table 5). Similar results were obtained in a previous study by Hirata et al., though the number of subjects was relatively small (n = 4439, male/female: 1498/2,941)²⁵. HRs of DM development in the T2 and T3 groups in the presence of IFG in the present study (male/female: 16.6–30.0/12.9–25.7) were higher than those in the previous study (male/female: 3.4–5.2/5.9–6.2)²⁵. Furthermore, HRs of DM development in the T3 group in the

	T1 (0.4–4.4)	T2 (4.5–12.6)	T3 (12.7–97.4)	
	n = 1458	n = 1455	n = 1452	P
Age (years)	43 ± 9	48 ± 9	51 ± 10	<0.001
Body mass index	19.2 ± 1.5	21.2 ± 1.7	24.8 ± 3.3	<0.001
Waist circumference (cm)	70.9 ± 4.7	78.0 ± 4.8	87.7 ± 7.9	<0.001
Systolic blood pressure (mmHg)	105 ± 13	110 ± 16	118 ± 16	<0.001
Diastolic blood pressure (mmHg)	66 ± 9	69 ± 10	75 ± 10	<0.001
Smoking habit	263 (18.6)	254 (17.9)	285 (20.2)	0.219
Alcohol drinking habit	211 (14.5)	257 (17.5)	261 (17.8)	0.001
Family history				
Diabetes mellitus	272 (18.7)	292 (20.0)	331 (22.8)	0.020
Comorbidity				
Hypertension	28 (1.9)	76 (5.2)	216 (14.8)	<0.001
Dyslipidemia	21 (1.4)	53 (3.6)	105 (7.2)	<0.001
Biochemical data				
Albumin (g/dL)	4.3 ± 0.2	4.3 ± 0.2	4.3 ± 0.2	0.234
eGFR (mL/min/1.73 m ²)	89.1 ± 15.1	85.9 ± 15.3	85.0 ± 15.6	<0.001
Uric acid (mg/dL)	4.1 ± 0.8	4.3 ± 0.9	4.8 ± 1.0	<0.001
FPG (mg/dL)	83 ± 7	86 ± 8	90 ± 9	<0.001
Hemoglobin A1c (%)	5.1 ± 0.3	5.2 ± 0.3	5.3 ± 0.4	<0.001
AST (U/L)	19 (16–20)	20 (16–22)	22 (18–24)	<0.001
ALT (U/L)	14 (11–16)	16 (12–18)	23 (15–27)	<0.001
γGTP (U/L)	16 (12–18)	21 (14–23)	41 (19–44)	<0.001
Total cholesterol (mg/dL)	191 ± 31	205 ± 32	216 ± 33	<0.001
LDL cholesterol (mg/dL)	106 ± 26	120 ± 30	131 ± 31	<0.001
HDL cholesterol (mg/dL)	75 ± 14	70 ± 14	63 ± 14	<0.001
Non-HDL cholesterol (mg/dL)	117 ± 27	134 ± 31	154 ± 34	<0.001
Triglycerides (mg/dL)	49 (40–60)	68 (54–85)	100 (76–134)	<0.001

Table 3. Characteristics of female subjects divided by tertiles of FLI at baseline (n = 4365). Variables are expressed as number (%), means ± SD or medians (interquartile ranges). *AST* aspartate aminotransferase, *ALT* alanine aminotransferase, *eGFR* estimated glomerular filtration rate, *FLI* fatty liver index, *FPG* fasting plasma glucose, *γGTP* γ-glutamyl transferase, *HDL* high-density lipoprotein, *LDL* low-density lipoprotein.

absence of IFG were significantly higher than those in the T1 group of both sexes in the present study, but there was no significant difference in HRs in the non-IFG group of both sexes in the previous study²⁵. Possible reasons for the difference were the number of study subjects (12,290 vs. 4439) and length of the follow-up period (mean: 9.5 vs. 3.0 years). Therefore, the results of the present study may accurately reflect the risk of DM development.

It has been reported that insulin resistance with visceral obesity causes compensatory hyperinsulinemia, leading to the development of NAFLD³². Furthermore, fat accumulation in the liver contributes to an increase of glucose production and high serum free fatty acid flux, which is caused by impaired insulin action³³. Conversely, insulin-resistant fatty liver overproduces glucose and very low-density lipoprotein, leading to exhaustion of the pancreatic β cell reserve and subsequent development of DM³². Since FLI has been reported to be strongly associated with reduced insulin sensitivity assessed by the euglycemic hyperinsulinemic clamp method³⁴, a possible mechanism for the association between FLI and DM development is insulin resistance. As another possibility of the mechanism, several hepatokines, secretory molecules from the liver, mediate the relationship between NAFLD and the development of DM. A steatotic and inflamed liver has been reported to secrete several hepatokines, including fetuin-A³⁵, fibroblast growth factor 21^{36,37}, selenoprotein P³⁸ and xanthine oxidase^{39–41}, which are known to have endocrine functions at extrahepatic sites to cause insulin resistance and other adverse effects on glucose homeostasis.

The present study has some limitations. First, FLI was used as a surrogate marker for NAFLD because of the absence of liver biopsy as a gold standard for diagnosis of NAFLD and imaging examinations including ultrasonography, computed tomography and magnetic resonance spectroscopy¹². However, it has recently been reported that FLI can predict NAFLD and overcome the limitation of diagnosis based on abdominal ultrasonography⁴². Second, the presence of hepatitis B and hepatitis C was unknown at baseline, though the prevalence of hepatitis B (0.63%) and hepatitis C (0.49%) was reported to be relatively low in the Japanese population^{43,44}. Third, accurate information on alcohol consumption was not obtained in the present study. However, when subjects who had an alcohol drinking habit were excluded from analyses, most of the results were similar (Supplementary Tables S3, S4). Fourth, the possibility of selection bias in the samples cannot be excluded since the study subjects were urban residents who received annual health checkups in a single clinic.




Figure 2. Prediction of new onset of DM by FLI and FPG at baseline. (A,B) Receiver operating characteristic (ROC) curves of fatty liver index (FLI) at baseline to predict new onset of diabetes mellitus (DM) in males (A) and females (B). (C,D) ROC curves of fasting plasma glucose (FPG) at baseline to predict new onset of DM in males (C) and females (D). AUC area under curve, CI confidence interval.

Finally, the relationship between change in FLI level and new onset of DM was not investigated in the present study, and this needs to be examined in the future.

In conclusion, an elevated FLI level can predict the development of DM during a 10-year period in a general population of both sexes. The pathophysiology underlying the association between factors that influence FLI level and DM development needs to be addressed in future basic and clinical studies.

Methods

Study subjects. A total of 28,990 subjects who received annual medical checkups at Keijinkai Maruyama Clinic, Sapporo, Japan in 2006 were enrolled in this registry^{45,46}. A flow chart of the study subjects is shown in Fig. 1. Exclusion criteria were the diagnosis of DM at baseline and absence of data for WC, BMI and laboratory data including FPG, HbA1c, triglycerides and γ GTP at baseline. After prespecified exclusion, a total of 12,290 subjects (male/female: 7925/4365) who received annual health examinations at least once from 2007 to 2015 were finally recruited in the present study. The study was performed with the approval of the institutional ethical committee of Sapporo Medical University (Numbers: 29-2-64, 30-2-32) and conformed to the principles of the Declaration of Helsinki. Written informed consent was obtained from the recruited subjects.

Measurements. Blood pressure measurements, medical examinations and samplings of blood were performed after an overnight fast. Body height and weight were measured, and BMI was calculated as body weight in kilograms divided by height in meters squared. HbA1c level was presented as the National Glycohemoglobin Standardization Program (NGSP) equivalent value. eGFR was calculated by an equation for Japanese: $eGFR \text{ (mL/min/1.73 m}^2\text{)} = 194 \times \text{serum creatinine}^{(-1.094)} \times \text{age}^{(-0.287)} \times 0.739 \text{ (if female)}$ ⁴⁷. A self-administered questionnaire survey was performed to obtain information on current smoking habit, alcohol drinking habit (≥ 3 times/week), family history of DM, and use of drugs for hypertension, dyslipidemia and DM.




Figure 3. Hazard ratio of the development of DM by FLI at baseline. (A,B) Hazard ratios (HRs) for the development of diabetes mellitus (DM) by fatty liver index (FLI) at baseline by multivariable Cox proportional hazard models with a restricted cubic spline in males (A) and females (B) after adjustment of age, fasting plasma glucose, habits of smoking and alcohol drinking, family history of DM, and diagnosis of hypertension and dyslipidemia at baseline during a 10-year follow-up period. Solid line: HR, dashed line: 95% confidence interval (CI). The reference values of FLI in males and females were 0.9 and 0.4 as minimum values, respectively.

DM was diagnosed in accordance with the guidelines of the American Diabetes Association⁴⁸: self-reported use of anti-diabetic drugs, FPG ≥ 126 mg/dL or HbA1c $\geq 6.5\%$. IFG was defined as FPG ≥ 100 mg/dL⁴⁸. Hypertension was diagnosed as self-reported use of anti-hypertensive drugs, systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg. Dyslipidemia was diagnosed as self-reported use of anti-dyslipidemic drugs, low-density lipoprotein cholesterol ≥ 140 mg/dL, triglycerides ≥ 150 mg/dL or high-density lipoprotein cholesterol < 40 mg/dL.

	Male (n = 7925)		Female (n = 4365)	
	HR (95% CI)	P	HR (95% CI)	P
FLI				
T1	Reference	–	Reference	–
T2	1.43 (1.05–1.94)	0.022	1.05 (0.57–1.92)	0.885
T3	2.44 (1.84–3.24)	<0.001	1.95 (1.13–3.36)	0.016
Age (per 1 year)	1.01 (1.00–1.02)	0.099	1.01 (0.99–1.03)	0.531
FPG (per 1 mg/dL)	1.13 (1.12–1.14)	<0.001	1.12 (1.10–1.14)	<0.001
Smoking habit	1.68 (1.41–2.01)	<0.001	1.42 (0.92–2.20)	0.115
Alcohol drinking habit	0.60 (0.50–0.71)	<0.001	0.82 (0.55–1.24)	0.356
Family history of DM	0.90 (0.65–1.24)	0.510	0.77 (0.39–1.53)	0.451
Hypertension	1.11 (0.91–1.36)	0.293	1.08 (0.68–1.71)	0.758
Dyslipidemia	1.52 (1.13–2.04)	0.006	2.31 (1.20–4.46)	0.013
	AIC = 7616		AIC = 1732	

Table 4. Multivariable Cox proportional hazard analyses for new onset of DM in tertiles of FLI.

	n	HR (95% CI)	P
Male subjects			
IFG (–)			
T1	2390	Reference	–
T2	2203	1.69 (1.06–2.68)	0.026
T3	1952	3.99 (2.63–6.04)	<0.001
IFG (+)			
T1	248	12.8 (7.74–21.2)	<0.001
T2	442	16.6 (10.8–25.6)	<0.001
T3	690	30.0 (20.2–44.6)	<0.001
Female subjects			
IFG (–)			
T1	1430	Reference	–
T2	1391	1.19 (0.60–2.39)	0.618
T3	1270	2.19 (1.16–4.14)	0.016
IFG (+)			
T1	28	9.51 (2.72–33.2)	<0.001
T2	64	12.9 (5.61–29.4)	<0.001
T3	182	25.7 (13.7–48.1)	<0.001

Table 5. Multivariable Cox proportional hazard regression analyses for new onset of DM in tertiles of FLI in the absence and presence of IFG. CI confidence interval, DM diabetes mellitus, FLI fatty liver index, HR hazard ratio, IFG impaired fasting glucose. IFG was defined as fasting plasma glucose ≥ 100 mg/dL. The model was adjusted for age, smoking habit, alcohol drinking habit, family history of DM, hypertension and dyslipidemia.

Fatty liver index. FLI was calculated using the following formula¹⁴: $FLI = [e^{(0.953 \times \ln(\text{triglycerides}) + 0.139 \times \text{BMI} + 0.718 \times \ln(\gamma\text{GTP}) + 0.053 \times \text{WC} - 15.745)}] / [1 + e^{(0.953 \times \ln(\text{triglycerides}) + 0.139 \times \text{BMI} + 0.718 \times \ln(\gamma\text{GTP}) + 0.053 \times \text{WC} - 15.745)}] \times 100$.

Statistical analysis. Numeric variables are expressed as means \pm SD for normally distributed parameters or medians (interquartile ranges) for skewedly distributed parameters. The distribution of each parameter was tested for its normality using the Shapiro–Wilk W test. Comparison between two groups was done with Mann–Whitney’s U test. Intergroup differences in demographic parameters were examined by the chi-square test. For detecting significant differences between data in multiple groups, one-way analysis of variance was used. The ability of FLI or FPG at baseline to predict new onset of DM was investigated using receiver operating characteristic (ROC) curves. The area under curve (AUC) was calculated, and cut-off values of FLI and FPG were obtained by the Youden index⁴⁹. The relationship between FLI and hazard ratio (HR) for the development of DM after adjustment of confounders including age, FPG, habits of alcohol drinking and current smoking, family history of DM and diagnosis of hypertension and dyslipidemia at baseline was analyzed by a multivariable Cox proportional hazard model with a restricted cubic spline. HRs and 95% confidence intervals in three subgroups according to tertiles of FLI level at baseline (T1–T3) in both males and females were calculated by adjustment of the covariates. HRs for new onset of DM among the T1–T3 groups in the absence and presence of IFG were also analyzed. A P value of <0.05 was considered statistically significant. All data were analyzed by using EZR⁵⁰

and R: A Language and Environment for Statistical Computing version 3.6.1 (R Core Team, R Foundation for Statistical Computing, Vienna, Austria, 2019, <https://www.R-project.org/>).

Data availability

The datasets analyzed during the current study are available from the corresponding author on reasonable request.

Received: 23 January 2021; Accepted: 8 June 2021

Published online: 18 June 2021

References

- Cosentino, F. *et al.* 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur. Heart J.* **41**, 255–323. <https://doi.org/10.1093/eurheartj/ehz486> (2020).
- Ogurtsova, K. *et al.* IDF Diabetes Atlas: global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes Res. Clin. Pract.* **128**, 40–50. <https://doi.org/10.1016/j.diabres.2017.03.024> (2017).
- Pertanen, J. *et al.* Natural history of peripheral neuropathy in patients with non-insulin-dependent diabetes mellitus. *N. Engl. J. Med.* **333**, 89–94. <https://doi.org/10.1056/NEJM199507133330203> (1995).
- Leon, B. M. & Maddox, T. M. Diabetes and cardiovascular disease: epidemiology, biological mechanisms, treatment recommendations and future research. *World J. Diabetes* **6**, 1246–1258. <https://doi.org/10.4239/wjcd.v6.i13.1246> (2015).
- Lonardo, A., Ballestri, S., Marchesini, G., Angulo, P. & Loria, P. Nonalcoholic fatty liver disease: a precursor of the metabolic syndrome. *Dig. Liver Dis.* **47**, 181–190. <https://doi.org/10.1016/j.dld.2014.09.020> (2015).
- Yang, K. C. *et al.* Association of non-alcoholic fatty liver disease with metabolic syndrome independently of central obesity and insulin resistance. *Sci. Rep.* **6**, 27034. <https://doi.org/10.1038/srep27034> (2016).
- Estes, C. *et al.* Modeling NAFLD disease burden in China, France, Germany, Italy, Japan, Spain, United Kingdom, and United States for the period 2016–2030. *J. Hepatol.* **69**, 896–904. <https://doi.org/10.1016/j.jhep.2018.05.036> (2018).
- Younossi, Z. M. *et al.* The economic and clinical burden of nonalcoholic fatty liver disease in the United States and Europe. *Hepatology* **64**, 1577–1586. <https://doi.org/10.1002/hep.28785> (2016).
- Chang, Y. *et al.* Cohort study of non-alcoholic fatty liver disease, NAFLD fibrosis score, and the risk of incident diabetes in a Korean population. *Am. J. Gastroenterol.* **108**, 1861–1868. <https://doi.org/10.1038/ajg.2013.349> (2013).
- Ballestri, S. *et al.* Nonalcoholic fatty liver disease is associated with an almost twofold increased risk of incident type 2 diabetes and metabolic syndrome. Evidence from a systematic review and meta-analysis. *J. Gastroenterol. Hepatol.* **31**, 936–944. <https://doi.org/10.1111/jgh.13264> (2016).
- Musso, G., Gambino, R., Cassader, M. & Pagano, G. Meta-analysis: natural history of non-alcoholic fatty liver disease (NAFLD) and diagnostic accuracy of non-invasive tests for liver disease severity. *Ann. Med.* **43**, 617–649. <https://doi.org/10.3109/07853890.2010.518623> (2011).
- Neuschwander-Tetri, B. A. & Caldwell, S. H. Nonalcoholic steatohepatitis: summary of an AASLD Single Topic Conference. *Hepatology* **37**, 1202–1219. <https://doi.org/10.1053/jhep.2003.50193> (2003).
- Papagianni, M., Sofogianni, A. & Tziomalos, K. Non-invasive methods for the diagnosis of nonalcoholic fatty liver disease. *World J. Hepatol.* **7**, 638–648. <https://doi.org/10.4254/wjh.v7.i4.638> (2015).
- Bedogni, G. *et al.* The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterol.* **6**, 33. <https://doi.org/10.1186/1471-230X-6-33> (2006).
- Otgonsuren, M. *et al.* Single non-invasive model to diagnose non-alcoholic fatty liver disease (NAFLD) and non-alcoholic steatohepatitis (NASH). *J. Gastroenterol. Hepatol.* **29**, 2006–2013. <https://doi.org/10.1111/jgh.12665> (2014).
- Yang, B. L. *et al.* External validation of fatty liver index for identifying ultrasonographic fatty liver in a large-scale cross-sectional study in Taiwan. *PLoS ONE* **10**, e0120443. <https://doi.org/10.1371/journal.pone.0120443> (2015).
- Huang, X. *et al.* Validation of the Fatty Liver Index for nonalcoholic fatty liver disease in middle-aged and elderly Chinese. *Medicine (Baltimore)* **94**, e1682. <https://doi.org/10.1097/MD.0000000000001682> (2015).
- Balkau, B. *et al.* Nine-year incident diabetes is predicted by fatty liver indices: the French D.E.S.I.R. study. *BMC Gastroenterol.* **10**, 56. <https://doi.org/10.1186/1471-230X-10-56> (2010).
- Jung, C. H. *et al.* Assessment of the fatty liver index as an indicator of hepatic steatosis for predicting incident diabetes independently of insulin resistance in a Korean population. *Diabet. Med.* **30**, 428–435. <https://doi.org/10.1111/dme.12104> (2013).
- Yadav, D. *et al.* Fatty liver index as a simple predictor of incident diabetes from the KoGES-ARIRANG study. *Medicine (Baltimore)* **95**, e4447. <https://doi.org/10.1097/MD.0000000000004447> (2016).
- Jager, S. *et al.* Association between the Fatty Liver Index and risk of type 2 diabetes in the EPIC-Potsdam Study. *PLoS ONE* **10**, e0124749. <https://doi.org/10.1371/journal.pone.0124749> (2015).
- Franch-Nadal, J. *et al.* Fatty liver index is a predictor of incident diabetes in patients with prediabetes: the PREDAPS study. *PLoS ONE* **13**, e0198327. <https://doi.org/10.1371/journal.pone.0198327> (2018).
- Wargny, M. *et al.* Fatty liver index is a strong predictor of changes in glycemic status in people with prediabetes: the IT-DIAB study. *PLoS ONE* **14**, e0221524. <https://doi.org/10.1371/journal.pone.0221524> (2019).
- Olubamwo, O. O., Virtanen, J. K., Pihlajamaki, J. & Tuomainen, T. P. Association of fatty liver index with risk of incident type 2 diabetes by metabolic syndrome status in an Eastern Finland male cohort: a prospective study. *BMJ Open* **9**, e026949. <https://doi.org/10.1136/bmjopen-2018-026949> (2019).
- Hirata, A. *et al.* Fatty liver index predicts incident diabetes in a Japanese general population with and without impaired fasting glucose. *Hepatol. Res.* **48**, 708–716. <https://doi.org/10.1111/hepr.13065> (2018).
- Khang, A. R., Lee, H. W., Yi, D., Kang, Y. H. & Son, S. M. The fatty liver index, a simple and useful predictor of metabolic syndrome: examination of the Korea National Health and Nutrition Examination Survey 2010–2011. *Diabetes Metab. Syndr. Obes.* **12**, 181–190. <https://doi.org/10.2147/DMSO.S189544> (2019).
- Examination Committee of Criteria for “Obesity Disease” in J. & Japan Society for the Study of, O. New criteria for “obesity disease” in Japan. *Circ. J.* **66**, 987–992. <https://doi.org/10.1253/circj.66.987> (2002).
- Satoh, H., Kishi, R. & Tsutsui, H. Body mass index can similarly predict the presence of multiple cardiovascular risk factors in middle-aged Japanese subjects as waist circumference. *Intern. Med.* **49**, 977–982. <https://doi.org/10.2169/internalmedicine.49.3006> (2010).
- Nakamura, K. *et al.* Optimal cutoff values of waist circumference and the discriminatory performance of other anthropometric indices to detect the clustering of cardiovascular risk factors for metabolic syndrome in Japanese men and women. *Environ. Health Prev. Med.* **16**, 52–60. <https://doi.org/10.1007/s12199-010-0165-y> (2011).
- Unno, M. *et al.* The utility of visceral fat level by bioelectrical impedance analysis in the screening of metabolic syndrome—the results of the Kyushu and Okinawa Population Study (KOPS). *J. Atheroscler. Thromb* **19**, 462–470. <https://doi.org/10.5551/jat.11528> (2012).

31. Kalia, H. S. & Gaglio, P. J. The prevalence and pathobiology of nonalcoholic fatty liver disease in patients of different races or ethnicities. *Clin. Liver Dis.* **20**, 215–224. <https://doi.org/10.1016/j.cld.2015.10.005> (2016).
32. Valenti, L., Bugianesi, E., Pajvani, U. & Targher, G. Nonalcoholic fatty liver disease: cause or consequence of type 2 diabetes?. *Liver Int.* **36**, 1563–1579. <https://doi.org/10.1111/liv.13185> (2016).
33. Seppala-Lindroos, A. *et al.* Fat accumulation in the liver is associated with defects in insulin suppression of glucose production and serum free fatty acids independent of obesity in normal men. *J. Clin. Endocrinol. Metab.* **87**, 3023–3028. <https://doi.org/10.1210/jcem.87.7.8638> (2002).
34. Gastaldelli, A. *et al.* Fatty liver is associated with insulin resistance, risk of coronary heart disease, and early atherosclerosis in a large European population. *Hepatology* **49**, 1537–1544. <https://doi.org/10.1002/hep.22845> (2009).
35. Pal, D. *et al.* Fetuin-A acts as an endogenous ligand of TLR4 to promote lipid-induced insulin resistance. *Nat. Med.* **18**, 1279–1285. <https://doi.org/10.1038/nm.2851> (2012).
36. Coskun, T. *et al.* Fibroblast growth factor 21 corrects obesity in mice. *Endocrinology* **149**, 6018–6027. <https://doi.org/10.1210/en.2008-0816> (2008).
37. Xu, J. *et al.* Fibroblast growth factor 21 reverses hepatic steatosis, increases energy expenditure, and improves insulin sensitivity in diet-induced obese mice. *Diabetes* **58**, 250–259. <https://doi.org/10.2337/db08-0392> (2009).
38. Misu, H. *et al.* A liver-derived secretory protein, selenoprotein P, causes insulin resistance. *Cell Metab.* **12**, 483–495. <https://doi.org/10.1016/j.cmet.2010.09.015> (2010).
39. Furuhashi, M. *et al.* Plasma xanthine oxidoreductase activity as a novel biomarker of metabolic disorders in a general population. *Circ. J.* **82**, 1892–1899. <https://doi.org/10.1253/circj.CJ-18-0082> (2018).
40. Furuhashi, M. *et al.* Annual change in plasma xanthine oxidoreductase activity is associated with changes in liver enzymes and body weight. *Endocr. J.* **66**, 777–786. <https://doi.org/10.1507/endocrj.EJ19-0053> (2019).
41. Furuhashi, M. New insights into purine metabolism in metabolic diseases: role of xanthine oxidoreductase activity. *Am. J. Physiol. Endocrinol. Metab.* **319**, E827–E834. <https://doi.org/10.1152/ajpendo.00378.2020> (2020).
42. Koehler, E. M. *et al.* External validation of the fatty liver index for identifying nonalcoholic fatty liver disease in a population-based study. *Clin. Gastroenterol. Hepatol.* **11**, 1201–1204. <https://doi.org/10.1016/j.cgh.2012.12.031> (2013).
43. Tanaka, J. *et al.* Sex- and age-specific carriers of hepatitis B and C viruses in Japan estimated by the prevalence in the 3,485,648 first-time blood donors during 1995–2000. *Intervirology* **47**, 32–40. <https://doi.org/10.1159/000076640> (2004).
44. Tanaka, J. *et al.* Total numbers of undiagnosed carriers of hepatitis C and B viruses in Japan estimated by age- and area-specific prevalence on the national scale. *Intervirology* **54**, 185–195. <https://doi.org/10.1159/000324525> (2011).
45. Higashiura, Y. *et al.* Low urine pH predicts new onset of diabetes mellitus during a 10-year period in men: BOREAS-DM1 study. *J. Diabetes Invest.* <https://doi.org/10.1111/jdi.13284> (2020).
46. Mori, K. *et al.* U-shaped relationship between serum uric acid level and decline in renal function during a 10-year period in female subjects: BOREAS-CKD2. *Hypertens. Res.* <https://doi.org/10.1038/s41440-020-0532-z> (2020).
47. Matsuo, S. *et al.* Revised equations for estimated GFR from serum creatinine in Japan. *Am. J. Kidney Dis.* **53**, 982–992. <https://doi.org/10.1053/j.ajkd.2008.12.034> (2009).
48. American Diabetes Association 2. Classification and diagnosis of diabetes. *Diabetes Care* **40**, S11–S24. <https://doi.org/10.2337/dc17-S005> (2017).
49. Schisterman, E. F., Perkins, N. J., Liu, A. & Bondell, H. Optimal cut-point and its corresponding Youden Index to discriminate individuals using pooled blood samples. *Epidemiology* **16**, 73–81. <https://doi.org/10.1097/01.ede.0000147512.81966.ba> (2005).
50. Kanda, Y. Investigation of the freely available easy-to-use software “EZ” for medical statistics. *Bone Marrow Transplant* **48**, 452–458. <https://doi.org/10.1038/bmt.2012.244> (2013).

Acknowledgements

This study was supported by Grants for Education and Research 2018 and 2019 from Sapporo Medical University.

Author contributions

Y.H., M.F., and M.T. designed the study, performed data analyses and wrote the paper. S.T. and N.M. performed data analyses. K.N., T.H., and N.H. performed data collection. M.K. and H.O. performed the statistical analyses. K.T. and T.M. supervised the analyses. All authors approved the final version of manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-92292-y>.

Correspondence and requests for materials should be addressed to M.F.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021