

REVIEW

Techniques and Emerging Trends in Flavor and Taste Development in Meat

Md Jakir Hossain^{1,†}, AMM Nurul Alam^{1,†}, So-Hee Kim¹, Chan-Jin Kim¹,
Seon-Tea Joo^{1,2}, and Young-Hwa Hwang^{2,*}

¹Division of Applied Life Science (BK 21 Four), Gyeongsang National University, Jinju 52828, Korea

²Institute of Agriculture & Life Science, Gyeongsang National University, Jinju 52828, Korea

 OPEN ACCESS

Received November 8, 2024

Revised December 3, 2024

Accepted December 3, 2024

*Corresponding author : Young-Hwa Hwang
Institute of Agriculture & Life Science,
Gyeongsang National University,
Jinju 52828, Korea
Tel: +82-55-772-1943
Fax: +82-55-772-1949
E-mail: philoria@gnu.ac.kr

*ORCID
Md Jakir Hossain
<https://orcid.org/0009-0008-7663-9202>
AMM Nurul Alam
<https://orcid.org/0000-0003-3153-3718>
So-Hee Kim
<https://orcid.org/0000-0003-3966-6160>
Chan-Jin Kim
<https://orcid.org/0000-0001-5020-6873>
Seon-Tea Joo
<https://orcid.org/0000-0002-5483-2828>
Young-Hwa Hwang
<https://orcid.org/0000-0003-3687-3535>

† These authors contributed equally to this work.

Abstract Flavor and taste are critical factors influencing consumer attraction for meat, shaping preferences and commercial demand. This review examines conventional and novel approaches to flavor and taste creation in the meat business, highlighting ways that improve sensory profiles and meet consumer demands. Conventional methods, such as aging and marination, are analyzed in conjunction with new technologies, including enzymatic treatment, fermentation, genetic treatments to alter texture and enhance umami. This study also emphasizes innovative methods to improve flavor of plant-based meat products, designed to meet the increasing demand for healthier, sustainable, and customizable meat products. The paper examines various methodologies and trends, offering a thorough grasp of flavor creation in the meat sector and highlighting the potential of creative approaches to transform meat flavor and taste profiles in response to evolving consumer and industry demands.

Keywords flavor and taste, plant based meat, genetic treatments, umami

Introduction

The sensory experience of meat intake is significantly affected by its flavor and taste, which are essential for consumer acceptance and market value. As customer preferences shift, influenced by heightened awareness of health, sustainability, and culinary diversity, the meat business encounters a rising demand for superior flavor development (Hossain et al., 2024; Zhang et al., 2021). The intricacy of meat flavor results from the interplay of several substances, including amino acids, peptides, lipids, and volatiles, which are affected by variables such as animal species, food, and processing techniques (Jiang et al., 2024). Understanding these complex interactions, researchers and industry experts have pursued novel methods to improve and alter the flavor profiles of meat products to satisfy contemporary customer expectations. Traditional methods for flavor development in meat include dry and wet aging, marination, and smoking (Alam et al.,

2024d). These traditional methods facilitate biological reactions that enhance the umami, sweetness, and overall complexity of meat's flavor. Nonetheless, recent developments in food technology are expanding the limits of flavor enhancement by offering innovative methods that utilize enzymatic activity, microbial fermentation, and improved flavor encapsulation technologies (Wang et al., 2022). Microbial transglutaminase (MTGase) has become a favored binding agent, enhancing texture and affecting flavor release in meat products (Chen et al., 2023).

The integration of artificial intelligence (AI) and big data analytics in the meat business is revolutionizing taste formulation methodologies. Data-driven models currently aid in forecasting consumer preferences, enhancing component interactions, and formulating tailored flavor profiles (Al-Ali et al., 2024). The integration of classic and contemporary techniques offers promising prospects to enhance and vary meat flavor profiles, improving both palatability and market attractiveness of meat products.

This review will examine contemporary techniques and upcoming trends in flavor and taste production in meat, highlighting developments in both classic and creative methodologies. This research seeks to elucidate how contemporary literature and industry practices are influencing the future of meat taste engineering in response to evolving customer needs.

Exploring the Biochemical Mechanisms of Flavor and Taste in Meat

Flavor and taste characteristics result from a complex interaction of chemical substances and sensory mechanisms, which collectively enhance a product's overall palatability. Taste, as conventionally defined, relates to the five fundamental gustatory sensations: sweet, sour, salty, bitter, and umami. Flavor is a composite of taste, scent, and supplementary sensations, including texture, mouthfeel, and visual cues, resulting in a multimodal experience that influences consumer satisfaction across various sensory dimensions (Shahidi and Zhong et al., 2005; Smith et al., 2018).

The flavor profile of meat is shaped by intrinsic characteristics, including species, age, and muscle type, alongside extrinsic ones such as diet, cooking style, and aging processes (Mottram, 1998). Principal factors influencing meat flavor encompass the Maillard reaction, lipid oxidation, and nucleotide degradation during cooking. The Maillard reaction, occurring between amino acids and reducing sugars at elevated temperatures, generates various volatile chemicals that provide cooked meat with its distinctive roasted and caramelized aromas (Nursten, 2005). Lipid oxidation is a crucial process that enhances flavor by producing aldehydes, ketones, and hydrocarbons, imparting meat its distinctive scent and sensory complexity (Mottram, 1998).

Fifth Fundamental Tastes “Umami”

Among the five fundamental tastes, umami is especially important in the flavor of meat. The umami flavor, mostly originating from amino acids like glutamate and nucleotides such as inosinate, contributes to the richness and complexity characteristic of premium meat products (Kurihara, 2009). Research indicates that aging processes, including wet and dry aging, might augment umami by decomposing proteins into glutamate and other umami-rich substances, hence enhancing the savory profile and overall sensory attractiveness of beef products (Ikeda, 2002; Son et al., 2024). Dry aging is recognized for augmenting the flavor of beef, yielding nutty and earthy undertones as a result of the enzymatic degradation of amino acids and lipids (Dashdorj et al., 2015). The ways of cooking significantly influence taste development. Grilling enhances the Maillard reaction and lipid oxidation, resulting in more pronounced flavors, while boiling or steaming produces a milder, less intricate flavor due to the restricted development of volatile chemicals (Zhong et al., 2021).

Numerous components, such as proteins, lipids, carbohydrates, and nucleotides, substantially influence the distinctive flavors experienced in meat. The influence of different compounds in the taste and flavor of food and meat are shown in Fig. 1. Meat is rich in proteins, and their decomposition during cooking or age liberates amino acids that are crucial for flavor enhancement. Amino acids, including glutamic acid, aspartic acid, and glycine, contribute to the umami flavor, whereas sulfur-containing amino acids such as methionine and cysteine produce appealing roasted or meaty odors upon heating (Herrera and Calkins, 2022; Mottram, 1998). Moreover, the Maillard reaction transpires between amino acids and reducing sugars during cooking, resulting in the creation of heterocyclic compounds that enhance the meat's browned exterior and intricate flavor. The oxidation of polyunsaturated fatty acids (PUFAs) during cooking generates chemicals including aldehydes, ketones, and alcohols, which are crucial for developing distinct meat tastes. For example, oleic acid produces agreeable fragrances, whereas linoleic and linolenic acids impart aromatic characteristics reminiscent of roasted meat (Resconi et al., 2013). Intramuscular fat, or marbling, is recognized for augmenting flavor intensity and juiciness, since it influences the retention and release of volatile chemicals during cooking (Wood, 2004). Despite their minimal presence, carbohydrates in meat, including glycogen, glucose, and ribose, significantly contribute to flavor production. The caramelization of sugars at high temperatures enhances the rich, sweet flavors of cooked meat, improving its overall appeal (Mottram, 1998).

Nucleotides like inosine monophosphate and guanosine monophosphate are essential elements in the umami flavor characteristic of meat. These molecules, when amalgamated with amino acids such as glutamate, synergistically augment the




Fig. 1. The influence of different compounds in the taste and flavor of food and meat. IMP, inosine monophosphate; GMP, guanosine monophosphate; AMP, adenosine monophosphate.

umami flavor and complexity in meat (Tikk, 2008). Free amino acids and peptides contribute to flavor; notably, glutamic acid enhances the delicious profile of aged meat (Dashdorj et al., 2015).

Umami, one of the five fundamental sensations with sweet, sour, salty, and bitter, is sometimes described as the “savory” or “meaty” flavor. Initially recognized by Japanese chemist Kikunae Ikeda in 1908, umami arises from the presence of glutamate, inosinate, and guanylate, chemicals prevalent in numerous foods, particularly meats. This flavor elevates the culinary profile of dishes, delivering a sumptuous experience and an enduring aftertaste. The umami flavor predominantly originates from amino acids, particularly glutamate, and nucleotides, including inosinate and guanylate, which are naturally found in meat (Ikeda, 2002). The umami and other taste perception mechanism is illustrated in Fig. 2. Meat processing techniques such as curing and boiling can augment umami levels. With aging, proteolytic enzymes degrade muscle proteins, liberating free amino acids and nucleotides that enhance umami flavor. In goods like restructured meat or meat replacements, using umami sources (e.g., mushroom extracts, yeast extracts, or soy sauce) enhances sensory perception and improves palatability, hence increasing their appeal to consumers seeking meat-like experiences (Samad et al., 2024).

The Application of Sensory Science in the Investigation of Flavor Profiles in Meat

Progress in sensory research and analytical methodologies has facilitated a more accurate identification and comprehension of the chemical components that influence beef flavor (Alam et al., 2024a). Gas chromatography-mass spectrometry (GC-MS) enables researchers to identify and quantify volatile molecules, elucidating the chemical foundation of meat flavors. Furthermore, consumer sensory panels and descriptive sensory analysis facilitate the correlation of specific flavor traits with consumer preferences, enabling the meat sector to customize goods to satisfy consumer wants (Lawless and Heymann, 2010).

Influence of Processing Methods on the Chemical Composition of Food Matrixes

The chemical composition of meat and its resultant flavor are influenced by processing processes. Dry aging facilitates the enzymatic degradation of proteins and fats, hence augmenting the content of amino acids, peptides, and fatty acids that enhance umami and savory flavors (Wang et al., 2020). Cooking methods such as grilling, smoking, and frying induce thermal




Fig. 2. The umami and other taste perception mechanism. Adapted from Schmidt and Mouritsen (2022) with CC-BY-NC-ND. GMP, guanosine monophosphate; IMP, inosine monophosphate.

breakdown of lipids and proteins, resulting in the creation of distinctive volatile compounds and enhancing meat flavor (Resconi et al., 2013).

The Maillard reaction is a primary mechanism of taste production during heat processing. This reaction transpires between reducing sugars and amino acids, resulting in a sequence of intricate reactions that yield various chemicals, including melanoidins, which impart the distinctive brown hue and umami flavors in meat. The Maillard reaction is significantly influenced by temperature, generally taking place at temperatures over 140°C (Lea and Swoboda, 1962). Research indicates that elevated temperatures result in a greater formation of Maillard reaction products, hence intensifying flavor and color in restructured meat (Samad et al., 2024).

The chemicals generated during the Maillard reaction differ based on the amino acids and sugars involved. Cysteine and ribose generate sulfur-containing compounds, including mercaptans and thiazoles, which provide savory and meaty flavors, whereas phenylalanine yields molecules with a roasted and nutty fragrance (van Boekel, 2006). These reactions are controlled in restructured meat processing to maximize flavor profiles and augment customer attraction.

Lipid oxidation is a significant element in the flavor characteristic of thermally cooked meat. Upon heating, unsaturated fats experience oxidation, resulting in the production of aldehydes, ketones, and alcohols, which enhance the aroma and flavor of the product (Shahidi and Zhong, 2005). Lipid oxidation can impart appealing aromas, such the distinctive “grilled” scent; nevertheless, excessive oxidation may lead to undesirable off-flavors that diminish the overall sensory quality of restructured meat.

Diverse fat sources and compositions markedly influence flavor results. Products rich in polyunsaturated fats typically exhibit more pronounced oxidation products, which can augment flavor complexity while also heightening the risk of rancidity. Consequently, meticulous selection of fats and regulated thermal processing are crucial in the manufacture of restructured meat to attain a harmonious flavor profile devoid of undesirable notes (Ren et al., 2024).

Gamma irradiation proved to enhance the flavor and taste characteristics by reducing the oxidative changes in meat (Sadakuzzaman et al., 2024). Thermal processing induces protein denaturation and amino acid degradation, which facilitate the formation of distinctive taste molecules. During protein denaturation, free amino acids are released, which then engage in the Maillard reaction or undergo Strecker degradation, resulting in the formation of new taste compounds, such as pyrazines and pyridines, so enhancing the meat’s flavor complexity (Xu and Yin, 2024). Heating elevates the release of glutamic acid, hence intensifying umami qualities in meat products (Yoshida and Ninomiya, 2023). The processing temperature and duration directly influence the degree of protein breakdown. Elevated temperatures and extended cooking durations augment protein degradation and amino acid liberation, hence intensifying umami qualities in restructured beef products. Enhancing these factors can consequently elevate the flavor profile and consumer approval of the final product.

Diverse thermal processing techniques, including roasting, grilling, and steaming, influence flavor development distinctively due to differing heat transmission mechanisms. Grilling delivers direct high heat, facilitating Maillard browning and lipid oxidation, which enhances the flavor profile. Conversely, steaming employs lower temperatures and moist heat, resulting in gentler aromas, preserving more subtle notes, and diminishing lipid oxidation (Parry, 2016).

Research comparing these techniques has indicated that whereas dry-heat procedures enhance flavor, they may also result in moisture loss, impacting texture and juiciness, which are essential for restructured meat products (Lawrie and Ledward, 2014). The selection of thermal processing method significantly influences the equilibrium between taste intensity and textural quality, affecting consumer preference and product satisfaction.

Impact of Animal Diet on Meat Flavor

The food of animals directly influences the accumulation of intramuscular fat and the composition of fatty acids, both of which are essential in the development of distinct flavors during cooking. Grass-fed and grain-fed animals exhibit distinct lipid profiles, with grass-fed diets yielding elevated concentrations of omega-3 fatty acids and conjugated linoleic acid (Rahman et al., 2009). These chemicals impart a unique grassy and gamey flavor typically associated with grass-fed meats, in contrast to the sweeter and buttery flavor of grain-fed meat (Jiang, 2011).

Certain dietary supplements, such as antioxidants (e.g., Vitamin E), can enhance oxidative stability, hence avoiding flavor degradation linked to lipid oxidation. Research indicates that administering specific natural antioxidants, such as rosemary, to cattle may improve the stability of meat flavor during storage (Rashidaie Abandansarie et al., 2019). Cattle provided with selenium-enriched diets have enhanced flavor profiles attributable to selenium's antioxidant properties, which mitigate off-flavors resulting from oxidation (Shin et al., 2021).

Genetic Influences on Meat Flavor

Genetics fundamentally determine muscle fiber types, fat deposition, and enzyme activity, all of which affect meat flavor. Various breeds demonstrate distinct flavor profiles influenced by hereditary characteristics. Wagyu beef is esteemed for its pronounced marbling, yielding a delicate texture and distinctive umami flavor profile (Smith et al., 2018). Iberian pork possesses a unique flavor, resulting from its genetics and a specialized acorn-rich diet during the finishing phase. Genetic factors influence the distribution of muscle fiber types, hence impacting meat flavor. Fast-twitch fibers (Type II) are associated with flavors related to lactate buildup, whereas slow-twitch fibers (Type I) possess a higher concentration of oxidative enzymes, resulting in a unique flavor profile (Purslow, 2022). Research indicates that the heritability of meat flavor is moderate, suggesting the feasibility of selective breeding for flavor-specific characteristics (Smith et al., 2003).

Comprehensive Sensory and Instrumental Approaches for Meat Flavor and Taste Assessment

Aging is a post-mortem process wherein meat experiences biochemical alterations that improve its flavor and softness. It is chiefly categorized into wet aging and dry aging, each conferring unique flavor profiles.

Wet aging entails vacuum-sealing the meat to preserve moisture, so facilitating the activity of proteolytic enzymes, such as calpains and cathepsins, which decompose muscle proteins into smaller, tasty peptides. Wet-aged meat possesses a subdued, more metallic flavor in contrast to dry-aged meat (Son et al., 2024). Dry aging involves storing meat at regulated temperatures and humidity levels without packaging, facilitating moisture evaporation, flavor concentration, and microbial activity that contributes nutty, earthy, and intricate umami characteristics. The Maillard process on the surface of the meat enhances the flavor profile (Kim et al., 2016). The length of the aging process influences taste strength, since prolonged aging provides additional time for enzymatic and microbiological activities, potentially resulting in complex flavors while also posing a spoiling risk if not well managed (Marcus, 2019).

The sensory quality of meat is a crucial determinant of consumer satisfaction. Attributes like flavor, texture, and juiciness drive consumer preferences and purchasing decisions. Meat flavor is a multi-faceted attribute, influenced by amino acids, peptides, and fat content, while the umami taste in particular enhances palatability (Kerth et al., 2024). This section discusses sensory evaluation methods to assess flavor and taste accurately, along with advanced instrumental techniques.

Descriptive analysis involves training panelists to characterize and quantify specific flavor attributes. It offers in-depth profiling and is widely used for understanding meat flavor (Lawless and Heymann, 2010). Techniques include flavor profile analysis uses trained panelists to evaluate attributes like “meaty,” “brothy,” or “metallic.” It provides a structured approach to understanding flavor nuances (Adam, 2021). Quantitative descriptive analysis applies statistical methods to quantify flavor intensities. It’s flexible for comparing different meat samples or treatments, such as aging or cooking methods (De Pilli et al., 2024). Spectrum™ Analysis is a highly standardized method enables cross-laboratory comparisons by using scales based on reference standards. It is particularly useful for research settings requiring consistency across sensory studies. Hedonic Rating is a widely used method, where participants rate samples on a scale (e.g., 9-point scale from “dislike extremely” to “like extremely”). This quick and simple test provides a direct measure of consumer acceptance (Lawless and Heymann, 2010). In Paired Preference Testing, consumers compare two samples to determine which they prefer, allowing direct comparisons between treatments like different binders or levels of additives (Meilgaard et al., 1999).

Instrumental techniques complement sensory analysis by providing objective measurements of flavor compounds in meat. GC-MS is a standard technique to analyze volatile flavor compounds (Shahidi and Cadwallader, 1997). It separates, identifies, and quantifies compounds contributing to meat aroma, like aldehydes and ketones. Electronic Nose (E-Nose) is sensor-based device mimics the human sense of smell to detect and differentiate flavor compounds (Peris and Escuder-Gilabert, 2009). It’s increasingly used in meat quality assessment due to its efficiency and non-destructive nature. Liquid chromatography-mass spectrometry is ideal for analyzing non-volatile compounds responsible for taste, such as amino acids and nucleotides that contribute to umami and savory tastes in meat (Koutsidis et al., 2009).

Traditional Techniques for Flavor Development in Meat

The formation of flavor in meat is a complex process affected by elements including animal species, muscle type, and cooking techniques. Conventional methods emphasize altering the inherent properties of meat to improve its flavor profile via aging, marination, fermentation, and curing. These procedures have been employed for ages, cultivating unique flavors desired in culinary practices. Fig. 3 illustrates the advantages and disadvantages of traditional flavor development techniques.




Fig. 3. Advantages and disadvantages of traditional flavor development techniques.

Aging, including dry and wet aging, is a prevalent method in the meat business to augment flavor. Dry aging entails the preservation of beef under regulated humidity and temperature for many weeks, facilitating the action of natural enzymes that decompose muscle fibers, resulting in a more intense, flavorful, and tender product (Dashdorj et al., 2016). In wet aging, meat is vacuum-sealed and refrigerated, permitting it to mature in its own juices. Wet aging often results in a more subdued flavor than dry age, as the meat preserves moisture; yet, enzyme activity continues to enhance a complex flavor profile (Kim et al., 2016).

Marination is a widely employed method to enhance the flavor, softness, and juiciness of meat. Marinades often comprise components such as acids (e.g., vinegar, citrus), oils, and seasonings. Acidic substances decompose proteins, resulting in a more tender product, while aromatic herbs and spices contribute supplementary flavors (Aminzare et al., 2016). An overview of using additives in the quality, taste and flavor development of meat and meat products are summarized in Table 1. Marinades function as vehicles for taste chemicals that engage with the meat, facilitating deeper flavor penetration into the muscle fibers. Moreover, the inclusion of sugar or salt in marinades might facilitate Maillard reactions during cooking,

Table 1. Overview of using additives in the quality, taste and flavor development of meat and meat products

Product type	Objectives	Natural additives	Additive concentration (%) in formulation	Salt type	Salt concentration (%)	Main effects	References
Pork ham	Flavor improvement	Seaweed extract	0.11	NaCl	1	Flavor improved	Barbieri et al. (2016)
Dry aged beef jerkey	Flavor and tenderness.	Soy sauce	2	NaCl	2	Flavor and tenderness improved	Lim et al. (2014)
Chicken nugget	Tenderness	Mushroom	2.5	NaCl	25	Tenderness improved	Akesowan (2021)
Beef patty	Flavor and lipid oxidation	Mushroom	2.5	NaCl	0.60%	Reduced lipid oxidation and improved flavor	Cerón-Guevara et al. (2020)
Dry-cured pork	Flavor	Glasswort extract	10	NaCl	90	Flavor improved	Ferreira et al. (2022)
	Tenderness and juiciness	Glasswort extract	10%	NaCl	1	Tenderness and juiciness improved	Lim et al. (2015)
Pork loin	Tenderness and juiciness	Red glasswort	2	NaCl	2	Tenderness and juiciness improved	Jeong et al. (2020)
Synthetic additives							
Fermented sausages	Tenderness	Calcium glutamate	0.03	NaCl, KCl, Calcium Ascorbate	2.25+0.42+0.3	Tenderness improved	Zhang et al. (2021)
	Flavor	K-lactate	10	NaCl, KCl	50+40	Flavor improved	Guàrdia et al. (2008)
Pork frankfurter	Texture and flavor	Glycine	20	NaCl, KCl	22.5+57.50	Texture and flavor improved	Wilailux et al. (2020)
Pork sausage	Texture	Maltodextrin+L-Lysine+L-Alanine, Citric acid, Ca-lactate	3.5+4+1+0.5+1%	NaCl, KCl	70+20	Texture improved	Chen et al. (2019)

resulting in caramelized tastes and enhanced coloration on the meat's surface (Latoch et al., 2023).

Fermentation is extensively employed in the production of cured foods such as sausages and salamis. This procedure entails the incorporation of particular bacterial cultures, such as *Lactobacillus* and *Staphylococcus*, into meat, facilitating lactic acid synthesis and reducing the pH level. The acidic environment preserves the meat and increases flavor through intricate biological interactions that release volatile molecules responsible for a tangy and umami-rich character (Lorenzo et al., 2018). Fermentation produces supplementary aromatic chemicals such as aldehydes and esters, enhancing the distinctive aromas of fermented meat products (Hugas and Monfort, 1997).

Smoking is a conventional method that imparts a unique, smoky flavor and preserves the meat. This approach involves exposing meat to smoke generated from wood combustion, which infuses it with phenols, carbonyls, and organic acids that provide aroma and antibacterial characteristics (Pintado et al., 2021). Various woods, including hickory, apple, and cherry, contribute various flavors, as each wood type emits a specific array of aromatic chemicals. Smoking is employed in goods such as ham, bacon, and smoked sausages, where the smoky scent and flavor are much valued (Aaslyng and Meinert, 2017).

Curing entails the application of salt, nitrates, or nitrites to meat, which suppresses bacterial proliferation and facilitates the emergence of distinctive flavors. Salt extracts moisture, intensifying the meat's flavor and fostering conditions for biological reactions that generate distinct taste and aroma profiles (Toldrá et al., 2012). Nitrate and nitrite curing is crucial for color stability in cured meats, imparting the distinctive pink hue and inhibiting oxidative rancidity, which can adversely affect flavor (Desmond, 2006). This process is crucial in the production of hams, prosciuttos, and other cured meats, where flavor intensity is a defining characteristic.

Biotechnological Methods for Flavor Enhancement

Biotechnology involves several techniques that alter biological creatures or systems to generate or improve flavors. These methods may encompass fermentation, metabolic engineering, and enzymatic procedures to generate preferred flavor molecules in food. The cutting-edge biotechnological methods are illustrated in Fig. 4.

Fermentation has been employed for centuries to enhance flavors in food and beverages. Recent advancements in microbial biotechnology have facilitated more precise fermentation procedures to enhance flavor qualities. Traditional fermentation methods utilizing lactic acid bacteria and yeast have been refined to produce flavors and fragrances characteristic of meat, dairy, or certain plant-based characteristics. Lactic acid bacteria, specifically *Lactobacillus* and *Pediococcus* species, can synthesize organic acids, esters, and alcohols that impart a meat-like flavor to plant-based products (Gänzle, 2015).

Metabolic engineering modifies the metabolic pathways of microorganisms to improve the synthesis of particular taste chemicals. In yeast, the metabolic pathways for ester synthesis can be modified to provide flavors akin to those present in cheese, wine, and other fermented goods. Genetically modifying *Saccharomyces cerevisiae*, a widely utilized yeast in taste manufacturing, to enhance the biosynthesis of isoamyl acetate or ethyl caproate can produce fruity and buttery flavors that are sought after in many plant-based dairy replacements (Ienczak et al., 2022).

Enzymes are crucial for catalyzing processes that generate taste chemicals. Biotechnology utilizes particular enzymes, such as lipoxigenase or peroxidase, to facilitate targeted reactions that produce desirable tastes. Lipoxigenase can transform linoleic acid into volatile chemicals linked to a "meaty" fragrance in plant-based meat. Improvements in enzyme immobilization and stabilization methods have significantly increased the commercial scalability of enzyme technology, facilitating reliable and efficient taste generation in food manufacturing (Hasan et al., 2006).




Fig. 4. The cutting-edge biotechnological methods for flavor development.

Genomics has elucidated the genetic foundations of flavor production, facilitating the identification and alteration of genes implicated in flavor pathways. By finding and manipulating these genes, scientists can genetically modify plants, yeast, and bacterial strains to yield various flavor profiles. Genomic research has pinpointed genes that govern the manufacturing of several taste chemicals in plants. In tomatoes, genes like LeHPL and LeCCD1 are recognized for their impact on volatile chemicals that enhance fruity and sweet aromas. Researchers can utilize CRISPR-Cas9 to accurately modify genes, so augmenting the natural tastes of plant components employed in food production (Tieman et al., 2017). Comparable research in soybeans has pinpointed genes that affect unwanted grassy or beany flavors, which can be suppressed or altered to yield a more neutral flavor profile appropriate for plant-based meat (Dixon, 2013).

Genetic alteration is a potent genomics instrument employed to augment flavor. By introducing or overexpressing particular flavor-producing genes, researchers can enhance the synthesis of desirable chemicals in microorganisms or plants. For instance, *S. cerevisiae* has been genetically modified to express flavor-related pathways from other species, such as *Pichia pastoris* or *Kluyveromyces lactis*, thereby augmenting its potential to generate intricate, dairy-like flavors in plant-based dairy substitutes (Steensels et al., 2014).

Epigenetic alterations entail altering the expression of flavor-associated genes without modifying the DNA sequence. This method can regulate gene expression to augment the synthesis of particular tastes in agricultural products. Methylation alterations in tomato plants have been linked to enhanced production of fruity smells (Guo et al., 2018). Epigenetic methodologies are

currently developing in flavor biotechnology but show potential for improving flavor profiles in diverse crops without direct genetic alteration.

Fundamental Sensory Attributes of Hybrid and Plant-Based Meat

Hybrid and plant-based meats must emulate essential sensory characteristics of conventional meat, including texture, flavor, appearance which are crucial determinant in consumer choice. In plant-based alternatives, attaining a meat-like texture frequently necessitates the use of binders, such as MTGase, which improves the structural integrity of plant proteins (Baugreet et al., 2018). Conventional meat flavors originate from amino acids and fatty acids, which are difficult to reproduce in plant-based formulations (Kumari et al., 2023). The incorporation of natural flavor enhancers and fermentation techniques can augment the meat-like flavor of alternative goods. Colorants or natural substances, such as beet juice or heme proteins, are employed to impart the distinctive red or pink hue in meat analogs (Kyriakopoulou et al., 2019). Use of scaffolding materials encapsulated with bioactive compounds may improve the taste and flavor of meat analogs (Alam et al., 2024c). Consumers are more inclined to embrace plant-based goods when their flavor closely resembles that of traditional meat (Elzerman et al., 2015). Sensory evaluation methods for any kinds of meat products has been illustrated in Fig. 5. A study conducted by Elzerman et al. (2015) revealed that consumers positively assessed the texture of plant-based meat analogs, while observing flavor discrepancies in comparison to traditional meat. In a separate study, Kyriakopoulou et al. (2019) investigated the impact of ingredient modification on the sensory attributes of restructured plant-based patties, observing that the incorporation of MTGase improved texture, resulting in increased chewiness and firmness. Baugreet et al. (2018) found that in consumer preference testing for hybrid meat products that integrate plant and animal proteins, those exhibiting enhanced juiciness and tenderness received better overall like scores. Alam et al. (2024b) assessed different type of plant ingredients in pork and beef products and hybrid products were well accepted by sensory panel in comparison to reference meat products. Additionally, the umami and richness values were higher in hybrid products. These studies indicate that hybrid products can attract consumers desiring meat substitutes that closely resemble the sensory characteristics of traditional meat.




Fig. 5. Sensory evaluation methods for any kinds of meat products.

Conclusion

The integration of traditional and contemporary approaches offers a viable foundation for augmenting the sensory attributes of meat. Although age, marination, and smoking are fundamental approaches, advancements in biotechnology and sensory research present novel potential for flavor enhancement. Enzymatic activity, microbial cultures, and data-driven insights are essential for creating customized flavor profiles that align with changing customer preferences. This collaborative strategy not only improves flavor but also corresponds with the market's transition towards sustainable and creative food solutions. Subsequent research ought to concentrate on enhancing these technologies to maximize flavor uniformity and user contentment.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was supported by a National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. 2020R1I1A2069379 & 2023R1A2C1004867).

Author Contributions

Conceptualization: Alam AN. Writing - original draft: Hossain MJ, Alam AN. Writing - review & editing: Hossain MJ, Alam AN, Kim SH, Kim CJ, Joo ST, Hwang YH.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Aaslyng MD, Meinert L. 2017. Meat flavour in pork and beef: From animal to meal. *Meat Sci* 132:112-117.
- Adam SSM. 2021. Sensory and instrumental methods of meat evaluation: A review. *Int J Food Sci Agric* 5:627-638.
- Akesowan A. 2021. Partial salt reduction in gluten-free chicken nugget extended with white button mushroom and quality development with eggplant flour. *J Food Sci Technol* 58:4738-4745.
- Al-Ali AR, Gupta R, Zualkernan I, Das SK. 2024. Role of IoT technologies in big data management systems: A review and Smart Grid case study. *Pervasive Mob Comput* 100:101905.
- Alam AN, Hashem MA, Matar AM, Ali MS, Monti JA, Hossain MJ, Yusuf HM, Mia N. 2024a. Cutting edge technologies for the evaluation of plant-based food and meat quality: A comprehensive review. *Meat Res* 4:1-12.
- Alam AN, Hossain MJ, Lee EY, Kim SH, Hwang YH, Joo ST. 2024b. Characterization of imitated hybrid cultured meat patty to compare with beef meat. *Food Sci Anim Resour* (in press). doi: 10.5851/kosfa.2024.e60
- Alam AN, Kim CJ, Kim SH, Kumari S, Lee EY, Hwang YH, Joo ST. 2024c. Scaffolding fundamentals and recent advances

- in sustainable scaffolding techniques for cultured meat development. *Food Res Int* 189:114549.
- Alam AN, Mia N, Monti JA, Hashem MA, Ali MS. 2024d. Enhancing the qualitative attributes of meat through processing and preservation techniques: A review. *Meat Res* 4:1-6.
- Aminzare M, Hashemi M, Hassanzadazar H, Hejazi J. 2016. The use of herbal extracts and essential oils as a potential antimicrobial in meat and meat products; a review. *J Hum Environ Health Promot* 1:63-74.
- Barbieri G, Barbieri G, Bergamaschi M, Francheschini M, Berizi E. 2016. Reduction of NaCl in cooked ham by modification of the cooking process and addition of seaweed extract (*Palmaria palmata*). *LWT-Food Sci Technol* 73:700-706.
- Baugreet S, Kerry JP, Allen P, Gallagher E, Hamill RM. 2018. Physicochemical characteristics of protein-enriched restructured beef steaks with phosphates, transglutaminase, and elasticised package forming. *J Food Qual* 2018:4737602.
- Cerón-Guevara MI, Rangel-Vargas E, Lorenzo JM, Bermúdez R, Pateiro M, Rodríguez JA, Sánchez-Ortega I, Santos EM. 2020. Reduction of salt and fat in frankfurter sausages by addition of *Agaricus bisporus* and *Pleurotus ostreatus* flour. *Foods* 9:760.
- Chen J, Hu Y, Wen R, Liu Q, Chen Q, Kong B. 2019. Effect of NaCl substitutes on the physical, microbial and sensory characteristics of Harbin dry sausage. *Meat Sci* 156:205-213.
- Chen Y, Wang Y, Liu R, Xiong S, Xu Y, Hu Y. 2023. Effects of microbial transglutaminase on the gelling property and *in vitro* digestibility of fish scale gelatin from grass carp. *Food Biosci* 53:102569.
- Dashdorj D, Amna T, Hwang I. 2015. Influence of specific taste-active components on meat flavor as affected by intrinsic and extrinsic factors: An overview. *Eur Food Res Technol* 241:157-171.
- Dashdorj D, Tripathi VK, Cho S, Kim Y, Hwang I. 2016. Dry aging of beef; review. *J Anim Sci Technol* 58:20.
- De Pilli T, Alessandrino O, Baiano A. 2024. Quantitative descriptive analysis as a strategic tool in research activities relating to innovative meat tenderisation technologies. *Heliyon* 10:e32618.
- Desmond E. 2006. Reducing salt: A challenge for the meat industry. *Meat Sci* 74:188-196.
- Dixon M. 2013. *Textbook on international law*. Oxford University Press, Oxford, UK.
- Elzerman JE, Hoek AC, van Boekel MJAS, Luning PA. 2015. Appropriateness, acceptance and sensory preferences based on visual information: A web-based survey on meat substitutes in a meal context. *Food Qual Prefer* 42:56-65.
- Ferreira I, Leite A, Vasconcelos L, Rodrigues S, Mateo J, Munekata PES, Teixeira A. 2022. Sodium reduction in traditional dry-cured pork belly using glasswort powder (*Salicornia herbacea*) as a partial NaCl replacer. *Foods* 11:3816.
- Gänzle MG. 2015. Lactic metabolism revisited: Metabolism of lactic acid bacteria in food fermentations and food spoilage. *Curr Opin Food Sci* 2:106-117.
- Guàrdia MD, Guerrero L, Gelabert J, Gou P, Arnau J. 2008. Sensory characterisation and consumer acceptability of small calibre fermented sausages with 50% substitution of NaCl by mixtures of KCl and potassium lactate. *Meat Sci* 80:1225-1230.
- Guo J, Zhou R, Ren X, Jia H, Hua L, Xu H, Lv X, Zhao J, Wei T. 2018. Effects of salicylic acid, epi-brassinolide and calcium on stress alleviation and Cd accumulation in tomato plants. *Ecotoxicol Environ Saf* 157:491-496.
- Hasan F, Shah AA, Hameed A. 2006. Industrial applications of microbial lipases. *Enzyme Microb Technol* 39:235-251.
- Herrera NJ, Calkins CR. 2022. Developments in meat flavor. In *New aspects of meat quality: From genes to ethics*. 2nd ed. Purslow P (ed). Woodhead, Sawston, UK. pp 195-235.
- Hossain MJ, Alam AMMN, Lee EY, Hwang YH, Joo ST. 2024. Umami characteristics and taste improvement mechanism of meat. *Food Sci Anim Resour* 44:515-532.

- Hugas M, Monfort JM. 1997. Bacterial starter cultures for meat fermentation. *Food Chem* 59:547-554.
- Ienczak JL, de Oliveira Pereira I, da Silveira JM. 2022. Utilization of *Saccharomyces cerevisiae* as a source of natural food additives. In *Natural additives in foods*. Valencia GA (ed). Springer, Cham, Switzerland. pp 185-214.
- Ikedo K. 2002. New seasonings. *Chem Senses* 27:847-849.
- Jeong TJ, Kim TK, Kim HW, Choi YS. 2020. Effects of red glasswort as sodium chloride substitute on the physicochemical properties of pork loin ham. *Asian-Australas J Anim Sci* 33:662-669.
- Jiang T. 2011. Palatability control points for grass-fed beef: Revealing the key compounds contributing to beef flavor or off-flavor. Ph.D. dissertation, Washington State Univ., Pullman, WA, USA.
- Jiang W, Yang X, Li L. 2024. Flavor of extruded meat analogs: A review on composition, influencing factors, and analytical techniques. *Curr Res Food Sci* 8:100747.
- Kerth CR, Legako JF, Woerner DR, Brooks JC, Lancaster JM, O'Quinn TG, Nair M, Miller RK. 2024. A current review of U.S. beef flavor I: Measuring beef flavor. *Meat Sci* 210:109437.
- Kim YHB, Kemp R, Samuelsson LM. 2016. Effects of dry-aging on meat quality attributes and metabolite profiles of beef loins. *Meat Sci* 111:168-176.
- Koutsidis G, Simons SPJ, Thong YH, Haldoupis Y, Mojica-Lazaro J, Wedzicha BL, Mottram DS. 2009. Investigations on the effect of amino acids on acrylamide, pyrazines, and Michael addition products in model systems. *J Agric Food Chem* 57:9011-9015.
- Kumari S, Alam AN, Hossain MJ, Lee EY, Hwang YH, Joo ST. 2023. Sensory evaluation of plant-based meat: Bridging the gap with animal meat, challenges and future prospects. *Foods* 13:108.
- Kurihara Y, Matsui A, Hanada K, Kawashima M, Ishida J, Morosawa T, Tanaka M, Kaminuma E, Mochizuki Y, Matsushima A, Toyoda T, Shinozaki K, Seki M. 2009. Genome-wide suppression of aberrant mRNA-like noncoding RNAs by NMD in Arabidopsis. *Proc Natl Acad Sci USA* 106:2453-2458.
- Kyriakopoulou K, Dekkers B, van der Goot AJ. 2019. Plant-based meat analogues. In *Sustainable meat production and processing*. Galanakis CM (ed). Academic Press, Cambridge, MA, USA. pp 103-126.
- Latoch A, Czarniecka-Skubina E, Moczowska-Wyrwiz M. 2023. Marinades based on natural ingredients as a way to improve the quality and shelf life of meat: A review. *Foods* 12:3638.
- Lawless HT, Heymann H. 2010. *Sensory evaluation of food: Principles and practices*. Springer, New York, NY, USA.
- Lawrie RA, Ledward D. 2014. *Lawrie's meat science*. Woodhead, Sawston, UK.
- Lea CH, Swoboda PAT. 1962. Simple vacuum distillation procedure for determination of the volatile carbonyl content of autoxidising edible fats. *J Sci Food Agric* 13:148-158.
- Lim HJ, Kim GD, Jung EY, Seo HW, Joo ST, Jin SK, Yang HS. 2014. Effect of curing time on the physicochemical and sensory properties of beef jerky replaced salt with soy sauce, red pepper paste and soybean paste. *Asian-Australas J Anim Sci* 27:1174-1180.
- Lim YB, Kim HW, Hwang KE, Song DH, Kim YJ, Ham YK, Jang SJ, Lee CH, He FY, Choi YS, Kim CJ. 2015. Effects of glasswort (*Salicornia herbacea* L.) hydrates on quality characteristics of reduced-salt, reduced-fat frankfurters. *Korean J Food Sci Anim Resour* 35:783-792.
- Lorenzo JM, Pateiro M, Domínguez R, Barba FJ, Putnik P, Kovačević DB, Shpigelman A, Granato D, Franco D. 2018. Berries extracts as natural antioxidants in meat products: A review. *Food Res Int* 106:1095-1104.
- Marcus JB. 2019. Aging, nutrition and taste: Nutrition, food science and culinary perspectives for aging tastefully. Academic

Press.

- Meilgaard MC, Carr BT, Civille GV. 1999. Sensory evaluation techniques. CRC Press, Boca Raton, FL, USA.
- Mottram DS. 1998. Flavour formation in meat and meat products: A review. *Food Chem* 62:415-424.
- Nursten HE. 2005. The Maillard reaction: Chemistry, biochemistry and implications. The Royal Society of Chemistry, London, UK.
- Parry DA. 2016. The organoleptic qualities of protein foods (with special reference to cooking procedures). In *Proteins as human food: Proceedings of the Sixteenth Easter School in Agricultural Science*, University of Nottingham, 1969. Lawrie RA (ed). Butterworth-Heinemann, Oxford, UK. pp 365-387.
- Peris M, Escuder-Gilabert L. 2009. A 21st century technique for food control: Electronic noses. *Anal Chim Acta* 638:1-15.
- Pintado T, Muñoz-González I, Salvador M, Ruiz-Capillas C, Herrero AM. 2021. Phenolic compounds in emulsion gel-based delivery systems applied as animal fat replacers in frankfurters: Physico-chemical, structural and microbiological approach. *Food Chem* 340:128095.
- Purslow PP. 2022. New aspects of meat quality: From genes to ethics. Woodhead, Sawston, UK.
- Rahman MA, Alam AMMN, Shahjalal M. 2009. Supplementation of urea-molasses-straw based diet with different levels of concentrate for fattening of emaciated bulls. *Pak J Biol Sci* 12:970-975.
- Rashidaie Abandansarie SS, Ariaii P, Charmchian Langerodi M. 2019. Effects of encapsulated rosemary extract on oxidative and microbiological stability of beef meat during refrigerated storage. *Food Sci Nutr* 7:3969-3978.
- Ren Z, Li Z, Hu Z, Xia W, Zhou M, Pan Z, Li J, Zhen Z. 2024. Recent insights into bonding technologies in restructured meat production: A review. *Food Chem X* 23:101712.
- Resconi VC, Escudero A, Campo MM. 2013. The development of aromas in ruminant meat. *Molecules* 18:6748-6781.
- Sadakuzzaman M, Rahman MM, Hashem MA, Ali MS, Alam AN. 2024. Synergistic effects of gamma irradiation and butylated hydroxyanisole on the sensory, physicochemical, and microbiological quality of beef. *Appl Food Res* 4:100547.
- Samad A, Alam AN, Kumari S, Hossain MJ, Lee EY, Hwang YH, Joo ST. 2024. Modern concepts of restructured meat production and market opportunities. *Food Sci Anim Resour* 44:284-298.
- Schmidt CV, Mouritsen OG. 2022. Umami taste as a driver for sustainable eating. *Int J Food Des* 7:187-203.
- Shahidi F, Cadwallader KR. 1997. Flavor and lipid chemistry of seafoods: An overview. ACS, Washington, DC, USA.
- Shahidi F, Zhong Y. 2005. Lipid oxidation: Measurement methods: Edible oil and fat products: Chemistry, properties, and safety aspects. In *Bailey's industrial oil and fat products*. Shahidi F (ed). John Wiley & Son, Hoboken, NJ, USA.
- Shin YG, Rathnayake D, Mun HS, Dilawar MA, Pov S, Yang CJ. 2021. Sensory attributes, microbial activity, fatty acid composition and meat quality traits of Hanwoo cattle fed a diet supplemented with stevioside and organic selenium. *Foods* 10:129.
- Smith RL, Cohen SM, Fukushima S, Gooderham NJ, Hecht SS, Guengerich FP, Rietjens IMCM, Bastaki M, Harman CL, McGowen MM, Taylor SV. 2018. The safety evaluation of food flavouring substances: The role of metabolic studies. *Toxicol Res* 7:618-646.
- Smith TPL, Thallman RM, Casas E, Shackelford SD, Wheeler TL, Koohmaraie M. 2003. Theory and application of genome-based approaches to improve the quality and value of beef. *Outlook Agric* 32:253-265.
- Son YM, Lee EY, Alam AMMN, Samad A, Hossain MJ, Hwang YH, Seo JK, Kim CB, Choi JH, Joo ST. 2024. The application of high-intensity ultrasound on wet-dry combined aged pork loin induces physicochemical and oxidative alterations. *Food Sci Anim Resour* 44:899-911.

- Steensels J, Snoek T, Meersman E, Nicolino MP, Voordeckers K, Verstrepen KJ. 2014. Improving industrial yeast strains: Exploiting natural and artificial diversity. *FEMS Microbiol Rev* 38:947-995.
- Tieman D, Zhu G, Resende MFR Jr, Lin T, Nguyen C, Bies D, Rambla JL, Beltran KSO, Taylor M, Zhang B, Ikeda H. 2017. A chemical genetic roadmap to improved tomato flavor. *Science* 355:391-394.
- Tikk M. 2008. Peptides and ribonucleotides in fresh meat as a function of aging in relation to sensory attributes of pork. Ph.D. dissertation, Swedish University of Agricultural Sciences, Uppsala, Sweden.
- Toldrá F, Aristoy MC, Mora L, Reig M. 2012. Innovations in value-addition of edible meat by-products. *Meat Sci* 92:290-296.
- van Boekel MAJS. 2006. Formation of flavour compounds in the Maillard reaction. *Biotechnol Adv* 24:230-233.
- Wang L, O'Connor D, Rinklebe J, Ok YS, Tsang DCW, Shen Z, Hou D. 2020. Biochar aging: Mechanisms, physicochemical changes, assessment, and implications for field applications. *Environ Sci Technol* 54:14797-14814.
- Wang Y, Tuccillo F, Lampi AM, Knaapila A, Pulkkinen M, Kariluoto S, Coda R, Edelmann M, Jouppila K, Sandell M, Piironen V, Katina K. 2022. Flavor challenges in extruded plant-based meat alternatives: A review. *Compr Rev Food Sci Food Saf* 21:2898-2929.
- Wilailux C, Sriwattana S, Chokumnoyporn N, Prinyawiwatkul W. 2020. Texture and colour characteristics, and optimisation of sodium chloride, potassium chloride and glycine of reduced-sodium frankfurter. *Int J Food Sci Technol* 55:2232-2241.
- Wood SN. 2004. Stable and efficient multiple smoothing parameter estimation for generalized additive models. *J Am Stat Assoc* 99:673-686.
- Xu C, Yin Z. 2024. Unraveling the flavor profiles of chicken meat: Classes, biosynthesis, influencing factors in flavor development, and sensory evaluation. *Compr Rev Food Sci Food Saf* 23:e13391.
- Yoshida R, Ninomiya Y. 2023. Umami and MSG. In *Umami: Taste for health*. Gabriel AS, Rains TM, Beauchamp G (ed). Springer, Cham, Switzerland. pp 7-42.
- Zhang Y, Zhang Y, Zhou X, Wang S, Li P. 2021. Salt replacement changed the bacterial community composition and physicochemical characteristics of sodium-reduced fermented sausages during fermentation and ripening. *Foods* 10:630.
- Zhong R, Lu X, Zhong J, Chen L, Cheng W, Liang P. 2021. Influence of Maillard reaction in volatile flavor compounds of blue round scad (*Decapterus maruadsi*) enzymatic hydrolysate. *J Aquat Food Prod Technol* 30:615-629.