



REVIEW ARTICLE

Stability of biomaterials used in adjunct to coronally advanced flap: A systematic review and network meta-analysis

Sourav Panda^{1,2}  | Shahnawaz Khijmatgar^{2,3}  | Heber Arbildo-Vega⁴  |
Abhaya Chandra Das¹  | Manoj Kumar¹  | Mohit Das¹  |
Leonardo Mancini⁵  | Massimo Del Fabbro^{2,6} 

¹Department of Periodontics and Oral Implantology, Institute of Dental Sciences, Siksha O Anusandhan (Deemed to be) University, Bhubaneswar, Odisha, India

²Department of Biomedical, Surgical and Dental Sciences, University of Milan, Milan, Italy

³Department of Oral Biology and Genomic Studies, Nitte (Deemed to be University), AB Shetty Memorial Institute of Dental Sciences, Mangalore, India

⁴Department of General Dentistry, Dentistry School, Universidad San Martín de Porres, Chiclayo, Peru

⁵Department of life health and environmental sciences, University of L'Aquila, Italy

⁶IRCCS Istituto Ortopedico Galeazzi, Milan, Italy

Correspondence

Shahnawaz Khijmatgar, Department of Biomedical, Surgical and Dental Sciences, University of Milan, Milan, Italy.
Email: khijmatgar@gmail.com

Abstract

Aim: The objective of this network meta-analysis was to rank different biomaterials used in adjunct to coronally advanced flap (CAF), based on their performance in root-coverage for Miller's Class I and II gingival recessions.

Materials and methods: An electronic database search was carried out in PUBMED, CENTRAL, SCOPUS, and EMBASE to identify the eligible articles and compiled into the citation manager to remove the duplicates. The primary outcome was keratinized gingival tissue width (KGW) and percentage of root coverage (%RC). The treatment effect of different biomaterials was estimated using predictive interval plots and ranked based on biomaterials performance, using multidimensional scale ranking.

Results: CAF + connective tissue graft (CTG), CAF + platelet concentrate matrix (PCM) and acellular dermal matrix (ADM) ranked at the top positions in performance in improving KGW. The highest ranked materials in improving percentage of root coverage in gingival recession were CAF + collagen matrix (CM) + gingival fibroblasts (GF), CAF + ADM + platelet rich plasma (PRP) and CAF + ADM, as compared to CAF alone.

Conclusion: CTG, ADM, platelet concentrates, and CM + GFs, when used in adjunct to CAF, showed improved stability over ≥ 12 months of follow-up, better percentage of root coverage, and improved keratinized gingival width.

KEYWORDS

CAF, clinical attachment level, keratinised tissue width, network meta-analysis, oral, recession height, recession width, regeneration, root coverage, soft tissue, systematic review

1 | INTRODUCTION

Gingival recession (GR) is characterized by displacement of the gingival margin below the level of cemento-enamel junction (Cortellini & Bissada, 2018). Several etiological factors like age, anatomical, physiological, pathological, trauma, hygiene, abnormal

frenal attachment, and so on, were identified for this condition (Fu et al., 2012) which may account for its relatively high incidence in the population (Rios et al., 2014; Susin et al., 2004). It affects more than 50% of population including healthy individuals. Recession of 1 mm or more is prevalent in aged 30 years and older. The risk increases with age. Root exposure leads to hypersensitivity, cervical caries, aesthetics complications, and non-carious cervical lesions (Jepsen et al., 2017).

PROSPERO registration number: CRD42020208010

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Clinical and Experimental Dental Research* published by John Wiley & Sons Ltd.

Root coverage procedures (RCP) have shown to be effective in treating single and multiple GRs (Cairo et al., 2014; Tavelli, Barootchi, et al., 2018; Tavelli, Barootchi, et al., 2019; Tavelli, Ravidà, et al., 2019), and in literature, several techniques were proposed. Nevertheless, the superiority of coronally advanced flap (CAF) combined with connective tissue graft (CTG) is clear. Indeed, CAF + CTG is a gold standard in RCP for the best outcomes achieved in terms of mean root coverage, keratinized tissue width, gingival thickness and aesthetics results (Zucchelli et al., 2020).

However, the presence of a second surgical site, patient morbidity, and limited availability are the main drawbacks that have been largely described for CTG (Tavelli, Asa'ad, et al., 2018; Tavelli, Barootchi, et al., 2019; Tavelli, Ravidà, et al., 2019). For this reason, several CTG substitutes were introduced including Platelet rich plasma (PRP) or Platelet rich fibrin (PRF), acellular dermal matrix (ADM), enamel matrix derivative (EMD), and xenogeneic collagen matrix (CMX). These biomaterials suffer from limitations in term of shape, consistency and size. PRF is a living cellular graft enriched with growth factors and it is a good alternative to CTG for the availability and the easy handling (Dohan et al., 2006; Miron et al., 2020). The ADM is soft CTG generated by a decellularization process preserving the extracellular skin matrix with high costs and it is not in use for ethical problems in diverse countries (McGuire et al., 2020; Tavelli et al., 2020). EMD is a porcine fetal tooth material extracted and manipulated as a gel used as an enhancer in oral regenerative procedures (McGuire & Nunn, 2003). CMX is another biomaterial, which has different layers of collagen fibers and porous layer facilitating blood clot formation and in-growth of tissue from adjacent sites (McGuire & Scheyer, 2016; Vignoletti et al., 2011).

Several studies revealed the comparison among CAF + CTG and each alternative therapy during a follow-up period of 6–12 months showing divergent results (Keceli et al., 2015; McGuire & Scheyer, 2010; Tonetti et al., 2018). According to the authors in the literature, a direct, indirect and mixed evidence for all these biomaterials contributing to the success of root coverage is not present and data extracted from non-systematic comparisons might be confusing and not well interpreted. In addition, the data from all studies are heterogeneous (differences in estimates of effect across studies that assessed the same comparison), which makes difficult to compare all materials. A conventional pairwise meta-analysis results in only one pooled effect estimate. Therefore, a novel method of weighing the effect estimate through network meta-analysis (NMA) has been proposed.

Previous network systematic reviews tried to collect data evaluating the clinical advantages for each CTG substitute with several limitations such as the follow-up period of 6-months which might be a limit, the inclusion of randomized clinical trials (RCTs) with a high risk of bias influencing results and the inclusion of RCTs with smoker patients or RCTs where the absence or presence of smoker patients was not reported (Buti et al., 2013; Moraschini et al., 2020). Thus, the purpose of this systematic review and NMA was to compare the clinical effects among patients who have one or more gingival recession sites and corrected with intervention of CTG substitutes and compared with controls or CAF alone or in combination for regeneration of keratinised gingival width (KGW), clinical attachment level (CAL), recession width (RW), recession height (RH), pocket depth outcomes during a long follow-up period.

2 | MATERIAL AND METHODS

This review was performed in accordance with of PRISMA guidelines. The protocol for this review and NMA was registered in PROSPERO with registration ID: CRD42020208010.

The eligibility of study was decided based on PICO format.

Type of Patients: Patients who had one or more than one site of gingival recession was considered for assessment and further review analysis.

Type of Intervention: CAF or/and CTG Substitutes.

Type of Comparator: Compared with Placebo, Control or CAF or/and combination of CAF + biomaterials or CTG substitutes.

Type of outcomes: KGW, CAL, Recession Height, Recession Width, Pocket Depth were the outcomes.

Type of Duration: More than 6 months' follow-up periods.

2.1 | Research question

What is the treatment effect of different biomaterials like CTG, EMD, ADM, PRP/PRF, CMX, and combination of these when used in adjunct to CAF for root coverage?

2.2 | Search strategy

An electronic database search was carried out in PUBMED, CENTRAL, SCOPUS, and EMBASE to identify the potentially eligible articles using the following strategy:

“(((Coronally advanced flap) OR (CAF)) OR (modified coronally advanced flap)) OR (coronally displaced flap) AND (((((((((((Enamel matrix derivative) OR (Connective tissue graft)) OR (Guided tissue regeneration)) OR (Collagen matrix)) OR (Acellular dermal matrix)) OR (platelet rich fibrin)) OR (platelet rich plasma)) OR (PRF)) OR (PRP)) OR (barrier membrane)) OR (amniotic membrane)) OR (hyaluronic acid)) OR (Emdogain)) OR (CTG).”

A manual search in periodontal journals like Journal of Clinical Periodontology, Journal of Periodontal Research, Journal of Periodontal and Implant Science, International Journal of Periodontics and Restorative Dentistry, and Journal of Periodontology. There were no limits or filters applied during the search. Both studies relevant to the topic in areas of systematic reviews and clinical trials were searched.

2.3 | Inclusion criteria

1. Randomized Clinical Trials (Both parallel and split mouth design)
2. Study follow up duration at least 12 months
3. Minimum sample size of 10 per group
4. CAF procedure should be employed both in test and control group.
5. The test group should have any of the biomaterials in adjunct to CAF compared to control group with a different biomaterial or none in adjunct to CAF.

6. Treatment of Class 1 and 2 gingival recessions only.
7. Both isolated and multiple recession

2.4 | Exclusion criteria

1. Studies not in English
2. Study participants under any medication which could influence the outcome of treatment.
3. Teeth with non-carious cervical lesions (NCCL)
4. Animal studies

2.5 | Study selection

The studies from the databases searches were compiled into citation manager to remove duplicates and screened for all titles and abstract by two independent reviewers (M.K and A.C.D). The eligible studies were then subjected to full text assessment and included for qualitative assessment. In case of disagreement or uncertainty while selecting the eligible articles, an expert third reviewer (M.D.F) was consulted until a consensus was reached. Detailed reasons were mentioned for all excluded studies.

2.6 | Data extraction

The qualitative data was extracted using excel spreadsheet. The data extraction was carried out by using two independent reviewers (M.

Dd, H.A.V.). In cases like missing or unclear data or need for additional data or raw data, the authors were approached through emails or telephone for enquiring the details of missing or unclear information.

2.7 | Outcomes

The primary outcomes that were assessed in this review were keratinized gingival width (KGW) and the percentage of root coverage (%RC). The secondary outcomes assessed included CAL, RW, and RD.

2.8 | Data synthesis

The data extracted were both qualitative and quantitative. The former were related to demographics of the study and type of publication. The quantitative data for the different outcomes allowed to undertake NMA. The NMA enables to develop a network geometry plot, where the number of studies and subjects between the comparators are projected. The risk of bias in each network was also estimated. The predictive interval plots (PrI) were calculated to predict the effects of future clinical studies incorporating heterogeneity. The surface under the cumulative ranking curve (SUCRA) was calculated for each treatment. Treatments were ranked based on their respective performances. Treatments with SUCRA values with higher percentage of being first were ranked higher and values with lower percentage were ranked lower (0–100%). The multi-dimensional scale ranking was employed to rank the biomaterials based on their dissimilarity. The network estimates for all comparisons are




FIGURE 1 PRISMA flow chart diagram

treated as proximity data aimed to reveal their latent structure. By this, the dissimilarity between any two treatments was distinguished. NMA was carried out using Stata version 16 (StataCorp, College Station, TX) by a single reviewer (S.K).

3 | RESULTS

This qualitative and NMA analysis was carried out by assessing 39 RCTs analyzing the stability of CAF when used alone or in combination with different biomaterials in treatment of class I and II gingival recession defects, over at least 12 months follow-up. The electronic database search and manual search of related journals and bibliographies yielded a total of 1223 articles. The searches from different databases were imported to a citation manager (ENDNOTE) to identify 938 articles after removing duplicates. All the articles were subjected to title and abstract screening, and were narrowed down to identify 56 potentially eligible studies. These studies were subjected to detailed full text assessment by two independent reviewers. Out of 56 eligible studies, 39 RCTs were included in this systematic review and 19 were considered for NMA. The detailed process of study selection is provided in the PRISMA flow chart (Figure 1). The rest of 17 articles were excluded with detailed reason for exclusion (refer Table 1).

3.1 | Characteristics of included studies

The demographic and interventional characteristics of all included studies are presented in Tables 2 and 3, respectively.

This systematic review analyzed a total of 1733 sites from 936 participants included in 39 RCTs. The trials included 546 females and 390 male participants and the age range of the participants recruited in the included RCTs were 18–74 years.

3.2 | Network meta-analysis

For the NMA, 19 RCTs with a total of 858 teeth in the entire network was considered.

3.2.1 | Keratinized gingival width

The most common comparator is between CAF, CAF + CMX, CAF + CTG and CAF + EMD. The risk of bias is found to be low between CAF and CAF + CM and high between other comparisons (Figures 2–5). A total of $N = 858$ patients are included in the entire network. There are three studies included in CAF and CAF + CTG and four studies in CAF and CAF + EMD. The percentage contribution of each direct comparison in the entire network CAF and CAF + CTG (13.33%) followed by CAF and CAF + rhPDGF-BB + TCP (10.33%) and that of each pairwise summary effect is from CAF and

TABLE 1 List of excluded studies

SI no.	Study	Reason of exclusion
1.	Tavelli, Barootchi, et al. (2019); Tavelli, Ravidà, et al. (2019)	Coronally advanced flap compared to tunnel technique
2.	Stefanini et al. (2018)	Not a randomized clinical trial but a case series
3.	Bellver-Fernández et al. (2016)	Test group has less than 10 sites
4.	Wang et al. (2015)	Same biomaterial used in both groups
5.	Wang et al. (2014)	Same biomaterial used in both groups
6.	Zucchelli et al. (2014)	Same biomaterial used in both groups
7.	Aroca et al. (2013)	Tunnel technique has been used in both the groups
8.	McGuire et al. (2012)	Less than 10 patients as sample size
9.	Aleksic et al. (2010)	Study in Russian language
10.	Aroca et al. (2009)	Follow up till 6 months
11.	Pourabbas et al. (2009)	Follow up till 6 months
12.	Moses et al. (2006)	Two separate procedures have been compared
13.	Dominiak et al. (2006)	Three separate procedures have been compared
14.	Nemcovsky et al. (2004)	Inclusion of cervically abraded teeth
15.	Wennström et al. (1996)	Non-randomized, prospective clinical study
16.	Pini-Prato et al. (2010)	Non randomized study
17.	Henriques et al. (2010)	Class III gingival recessions

CAF + ADM (100%) followed by CAF and CAF + CTG (99.0%). CAF + GTR(NonBio) (11) and CAF + CMX (12) have significant variance and inconsistency. CAF + GTR-Bio and CAF + GTR-NonBio more likely to perform poor in future studies in gaining width thickness of gingival recession cases. A favorable outcome is expected through CAF + PCM and CAF + CTG in present (CrI) and future studies (PrI). CAF + CTG and CAF + PCM are ranked higher in SUCRA ranking. GTR Bio and Non-Bio most distinct and dissimilar among other materials as illustrated in MDS rank.

3.2.2 | Percentage of root coverage

The network geometry plot illustrates the total number of subjects ($N = 826$) and comparisons between each intervention (Figures 6–9). The number of RCT's in CAF and CTG are two and CAF + EMD are three. The majority of direct evidence contribution was by CAF + PCM and CAF + CTG (99.04) followed by CAF + ADM and CAF + CMX (98.36). The Majority of indirect evidence contribution is

TABLE 2 Demographic characteristics of included studies

SI no.	Author and year	Study design	Age in years (range, mean SD)	Type of recession	No. of patients	No. of sites	Test procedure	Control procedure	Follow-up
1.	Barakat and Dayoub (2020)	RCT split-mouth	20–45	Single	20	40	CAF + PCM	CAF + CTG	12 months
2.	Rotundo et al. (2021)	RCT parallel	> 18	Multiple	24	61	CAF + CMX	CAF	12 months
3.	Aydinyurt et al. (2019)	RCT split-mouth	18–55	Single	19	38	CAF + CTG + EMD	CAF + CTG	12 months
4.	Pilloni et al. (2019)	RCT parallel	21–47	Single	30	30	CAF + HA	CAF	18 months
5.	Rotundo et al. (2019)	RCT parallel	> 18	Multiple	24	61	CAF + CMX	CAF	12 months
6.	França-Grohmann et al. (2019)	RCT parallel	29.74	Single	30	30	CAF + EMD	CAF	12 months
7.	Gürlek et al. (2020)	RCT split-mouth	> 18	Multiple	15	82	CAF + ADM	CAF + CTG	18 months
8.	Matoh et al. (2019)	RCT split-mouth	21–52	Single	10	20	CAF + CM	CAF + CTG	12 months
9.	Francetti et al. (2018)	RCT parallel	> 18	Single	20	NR	CAF + CTG	CAF	60 months
10.	Kuka et al. (2018)	RCT parallel	21–41	Multiple	24	52	CAF + PRF	CAF	12 months
11.	Rasperini et al. (2018)	RCT parallel	37–63	Single	25	25	CAF + CTG	CAF	108 months
12.	Çetiner et al. (2018)	RCT split-mouth	20–67	Multiple	12	84	CAF + ADM + PRP	CAF + ADM	12 months
13.	Abou-Arraj et al. (2017)	RCT parallel	24–74	Single	17	17	CAF + ADM-A	CAF + ADM-B	12 months
14.	Jepsen et al. (2017)	RCT split-mouth	20–73	Single	18	36	CAF + CMX	CAF	36 months
15.	Cairo et al. (2016)	RCT parallel	20–53	Multiple	32	74	CAF + CTG	CAF	12 months
16.	McGuire & Scheyer, (2016)	RCT split-mouth	51.3 ± 13.9	Single	17	34	CAF + CMX	CAF + CTG	60 months
17.	Godavarthi et al. (2016)	RCT split-mouth	41.4 ± 7.6	Single	14	28	CAF + PPG	CAF + ADM	12 months
18.	Stefanini et al. (2016)	RCT split-mouth	39.5 ± 13.8	Multiple	45	41	CAF + CMX	CAF	12 months
19.	Lops et al. (2015)	RCT parallel	46	Single	28	28	CAF + CTG	CAF	12 months
20.	Cairo et al. (2015)	RCT parallel	45.9 ± 10.3	Single	24	24	CAF + CTG	CAF	36 months
21.	Milinkovic et al. (2015)	RCT split-mouth	18–55	Both	18	36	CAF + AFCC	CAF + CTG	12 months
22.	McGuire et al. (2014)	RCT split-mouth	29–68	Both	30	60	CAF + rhPDGF-BB	CAF + CTG	60 months
23.	Ahmedbeyli et al. (2014)	RCT parallel	22–40	Multiple	24	48	CAF + ADM	CAF	12 months
24.	Zucchelli et al. (2014)	RCT parallel	>18	Multiple	50	50	CAF + CTG	CAF	60 months
25.	Cardaropoli et al. (2014)	RCT parallel	38.4 ± 11.1	Multiple	32	113	mCAF+CM	mCAF	12 months
26.	Alkan and Parlar (2013)	RCT split-mouth	35–53	Multiple	12	56	CAF + EMD	CAF + CTG	12 months
27.	Kuis et al. (2013)	RCT split-mouth	20–52	Multiple	37	114	CAF + CTG	CAF	60 months
28.	Köseoğlu et al. (2013)	RCT split-mouth	19–41	Single	11	22	CAF + CMX + GF	CAF + CMX	12 months
29.	Roman et al. (2013)	RCT split-mouth	21–28	Both	42	57	CAF + CTG + EMD	CAF + CTG	12 months
30.	Kumar and Murthy (2013)	RCT split-mouth	18–60	Single	12	24	CAF + PCG	CAF + CTG	12 months
31.	Cordaro et al. (2012)	RCT split-mouth	18–60	Multiple	10	58	CAF + EMD	CAF	24 months
32.	Cardaropoli et al. (2012)	RCT split-mouth	21–59	Multiple	18	22	CAF + CMX	CAF + CTG	12 months
33.	Alkan and Parlar (2011)	RCT split-mouth	23–42	Single	12	24	CAF + EMD	CAF + CTG	12 months
34.	Jankovic et al. (2010)	RCT split-mouth	21–48	Single	20	40	CAF + EMD	CAF + PRF	12 months
35.	Del Pizzo et al. (2005)	RCT split-mouth	18–56	Single	15	30	CAF + EMD	CAF	24 months

(Continues)

TABLE 2 (Continued)

SI no.	Author and year	Study design	Age in years (range, mean SD)	Type of recession	No. of patients	No. of sites	Test procedure	Control procedure	Follow-up
36.	Spahr et al. (2005)	RCT split-mouth	23–62	Single	30	60	CAF + EMD	CAF + Placebo	24 months
37.	McGuire and Nunn (2003)	RCT split-mouth	23–62	Single	20	40	CAF + EMD	CAF + CTG	12 months
38.	Hägewald et al. (2002)	RCT split-mouth	22–62	Single	36	72	CAF + EMD	CAF + Placebo	12 months
39.	Zucchelli et al. (1998)	RCT parallel	23–33	Single	54	54	CAF + GTR(R) CAF + GTR (NR)	CAF + CTG	12 months

Abbreviations: RCT- Randomized controlled trial, CAF- Coronally Advanced Flap, mCAF- modified Coronally Advanced Flap, PCM- Platelet concentrate matrix, CMX- Xenogenic Collagen Matrix, CTG- Connective Tissue Graft, EMD- Enamel Matrix Derivative, HA- Hyaluronic Acid, ADM- Acellular Dermal Matrix, PRF- Platelet Rich Fibrin, PRP- Platelet Rich Plasma, ADMA- Acellular Dermal Matrix(AlloDerm BioHorizons), ADMB- Acellular Dermal Matrix(Puros Dermis Zimmer Biomet), PPG- Periosteal Pedicle Graft, AFCC- Autologous Fibroblast Cell Culture, rh-PDGF-BB - recombinant human Platelet Derived Growth Factor- BB, GF- Gingival Fibroblasts, PCG- Platelet Concentrate Graft, GTR(R)- Resorbable Guided Tissue Regeneration Membrane, GTR(NR)- None Resorbable Guided Tissue Regeneration Membrane.

from CAF and CAF + CTG (41.15%) followed by CAF + CTG and CAF + PRF (20.54%). The evidence from entire network contribution is between CAF and CAF + ADM (9.87%) and CAF and CAF + CMX (8.34%).

There is no statistically significant inconsistency in the loop formed by CAF (01), CAF + PCM (03) and CAF + CMX (08) as the lower limit confidence intervals reaches the zero line. CAF + rhPDGF-BB + TCP and CAF + CTG + EMD more likely to perform poor in future studies. It is predicted that CAF + CM + GF treatment would be 73.86% successful followed by CAF + ADM + PRP (45.83%).

CAF + CM + GF and CAF + ADM + PRP ranked higher in SUCRA ranking. Multidimensional scale (MDS) ranking shows that CAF + CTG and CAF + ADM + PRP are the most distinct interventions in improving the outcomes of root coverage in gingival recessions.

3.2.3 | Clinical attachment level

The highest ranked materials for this specific outcome in CAL were CAF + ADM and CAF + PRF in gingival tissue reconstruction. The effect estimate for CAF + ADM was 1.79(1.21, 2.84) and CAF + PRF was 1.43(0.99, 2.08) (Supplementary Figures 1(a)–(d)).

3.2.4 | Recession width

The highest ranked materials for RW gain in gingival recession were CAF + CTG and CAF + ADM. The effect estimate was 0.75(0.40, 1.39) and 0.37(0.11, 1.23) for CAF + CTG and CAF + ADM, respectively (Supplementary Figures 2(a)–(d)).

3.2.5 | Recession height

The highest ranked materials for RD gain were CAF + ADM and CAF + PCM. The effect estimate was found to be 2.03(1.38, 3.00) for CAF + ADM and 1.61 (1.04, 2.49) for CAF + PCM (Supplementary Figures 3(a)–(d)).

3.3 | GRADE analysis

The overall level of evidence for the regenerative material is moderate. We predominantly downgraded the rating of the evidence due to different levels of risk of bias and imprecision (Table 4).

4 | DISCUSSION

Our results from NMA explored the effectiveness of different biomaterials in periodontal regeneration (i.e., Gingival Recession). KGW and percentage of root coverage are considered as primary outcomes for validating the successful periodontal regeneration by a biomaterial (Lang & Löe, 1972). We found that, CAF + CTG ranked with highest probability followed by CAF + PCM in keratinised tissue width regeneration (Figure 5). A favorable outcome is expected through CAF + PCM and CAF + CTG as indicated from effect sizes, CrI and PrI. Similarly, CAF + CTG ranked with highest probability followed by CAF + ADM + PRP in favorable outcomes in percentage of root coverage. It is predicted that, CAF + CM + GF treatment would be 73.86% more successful followed by CAF + ADM + PRP (45.83%) (Figure 9).

The ROB among studies that included CAF and CAF + CMX intervention was high and ROB between CAF and CAF + CM was low. Other studies were unclear or had moderate ROB in keratinised gingival width (KGW) outcome (Figure 1(a)). Similarly, ROB was high among majority of the comparisons except the comparison between CAF Vs CAF + CM for percentage of root coverage (Figure 2). The ROB for all the included studies was illustrated in Figure 10. According to the inconsistency plot, the loop formed by CAF, CAF + CTG, CAF + EMD, CAF + GTR (Nonbio) and CAF + CM had significant inconsistency in KGW outcome (Figure 1(b)). Similarly, CAF, CAF + PCM, CAF + CM, CAF + CTG + EMD and CAF + EMD had significant inconsistency in percentage of root coverage outcome (Figure 3). The overall evidence from all the comparisons for KGW and percentage of root coverage was found to be moderate (Table 3). The ranking of materials rated highest and lowest should be interpreted carefully by taking ROB and inconsistencies factors between these comparisons and dissimilarity between the materials illustrated by multidimensional scaling (Figure 1(e) and 5(b)).

TABLE 3 Interventional characteristics of included studies

Author and year	Procedure	Test biomaterial	Type of biomaterial	Trade name - test	Control biomaterial	Type of biomaterial	Trade name - control	Surgery time(test)	Surgery time (control)	Outcomes assessed
Barakat and Davoub (2020)	CAF	CMX	XENO	Mucograft collagen matrix; Geistlich Pharma	CTG	AUTO	Autologous	NR	NR	RD, PD,CAL, WKG, RC%, CRC%, aesthetic score(RES) and pain intensity
Rotundo et al. (2021)	CAF	CMX	XENO	Geistlich Mucograft®, Geistlich Pharma AG	None	-	None	NR	NR	PI, FMBS, RD, WKG, GT, PD
Aydinyurt et al. (2019)	CAF	EMD	XENO	Emdogain®, Straumann, Basel, Switzerland	CTG	AUTO	Autologous	NR	NR	RD, RW, RC%, CRC%, aesthetic score (RES)
Pilloni et al. (2019)	CAF	HA	ALLP	HyADENT BG, Bioscience, Germany	None	-	None	NR	NR	RD, CAL, PD, WKG, CRC%, RC%, pain intensity
Rotundo et al. (2019)	CAF	CMX	XENO	Geistlich Mucograft®, Geistlich Pharma AG	None	-	None	47.3 ± 5.8	36.1 ± 4.6	RD, RW, BOP, PI, PD, CAL, WKG, GT
França-Grohmann et al. (2019)	CAF	EMD	XENO	Emdogain® Institut Straumann AG, Basel, Switzerland	None	-	None	NR	NR	RD, RW, WKG, GT, PD, CAL
Gürlek et al. (2020)	Modified CAF	ADM	ALLO	Mucoderm, Botiss GmbH, Berlin, Germany	CTG	AUTO	Autologous	NR	NR	RD, RW, WKG, PD, CAL
Match et al. (2019)	CAF	CM	XENO	OsteoBiol Derma, Tecness	CTG	AUTO	Autologous	NR	NR	RD, PD, CAL, WKG, GT, RC%
Francetti et al. (2018)	CAF	CTG	AUTO	Autologous	None	-	None	NR	NR	RD, CRC%, WKG, CAL, RC%
Kukka et al. (2018)	CAF	PRF	BIO	Autologous	None	-	None	NR	NR	RD, RW, PD, CAL, GT, WKG, PI, GI and BOP
Rasperini et al. (2018)	CAF	CTG	AUTO	Autologous	None	-	None	NR	NR	PD, RD, CAL, WKG, CRC%
Çetiner et al. (2018)	CAF	PRP	BIO	Curasan, Pharma GmbH AG, Lindigstrab, Germany	ADM	ALLO	SureDerm, Seoul, Korea	NR	NR	RD, RW, WKG, PI, GI, PD, CAL, RC%
Abou-Arraj et al. (2017)	CAF	ADM-A	ALLO	AlloDerm BioHorizons	ADM-B	ALLO	Puros Dermis Zimmer Biomet	NR	NR	WKG, RD, GT
Jepson et al. (2017)	CAF	CMX	XENO	NR	None	-	None	NR	NR	RD, CRC%, RD, WKG, GT, CAL, PD
Cairo et al. (2016)	CAF	CTG	AUTO	Autologous	None	-	None	NR	NR	RD, CRC%, PD, CAL, WKG, GT
McGuire and Scheyer (2016)	CAF	CMX	XENO	NR	CTG	AUTO	None	NR	NR	RD, CRC%, WKG, PD and CAL
Godavarthi et al. (2016)	CAF	PPG	BIO	Autologous	ADM	ALLO	AlloDerm BioHorizons	NR	NR	RD, CRC%, WKG, PD, CAL
Stefanini et al. (2016)	CAF	CMX	XENO	Geistlich Mucograft®, Geistlich Pharma AG	None	-	None	NR	NR	RD, RW, WKG, GT, CAL, PD
Lops et al. (2015)	CAF	CTG	AUTO	Autologous	None	-	None	NR	NR	RD, WKG, PD, CAL
Cairo et al. (2015)	CAF	CTG	AUTO	Autologous	None	-	None	NR	NR	RD, WKG, PD, CAL
Milinkovic et al. (2015)	CAF	AFCC	AUTO	Autologous	CTG	AUTO	Autologous	NR	NR	RD, WKG, CAL
McGuire et al. (2014)	CAF	rhPDGF-BB	BIO	GEM21-S	CTG	AUTO	Autologous	NR	NR	RD, CAL, PD, WKG, RC%, CRC%, Pain intensity
Ahmedbeyli et al. (2014)	CAF	ADM	ALLO	AlloDerm, BioHorizon, USA	None	-	None	NR	NR	RD, WKG, GT
Zucchelli et al. (2014)	CAF	CTG	AUTO	Autologous	None	-	None	29.8 ± 3.2	40.2 ± 6.8	RD, CRC%, PD, CAL, WKG, Pain Intensity

(Continues)

TABLE 3 (Continued)

Author and year	Procedure	Test biomaterial	Type of biomaterial	Trade name - test	Control biomaterial	Type of biomaterial	Trade name - control	Surgery time(test)	Surgery time (control)	Outcomes assessed
Cardaropoli et al. (2014)	Modified CAF	CMX	XENO	Mucograft, Geistlich	None	-	None	NR	NR	RD, PD, CAL, WKG, GT
Alkan and Parlar (2013)	CAF	EMD	XENO	Emdogain [®] , Institut Straumann AG, Basel, Switzerland	CTG	AUTO	Autologous	NR	NR	RD, RW, PD, CAL, RC%, WKG, GT
Kuis et al. (2013)	CAF	CTG	AUTO	Autologous	None	-	None	NR	NR	PI, FMBS, PD, RD, WKG, CAL, RC%, CRC%
Köseoğlu et al. (2013)	CAF	GF	AUTO	Autologous	CMX	XENO	Collagen AD	NR	NR	PI, GI, PD, CAL, RD, RW, WKG, GT
Roman et al. (2013)	CAF	EMD	XENO	Emdogain [®] , Institut Straumann AG, Basel, Switzerland	CTG	AUTO	Autologous	NR	NR	RD, RC%, CRC%, WKG
Kumar and Murthy (2013)	CAF	PCG	BIO	Autologous	CTG	AUTO	Autologous	NR	NR	PI, GI, PD, CAL, RD, Pain Intensity, WKG
Cordaro et al. (2012)	CAF	EMD	ALLO	Emdogain [®] , Institut Straumann AG, Basel, Switzerland	None	-	None	NR	NR	RD, PD, CAL, WKG, FMBS, PI
Cardaropoli et al. (2012)	CAF	CMX	XENO	Mucograft, Geistlich	CTG	AUTO	Autologous	NR	NR	RD, PD, CAL, WKG
Alkan and Parlar (2011)	CAF	EMD	ALLO	Emdogain [®] , Institut Straumann AG, Basel, Switzerland	CTG	AUTO	Autologous	NR	NR	RD, CAL, PD
Janković et al. (2010)	CAF	EMD	ALLO	Emdogain [®] , Institut Straumann AG, Basel, Switzerland	PRF	BIO	Autologous	NR	NR	RD, PD, CAL, Pain Intensity
Del Pizzo et al. (2005)	CAF	EMD	ALLO	Emdogain Biora AB	None	-	None	NR	NR	RD, RW, PD, CAL, WKG
Spahr et al. (2005)	CAF	EMD	AUTO	Emdogain Biora AB	Placebo	-	PGA	NR	NR	RD, RW, WKG, CAL, PD
McGuire and Nunn (2003)	CAF	EMD	ALLO	Emdogain Biora AB	CTG	AUTO	Autologous	NR	NR	RD, RW, PD, CAL, WKG
Hägewald et al. (2002)	CAF	EMD	ALLO	Emdogain Biora AB	Placebo	-	PGA	NR	NR	RD, RW, WKG, CAL, PD
Zucchelli et al. (1998)	CAF	GTR	ALLP	NR	CTG	AUTO	Autologous	NR	NR	PD, CAL, RD, WKG, RC%

Abbreviations: CAF- Coronally Advanced Flap, mCAF- modified Coronally Advanced Flap, PCM- Platelet concentrate matrix, CMX- Xenogenic Collagen Matrix, CTG- Connective Tissue Graft, EMD- Enamel Matrix Derivative, HA- Hyaluronic Acid, ADM- Acellular Dermal Matrix, PRF- Platelet Rich Fibrin, PRP- Platelet Rich Plasma, ADMA- Acellular Dermal Matrix(AlloDerm BioHorizons), ADMB- Acellular Dermal Matrix(PuroS Dermis Zimmer Biomet), PPG- Periosteal Pedicle Graft, AFCC- Autologous Fibroblast Cell Culture, rh-PDGF-BB – recombinant human Platelet Derived Growth Factor- BB, GF- Gingival Fibroblasts, PCG- Platelet Concentrate Graft, GTR(R)- Resorbable Guided Tissue Regeneration Membrane, GTR(NR)- None Resorbable Guided Tissue Regeneration Membrane, AUTO- Autologous Biomaterial, ALLO – Allogenic Biomaterial, XENO- Xenogenic Biomaterial, ALLP – Allogenic Biomaterial, BIO- Biologic Biomaterial, RD- Recession Depth, RW- Recession Width, PD- Probing Depth, CAL- Clinical Attachment Level, WKG- Width of Keratinised Gingiva, RC% - Percentage of Root Coverage, FMBS – Full Mouth Bleeding Score, CRC% - Complete Root Coverage %, GT- Gingival Thickness, PI- Plaque index, GI- Gingival Index, BOP- Bleeding on Probing, RES- Aesthetic score, NR – Not Reported.




FIGURE 2 Network geometry plot for keratinized gingival width

CTG has always been considered as gold standard intervention for root coverage and for the modification of periodontal phenotype (Barootchi et al., 2020; Chambrone & Tatakis, 2015; Tatakis et al., 2015) because it demonstrates the best long-term maintenance of treatment (Pini Prato, Franceschi, et al., 2018). However, it presents with limitations such as increased surgical morbidity, bleeding and postoperative pain. Therefore, clinicians and the patients look for alternatives that can meet the clinical need and also improve the post treatment quality of life (Moraschini et al., 2019). In contrast, the use of CTG substitutes does not affect postoperative pain and quality of life (Rotundo et al., 2019; Tonetti et al., 2018). For this reason, the decision making to choose among different biomaterial substitutes in adjunct to CAF for root coverage of single or multiple gingival recessions must be based on scientific evidence.





FIGURE 3 The inconsistency plot for KGW

FIGURE 4 Predictive interval (PrI) and confidence interval plot (CrI) for KGW






FIGURE 5 SUCRA ranking; 2E. Multi-dimensional scale ranking (MDS) for keratinized gingival width




FIGURE 6 Network geometry plot for percentage of root coverage

Recently, the effect of time on the stability of postsurgical results emerged as an important factor for root coverage and periodontal procedures (Cortellini et al., 2017; Pini Prato, Magnani, et al., 2018; Wu et al., 2017). A duration of 6 months is considered as a sufficient time for healing and tissue stability after mucogingival surgery (Cheng et al., 2007) and some authors have shown that the data obtained at this time can already predict the results of 3 years of the RCP (Cairo et al., 2015; Jepsen et al., 2017), and at 12 months the maturation of the tissue after the procedure is already complete (Gurtner et al., 2008; Smith et al., 2015).

A data on long term effects of different RCP has been recently being reported (Moslemi et al., 2011; Pini Prato et al., 2011; Pini Prato, Franceschi, et al., 2018; Pini Prato, Magnani, et al., 2018; Rasperini et al., 2018) and although there are still some controversies, CTG-based techniques show the least changes over time (Pini Prato, Franceschi, et al., 2018; Rasperini et al., 2018). But, despite the fact that the evidence provides favorable results of early

treatment (6 or 12 months) for gingival recessions (Francesco Cairo et al., 2014; Tavelli, Barootchi, et al., 2018), whether they persist for a longer time, has not yet been determined (Chambrone et al., 2019). In addition, a definitive conclusion cannot be drawn individually because of limited sample size and high drop outs (Chambrone et al., 2019; McGuire et al., 2012; Rasperini et al., 2018). Therefore, a time greater than 12 months, as a variable of the obtained treatment results, has never been explored. For this reason, this SR and NMA made direct and indirect comparisons between possible CTG substitutes with a minimum follow-up time of 12 months, avoiding any influence of extrinsic factors in the healing process. Furthermore, the incorporation of an NMA can provide information on the effect of time on the changes that occur in the results obtained postoperatively and at the same time the different substitutes of the CTG are compared.

Although the treatment of choice, in terms of flap design, remains controversial (Santamaria et al., 2017; Zuhr et al., 2014), in order to guarantee homogeneity in the analyses of the present study, all included studies used the CAF as the flap design. All biomaterials had superior performance compared to CAF alone, for PD, KGW, CAL, RW, and RH parameters. These results are similar to those found by several systematic reviews aimed at evaluating the efficacy of RCP (Francesco Cairo et al., 2014; Roman et al., 2013; Tavelli, Asa'ad, et al., 2018).

One of the objectives of this article was to evaluate the effect of time on gingival recessions using the CAF as a flap design and comparing it with other biomaterials. Although it was found that the CAF + CTG and CAF + ADM + PRP approaches showed the best results in time for the percentage of root coverage, the CAF + CTG approach showed a greater difference in relation to the other approaches. These results are similar to those reported by other authors (Rasperini et al., 2018; Cairo et al., 2014; Dai et al., 2019) where they found that CAF + CTG have a tendency to displace gingival margin coronally, while CAF alone had a tendency towards apical relapse. It is





FIGURE 7 Inconsistency plots for Keratinized tissue width; Percentage of Root coverage and Recession Height

FIGURE 8 Predictive interval and confidence interval plot for percentage of root coverage



reported that, due to biological filler content of CTG, it has the ability to adapt flap on the root surface (Francesco Cairo et al., 2016) and increases the marginal thickness of soft tissue. This enables greater chance of achieving root coverage (Rebele et al., 2014). This is fundamentally crucial also for the stability of the gingival margin, since an increase in the thickness of the gingival tissue after a CTG has been associated with the effect of progressive adhesion

over the years (Pini-Prato et al., 2010; Rasperini et al., 2018). Furthermore, it is also similar to that reported by Chambrone et al. (2019)) and Mehta et al. (2019)), where authors mention that there is, evidence suggesting that ADMs appear as the soft tissue surrogate that can provide the most similar results to those achieved by CTG for single or multiple recessions (Lee et al., 2002). On contrary, Leknes et al. (2005)) did not find any difference in




FIGURE 9 SUCRA ranking; 3E: Multi-dimensional scale ranking (MDS) for percentage of root coverage

TABLE 4 GRADE analysis

Comparison	Network meta-analysis (quality evidence)					Overall
	Pocket depth	Keratinized gingival width	Clinical attachment level	Recession width	Recession height	
CAF vs. CAF + PCM	Low	Low	Low	–	Low	Low
CAF vs. CAF + CTG	Low	Moderate	Low	Moderate	Low	Moderate
CAF vs. CAF + EMD	Low	Low	Moderate	Moderate	Low	Moderate
CAF vs. CAF + XADM	Low	Low	Low	Moderate	Low	Moderate
CAF vs. CAF + PRF	Low	Low	Moderate	Low	Low	Moderate
CAF vs. CAF + CMX	Moderate	Moderate	Moderate	Low	Low	Moderate
CAF vs. CAF + rhPDGF-BB + TCP	Moderate	Moderate	Moderate	–	Moderate	Moderate
CAF vs. CAF + ADM	Moderate	Moderate	Moderate	–	Moderate	Moderate
CAF vs. CAF + GTR(Bio)	–	Moderate	–	–	–	Moderate
CAF vs. CAF + GTR(Non Bio)	–	Moderate	–	–	–	Moderate
CAF vs. CAF + CM	Moderate	Moderate	Moderate	–	Low	Moderate
Overall						Moderate

time intervals between CAF and GTR for root coverage. It is suggested that, a strict oral hygiene maintenance after each appointment, after the root coverage procedure was recommended (McGuire et al., 2014; Pini Prato et al., 2011; Zucchelli et al., 2018).

The importance of at least 2 mm KTW has been demonstrated as an important factor for the stability of the gingival margin over time (Pini Prato, Magnani, et al., 2018). Furthermore, it has also been suggested that KTW plays a crucial role in facilitating long-term maintenance of the patients themselves and reducing the risk of soft tissue relapse (Stefanini et al., 2018; Zucchelli et al., 2014). In our analysis, we found that KTW was a significant predictor that greatly affected treatment slopes, which is also mentioned by Tavelli, Barootchi, et al. (2019) and Tavelli, Ravidà, et al. (2019).

Among treatment approaches, CAF + PCM exhibited positive slopes for KTW increase in future recessions. A possible explanation could be the potential of this material to increase the width of the keratinized tissue (Yu et al., 2018). Despite all this, our results confirm that a CTG was the best treatment to increase KTW over time.

This SR included only data from RCTs, analyzing the best available evidence where different biomaterials were used as a complement to CAF (PCM, EMD, XADM, PRF, CMX, rhPDGF-BB + TCP, ADM, GTR (Bio), GTR (Non Bio), and CM). Furthermore, studies where there were smokers were not included, as smokers may have greater gingival margin instability than non-smokers (Raes et al., 2015) due to ecological, immunological and vascular deficiencies caused by tobacco use (Palmer et al., 2005).

Study	Risk of bias domains							Overall
	D1	D2	D3	D4	D5	D6	D7	
Barakat H et al. (2020)	+	+	-	×	+	+	+	×
Rotundo R et al (2020)	+	+	+	×	+	+	+	-
Aydinyurt HS et al (2019)	+	+	+	×	+	×	+	×
Pilloni A et al (2019)	+	+	+	+	+	+	+	+
Rotundo R et al (2019)	+	+	+	+	+	+	+	+
França-Grohmann IL et al (2018)	+	+	+	+	+	+	+	+
Gürlek Ö et al (2019)	-	+	+	×	+	+	+	×
Match U et al (2019)	+	+	+	×	+	+	+	-
Francetti L et al (2018)	×	+	+	+	+	+	+	-
Kuka S et al (2018)	+	+	-	+	+	+	+	-
Rasperini G et al (2018)	+	+	+	+	+	+	+	+
Çetiner D et al (2018)	+	+	+	+	+	+	+	+
Abou-Arraj RV et al (2017)	-	+	-	+	+	+	+	×
Jepsen K et al (2017)	+	+	-	-	+	+	+	×
Cairo F et al (2016)	+	+	+	+	+	+	+	+
McGuire MK et al (2016)	+	+	+	+	+	+	+	+
Lalasa Godavarthi et.al (2016)	+	+	×	+	+	+	+	-
Stefanini M et al (2016)	+	+	+	+	+	+	+	+
Lops D et al (2015)	+	+	+	×	+	+	+	-
Cairo F et al (2015)	+	+	+	+	+	+	+	+
Milinkovic I et al (2015)	+	+	-	-	+	+	+	×
McGuire MK et al (2014)	-	+	+	×	+	+	+	×
Ahmedbeyli C et al (2014)	+	+	+	×	+	+	+	-
Zucchelli G et al (2014)	+	+	+	+	+	+	+	+
Cardaropoli D et al (2014)	+	+	+	+	+	+	+	+
Alkan EA et al (2013)	+	+	-	×	+	+	+	×
Kuis D et al (2013)	+	+	+	×	×	+	+	×
Koseoglu S et al (2013)	+	+	+	-	+	+	+	-
Roman A et al (2013)	+	+	+	+	+	+	+	+
Kumar G et al (2020)	-	+	×	+	+	+	+	×
Cordaro L et al (2012)	-	+	-	+	+	+	+	×
Cardaropoli D et al (2012)	+	+	-	+	+	+	+	-
Alkan EA et al (2011)	-	+	×	×	+	+	+	×
Jankovic S et al (2010)	+	+	-	+	+	×	+	×
Del Pizzo M et al (2005)	-	+	×	+	+	+	+	×
Spahr A et al (2005)	+	+	×	+	+	+	+	-
McGuire MK et al (2003)	+	+	+	×	+	-	+	×
Hagewald S et al (2002)	+	+	-	+	+	+	+	-
Zucchelli G et al (1998)	+	-	+	-	+	+	+	×

Domains:
D1: Random allocation
D2: Inclusion/exclusion criteria clearly defined
D3: Blinding of participants
D4: Blinding of examiners
D5: Balanced experimental groups
D6: Identical treatment between the groups
D7: Reporting of follow-up

Judgement
× Serious
- Moderate
+ Low

FIGURE 10 Risk of bias summary for all included studies

Some weaknesses of this study should be highlighted. First, most of the studies presented a high or moderate risk of bias, which increases the inconsistency of the results, leading to the fact that in the comparisons between CAF only with CAF with a biomaterial they presented a moderate GRADE analysis, thus decreasing, the recommendation of clinical results.

Another problem could be the differences in the process of making platelet concentrates. Variations in centrifuge type, speed, and G-force could change membrane patterns and, consequently, the number of growth factors and cytokines (Miron et al., 2019). Furthermore, the limited research and high risk of bias in these studies, mentioned by Li et al. (2019) and Moraschini and Barboza (2016)), can make the interpretation of the results difficult.

Among the limitations of the literature, we observed that included RCTs provided no information regarding gingival phenotype of the patient at the start of the study or at follow-up intervals. Gingival phenotype was suggested to play a key role in determining future graft procedures, and this could not be explored with NMA. There was no significant added information from the analysis about gingival thickness (i.e., gingival thickness \geq 0.8 mm or 1.2 mm has shown to be associated with greater chance of complete root coverage) (Baldi et al., 1999; Cairo et al., 2016). In addition, due to the lack of individual patient data, the impact of age and gender on the stability of the results was not investigated. However, in a recent article, age and gender were not found to be relevant factors in maintaining the stability of the gingival margin (Rasperini et al., 2018).

5 | IMPLICATIONS FOR CLINICAL PRACTICE AND FUTURE DIRECTIONS

Our SR and NMA found that ADM + PRP and PCM have the better clinical performance as an adjunct to CAF, for the percentage of root coverage and KTW, respectively, in the treatment of Miller's class I and II gingival recessions (Cairo RT I). Based on the ranking of biomaterials, clinician will be able to make informed decisions in daily clinical practice. Standardization of the methods for using these biomaterials is essential to ensure that results are reproducible and predictable for monitoring long-term tissue stability and behavior.

6 | CONCLUSION

Our NMA found that CAF + CTG and CAF + PCM for KGW and CAF + CM + GF AND CAF + ADM + PRP for percentage of root coverage were ranked higher and would perform better in future clinical studies. The highest ranked material in improving CAL was CAF + ADM and CAF + PRF. In conclusion, CTG, ADM, and CM along with GFs showed improved stability for \geq 12 months follow-up, when used in adjunct to CAF in terms of better percentage of root coverage and improved KGW. However, the overall evidence was moderate and therefore, well designed clinical trials are needed.

CONFLICT OF INTEREST

The authors declare no conflicts of interest related to the contents of this study.

AUTHORS CONTRIBUTION

Concepts, design, definition of intellectual content, literature search, clinical studies, experimental studies, data acquisition, data analysis, statistical analysis, manuscript preparation, manuscript editing, manuscript review, and guarantor: Sourav Panda, Shahnawaz Khijmatgar, Heber Arbildo-Vega, Abhaya Chandra Das, Manoj Kumar, Mohit Das, Leonardo Mancini, and Massimo Del Fabbro.

SOURCES OF FUNDING STATEMENT

No external funding, apart from the support of the authors' institution, was available for this study.

DATA AVAILABILITY STATEMENT

Data sharing not applicable - no new data generated, or the article describes entirely theoretical research.

ORCID

Sourav Panda  <https://orcid.org/0000-0002-0198-748X>

Shahnawaz Khijmatgar  <https://orcid.org/0000-0002-9177-3505>

Heber Arbildo-Vega  <https://orcid.org/0000-0003-3689-7502>

Abhaya Chandra Das  <https://orcid.org/0000-0001-7832-3406>

Manoj Kumar  <https://orcid.org/0000-0002-8775-0589>

Mohit Das  <https://orcid.org/0000-0001-8224-5500>

Leonardo Mancini  <https://orcid.org/0000-0002-7030-155X>

Massimo Del Fabbro  <https://orcid.org/0000-0001-7144-0984>

REFERENCES

- Abou-Arraj, R. V., Kaur, M., Vassilopoulos, P. J., & Geurs, N. C. (2017). Creation of a zone of immobile connective tissue with acellular dermal matrix allografts. *International Journal of Periodontics & Restorative Dentistry*, 37(4), 571–579.
- Ahmedbeyli, C., İpçi, Ş. D., Cakar, G., Kuru, B. E., & Yılmaz, S. (2014). Clinical evaluation of coronally advanced flap with or without acellular dermal matrix graft on complete defect coverage for the treatment of multiple gingival recessions with thin tissue biotype. *Journal of Clinical Periodontology*, 41(3), 303–310.
- Aleksić, Z., Janković, S., Dimitrijević, B., Divnić-Resnik, T., Milinković, I., & Leković, V. (2010). The use of platelet-rich fibrin membrane in gingival recession treatment. *Srpski arhiv za celokupno lekarstvo*, 138(1-2), 11–18.
- Alkan, E. A., & Parlar, A. (2011). EMD or subepithelial connective tissue graft for the treatment of single gingival recessions: A pilot study. *Journal of Periodontal Research*, 46(6), 637–642.
- Alkan, E. A., & Parlar, A. (2013). Enamel matrix derivative (emdogain) or subepithelial connective tissue graft for the treatment of adjacent multiple gingival recessions: A pilot study. *International Journal of Periodontics & Restorative Dentistry*, 33(5), 619–625.
- Aroca, S., Molnár, B., Windisch, P., Gera, I., Salvi, G. E., Nikolidakis, D., & Sculean, A. (2013). Treatment of multiple adjacent Miller class I and II gingival recessions with a Modified Coronally Advanced Tunnel (MCAT) technique and a collagen matrix or palatal connective tissue graft: a randomized, controlled clinical trial. *Journal of clinical periodontology*, 40(7), 713–720.

- Aroca, S., Keglevich, T., Barbieri, B., Gera, I., & Etienne, D. (2009). Clinical evaluation of a modified coronally advanced flap alone or in combination with a platelet-rich fibrin membrane for the treatment of adjacent multiple gingival recessions: A 6-month study. *Journal of periodontology*, *80*(2), 244–252.
- Aydinyurt, H. S., Tekin, Y., & Ertugrul, A. S. (2019). The effect of enamel matrix derivatives on root coverage: A 12-month follow-up of a randomized clinical trial. *Brazilian Oral Research*, *33*, e006.
- Baldi, C., Pini-Prato, G., Pagliaro, U., Nieri, M., Saletta, D., Muzzi, L., & Cortellini, P. (1999). Coronally advanced flap procedure for root coverage. Is flap thickness a relevant predictor to achieve root coverage? A 19-case series. *Journal of Periodontology*, *70*(9), 1077–1084.
- Barakat, H., & Dayoub, S. (2020). Treatment of miller type I and II gingival recession defects using three-dimensional porcine collagen matrix with coronally advanced flap: A randomized clinical split-mouth trial (a 1-year follow-up). *Indian Journal of Dental Research*, *31*(2), 209–216.
- Barootchi, S., Tavelli, L., Zucchelli, G., Giannobile, W. V., & Wang, H. L. (2020). Gingival phenotype modification therapies on natural teeth: A network meta-analysis. *Journal of Periodontology*, *91*(11), 1386–1399.
- Bellver-Fernández, R., Martínez-Rodríguez, A. M., Gioia-Palavecino, C., Caffesse, R. G., & Peñarocha, M. (2016). Surgical treatment of localized gingival recessions using coronally advanced flaps with or without subepithelial connective tissue graft. *Medicina oral, patología oral y cirugía bucal*, *21*(2), p.e222.
- Buti, J., Baccini, M., Nieri, M., La Marca, M., & Pini-Prato, G. P. (2013). Bayesian network meta-analysis of root coverage procedures: Ranking efficacy and identification of best treatment. *Journal of Clinical Periodontology*, *40*(4), 372–386.
- Cairo, F., Cortellini, P., Pilloni, A., Nieri, M., Cincinelli, S., Amunni, F., Pagavino, G., & Tonetti, M. S. (2016). Clinical efficacy of coronally advanced flap with or without connective tissue graft for the treatment of multiple adjacent gingival recessions in the aesthetic area: A randomized controlled clinical trial. *Journal of Clinical Periodontology*, *43*(10), 849–856.
- Cairo, F., Cortellini, P., Tonetti, M., Nieri, M., Mervelt, J., Pagavino, G., & Pini-Prato, G. P. (2015). Stability of root coverage outcomes at single maxillary gingival recession with loss of interdental attachment: 3-year extension results from a randomized, controlled, clinical trial. *Journal of Clinical Periodontology*, *42*(6), 575–581.
- Cairo, F., Nieri, M., & Pagliaro, U. (2014). Efficacy of periodontal plastic surgery procedures in the treatment of localized facial gingival recessions. A systematic review. *Journal of clinical periodontology*, *41*, S44–S62.
- Cardaropoli, D., Tamagnone, L., Roffredo, A., & Gaviglio, L. (2012). Treatment of gingival recession defects using coronally advanced flap with a porcine collagen matrix compared to coronally advanced flap with connective tissue graft: A randomized controlled clinical trial. *Journal of Periodontology*, *83*(3), 321–328.
- Cardaropoli, D., Tamagnone, L., Roffredo, A., & Gaviglio, L. (2014). Coronally advanced flap with and without a xenogenic collagen matrix in the treatment of multiple recessions: A randomized controlled clinical study. *International Journal of Periodontics & Restorative Dentistry*, *34*, s97–s102.
- Çetiner, D., Kalabay, P. G., Özdemir, B., & Çankaya, Z. T. (2018). Efficiency of platelet-rich plasma on acellular dermal matrix application with coronally advanced flap in the treatment of multiple adjacent gingival recessions: A randomized controlled clinical trial. *Journal of dental sciences*, *13*(3), 198–206.
- Chambrone, L., Ortega, M. A. S., Sukekava, F., Rotundo, R., Kalemaj, Z., Buti, J., & Prato, G. P. P. (2019). Root coverage procedures for treating single and multiple recession-type defects: An updated Cochrane systematic review. *Journal of Periodontology*, *90*(12), 1399–1422.
- Chambrone, L., & Tatakis, D. N. (2015). Periodontal soft tissue root coverage procedures: a systematic review from the AAP Regeneration Workshop. *Journal of periodontology*, *86*, S8–S51.
- Cheng, Y. F., Chen, J. W., Lin, S. J., & Lu, H. K. (2007). Is coronally positioned flap procedure adjunct with enamel matrix derivative or root conditioning a relevant predictor for achieving root coverage? A systematic review. *Journal of Periodontal Research*, *42*(5), 474–485.
- Cordaro, L., di Torresanto, T. V. M., & Torsello, F. (2012). Split-mouth comparison of a Coronally advanced flap with or without enamel matrix derivative for coverage of multiple gingival recession defects: 6-and 24-month follow-up-online. *International Journal of Periodontics and Restorative Dentistry*, *32*(1), 59.
- Cortellini, P., & Bissada, N. F. (2018). Mucogingival conditions in the natural dentition: Narrative review, case definitions, and diagnostic considerations. *Journal of Periodontology*, *89*, S204–S213.
- Cortellini, P., Buti, J., Pini Prato, G., & Tonetti, M. S. (2017). Periodontal regeneration compared with access flap surgery in human intra-bony defects 20-year follow-up of a randomized clinical trial: Tooth retention, periodontitis recurrence and costs. *Journal of Clinical Periodontology*, *44*(1), 58–66.
- Dai, A., Huang, J. P., Ding, P. H., & Chen, L. L. (2019). Long-term stability of root coverage procedures for single gingival recessions: A systematic review and meta-analysis. *Journal of Clinical Periodontology*, *46*(5), 572–585.
- Del Pizzo, M., Zucchelli, G., Modica, F., Villa, R., & Debernardi, C. (2005). Coronally advanced flap with or without enamel matrix derivative for root coverage: A 2-year study. *Journal of Clinical Periodontology*, *32*(11), 1181–1187.
- Dohan, D. M., Choukroun, J., Diss, A., Dohan, S. L., Dohan, A. J., Mouhyi, J., & Gogly, B. (2006). Platelet-rich fibrin (PRF): A second-generation platelet concentrate. Part II: Platelet-related biologic features. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, *101*(3), e45–e50.
- Dominiak, M., Kalecińska, E., Krzysztosiński, E., Dominiak, P., Mierzwa-Dudek, D., & Sulka, A. (2006). Correlation between temporomandibular dysfunction, disturbances of occlusion and gingival recession in a group of youth students. *Bulletin du Groupement international pour la recherche scientifique en stomatologie & odontologie*, *47*(1), 40–46.
- França-Grohmann, I. L., Sangiorgio, J. P. M., Bueno, M. R., Casarin, R. C. V., Silvério, K. G., Nociti, F. H., Jr., Casati, M. Z., & Sallum, E. A. (2019). Does enamel matrix derivative application improve clinical outcomes after semilunar flap surgery? A randomized clinical trial. *Clinical Oral Investigations*, *23*(2), 879–887.
- Francetti, L., Weinstein, R., Taschieri, S., & Corbella, S. (2018). Coronally advanced flap with or without subepithelial connective tissue graft for the treatment of single recession: 5-year outcomes from a comparative study. *International Journal of Periodontics & Restorative Dentistry*, *38*(6), 819–825.
- Fu, J. H., Su, C. Y., & Wang, H. L. (2012). Esthetic soft tissue management for teeth and implants. *Journal of Evidence Based Dental Practice*, *12*(3), 129–142.
- Godavarthi, L., Murthy, K. R., & Pavankumar, S. (2016). A comparison of acellular dermal matrix allograft and periosteal pedicle graft covered by coronally advanced flap in the treatment of gingival recession: 1-year follow-up study. *The International Journal of Periodontics & Restorative Dentistry*, *36*(4), e67–e75.
- Gürlek, Ö., Gümüş, P., Nizam, N., & Buduneli, N. (2020). Coronally advanced flap with connective tissue graft or xenogeneic acellular dermal matrix in the treatment of multiple gingival recessions: A split-mouth randomized clinical trial. *Journal of Esthetic and Restorative Dentistry*, *32*(4), 380–388.
- Gurtner, G. C., Werner, S., Barrandon, Y., & Longaker, M. T. (2008). Wound repair and regeneration. *Nature*, *453*(7193), 314–321.

- Hägeward, S., Spahr, A., Rompolo, E., Haller, B., Heijl, L., & Bernimoulin, J. P. (2002). Comparative study of Emdogain® and coronally advanced flap technique in the treatment of human gingival recessions: A prospective controlled clinical study. *Journal of Clinical Periodontology*, 29(1), 35–41.
- Henriques, P. S., Pelegrine, A. A., Nogueira, A. A., & Borghi, M. M. (2010). Application of subepithelial connective tissue graft with or without enamel matrix derivative for root coverage: a split-mouth randomized study. *Journal of oral science*, 52(3), 463–471.
- Jankovic, S., Zoran, A., Iva, M., & Bozidar, D. (2010). The coronally advanced flap in combination with platelet-rich fibrin (PRF) and enamel matrix derivative in the treatment of gingival recession: A comparative study. *The European Journal of Esthetic Dentistry*, 5(3), 260–273. <https://pubmed.ncbi.nlm.nih.gov/20820456/>.
- Jepsen, K., Stefanini, M., Sanz, M., Zucchelli, G., & Jepsen, S. (2017). Long-term stability of root coverage by coronally advanced flap procedures. *Journal of Periodontology*, 88(7), 626–633.
- Keceli, H. G., Kamak, G., Erdemir, E. O., Evginer, M. S., & Dolgun, A. (2015). The adjunctive effect of platelet-rich fibrin to connective tissue graft in the treatment of buccal recession defects: Results of a randomized, parallel-group controlled trial. *Journal of Periodontology*, 86(11), 1221–1230.
- Köseoğlu, S., Duran, İ., Sağlam, M., Bozkurt, S. B., Kırtıoğlu, O. S., & Hakkı, S. S. (2013). Efficacy of collagen membrane seeded with autologous gingival fibroblasts in gingival recession treatment: A randomized, controlled pilot study. *Journal of Periodontology*, 84(10), 1416–1424.
- Kuis, D., Sciran, I., Lajnert, V., Snjarić, D., Prpic, J., Pezelj-Ribaric, S., & Bosnjak, A. (2013). Coronally advanced flap alone or with connective tissue graft in the treatment of single gingival recession defects: A long-term randomized clinical trial. *Journal of Periodontology*, 84(11), 1576–1585.
- Kuka, S., Ipci, S. D., Cakar, G., & Yılmaz, S. (2018). Clinical evaluation of coronally advanced flap with or without platelet-rich fibrin for the treatment of multiple gingival recessions. *Clinical Oral Investigations*, 22(3), 1551–1558.
- Kumar, G. N. V., & Murthy, K. R. V. (2013). A comparative evaluation of subepithelial connective tissue graft (SCTG) versus platelet concentrate graft (PCG) in the treatment of gingival recession using coronally advanced flap technique: A 12-month study. *Journal of Indian Society of Periodontology*, 17(6), 771–776.
- Lang, N. P., & Löe, H. (1972). The relationship between the width of keratinized gingiva and gingival health. *Journal of Periodontology*, 43(10), 623–627.
- Lee, E. J., Meraw, S. J., Oh, T. J., Giannobile, W. V., & Wang, H. L. (2002). Comparative histologic analysis of coronally advanced flap with and without collagen membrane for root coverage. *Journal of Periodontology*, 73(7), 779–788.
- Leknes, K. N., Amarante, E. S., Price, D. E., Bøe, O. E., Skavland, R. J., & Lie, T. (2005). Coronally positioned flap procedures with or without a biodegradable membrane in the treatment of human gingival recession: A 6-year follow-up study. *Journal of Clinical Periodontology*, 32(5), 518–529.
- Li, R., Liu, Y., Xu, T., Zhao, H., Hou, J., Wu, Y., & Zhang, D. (2019). The additional effect of autologous platelet concentrates to coronally advanced flap in the treatment of gingival recessions: A systematic review and meta-analysis. *BioMed Research International*, 2019, 2587245.
- Lops, D., Gobbato, L., Nart, J., Guazzo, R., Ho, D. K., & Bressan, E. (2015). Evaluation of root coverage with and without connective tissue graft for the treatment of single maxillary gingival recession using an image analysis system: A randomized controlled clinical trial. *International Journal of Periodontics & Restorative Dentistry*, 35(2), 247–254.
- Matoh, U., Petelin, M., & Gašperšič, R. (2019). Split-mouth comparison of Coronally advanced flap with connective tissue graft or collagen matrix for treatment of isolated gingival recessions. *International Journal of Periodontics & Restorative Dentistry*, 39(3), 439–446.
- McGuire, M. K., & Nunn, M. (2003). Evaluation of human recession defects treated with coronally advanced flaps and either enamel matrix derivative or connective tissue. Part 1: Comparison of clinical parameters. *Journal of Periodontology*, 74(8), 1110–1125.
- McGuire, M. K., & Scheyer, E. T. (2010). Xenogeneic collagen matrix with coronally advanced flap compared to connective tissue with coronally advanced flap for the treatment of dehiscence-type recession defects. *Journal of Periodontology*, 81(8), 1108–1117.
- McGuire, M. K., & Scheyer, E. T. (2016). Long-term results comparing xenogeneic collagen matrix and autogenous connective tissue grafts with coronally advanced flaps for treatment of dehiscence-type recession defects. *Journal of Periodontology*, 87(3), 221–227.
- McGuire, M. K., Scheyer, E. T., & Nunn, M. (2012). Evaluation of human recession defects treated with coronally advanced flaps and either enamel matrix derivative or connective tissue: Comparison of clinical parameters at 10 years. *Journal of Periodontology*, 83(11), 1353–1362.
- McGuire, M. K., Scheyer, E. T., & Snyder, M. B. (2014). Evaluation of recession defects treated with coronally advanced flaps and either recombinant human platelet-derived growth factor-BB plus β -tricalcium phosphate or connective tissue: Comparison of clinical parameters at 5 years. *Journal of Periodontology*, 85(10), 1361–1370.
- McGuire, M. K., Tavelli, L., Feinberg, S. E., Rasperini, G., Zucchelli, G., Wang, H. L., & Giannobile, W. V. (2020). Living cell-based regenerative medicine technologies for periodontal soft tissue augmentation. *Journal of Periodontology*, 91(2), 155–164.
- Mehta, V., Bagwe, S., Gopalakrishnan, D., & Mathu, A. (2019). Comparison of effectiveness between acellular dermal matrix allograft and subepithelial connective tissue graft with Coronally advanced flap in recession treatment: A systematic review. *Acta Scientific Dental Sciences*, 3(10), 2–7.
- Milinkovic, I., Aleksic, Z., Jankovic, S., Popovic, O., Bajic, M., Cakic, S., & Lekovic, V. (2015). Clinical application of autologous fibroblast cell culture in gingival recession treatment. *Journal of Periodontal Research*, 50(3), 363–370.
- Miron, R. J., Moraschini, V., Del Fabbro, M., Piattelli, A., Fujioka-Kobayashi, M., Zhang, Y., Saulacic, N., Schaller, B., Kawase, T., Cosgarea, R., & Jepsen, S. (2020). Use of platelet-rich fibrin for the treatment of gingival recessions: A systematic review and meta-analysis. *Clinical Oral Investigations*, 24(8), 2543–2557.
- Miron, R. J., Pinto, N. R., Quirynen, M., & Ghanaati, S. (2019). Standardization of relative centrifugal forces in studies related to platelet-rich fibrin. *Journal of Periodontology*, 90(8), 817–820.
- Moraschini, V., & Barboza, E. D. S. P. (2016). Use of platelet-rich fibrin membrane in the treatment of gingival recession: A systematic review and meta-analysis. *Journal of Periodontology*, 87(3), 281–290.
- Moraschini, V., Calasans-Maia, M. D., Dias, A. T., de Carvalho Formiga, M., Sartoretto, S. C., Sculean, A., & Shibli, J. A. (2020). Effectiveness of connective tissue graft substitutes for the treatment of gingival recessions compared with coronally advanced flap: A network meta-analysis. *Clinical Oral Investigations*, 24(10), 3395–3406.
- Moraschini, V., de Almeida, D. C. F., Sartoretto, S., Bailly Guimarães, H., Chaves Cavalcante, I., & Diuana Calasans-Maia, M. (2019). Clinical efficacy of xenogeneic collagen matrix in the treatment of gingival recession: A systematic review and meta-analysis. *Acta Odontologica Scandinavica*, 77(6), 457–467.
- Moses, O., Artzi, Z., Sculean, A., Tal, H., Kozlovsky, A., Romanos, G. E., & Nemcovsky, C. E. (2006). Comparative study of two root coverage procedures: A 24-month follow-up multicenter study. *Journal of periodontology*, 77(2), 195–202.
- Moslemi, N., Mousavi Jazi, M., Haghghati, F., Morovati, S. P., & Jamali, R. (2011). Acellular dermal matrix allograft versus subepithelial connective tissue graft in treatment of gingival recessions: A 5-year randomized clinical study. *Journal of Clinical Periodontology*, 38(12), 1122–1129.

- Nemcovsky, C. E., Artzi, Z., Tal, H., Kozlovsky, A., & Moses, O. (2004). A multi-center comparative study of two root coverage procedures: coronally advanced flap with addition of enamel matrix proteins and subpedicle connective tissue graft. *Journal of periodontology*, 75(4), 600–607.
- Palmer, R. M., Wilson, R. F., Hasan, A. S., & Scott, D. A. (2005). Mechanisms of action of environmental factors-tobacco smoking. *Journal of Clinical Periodontology*, 32(Suppl 6), 180–195.
- Pilloni, A., Schmidlin, P. R., Sahrman, P., Sculean, A., & Rojas, M. A. (2019). Effectiveness of adjunctive hyaluronic acid application in coronally advanced flap in miller class I single gingival recession sites: A randomized controlled clinical trial. *Clinical Oral Investigations*, 23(3), 1133–1141.
- Pini Prato, G., Rotundo, R., Franceschi, D., Cairo, F., Cortellini, P., & Nieri, M. (2011). Fourteen-year outcomes of coronally advanced flap for root coverage: Follow-up from a randomized trial. *Journal of Clinical Periodontology*, 38(8), 715–720.
- Pini Prato, G. P., Franceschi, D., Cortellini, P., & Chambrone, L. (2018). Long-term evaluation (20 years) of the outcomes of subepithelial connective tissue graft plus coronally advanced flap in the treatment of maxillary single recession-type defects. *Journal of Periodontology*, 89(11), 1290–1299.
- Pini Prato, G. P., Magnani, C., & Chambrone, L. (2018). Long-term evaluation (20 years) of the outcomes of coronally advanced flap in the treatment of single recession-type defects. *Journal of Periodontology*, 89(3), 265–274.
- Pini-Prato, G. P., Cairo, F., Nieri, M., Franceschi, D., Rotundo, R., & Cortellini, P. (2010). Coronally advanced flap versus connective tissue graft in the treatment of multiple gingival recessions: A split-mouth study with a 5-year follow-up. *Journal of Clinical Periodontology*, 37(7), 644–650.
- Pourabbas, R., Chitsazi, M. T., Kosarieh, E., & Olyae, P. (2009). Coronally advanced flap in combination with acellular dermal matrix with or without enamel matrix derivatives for root coverage. *Indian Journal of Dental Research*, 20(3), 320.
- Raes, S., Rocci, A., Raes, F., Cooper, L., De Bruyn, H., & Cosyn, J. (2015). A prospective cohort study on the impact of smoking on soft tissue alterations around single implants. *Clinical Oral Implants Research*, 26(9), 1086–1090.
- Rasperini, G., Acunzo, R., Pellegrini, G., Pagni, G., Tonetti, M., Pini Prato, G. P., & Cortellini, P. (2018). Predictor factors for long-term outcomes stability of coronally advanced flap with or without connective tissue graft in the treatment of single maxillary gingival recessions: 9 years results of a randomized controlled clinical trial. *Journal of Clinical Periodontology*, 45(9), 1107–1117.
- Rebele, S. F., Zuhr, O., Schneider, D., Jung, R. E., & Hürzeler, M. B. (2014). Tunnel technique with connective tissue graft versus coronally advanced flap with enamel matrix derivative for root coverage: A RCT using 3D digital measuring methods. Part II. Volumetric studies on healing dynamics and gingival dimensions. *Journal of Clinical Periodontology*, 41(6), 593–603.
- Rios, F. S., Costa, R. S. A., Moura, M. S., Jardim, J. J., Maltz, M., & Haas, A. N. (2014). Estimates and multivariable risk assessment of gingival recession in the population of adults from Porto Alegre, Brazil. *Journal of Clinical Periodontology*, 41(11), 1098–1107.
- Roman, A., Soancă, A., Kasaj, A., & Stratul, S. I. (2013). Subepithelial connective tissue graft with or without enamel matrix derivative for the treatment of miller class I and II gingival recessions: A controlled randomized clinical trial. *Journal of Periodontal Research*, 48(5), 563–572.
- Rotundo, R., Genzano, L., Nieri, M., Covani, U., Peñarrocha-Oltra, D., & Peñarrocha-Diago, M. (2021). Smile esthetic evaluation of mucogingival reconstructive surgery. *Odontology*, 109(1), 295–302.
- Rotundo, R., Genzano, L., Patel, D., D'Aiuto, F., & Nieri, M. (2019). Adjunctive benefit of a xenogenic collagen matrix associated with coronally advanced flap for the treatment of multiple gingival recessions: A superiority, assessor-blind, randomized clinical trial. *Journal of Clinical Periodontology*, 46(10), 1013–1023.
- Santamaria, M. P., Neves, F. L. D. S., Silveira, C. A., Mathias, I. F., Fernandes-Dias, S. B., Jardim, M. A. N., & Tatakis, D. N. (2017). Connective tissue graft and tunnel or trapezoidal flap for the treatment of single maxillary gingival recessions: A randomized clinical trial. *Journal of Clinical Periodontology*, 44(5), 540–547.
- Smith, P. C., Cáceres, M., Martínez, C., Oyarzún, A., & Martínez, J. (2015). Gingival wound healing: An essential response disturbed by aging? *Journal of Dental Research*, 94(3), 395–402.
- Spahr, A., Haegewald, S., Tsoulfidou, F., Rompola, E., Heijl, L., Bernimoulin, J. P., Ring, C., Sander, S., & Haller, B. (2005). Coverage of miller class I and II recession defects using enamel matrix proteins versus coronally advanced flap technique: A 2-year report. *Journal of Periodontology*, 76(11), 1871–1880.
- Stefanini, M., Jepsen, K., de Sanctis, M., Baldini, N., Greven, B., Heinz, B., Wennström, J., Cassel, B., Vignoletti, F., Sanz, M., & Jepsen, S. (2016). Patient-reported outcomes and aesthetic evaluation of root coverage procedures: A 12-month follow-up of a randomized controlled clinical trial. *Journal of Clinical Periodontology*, 43(12), 1132–1141.
- Stefanini, M., Zucchelli, G., Marzadori, M., & de Sanctis, M. (2018). Coronally advanced flap with site-specific application of connective tissue graft for the treatment of multiple adjacent gingival recessions: A 3-year follow-up case series. *International Journal of Periodontics & Restorative Dentistry*, 38(1), 25–33.
- Susin, C., Haas, A. N., Oppermann, R. V., Haugejorden, O., & Albandar, J. M. (2004). Gingival recession: Epidemiology and risk indicators in a representative urban Brazilian population. *Journal of Periodontology*, 75(10), 1377–1386.
- Tatakis, D. N., Chambrone, L., Allen, E. P., Langer, B., McGuire, M. K., Richardson, C. R., Zabalegui, I., & Zadeh, H. H. (2015). Periodontal soft tissue root coverage procedures: A consensus report from the AAP regeneration workshop. *Journal of Periodontology*, 86, S52–S55.
- Tavelli, L., Asa'ad, F., Acunzo, R., Pagni, G., Consonni, D., & Rasperini, G. (2018). Minimizing patient morbidity following palatal gingival harvesting: A randomized controlled clinical study. *The International Journal of Periodontics & Restorative Dentistry*, 38(6), e127–e134.
- Tavelli, L., Barootchi, S., Cairo, F., Rasperini, G., Shedden, K., & Wang, H. L. (2019). The effect of time on root coverage outcomes: A network meta-analysis. *Journal of Dental Research*, 98(11), 1195–1203.
- Tavelli, L., Barootchi, S., Nguyen, T. V. N., Tattan, M., Ravidà, A., & Wang, H.-L. (2018). Efficacy of tunnel technique in the treatment of localized and multiple gingival recessions: A systematic review and meta-analysis. *Journal of Periodontology*, 89(9), 1075–1090.
- Tavelli, L., McGuire, M. K., Zucchelli, G., Rasperini, G., Feinberg, S. E., Wang, H.-L., & Giannobile, W. V. (2020). Extracellular matrix-based scaffolding technologies for periodontal and peri-implant soft tissue regeneration. *Journal of Periodontology*, 91(1), 17–25.
- Tavelli, L., Ravidà, A., Saleh, M. H. A., Maska, B., Del Amo, F. S.-L., Rasperini, G., & Wang, H.-L. (2019). Pain perception following epithelialized gingival graft harvesting: A randomized clinical trial. *Clinical Oral Investigations*, 23(1), 459–468.
- Tonetti, M. S., Cortellini, P., Pellegrini, G., Nieri, M., Bonaccini, D., Allegri, M., Bouchard, P., Cairo, F., Conforti, G., Fourmousis, I., & Graziani, F. (2018). Xenogenic collagen matrix or autologous connective tissue graft as adjunct to coronally advanced flaps for coverage of multiple adjacent gingival recession: Randomized trial assessing non-inferiority in root coverage and superiority in oral health-related quality of life. *Journal of Clinical Periodontology*, 45(1), 78–88.
- Vignoletti, F., Nuñez, J., Discepoli, N., De Sanctis, F., Caffesse, R., Muñoz, F., Lopez, M., & Sanz, M. (2011). Clinical and histological healing of a new collagen matrix in combination with the coronally advanced flap for the treatment of miller class-I recession defects: An experimental study in the minipig. *Journal of Clinical Periodontology*, 38(9), 847–855.

- Wang, H. L., Romanos, G. E., Geurs, N. C., Sullivan, A., Suárez-López del Amo, F., & Eber, R. M. (2014). Comparison of two differently processed acellular dermal matrix products for root coverage procedures: a prospective, randomized multicenter study. *Journal of periodontology*, 85(12), 1693–1701.
- Wang, H. L., Suarez-Lopez Del Amo, F., Layher, M., & Eber, R. (2015). Comparison of freeze-dried and solvent-dehydrated acellular dermal matrix for root coverage: a randomized controlled trial. *Int J Periodontics Restorative Dent*, 35(6), 811–817.
- Wennström, J. L. (1996). Mucogingival therapy. *Annals of periodontology*, 1(1), 671–701.
- Wu, Y. C., Lin, L. K., Song, C. J., Su, Y. X., & Tu, Y. K. (2017). Comparisons of periodontal regenerative therapies: A meta-analysis on the long-term efficacy. *Journal of Clinical Periodontology*, 44(5), 511–519.
- Yu, S.-H., Tseng, S.-C., & Wang, H.-L. (2018). Classification of soft tissue grafting materials based on biologic principles. *The International Journal of Periodontics & Restorative Dentistry*, 38(6), 849–854.
- Zucchelli, G., Clauser, C., De Sanctis, M., & Calandriello, M. (1998). Mucogingival versus guided tissue regeneration procedures in the treatment of deep recession type defects. *Journal of Periodontology*, 69(2), 138–145.
- Zucchelli, G., Felice, P., Mazzotti, C., Marzadori, M., Mounssif, I., Monaco, C., & Stefanini, M. (2018). 5-year outcomes after coverage of soft tissue dehiscence around single implants: A prospective cohort study. *European Journal of Oral Implantology*, 11(2), 215–224.
- Zucchelli, G., Mounssif, I., Mazzotti, C., Stefanini, M., Marzadori, M., Petracci, E., & Montebugnoli, L. (2014). Coronally advanced flap with and without connective tissue graft for the treatment of multiple gingival recessions: A comparative short- and long-term controlled randomized clinical trial. *Journal of Clinical Periodontology*, 41(4), 396–403.
- Zucchelli, G., Tavelli, L., McGuire, M. K., Rasperini, G., Feinberg, S. E., Wang, H. L., & Giannobile, W. V. (2020). Autogenous soft tissue grafting for periodontal and peri-implant plastic surgical reconstruction. *Journal of Periodontology*, 91(1), 9–16.
- Zuhr, O., Rebele, S. F., Schneider, D., Jung, R. E., & Hürzeler, M. B. (2014). Tunnel technique with connective tissue graft versus coronally advanced flap with enamel matrix derivative for root coverage: A RCT using 3D digital measuring methods. Part I. clinical and patient-centred outcomes. *Journal of Clinical Periodontology*, 41(6), 582–592.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Panda, S., Khijmatgar, S., Arbildo-Vega, H., Das, A. C., Kumar, M., Das, M., Mancini, L., & Del Fabbro, M. (2022). Stability of biomaterials used in adjunct to coronally advanced flap: A systematic review and network meta-analysis. *Clinical and Experimental Dental Research*, 8, 421–438. <https://doi.org/10.1002/cre2.461>