

RESEARCH

Open Access

A locus-specific database for mutations in *GDAP1* allows analysis of genotype-phenotype correlations in Charcot-Marie-Tooth diseases type 4A and 2K

Julien Cassereau^{1,2,3,4}, Arnaud Chevrollier^{1,2,5}, Dominique Bonneau^{1,2,3,5}, Christophe Verny^{1,2,3,4}, Vincent Procaccio^{1,2,3,5}, Pascal Reynier^{1,2,3,5} and Marc Ferre^{1,2,5*}

Abstract

Background: The ganglioside-induced differentiation-associated protein 1 gene (*GDAP1*), which is involved in the Charcot-Marie-Tooth disease (CMT), the most commonly inherited peripheral neuropathy, encodes a protein anchored to the mitochondrial outer membrane. The phenotypic presentations of patients carrying *GDAP1* mutations are heterogeneous, making it difficult to determine genotype-phenotype correlations, since the majority of the mutations have been found in only a few unrelated patients. Locus-specific databases (LSDB) established in the framework of the Human Variome Project provide powerful tools for the investigation of such rare diseases.

Methods and Results: We report the development of a publicly accessible LSDB for the *GDAP1* gene. The *GDAP1* LSDB has adopted the Leiden Open-source Variation Database (LOVD) software platform. This database, which now contains 57 unique variants reported in 179 cases of CMT, offers a detailed description of the molecular, clinical and electrophysiological data of the patients. The usefulness of the *GDAP1* database is illustrated by the finding that *GDAP1* mutations lead to primary axonal damage in CMT, with secondary demyelination in the more severe cases of the disease.

Conclusion: Findings of this nature should lead to a better understanding of the pathophysiology of CMT. Finally, the *GDAP1* LSDB, which is part of the mitodyn.org portal of databases of genes incriminated in disorders involving mitochondrial dynamics and bioenergetics, should yield new insights into mitochondrial diseases.

Keywords: inherited peripheral neuropathy, Charcot-Marie-Tooth disease (CMT), ganglioside-induced differentiation-associated protein 1 (*GDAP1*), locus-specific database (LSDB)

Background

Charcot-Marie-Tooth (CMT) disease is one of the commonest inherited peripheral neuropathies with a prevalence of about 10–30 per 100,000, depending on the country of origin [1]. CMT neuropathies are characterized clinically by progressive sensory loss, weakness, muscle atrophy, loss of deep tendon reflexes, and foot deformities. To date, about forty genes have been incriminated in CMT. Some of these genes encode mitochondrial proteins, i.e. mitofusin 2 (*MFN2*) [MIM:608507]

and the ganglioside-induced differentiation-associated protein 1 (*GDAP1*) [MIM:606598]. *GDAP1* mutations are responsible: for CMT4A [MIM:214400] [2], the most frequent recessive subtype of demyelinating CMT; for AR-CMT2 [MIM:607706] [3], the axonal recessive subtype; for CMTRIA [MIM:608340] [4], the intermediate recessive CMT; and for CMT2K [MIM:607831] [5–7], the rare dominant subtype.

The gene locus, first identified in cases of CMT4A in Tunisian families on chromosome 8q13-q21 (CMT4A), is associated with slow motor nerve conduction velocity, and hypomyelination as detected by nerve biopsy [8]. The corresponding gene, *GDAP1*, was identified by

* Correspondence: MaFerre@chu-angers.fr

¹INSERM, U771, Angers, F-49000, France

Full list of author information is available at the end of the article

Baxter et al. (2002). *GDAP1* was previously identified as one of ten cDNAs expressed in a differentiated Neuro2a mouse neuroblastoma cell line, in which cholinergic differentiation with neurite sprouting had been induced by transfection of GD3 synthase (α2,8 sialyltransferase) cDNA [9].

The *GDAP1* gene spans ~17 kilobases of genomic DNA, containing six exons and five introns. Two transcript variants of *GDAP1* have been identified. Transcript variant 1, representing the longer variant, encodes the longer isoform, designated as isoform a (358 amino acids). Transcript variant 2, encoding isoform b, contains an alternate in-frame exon, uses an alternate splice site in the 5'-coding region and a downstream start codon, compared to transcript variant 1. Isoform b (290 amino acids) has a shorter N-terminal sequence compared to isoform a. mRNA studies using RT-PCR have demonstrated the ubiquitous expression of *GDAP1*, predominantly in nervous tissues [3].

Phylogenetic and structural analyses suggest that *GDAP1* belongs to a subfamily of glutathione-S-transferases (GSTs) [3,10,11]. *GDAP1* has two typical GST domains, domain I (GST-N) at the N-terminal region and domain II (GST-C) at the C-terminal region. The existence of these domains was predicted by using bioinformatics tools [3,10] but no functional GST activity has been demonstrated so far [11,12]. It has also been shown that *GDAP1* has a single transmembrane domain (TMD) at the extremity of the C-terminal, and a hydrophobic domain (HD) in the flanking C-terminal region [10,13]. Moreover, *GDAP1* has two additional regions between amino acids 152–164 and 169–195 that are predicted to represent two helices, α4 and α5, constituting the α4–α5 loop.

Several studies have suggested that *GDAP1* plays a role in mitochondrial fission [14,15]. It was shown that the TMD plays a key role in mitochondrial targeting and mitochondrial membrane insertion [13]. Positively charged amino acids flanking the TMD at the C-terminal are involved in mitochondrial targeting and the fission function of *GDAP1*. The HD seems essential for the mitochondrial fission mediated by *GDAP1*. We have recently shown that *GDAP1* mutations in fibroblasts from CMT2K patients are associated with defective mitochondrial complex I activity [5].

The examination of the relationship between the genotype and the phenotype of the various forms of CMT disease involving *GDAP1* mutations shows that the recessive forms are generally far more severe than the dominant forms. In particular, the truncating mutations have a more severe phenotype in the recessive forms with early onset and rapid progression of the disease leading to the inability to walk by the age of 10,

generally followed by wheelchair-dependence before the end of the third decade [16].

However, the phenotypic presentations of patients carrying *GDAP1* mutations are heterogeneous. There is not only a large clinical variability concerning the ages of onset and functional disability among patients carrying the same mutations [17] but also a significant intra-familial variability [18]. Since the majority of mutations have been described in only a few unrelated patients, it is difficult to determine phenotype-genotype correlations for *GDAP1* gene mutations. The analysis of a larger number of cases of CMT involving *GDAP1* mutations, along with detailed clinical, electrophysiological and pathological data, will be required to establish a reliable relationship between the genotype and phenotype of the different forms of the disease.

The increasing importance of *GDAP1* in CMT prompted us to develop a web-based locus-specific database (LSDB) for *GDAP1* sequence variations with short publication delays. To our knowledge, the only other database collecting *GDAP1* mutations is the Inherited Peripheral Neuropathies Mutation Database (IPNMDB) [19], which includes 29 *GDAP1* mutations among the 953 mutations in 44 genes related to inherited peripheral neuropathies. Our LSDB aims at collecting all the *GDAP1* mutations incriminated in disorders involving mitochondrial dynamics and bioenergetics, with a full record of clinical, electrophysiological and biochemical data. We here describe the construction of the database, the procedure for data submission, and the presentation of the data.

Methods

We have established a locus-specific database for *GDAP1* using the Leiden Open-source Variation Database (LOVD) version 2 build 31 system [20]. This database includes a total of 20 variant items and 18 clinical patient items. Figure 1 shows a typical entry. A standardized description of the clinical and molecular items is set up using drop-down lists with predefined variables. The clinical features are based on the usual symptoms of mitochondrial diseases.

The data is openly accessible and should prove a valuable tool for clinicians and researchers alike since it will contain published as well as unpublished sequence variants. Contributors can submit their variants online to the database after registering for a login and password. This contact information is collected for reference purposes and clarification of the data submitted. The description of mutational data is confirmed using the web-based software Mutalyzer, to which the LOVD software is currently also linked [21]. Variations submitted directly to the LSDB are re-checked with these tools.

a Patient data (#0000025)	
Gender	Female
Disease	CMT2K
Age of onset	21-30 years
Age at last examination	52 years
Duration of disease	21-30 years
Additional symptoms	-
Functional disability	LL: wheelchair bound (disease duration 21-30 years)
Clinical score	-
Electroneuromyography	Sensori-motor neuropathy, Axonal, Median or ulnar nerves: MNCV > 40 m/s, Median or ulnar nerves: CMAP > 6 mV
Ocular electrophysiology	-
Histology	-
Radiology	-
Geographic origin	France
Reference	Cassereau et al. (2009)
Remarks	-
# Reported	1
Submitter	Julien Cassereau (Curator, according to published data)

b Variant data	
Allele	Maternal (confirmed)
Reported pathogenicity	Pathogenic
Concluded pathogenicity	Pathogenic
DB-ID	GDAP1_00008
DNA change	c.719G>A (View in UCSC Genome Browser)
Type	Substitution
Location	Exon
Exon	6
Affected domain	GST-C (C-term Glutathione-S-transferase)
RNA change	-
Protein	p.(Cys240Tyr)
Reference	Cassereau et al. (2009)
Technique	SEQ
Template	DNA
Tissue	Blood
Re-site	-
DNA published	-
Variant remarks	-
Frequency	-

Figure 1 Sample record in the *GDAP1* database. **a:** clinical items; and **b:** molecular items. MNCV: Motor Nervous Conduction Velocity (in m/s); CMAP: Compound Motor Action Potential (in mV); LL: lower limbs.

The *GDAP1* database reviews all the clinical and molecular data of patients carrying *GDAP1* mutations published in peer-reviewed literature, as well as unpublished contributions that may be directly submitted. While most variants can be described in terms of the

latest update of the standard nomenclature [22,23], some inaccuracies may persist because gene anomalies discovered earlier might have been named according to a convention now out of use. Eventually, the *DNA published* field of the page dedicated to each variant (Figure

1b) indicates whether the published name of the mutation has been modified by the curator. The *GDAP1* LSDB website requires absolute compliance with the rules set out below to describe sequence variants in order to provide uniform and comparable data.

The *GDAP1* mutations are described according to the *GDAP1* transcript variant 1 [RefSeq:NM_018972.2], containing 6 exons and encoding isoform a containing 358 amino acids. The nucleotide numbering of nuclear genes reflects cDNA numbering with +1 corresponding to the A of the ATG translation initiation codon in the reference sequence, according to the guidelines of the Human Genome Variation Society (HGVS). The initiation codon is codon 1.

The criteria of pathogenicity, dependent on the clinical context and molecular findings, are stated in the *Variant data* fields: *Reported pathogenicity* and *Concluded pathogenicity* (Figure 1b). Putative novel mutations detected in affected patients must segregate according to disease status and not be present in control individuals. Putative mutations are graded according to type of mutation: frameshift and nonsense mutations are considered to be pathogenic; missense mutations are described as being of *unknown pathogenicity* when detected in single families, or as *probably pathogenic* when detected in a group of families; the mutations are considered to be pathogenic when so proven by experimental evidence or detected in multiple families.

Results and discussion

Type of mutations and affected domains

To date, the *GDAP1* database contains 57 unique variants, four of which are considered as non-pathogenic sequence variants (NPSV). Pathogenic mutations, which affect the coding sequence of the gene, are more frequently found in exon 3 (17 mutations) and exon 6 (14 mutations) (Figure 2a). The most frequent mutational protein consequences are missense mutations observed in 34 of the 53 variant mutations. In addition, nine frameshifts (18%), seven nonsense (14%) and three splice-site (6%) mutations have also been reported (Figure 2b).

Whereas few mutations are recurrent, some have been frequently reported: the c.581C>G mutation (p.Ser194X) reported 38 times in Spain, Morocco and Tunisia; the c.487C>T mutation (p.Gln163X) reported 34 times, mainly in Spain; the c.715C>T mutation (p.Leu239Phe) reported 28 times, mainly in Eastern Europe; and the c.358C>T mutation (p.Arg120Trp) 22 times, mainly in Spain.

Among all the pathogenic variants mentioned in the database, 56% are located in the glutathione S-transferase (GST) domains of the protein (Figure 2c), highlighting the important role of these domains in protein function, as recently reported [24]. Among the 34

missense mutations, 21 (62%) affect GST domains: 6 in GST-N and 15 in GST-C; 5 (14.5%) affect the α 4- α 5 loop; one affects the hydrophobic domain, one affects the transmembrane domain; and 6 (17.5%) affect no particular domain of the protein. Interestingly, 5 of the 9 missense mutations involved in CMT2K are located in GST domains (p.Ser34Cys in GST-N, and p.Gln218Glu, p.Arg226Ser, p.Cys240Tyr and p.Pro274Leu in GST-C).

Similarly, 35 of the 45 patients (78%) presenting with the recessive form of the disease associated with homozygous missense mutations, harbor mutations located in GST domains (12 in GST-N and 23 in GST-C). In addition, truncating mutations located upstream of amino acid 287 affect at least one GST domain, as is the case in almost all truncating mutations reported.

Clinical data

To date, the database includes 179 patients (93 females, 75 males and 11 not specified): 47 with a dominant form CMT2K, and 126 with a recessive form (CMT4A, AR-CMT2 or CMTRIA). Six patients were unclassified, either because the authors suspected mutations on the second allele, or because the pathogenicity of the second variant was uncertain. The database shows the clinical data for all the patients reported, with the exception of 28 patients for whom the data was unavailable or insufficient for classification of the phenotype.

Clinical severity was evaluated according to the age of onset of the disease, i.e. before age 10, between ages 11–30, and after age 30, and by walking ability, i.e. independent, requiring technical assistance with a cane or a crutch, or wheelchair-bound, before or after age 30. The phenotypic data correlated with the mutational data in patients with sufficient clinical data for each item:

- age of onset according to: (1) the mode of inheritance (dominant or recessive); and (2) the domains affected in the dominant forms of CMT (GST domains, α 4- α 5 loop, transmembrane domain, hydrophobic domain or non-specific domain);

- disability in walking according to: (1) the mode of inheritance (dominant or recessive); and (2) the predicted effect at the protein level: truncating mutations or missense mutations.

Data analysis shows that patients with recessive forms of CMT, i.e. CMT4A and AR-CMT2, presented a more severe phenotype than those with the dominant form, i.e. CMT2K, with a more early onset and a more rapid evolution (Table 1). Interestingly, nineteen percent of patients with the dominant form of CMT were asymptomatic until age 30, whereas all patients with a recessive form had symptoms before the end of the adolescence. In addition, mutations affecting GST domains were responsible for a more severe phenotype probably produced by a different pathophysiological mechanism.




Figure 2 Distribution of the 53 pathogenic mutations in the *GDAP1* gene. **a:** exons involved in the mutations are shown as blue bars; the mutations in the intronic neighbourhood of the exons are shown as red bars; **b:** type; and **c:** domain. bp: base pairs; GST: glutathione S-transferase; TMD: transmembrane domain; HD: hydrophobic domain.

Almost all patients with mutations in GST domains were symptomatic after age 50 whereas 17% of patients with mutations in non-specific domains were asymptomatic after the age of 50. These findings suggest that mutations affecting GST domains are responsible for a more severe phenotype probably due to a different pathophysiological mechanism.

The course of CMT associated with *GDAP1* mutations was more rapid in the recessive form than in the dominant forms (Table 2). All AR-CMT patients required technical support for walking after age 30. In contrast, among the CMT2K patients reported in the database, 78% walked independently after age 30. Nonsense and frameshift mutations were more often associated with a

Table 1 Age of onset of CMT

	Age of onset				Total
	< 10 y	10-30 y	> 30 y	asymptomatic > 50 y	
Recessive forms	120 (98%)	3 (2%)	0	0	123
Dominant forms	11 (25%)	18 (41%)	9 (20%)	6 (14%)	44
Domain affected:					
GST	3 (33%)	2 (22%)	4 (44%)	0	9
Others	8 (23%)	16 (36%)	5 (14%)	6 (17%)	35

Genotype-phenotype correlation analysis using *GDAP1* locus-specific database data evaluating the age of onset according to mode of inheritance and affected domains in dominant forms. Values represent the number of patients for each situation; the corresponding percentages are shown in brackets. y: years; GST: glutathione S-transferase.

severe phenotype of CMT than missense mutations. Although in each case all patients needed technical assistance with a cane or a crutch for walking after age 30, the course of the disease was more rapid in patients with homozygous truncating mutations: almost all patients over 30 (93%) were wheelchair-bound *versus* 50% of patients harboring missense mutations (Table 2).

A greater number of cases of CMT disease involving *GDAP1* mutations, along with detailed clinical, electrophysiological and pathological data, will be required to determine the relationship between the genotype and phenotype of the different forms of the disease. This is precisely the aim of our database.

Electrophysiological data

Major axonal loss reduces the motor median nerve conduction velocity (MNCV) so that the cut-off value of MNCV should be adjusted according to the compound motor action potential (CMAP), such that the MNCV is

32 m/s when the CMAP is 4 mV, as proposed for acquired demyelinating neuropathies [25]. The demyelinating forms of CMT may thus be defined by an MNCV slower than 30 m/s, the axonal forms by an MNCV above 40 m/s, and the intermediate forms by an MNCV between 30-40 m/s [26]. We have therefore included these electrophysiological categories in our database in order to characterize the neuropathies associated with *GDAP1* mutations. To determine the type of neuropathy, we need to report not only the MNCV but also the CMAP reflecting the axonal loss. *GDAP1* mutations have been described as responsible for CMT4A, the demyelinating form; AR-CMT2 and CMT2K, the axonal forms; and CMTRIA, the intermediate form of CMT.

Our database contains the electrophysiological parameters for all the CMT patients reported, with the exception of 48 patients for whom the data were insufficient. Thus, our analysis concerns 131 patients among whom 30 (23%) had no recordable median nerve parameters suggesting severe axonal loss (Table 3). The majority of patients are reported to have MNCVs greater than 40 m/s, corresponding to the axonal forms of CMT and only 9 patients (7%) with MNCVs less than 30 m/s, corresponding to the demyelinating forms of CMT. However, all MNCVs less than 30 m/s (100%) were associated with major reductions of amplitude, with values less than 0.5 mV. These electrophysiological findings underscore the presence of severe axonal neuropathies with secondary demyelination in CMT associated with *GDAP1* mutations.

Indeed, in demyelinating forms of CMT disease affecting myelinic proteins, such as CMT1A [MIM:118220], CMT1B [MIM:118200] [27-29] as well as in other forms of the disease, such as CMT4 [30], the conduction velocity is lower than 30 m/s even in the early phases of myelination and is definitely established between three

Table 2 Walking ability

	Independent	Walking disability		Total
		Technical assistance	Wheelchair-bound	
Recessive forms ≤ 30 y	23 (32.9%)	28 (40%)	19 (27.1%)	70
> 30 y	0	6 (24%)	19 (76%)	25
missense/missense ≤ 30 y	16 (45.7%)	10 (28.6%)	9 (25.7%)	35
> 30 y	0	5 (50%)	5 (50%)	10
truncating/truncating ≤ 30 y	6 (22.2%)	13 (48.2%)	8 (29.6%)	27
> 30 y	0	1 (6.7%)	14 (93.3%)	15
missense/truncating ≤ 30 y	1 (12.5%)	5 (62.5%)	2 (25%)	8
> 30 y	0	0	0	0
Dominant forms ≤ 30 y	8 (73%)	3 (27%)	0	11
> 30 y	28 (78%)	5 (14%)	3 (8%)	36

Genotype-phenotype correlation analysis using *GDAP1* locus-specific database data evaluated walking ability according to the mode of inheritance and the predicted effect on the protein in the recessive forms of CMT. Values represent the number of patients for each situation; the corresponding percentages are shown in brackets. y: years.

Table 3 Analysis of electrophysiological parameters using *GDAP1* locus-specific database data

Amplitude (mV)	Conduction Velocity (m/s)			Total
	MNCV > 40	MNCV 31-40	MNCV ≤ 30	
CMAP > 6	35 (45%)	2 (15.4%)	0	/ 37
CMAP 1.1-6	30 (39%)	5 (23.1%)	0	/ 35
CMAP 0.5-1	6 (8%)	3 (23.1%)	0	/ 9
CMAP < 0.5	6 (8%)	5 (38.5%)	9 (100%)	/ 20
Not recordable	/	/	/	30/30
Total	77	15	9	131

Values represent the number of patients for each situation; the corresponding percentages are mentioned in brackets. MNCV: motor nerve conduction velocity; CMAP: compound motor action potential.

to five years of age. This conduction velocity is independent of the clinical severity and progression of CMT, and could therefore serve as a marker of the disease even in asymptomatic patients. In contrast, the amplitudes that were initially normal decreased progressively over time following secondary axonal loss. Moreover, histopathological analyses of nerve biopsies described in cases of CMT involving *GDAP1* mutations have revealed a loss of large myelinated fibres, which is known to reduce the MNCV [16].

Finally, the analysis of the database presented here shows that *GDAP1* mutations are responsible for primary axonal damage in CMT, with secondary demyelination in the more severe cases of the disease. Findings of this nature should lead to a better understanding of the pathophysiology of CMT.

Conclusions

One of the patients with a *GDAP1* mutation referred to our database also carried a mutation in the *MFN2* gene [31]. Such cases of multiple gene mutations are now being brought to light more often with the rapid advance of genetic investigations. The *GDAP1* database is part of the MITochondrial DYNamics variation portal (mitodyn.org), aimed at specializing in genes involved in disorders of mitochondrial dynamics [32]. The mitodyn.org will therefore incorporate other genes involved in diseases affecting mitochondrial dynamics and bioenergetics. Thus, the *OPA1* LSDB (*eOPA1*) [33], curated by our laboratory, will soon be transferred to mitodyn.org. We will then integrate *MFN2*, responsible for CMT2A2 [MIM:609260] [34], and *DNM1L* associated with encephalopathy with lactic acidosis [MIM:603850] [35]. Since the same patients may be affected by these different diseases involving mitochondrial dysfunction, the data collected should interest physicians and researchers alike. Eventually, we hope to be able to refine genotype-phenotype correlations in diseases involving mitochondrial dynamics by comparing and cross-checking the clinical, electrophysiological and biochemical data recorded in the mitodyn.org databases.

Acknowledgements

We are grateful to Marie-Claire Malinge for performing molecular diagnosis, and Kanaya Malkani for critical reading and comments on the manuscript.

Author details

¹INSERM, U771, Angers, F-49000, France. ²CNRS, 6214, F-49000 Angers, France. ³Univ Angers, Angers, F-49000, France. ⁴CHU Angers, Département de Neurologie, Angers, F-49000, France. ⁵CHU Angers, Département de Biochimie et Génétique, Angers, F-49000, France.

Authors' contributions

JC and MF initiated the project, created the database, monitored data collection and drafted the manuscript. AC participated in writing the article and the analysis of results. DB, CV, VP, PR and MF supervised the project and revised the manuscript. All authors have given final approval of the version to be published.

Competing interests

The authors declare that they have no competing interests.

Received: 31 August 2011 Accepted: 26 December 2011

Published: 26 December 2011

References

1. Emery AE: Population frequencies of inherited neuromuscular diseases—a world survey. *Neuromuscul Disord* 1991, 1:19-29.
2. Baxter RV, Ben Othmane K, Rochelle JM, Stajich JE, Hulette C, Dew-Knight S, Hentati F, Ben Hamida M, Bel S, Stenger JE, et al: Ganglioside-induced differentiation-associated protein-1 is mutant in Charcot-Marie-Tooth disease type 4A/8q21. *Nat Genet* 2002, 30:21-22.
3. Cuesta A, Pedrola L, Sevilla T, Garcia-Planells J, Chumillas MJ, Mayordomo F, LeGuern E, Marin I, Vilchez JJ, Palau F: The gene encoding ganglioside-induced differentiation-associated protein 1 is mutated in axonal Charcot-Marie-Tooth type 4A disease. *Nat Genet* 2002, 30:22-25.
4. Senderek J, Bergmann C, Ramaekers VT, Nelis E, Bernert G, Makowski A, Zuchner S, De Jonghe P, Rudnik-Schoneborn S, Zerres K, Schroder JM: Mutations in the ganglioside-induced differentiation-associated protein-1 (*GDAP1*) gene in intermediate type autosomal recessive Charcot-Marie-Tooth neuropathy. *Brain* 2003, 126:642-649.
5. Cassereau J, Chevrollier A, Gueguen N, Malinge MC, Letournel F, Nicolas G, Richard L, Ferre M, Verney C, Dubas F, et al: Mitochondrial complex I deficiency in *GDAP1*-related autosomal dominant Charcot-Marie-Tooth disease (CMT2K). *Neurogenetics* 2009, 10:145-150.
6. Chung KW, Kim SM, Sunwoo IN, Cho SY, Hwang SJ, Kim J, Kang SH, Park KD, Choi KG, Choi IS, Choi BO: A novel *GDAP1* Q218E mutation in autosomal dominant Charcot-Marie-Tooth disease. *J Hum Genet* 2008, 53:360-364.
7. Claramunt R, Pedrola L, Sevilla T, Lopez de Munain A, Berciano J, Cuesta A, Sanchez-Navarro B, Millan JM, Saifi GM, Lupski JR, et al: Genetics of Charcot-Marie-Tooth disease type 4A: mutations, inheritance, phenotypic variability, and founder effect. *J Med Genet* 2005, 42:358-365.
8. Ben Othmane K, Hentati F, Lennon F, Ben Hamida C, Blel S, Roses AD, Pericak-Vance MA, Ben Hamida M, Vance JM: Linkage of a locus (CMT4A)

- for autosomal recessive Charcot-Marie-Tooth disease to chromosome 8q. *Hum Mol Genet* 1993, **2**:1625-1628.
- 9. Liu H, Nakagawa T, Kanematsu T, Uchida T, Tsui S: Isolation of 10 differentially expressed cDNAs in differentiated Neuro2a cells induced through controlled expression of the GD3 synthase gene. *J Neurochem* 1999, **72**:1781-1790.
 - 10. Marco A, Cuesta A, Pedrola L, Palau F, Marin I: Evolutionary and structural analyses of GDAP1, involved in Charcot-Marie-Tooth disease, characterize a novel class of glutathione transferase-related genes. *Mol Biol Evol* 2004, **21**:176-187.
 - 11. Shield AJ, Murray TP, Board PG: Functional characterisation of ganglioside-induced differentiation-associated protein 1 as a glutathione transferase. *Biochem Biophys Res Commun* 2006, **347**:859-866.
 - 12. Pedrola L, Espert A, Wu X, Claramunt R, Shy ME, Palau F: GDAP1, the protein causing Charcot-Marie-Tooth disease type 4A, is expressed in neurons and is associated with mitochondria. *Hum Mol Genet* 2005, **14**:1087-1094.
 - 13. Wagner KM, Ruegg M, Niemann A, Suter U: Targeting and function of the mitochondrial fission factor GDAP1 are dependent on its tail-anchor. *PLoS One* 2009, **4**:e5160.
 - 14. Niemann A, Ruegg M, La Padula V, Schenone A, Suter U: Ganglioside-induced differentiation associated protein 1 is a regulator of the mitochondrial network: new implications for Charcot-Marie-Tooth disease. *J Cell Biol* 2005, **170**:1067-1078.
 - 15. Pedrola L, Espert A, Valdes-Sanchez T, Sanchez-Piris M, Sirkowski EE, Scherer SS, Farinas I, Palau F: Cell expression of GDAP1 in the nervous system and pathogenesis of Charcot-Marie-Tooth type 4A disease. *J Cell Mol Med* 2008, **12**:679-689.
 - 16. Cassereau J, Chevrollier A, Gueguen N, Desquiret V, Verny C, Nicolas G, Dubas F, Amati-Bonneau P, Reynier P, Bonneau D, Procaccio V: Mitochondrial dysfunction and pathophysiology of Charcot-Marie-Tooth disease involving GDAP1 mutations. *Experimental neurology* 2011, **227**:31-41.
 - 17. Moroni I, Morbin M, Milani M, Ciano C, Bugiani M, Pagliano E, Cavallaro T, Pareyson D, Taroni F: Novel mutations in the GDAP1 gene in patients affected with early-onset axonal Charcot-Marie-Tooth type 4A. *Neuromuscul Disord* 2009, **19**:476-480.
 - 18. Azzedine H, Ruberg M, Ente D, Gilardeau C, Perie S, Wechsler B, Brice A, LeGuern E, Dubourg O: Variability of disease progression in a family with autosomal recessive CMT associated with a S194X and new R310Q mutation in the GDAP1 gene. *Neuromuscul Disord* 2003, **13**:341-346.
 - 19. The Inherited Peripheral Neuropathies Mutation Database. [http://www.molgen.ua.ac.be/CMTMutations/].
 - 20. Fokkema IF, Taschner PE, Schaafsma GC, Celli J, Laros JF, den Dunnen JT: LOVD v.2.0: the next generation in gene variant databases. *Human mutation* 2011, **32**:557-563.
 - 21. Wildeman M, van Ophuizen E, den Dunnen JT, Taschner PE: Improving sequence variant descriptions in mutation databases and literature using the Mutalyzer sequence variation nomenclature checker. *Human mutation* 2008, **29**:6-13.
 - 22. den Dunnen JT, Antonarakis SE: Mutation nomenclature extensions and suggestions to describe complex mutations: a discussion. *Hum Mutat* 2000, **15**:7-12.
 - 23. den Dunnen JT, Paalman MH: Standardizing mutation nomenclature: why bother? *Hum Mutat* 2003, **22**:181-182.
 - 24. Crimella C, Tonelli A, Airoldi G, Baschirotto C, D'Angelo MG, Bonato S, Losito L, Trabacca A, Bresolin N, Bassi MT: The GST domain of GDAP1 is a frequent target of mutations in the dominant form of axonal Charcot Marie Tooth type 2K. *J Med Genet* 2010, **47**:712-716.
 - 25. AAN: Research criteria for diagnosis of chronic inflammatory demyelinating polyneuropathy (CIDP). *Neurology* 1991, **41**:617-618.
 - 26. Dubourg O: Charcot-Marie-Tooth disease: from phenotype to genotype. *Revue neurologique* 2004, **160**:1221-1229.
 - 27. Birouk N, Gouider R, Le Guern E, Gugenheim M, Tardieu S, Maisonobe T, Le Forestier N, Agid Y, Brice A, Bouche P: Charcot-Marie-Tooth disease type 1A with 17p11.2 duplication. Clinical and electrophysiological phenotype study and factors influencing disease severity in 119 cases. *Brain* 1997, **120**(Pt 5):813-823.
 - 28. Hattori N, Yamamoto M, Yoshihara T, Koike H, Nakagawa M, Yoshikawa H, Ohnishi A, Hayasaka K, Onodera O, Baba M, et al: Demyelinating and axonal features of Charcot-Marie-Tooth disease with mutations of myelin-related proteins (PMP22, MPZ and Cx32): a clinicopathological study of 205 Japanese patients. *Brain* 2003, **126**:134-151.
 - 29. Lewis RA, Sumner AJ, Shy ME: Electrophysiological features of inherited demyelinating neuropathies: A reappraisal in the era of molecular diagnosis. *Muscle Nerve* 2000, **23**:1472-1487.
 - 30. Dubourg O, Azzedine H, Verny C, Durosier G, Birouk N, Gouider R, Salih M, Bouhouche A, Thiam A, Grid D, et al: Autosomal-recessive forms of demyelinating Charcot-Marie-Tooth disease. *Neuromolecular Med* 2006, **8**:75-86.
 - 31. Cassereau J, Casanovas C, Gueguen N, Malinge MC, Guillet V, Reynier P, Bonneau D, Amati-Bonneau P, Banchs I, Volpini V, et al: Simultaneous MFN2 and GDAP1 mutations cause major mitochondrial defects in a patient with CMT. *Neurology* 2011, **76**:1524-1526.
 - 32. The MitoOchondrial DYNAmics variation pages. [http://mitodyn.org].
 - 33. Ferre M, Amati-Bonneau P, Tourmen Y, Malthiery Y, Reynier P: eOPA1: an online database for OPA1 mutations. *Human mutation* 2005, **25**:423-428.
 - 34. Zuchner S, Mersiyanova IV, Muglia M, Bissar-Tadmouri N, Rochelle J, Dadali EL, Zappia M, Nelis E, Putitucci A, Senderek J, et al: Mutations in the mitochondrial GTPase mitofusin 2 cause Charcot-Marie-Tooth neuropathy type 2A. *Nat Genet* 2004, **36**:449-451.
 - 35. Waterham HR, Koster J, van Roermund CW, Mooyer PA, Wanders RJ, Leonard JV: A lethal defect of mitochondrial and peroxisomal fission. *N Engl J Med* 2007, **356**:1736-1741.

doi:10.1186/1750-1172-6-87

Cite this article as: Cassereau et al.: A locus-specific database for mutations in GDAP1 allows analysis of genotype-phenotype correlations in Charcot-Marie-Tooth diseases type 4A and 2K. *Orphanet Journal of Rare Diseases* 2011 **6**:87.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

