




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



# RdRp (RNA-dependent RNA polymerase): A key target providing anti-virals for the management of various viral diseases

Shelly Pathania<sup>a</sup>, Ravindra K. Rawal<sup>b,c,\*</sup>, Pankaj Kumar Singh<sup>d,\*</sup>

<sup>a</sup> Department of Pharmaceutical Chemistry, ISF College of Pharmacy, Moga-142001, Punjab, India

<sup>b</sup> Department of Chemistry, Maharishi Markandeshwar (Deemed to be University), Mullana-133207, Haryana, India

<sup>c</sup> CSIR-North East Institute of Science and Technology, Jorhat-785006, Assam, India

<sup>d</sup> Faculty of Medicine, Integrative Physiology and Pharmacology, Institute of Biomedicine, University of Turku, Turku, FI-20014, Finland



## ARTICLE INFO

### Article history:

Received 30 July 2021

Revised 13 October 2021

Accepted 15 October 2021

Available online 17 October 2021

### Keywords:

RdRp

HCV (Hepatitis)

Influenza

Dengue

BVDV

Covid 19

RdRp inhibitors

## ABSTRACT

With the arrival of the Covid-19 pandemic, anti-viral agents have regained center stage in the arena of medicine. Out of the various drug targets involved in managing RNA-viral infections, the one that dominates almost all RNA viruses is RdRp (RNA-dependent RNA polymerase). RdRp are proteins that are involved in the replication of RNA-based viruses. Inhibition of RdRps has been an integral approach for managing various viral infections such as dengue, influenza, HCV (Hepatitis), BVDV, etc. Inhibition of the coronavirus RdRp is currently rigorously explored for the treatment of Covid-19 related complications. So, keeping in view the importance and current relevance of this drug target, we have discussed the importance of RdRp in developing anti-viral agents against various viral diseases. Different reported inhibitors have also been discussed, and emphasis has been laid on highlighting the inhibitor's pharmacophoric features and SAR profile.

© 2022 The Authors. Published by Elsevier B.V.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

## 1. Introduction

Viruses are the most common infectious diseases causing agents. They typically contain RNA (Ribonucleic acid) or DNA (Deoxyribonucleic acid) as the core and source of genetic information encapsulated in a protein coat. One of the functions of the protein coats is to assist the virus in attaching to the host cell surface, followed by incursion and utilization of the host's bio-machinery for replication. Therefore, based on genetic material, the viruses are of two types, RNA and DNA viruses, out of which RNA-viruses result in critical affliction on healthcare systems throughout the globe [1, 2]. Further, depending on the type of genome, RNA viruses can be categorized into double-stranded (dsRNA), positive sense single-strand (RNA), and negative sense single-strand (ssRNA) [3]. The most recent RNA virus causing a burden on the global health care system is severe acute respiratory syndrome virus 2, the causative agent of COVID-19 [4–8]. Targeting RNA-viruses and their management is complicated due to an intrinsic viral property of integrating the viral genetic material with the host cell genome, which helps them in bypassing the human immune system [9]. Further

utilizing the host machinery, RNA viruses increase their genetic content [10]. Also, RNA viruses maintain mutation rates, which further complicates the management of viral diseases [11,12].

With the advancements in current tools and techniques, all the essential proteins targeted to develop anti-virals are well-known. Still, RNA-dependent RNA polymerases (RdRps) remain among the most critical viral drug targets [13, 14]. It is involved in the genome replication process of an RNA from an RNA template, which is then involved in the encoding of some of the important proteins for the proper functioning and survival of viruses [15]. Following their functional parallel, RdRps also share structural resemblance with DNA polymerases and reverse transcriptases, which also require them to preserve conserved sequences and motifs through their span. Recently there are few reviews published on RdRps as potent anti-viral target [16–18] however they focus on either the structure of RdRp or on the chemical classes of the reported inhibitors, while this work provides the exploration of RdRp as target for different viral infection of positive-sense single-stranded RNA viruses along with the rationales and architecture on which different RdRp inhibitors are designed in the last five years. This work provides the binding mode analysis of different RdRp inhibitors which could be utilized for further designing of potent inhibitors.

\* Corresponding authors.

E-mail addresses: [rawal.ravindra@gmail.com](mailto:rawal.ravindra@gmail.com) (R.K. Rawal), [pankaj.singh@utu.fi](mailto:pankaj.singh@utu.fi) (P.K. Singh).




Fig. 1. 3-D structure of RdRp along with the motifs of palm, finger and thumb domains (PDB ID: 6M71).

## 2. Structure of viral RNA-dependent RNA polymerases

The core domain of RdRp, consisting of fingers, thumb and palm sub-domains, mainly performs the catalytic function involving binding of template, entry of nucleoside triphosphate (NTP), and polymerization (Fig. 1). The palm sub-domain, present at the intersection of the thumb and fingers sub-domains, possess the majority of the conserved structural elements which are essential for the catalytic activity. In the palm sub-domain, a RNA Recognizing Motif (RRM) comprising of three-strands and catalytic aspartates is present [19]. The sub-domain is responsible for the identification of NTPs over deoxy NTPs and catalysis of phosphorylation reaction via metal ion coordination [20]. The palm sub-domain is conformationally dynamic and following the binding of NTP, it undergoes significant restructuring to facilitate catalysis [21]. In some RdRp proteins, belonging to different RNA-viruses, shuffling in the sequence of the catalytic sub-domain results in an uncommon topology [22].

The residues in the thumb sub-domain are responsible for padding alongside the template RNA, which stabilizes the starting NTPs on the template. Following polymerization, thumb sub-domain is involved in the migration of newly formed templates [23, 24]. The fingers sub-domain has a vital function of maintaining the architecture of the active site and holding the RNA template to facilitate polymerization [25]. This sub-domain aids in recognition of the template RNA via binding with the major groove. It consists of four intertwined fingers, which upon interacting with the thumb sub-domain, close the active site [26, 27].

Structurally, seven motifs (A-G) are present in viral RdRps which epitomizes the preserved structural constituents of RdRps [28]. The palm sub-domain contains motifs A to E and motifs G and F are present in the fingers sub-domain. In rare cases, RNA viruses possess an added motif H in the fingers sub-domains. A catalytic motif DX2-4D (an aspartate motif) having invariant first aspartate across numerous RdRps, is present in Motif A. Another aspartate, second in sequence, forms H-bond with 2'-OH of the in-bound NTP and therefore, along with a preserved asparagine of motif B, assists in the discernment of NTPs over deoxy NTPs [6]. In

some RNA viruses, the concerned aspartate is replaced by a conserved lysine permitting them to utilize manganese as the cofactor, in place of magnesium [29]. While in Motif B, a threonine is present in the palm sub-domain, within the  $\alpha$ -helix facing the active site, which contribute to the binding of the template RNA. It also links the N-terminal helix rising from the palm sub-domain to a strand of the fingers sub-domain. Acting as a pivot, it accommodates conformational deviations linked to the binding of the template and the substrate [30]. Whereas in Motif C, a preserved GDD (Gly-Asp-Asp) motif indispensable for complexation with the metal ions is present. Structurally, motifs A and C are juxtaposed and help in forming the RRM. Also, the aspartate from DX2-4D and the preserved aspartates of GDD align with each other at the tip of the RRM to aid in catalysis [31]. Further, motif D possess conserved glycine which participate in the adjustment of the conformation due to proper NTP binding. Detailed studies disclosed that motif D, responsible to select NTPs, is one of the most dynamic elements with variation of around 6 Å [32]. It is also reported to be responsible for the migration of the thumb sub-domain in the course of elongation. While a preserved lysine of motif D performs the key job of deprotonating the leaving pyrophosphate group [33]. Moreover, there is a primer grip formed by motif E, which maintains the proper orientation of 3'-hydroxyl of the primer during catalysis. An aromatic residue present on the face of the N-terminal side pointing at motif C remains conserved in this motif [34]. Furthermore, there is a motif F which consists of positively charged amino acid residues that protect the negative charges of the incoming NTP's phosphate groups by positioning themselves right above the palm sub-domain. This motif also contains a key conserved arginine in its C-terminal [28]. Lastly, motif G, consisting of loop, forms a part of the entrance tunnel for the template and also interacts with the priming NTPs [35].

A lot of channels/tunnels which pass through in various directions are also part of RdRps. They connect the catalytic sites with the external surface of the protein [36]. These channels are also considered as key targets in the development of RdRp inhibitors. The template entry channels possess positively charged residues which favor the inward movement of template RNA and NTPs into






Fig. 3. Compound 1 as anti-BVDV agent with SAR analysis.




Fig. 4. SAR profile of compound 2 as anti-BVDV agent.




Fig. 5. Substituted 9-aminoacridine as potent molecule against BVDV RdRp target.

In 2018, Loddo et al., described the synthesis of 9-aminoacridine derivatives as anti-viral agents *via* RdRp inhibition. The molecules were found able to selectively inhibit BVDV replication. Most potent compound **3** (Fig. 5) with EC<sub>50</sub> value of 1.2 μM, showed favourable selectivity index in comparison to ribavirin (standard drug). All the derivatives had high CC<sub>50</sub> values establishing the safety profile. Further the top molecules were

subjected to biochemical assay for BVDV RdRp inhibitory potential. The analysis disclosed the compounds to have sub-micromolar level potency against RdRp. Further, ITC (Isothermal titration calorimetry) measurements were performed to determine their binding affinities, which were found to be in the range of 0.57–48 μM (K<sub>d</sub> values). Detailed experimental analysis established that the 9-aminoacridine derivatives showed their action by inhibiting




Fig. 6. 1-indanone and *N*4-aryl substituted thiosemicarbazones as potent anti-BVDV agent.




Fig. 7. Imidazodipyridinylaniline derivative as inhibitor of the replication of pestiviruses.




Fig. 8. Imidazoquinolines derivative as anti-BVDV agent with SAR analysis.

the replication of virus assembly via inhibition of RdRp of BVDV [47].

In 2017, Santacruz et al., presented a study involving development of a series of thiosemicarbazones and *N*4-aryl substituted thiosemicarbazones as probable anti-BVDV compounds. The cell-based biological validation established compound **4** as active derivative, with  $EC_{50}$  value of  $0.7 \pm 0.1 \mu\text{M}$ . For *N*4-thiosemicarbazones, the SAR analysis disclosed that di-methoxy substitution on the indanic ring is crucial for the activity. Additionally, *N*4-aryl substitution on the thiosemicarbazone moiety is also a critical feature for the anti-viral potential. Furthermore, an electron-withdrawing group on the aryl ring linked to *N*4 site on the thiosemicarbazone improved the anti-BVDV activity (Fig. 6) [48].

Previously in 2016, the same group reported the identification of **5** (3-(imidazo[1,2-*a*:5,4-*b'*]dipyridin-2-yl)aniline) for the preferential inhibition of CPE (cytopathic effect) produced by BVDV using virus-cell-based assay, with the  $EC_{50}$ -values of  $13.0 \pm 0.6 \text{ mM}$ . Surprisingly, this molecule showed no *in vitro* inhibitory potential against RdRp of recombinant BVDV, but exhibited inhibitory action against BVDV replication complexes in a dose-dependent manner. Through docking studies, it was recognized that the fingertip area in pestivirus RdRp is very vital for the inhibition of its function (Fig. 7) [49].

In 2011, Carta et al., expanded the quinoline ring with 1,2,3-triazole/imidazole/pyrazine to explore the cytotoxicity and antiviral potential of linear *N*-tricyclic ring systems. Upon testing against representatives of different RNA viruses, these quinoline-based molecules could only potently inhibit BVDV. Out of the three classes, imidazoquinolines and pyridoquinoxalines had more potent anti-BVDV activity than triazoloquinolines. Overall, imidazoquinoline **6** (Fig. 8), was more potent with selective activity against BVDV ( $EC_{50} = 1.2 \mu\text{M}$ ) and potency comparable to that of 20-C-methyl-guanosine, a known Flaviviridae RdRp inhibitor. In addition, compound **6** also showed anti-HCV potential with  $EC_{50} = 3.1 \mu\text{M}$  in a replicon assay [50].

In continuation to the previous study, Carta et al., designed, synthesis and evaluated a plethora of molecules with linear aromatic *N*-tricyclic scaffold. Following their previous work, whereby derivatives of imidazo[4,5-*g*]quinoline were found to be potent anti-BVDV agents, they extended the SAR via rational alterations of lead molecule, 2-phenyl-3-*H*-imidazo[4,5-*g*]quinoline, having  $EC_{50}$  value of  $4 \mu\text{M}$  and no cellular toxicity. Various substitutions at multiple sites and replacing aryl ring with a furan ring resulted in 12 imidazo[4,5-*g*]quinoline based compounds. SAR analysis disclosed that chlorine/cyano at 4' position of phenyl ring improves the anti-viral potential of the respective compound. Overall, **7** ( $EC_{50} = 0.3 \mu\text{M}$ ) was found to be most potent amongst the whole series of compounds. Further *in-silico* experiments established the probable mechanism of action of the designed compounds. A 3D pharmacophore model was also developed, which disclosed that one aromatic hydrophobic feature and two H-bond acceptors were basic features required in a potent inhibitor. It was followed by MM/PBSA scoring, which disclosed the significance of a couple of RdRp amino acid residues, i.e., Arg295 and Tyr674, as the essential H-bond contacts (Fig. 9), on the basis of enthalpic component of binding Gibbs free energy change [51].

#### 4.2. For hepatitis C virus (HCV)

In 2016, Meguellat et al., developed pseudodimeric aurones, considering both structure-based and ligand-based approaches of drug discovery, as inhibitors of NS5B (Nonstructural protein 5B) RdRp. Overall, they synthesized 14 compounds having a linker between the benzofuranone moieties and examined their HCV RdRp inhibitory potential using an *in vitro* assay. The most potent compound was found to be pseudodimeric aurone **8**, with inhibitory activity of  $IC_{50} = 1.3 \mu\text{M}$ . The good inhibitory activity was attributed to the presence of phenyl ring as a spacer and -OH group at 4th and 6th position of benzofuran moiety (Fig. 10). The biological study also revealed that the distance between the two benzofuran moiety and shape of the molecule is also very important




Fig. 9. Imidazoquinoline derivative bearing cyano group as potent anti-BVDV agent.




Fig. 10. Pseudodimeric auranones with potential HCV RdRp inhibitory activity.




Fig. 11. Pyrrolo[2,3-d]pyrimidine derivative with potential HCV inhibitory activity.

for its inhibitory action. Molecular docking analysis also highlight the significance of -OH groups as they form H-bond with Asn291. Hydrophobic interaction of benzofuran moiety with residues like Pro197, Met414, Tyr415, Ile447 and Tyr448 was also observed. Anti-HCV activity of pseudodimeric auranones was also evaluated using infectious model which signifies the correlation between the result of both HCV RdRp inhibitory activity and anti-HCV activity [52].

In 2016, Mohamed et al., synthesized series of pyrrole, pyrrolo[2,3-d]pyrimidine and pyrrolo[3,2-e][1,2,4]triazolo[4,3-c]pyrimidine derivatives and evaluated them for HCV inhibitory activity. Pyrrole and pyrrolo[3,2-e][1,2,4]triazolo[4,3-c]pyrimidine derivatives were found inactive against HCV genotype 4, while pyrrolo[2,3-d]pyrimidine derivatives exhibited substantial inhibitory potential. Compound **9** with  $IC_{50}$  value of  $82.1 \pm 10.3 \mu M$ , showed best potency amongst the pool, which indicated that pyrrolo[2,3-d]pyrimidine is crucial nucleus for anti-HCV activity. Incorporation of arylamino group at 4th position also increased the anti-viral activity (Fig. 11). Molecular docking studies revealed that compound **9** formed H-bond with Gln446 and exhibited a docking score of  $-24.92$ , kcal/mol. Overall findings concluded that this molecule can be used as lead compound for the designing of more potent anti-HCV agents [53].

In 2016, Wei et al., described the utilization of random forest based virtual screening (RB-VS) followed by e-pharmacophore (PB-VS), and molecular docking (DB-VS) based screening to filter HCV NS5B polymerase inhibitors. The protocol involved sequential use of the three methods (RB-VS, PB-VS and DB-VS) to filter InterBio-




Fig. 12. Benzoxazole derivative as potent HCV NS5B polymerase inhibitor.

Screen database. Finally, 5 compounds were selected for further validation via biological assays. Upon biochemical and cell-based evaluation, all 5 hits inhibited NS5B polymerase with the  $IC_{50}$  values in low micro molar range ( $2.01$ – $23.84 \mu M$ ) and exhibited anti-HCV activity with  $EC_{50}$  values of  $1.61$  to  $21.88 \mu M$ , without any cytotoxicity ( $CC_{50} > 100 \mu M$ ). Out of all, **10** showed top anti-viral potential for HCV inhibition (Fig. 12) [54].

In 2015, Cakır et al., identified 5-arylidene-4-thiazolidinones as effective HCV NS5B polymerase inhibitors. A pool of 5-arylidene-4-thiazolidinones was synthesized and subjected to NS5B RdRp biochemical assay. The arylidene derivatives showed inhibition potential of 71–91% against NS5B RdRp while compounds without arylidene group showed only 6–31% inhibition. The screening result of NS5B inhibitory activity yielded **11** ( $IC_{50} = 25.3 \mu M$ ) as most active. During the high throughput cell-based anti-HCV screening, **11** showed inhibition of  $\geq 70\%$  in HCV RNA replication process with cell survival of  $\leq 45\%$  at  $100 \mu M$  concentration. Molecular docking studies disclosed the various binding interactions including cation- $\pi$  contact of Arg501 and 3-fluorophenyl ring, and electrostatic interaction of 2-chloro-6-fluorophenyl ring and Asn527/Arg422 residues in the thumb pocket-II of NS5B (Fig. 13). [55].

In 2014, Therese et al., performed a comparative pharmacophore analysis studying the co-crystallized HCV NS5B structures to explore the active conformation of the ligands inside the allosteric pockets of palm and thumb sub-domains. In the next step, a high-throughput virtual screening along with molecular docking exercise was performed to screen varied non-peptide hits. The resulting top 10 hits, exhibiting different scaffolds, were then evaluated using NS5B and anti-HCV inhibition assays. Out of all, **12** was the most promising lead compound with benzimidazole scaffold (Fig. 14), with  $IC_{50}$  value of  $28.8 \mu M$  in NS5B biochemical assay and inhibition of 97% at  $50 \mu M$  in anti-HCV cell-based assay [56].




Fig. 13. 5-arylidene-4-thiazolidinone derivative as potent inhibitor of HCV NS5B polymerase.




Fig. 14. Benzimidazole derivative with HCV NS5B inhibitory activity.




Fig. 15. Anilinobenzothiazole derivative with HCV RdRp inhibitory activity.

In 2013, Peng et al., synthesized many unique derivatives of anilinobenzothiazole and evaluated them for their inhibitory potential on viral genome replication as HCV RdRp inhibitors. The compounds were *in vitro* evaluated for HCV RdRp inhibitory activity using Ava5 cells (cell with HCV replicon). They disclosed that compound **13** showed 76% inhibition of HCV RdRp activity in non-competitive manner and exhibited  $EC_{50}$  value of  $8 \pm 0.5 \mu\text{M}$ . SAR analysis revealed that substituents at 4th position of phenyl ring were important for inhibitory action. Molecular docking experiment performed considering Thumb II Pocket of HCV RdRp suggested that compound **13** well occupied the pocket and was found to be surrounded by four main residues Leu419, Leu497, Arg501, and Trp528 which are responsible for its good inhibitory activity (Fig. 15) [57].

#### 4.3. For zika virus (ZIKV)

In 2020, Wang et al. designed and synthesized novel derivatives of 1,4-dibenzylsubstituted piperazine on the basis of an *in-silico* screening of an *in-built* small molecule library. In the next step,




Fig. 16. 1, 4-benzylsubstituted piperazine derivatives as Zika inhibitors.

screened hits were evaluated for their cytopathic protection effect (CPE) using a standard ZIKV-infected Vero E6 cellular assay. Initial examination disclosed five unique 4-amino-2-(4-benzylpiperazin-1-yl)methylbenzotriazole analogues having CPE lessening effects at micro-molar level. Furthermore, best hit **14** (Fig. 16), at low micro-molar concentrations, showed a substantial dose-dependent antiviral activity by inhibiting both RNA replication of Zika and virus protein expression [58].

In 2017, Pattnaik et al., identified potent RdRp inhibitors for ZIKV *via* structure-based approach. The study involved evaluation of virus replication inhibitory potential of a library of small molecules via virtual screening, followed by validation of the screened hits using cell-based *in vitro* assays. Out of all, **15** was found to effectively inhibit ZIKV replication at nano molar concentrations with  $IC_{50}$  of 94 nM. Further, docking experiments were implemented to study the conformation and binding pose of **15** (Fig. 17), within the catalytic pocket of RdRp, which disclosed the allosteric mechanism involved in blocking the viral RNA synthesis. The two H-bond formed with residues Asp535 and Asp665 were key interactions as these coordinate with divalent metal ions ( $Mg^{2+}$ ) [59].






Fig. 17. Binding pattern of 32 as potential drug candidate for ZIKV infections.




Fig. 18. 6'-fluorinated aristeromycin analogues with inhibitory effect against Chikungunya virus.

#### 4.4. For Chikungunya virus (CHIKV)

In 2020, Kovacicova and Hemert discussed small molecule inhibitors for chikungunya virus, focussing on viral targets, mode of action, and mechanisms of antiviral drug resistance along with associated mutations [60]. In 2019, Yoon et al., designed and developed 6'-fluorinated analogues of aristeromycin, targeting viral RdRp along with SAH hydrolase of the host. The SAR analysis of the 6'-fluorinated aristeromycin analogues disclosed that incorporating fluorine on 6'-position upsurges the SAH hydrolase inhibitory activity along with inhibition of replication in certain +RNA viruses. It was also found that the 6'-fluorinated analogues more efficiently inhibited SAH hydrolase activity than the 6'-unsubstituted derivatives. Amongst all, 6'- $\beta$ -fluoroaristeromycin, **16** (Fig. 18), proved to be better inhibitor for MERS-CoV replication, having an EC<sub>50</sub> value of 0.20  $\mu$ M. [61].

Wada et al., in 2017, screened chemical libraries virtually and found one candidate, a benzimidazole-based compound (**17**, Fig. 19), which at nano molar range, might potentially manage the infection of numerous CHIKV strains. Further to study the mechanism of inhibition, a resistant CHIKV (res-CHIKV) was isolated, which was then subjected to reverse-genetic recombinant CHIKVs to identify the key mutation associated with resistance. This detailed analysis disclosed that at the target site in the RdRp, the Met2295 residue was crucial for the binding of **9**. They also confirmed that **9** used CHIKV replicons to inhibit the function of CHIKV RdRp [62].

#### 4.5. For covid-19

In 2020, Khan et al., screened NNPDB database for SARS-CoV-2 utilizing molecular docking approach. Out of all, 42 molecules showed improved docking score than Remdesivir. Top 5 hits along

with top hit, **18**, were subjected to MD simulation and free binding energy calculations, predicting better binding affinity than Remdesivir. Compound **18** (Diosmetin-7-O-Beta-D-apiofuranoside) exhibited high binding affinity score of  $-10.4$  kcal/mol when compared to Remdesivir ( $-7.1$  kcal/mol). It overall formed several H-bonds with different amino acid residues like Trp617, Tyr619, Lys621, Cys622, Asp623, Asp760, Asp761, Ala762, and Trp800, along with hydrophobic contacts with residues such as Asp618, Lys798, Glu811. Molinspiration Cheminformatics tool revealed the bioactivity score of **18** (Fig. 20) as 0.36 which is quite similar to Remdesivir (0.38). Finally, compound **18** was also found to comply pharmacokinetic rules such as Log P, hydrogen bond donor and acceptors [63].

In 2020, Athar and Beg elaborated the function of anti-HIV and anti-HCV drugs in inhibiting RdRp of SARS CoV 2. The study involved homology modeling followed by validation of the obtained protein structure and docking experiments to study the binding potential of HIV/HCV drugs with RdRp. Results suggested that Nelfinavir, Raltegravir and Delavirdine, out of the studied anti-HIV, and Paritaprevir, Beclabuvir and Ledipasvir, out of the studied anti-HCV, might has potential to inhibit RdRp of SARS CoV-2 [64].

Procacci et al., in the year 2020 disclosed that HCQ (hydroxychloroquine) might behave as a moderate inhibitor of some SARS-CoV2 replication functional proteins. They utilized both enhanced sampling MD techniques and non-equilibrium alchemical transformations together. The  $\Delta G_0$  value for PLpro was found to be  $7.7 \pm 0.9$  kcal/mol, while for 3CLpro, it was  $8.5 \pm 1.1$  kcal/mol, and for RdRp, it was  $9.1 \pm 1.4$  kcal/mol. Further by analysing the state in bound configurations, they could enhance the affinity for 3CLpro, leading to a HCQ-based, **19**, with  $\Delta G_0$  value =  $9.8 \pm 1.4$  kcal/mol. The binding potential increased upon introducing methyl-naphthyl moiety which enhances the hydrophobic interactions (Fig. 21). Also, the -OH of **19** made H-bond with Ser46 residue [65].

Gordon et al., described the mechanistic study of RdRp inhibition via remdesivir in MERS-CoV. Study involved expression of non-structural proteins including RdRp, as a part a polyprotein in insect cells. Further analysis demonstrated that an active complex involves both nsp8 and nsp12 (RdRp) along with remdesivir (triphosphate), which contests with ATP, an RdRp substrate. Importantly, it was observed that active complex had preference for remdesivir-triphosphate over ATP and two other nucleotide analogues which reflects that it is more proficiently assimilated. The mechanism of remdesivir was found to be delay in RNA chain termination. Overall, this study helped in explaining the affinity potential of remdesivir against RNA viruses [66].

#### 4.6. For Dengue virus (DENV)

In 2020, Wan et al., utilized the five-component reaction to synthesize derivatives of octahydroquinazoline-5-one (OHQs). The compounds were tested for anti-dengue activity by lactate dehydrogenase release assay in DENV-2 infected BHK-21 cells. Results suggested that octahydroquinazoline-5-one bearing 4-BrPh at 2nd position exhibited maximum activity (EC<sub>50</sub> = 2.85  $\mu$ M) amongst all in the series. Further they performed SAR analysis by introducing different substituents at 2nd position. It was observed that at 2nd position of OHQs, phenyl substituent with electron withdrawing group at its para position showed good activity (Fig. 22). In addition, OHQs with 3-BrPh resulted into a most potent compound **20** (EC<sub>50</sub> =  $1.31 \pm 0.21$   $\mu$ M) [67].

In 2019, Shimizu et al., employed a HTS (high throughput screen) to filter a library of 16,240 molecules and identified a hit molecule **21**. The molecule was tested against four types of dengue i.e. DENV1-4 and displayed EC<sub>50</sub> values from 6.0 to 31.9  $\mu$ M. Considering best for DENV2 (EC<sub>50</sub> = 6.0  $\mu$ M), **21** was docked in DENV2




Fig. 19. Benzimidazole related compound as anti-CHIKV agent.




Fig. 20. Docking interactions of Diosmetin-7-O-Beta-D-apiofuranoside.




Fig. 21. A novel HCQ-inspired compound with docking interactions.




Fig. 22. Octahydroquinazoline-5-ones with potential anti-dengue activity.

RdRp protein. Docking analysis identified two binding sites (Site 1 on thumb sub-domain and site 2 on palm-subdomain) for **21** within the RdRp. The residues like Cys780, Tyr882, Met809 surrounds benzoxathiole ring of **21** and acetate group showed interactions with Trp833, Arg773 (Fig. 23). While in site 2 benzoxathiole ring is surrounded by Glu510, Gly511, Ser661, and Cys709. Cysteine residue was the common residue for site 1 and site 2 and have close binding contact with **21**. Also, the mutation analysis of

Cys-to Ala-revealed that interaction of compound with cysteine is important for its anti-viral action [68].

In 2018, Lim et al., identified a biphenyl acetic acid fragment targeting unique binding site, present at the thumb/palm boundary near to its N pocket, in the RdRp of dengue virus (DENV). They utilized a fragment-based screening via co-crystallized structure. Structure-directed modification resulted in a pool of nano molar range RdRp inhibitors (**22**; Fig. 24), which also showed sub-micro molar level EC<sub>50</sub> values in cell-based assays [69].

In 2017, Pelliccia et al., reported putative anti-DENV inhibitor which targets the NS5 RdRp. They recognized five lead molecules which showed potent anti-DENV activity. Out of five lead molecules, compound **23** (Fig. 25), EC<sub>50</sub> = 5.99 ± 0.30 μM, exhibited most prominent anti-RdRp activity in cell-based assay. Further, enzymatic assays also confirmed the anti-RdRp activity of **23** with EC<sub>50</sub> of 4.87 ± 0.24 μM. *In-vivo* DENV-infected ICR-suckling mouse model was used to evaluate the anti-DENV activity, which established that compound **23** was able to treat the DENV infection of mice [70].

In 2016, Tarantino et al., identified a pyridobenzothiazole compound, **24**, via an *in silico* docking approach. The compound was found to exhibit dengue RdRp inhibitory activity with IC<sub>50</sub> = 1.5 ± 0.2 μM. Crystallographic analysis displayed that **24** was found to interact in the binding domain of fingers and priming loop of DENV 3 RdRp. Phenolic ring of **16** exhibited π-π stacking




Fig. 23. Compound 21 showing the interacting residues in the active domain of DENV2 RdRp.




Fig. 24. Compound 14 with potential DENV RdRp inhibitory activity.




Fig. 25. Pyrazole derivative with potential RdRp inhibitory activity.




Fig. 26. Pyridobenzothiazole derivative with dengue RdRp inhibitory activity.




Fig. 27. 1,2,4-oxadiazole derivative as potent DENV-2 RdRp inhibitor.

interaction with Trp795 in the priming loop which stabilized the molecule. Though, the bulky cyclohexane ring lead to loss of H-bonding with side chain Asn492 (Fig. 26). Further, mutational analyses identified a new binding cavity i.e. site 2, out of the primary loop. Interactions like H-bond with Cys753 was also observed, suggesting the importance of site 2 in RdRp inhibition [71].

In 2016, Benmansour et al., reported various non-nucleoside based inhibitors of viral polymerase of dengue by modifying the *N*-phenyl-3- arylacrylamide scaffold of hit molecule ((*2E*)-*N*-(2- hydroxy-4-nitrophenyl)-3-(5-nitrothiophen-2-yl)prop-2-enamide). The 1,3,4-oxadiazoles and 1,2,4-oxadiazoles derivatives

of the same hit compound were developed and evaluated for DENV-2 RdRp. It resulted in a potent compound **25** (bearing 1,2,4-oxadiazole moiety) with  $IC_{50}$  value of  $2.2 \pm 0.1 \mu M$ . The compound also exhibited anti-viral activity in micromolar concentration, when tested against DENV-infected Vero cells. SAR analysis indicated that presence of 4-chlorophenyl at 3rd position of oxadiazole moiety is important for DENV-2 RdRp activity (Fig. 27). In addition, bromo group at thiophene is also crucial for anti-viral activity [72].

In 2015, Manvar et al., described the development of coupled thiazolidinone-thiadiazole molecules as novel inhibitors of DENV-2 NS5 RdRp. Around 40 molecules bearing different substituents on thiazolidinone and thiadiazole rings were synthesized. Out of them, **26** ( $IC_{50} = 2.1 \mu M$ ) emerged as a potent compound. SAR study showed that 3-fluorobenzylidene at C5 of the thiazolidinone nucleus is important for activity (Fig. 28). Also, the existence of 2-chlorophenyl at C5 in the 1,3,4-thiadiazole ring is crucial. Molecular docking studies identified five binding sites (S1-S5). S1 is recognized as palm, S2 and S3 covers thumb pocket I and II, and




Fig. 28. Hybrid thiazolidinone-thiadiazole derivative as DENV-2 NS5 RdRp inhibitor.




Fig. 29. 5-(5-fluoro-1H-pyrrolo[2,3-b]pyridin-3-yl)pyrazin-2(1H)-one as potent PB2 inhibitor for the treatment of influenza.

S5 is located at fingers of protein. Binding energy score revealed site 3 with thumb pocket-II as desirable binding domain for compound **26** [73]. The H-bond interactions were observed between NH of thiazolidinone ring and Pro884, and between imine group and Ser885. The fluorobenzylidene group showed aromatic interaction with Tyr882 whereas chloro group exhibited electrostatic interaction with Asn777 [73].

#### 4.7. For influenza virus

In 2019, Yang et al., detailed the discovery of novel derivatives containing 5-(5-fluoro-1H-pyrrolo[2,3-b]pyridin-3-yl)pyrazin-2(1H)-one skeleton as inhibitors of PB2 (a significant subunit of influenza RdRp). Biological evaluation results revealed compound **27** as one of the most potent compound among the series with sub-micro molar  $K_d$  values in surface plasmon resonance (SPR;  $0.11 \mu\text{M}$ ) and isothermal titration calorimetry (ITC;  $0.19 \mu\text{M}$ ) assays. Compound **27**, when examined for *in-vitro* anti-viral activity, showed higher  $\text{EC}_{50}$  value ( $1.025 \mu\text{M}$ ) than standard drug Ribavirin ( $9.116 \mu\text{M}$ ). Molecular docking analysis disclosed that scaffold of compound **27** formed many important interactions like H-bond with Lys376 and Glu361 residues, electrostatic interaction as well as H-bond with Arg355 and  $\pi$ - $\pi$  interactions with His357. SAR studies indicated the importance of ethyl group at N-1 of pyrazine ring of compound **27** (Fig. 29), if replaced by higher bulky group, a decrease in the anti-viral activity was observed [74].

In 2019, Zhao et al., worked on identifying inhibitors of influenza A via screening strategy focussed on the transcription/replication stages by means of an RdRp-focussed filter. Followed by employing those same molecules in a replication-competent reporter virus-based approach. A library of natural product derived compounds (891 molecules) was subjected to the same approach to validate its utility, which established the primary

screen to be rigorous for screening RdRp inhibitors for influenza A virus. In conclusion, out of the two screened hits, one was confirmed as a RdRp inhibitor [75].

In 2014, Lepri et al., designed various molecules as RdRp inhibitors aiming at PA-PB1 (polymerase acidic protein-basic protein 1) complex by modifying the chemical space of a molecule containing thiophene-3-carboxamide and pyrazolo[1,5-a]pyrimidine scaffolds [76], which was previously proved to possess RdRp inhibitory activity. Two approaches were utilized for the design of new compounds but thiophene-3-carboxamide moiety was preserved as an important moiety responsible for RdRp inhibition. Firstly, various new structural features were introduced to explore the interactions capable of interrupting of PA-PB1 complex. The second step utilized scaffold hopping to find new hits with different scaffolds as PA-PB1 complex inhibitor. When pyrazolo[1,5-a]pyrimidine was substituted with triazolo-pyrimidine ring, the PA-PB1 inhibitory activity increased three times. As a result, compound **28** containing triazolo-pyrimidine moiety with  $\text{CF}_3$  and cyclohexane substituents exhibited  $\text{IC}_{50}$  value of  $7.5 \pm 0.7 \mu\text{M}$  in ELISA PA-PB1 interaction assay. These results concluded that derivatives with linear shape are responsible for the good inhibitory activity. Scaffold hopping also resulted in a compound containing sulphonamide moiety with similar inhibitory activity as that of compound **28** (Fig. 30). Compound **28** was also tested against FluA and FluB strains and displayed significant anti-viral potential. [77].

#### 4.8. For norovirus

In 2020, Yi et al., explored the N-pocket in the RdRp of dengue virus (DENV) and its comparable B-site in the RdRp of human norovirus (hNV). They executed a structure-based virtual screening in parallel, considering both the binding sites, to identify hits which can hamper RdRps of hNV and DENV. They positively recognized Entrectinib (**29**) as an effective inhibitor of RdRps of both hNV and DENV. Explicitly, **29** displayed dose-responsive inhibition of both MNV (murine norovirus) and DENV2 with the  $\text{IC}_{50}$  values of 2.01 and 2.43 mM, respectively. It is reported to bind to respective RdRps and efficiently inhibit the catalytic function in the biological assays. The indazole ring of compound **29** exhibit cation- $\pi$  interaction with Arg392 (Fig. 31). It also interacts with other residues like Val504, Glu506, and Asp507 in C-terminal of NV-RdRp. [78].

In 2019, Harmalkar et al., designed various stilbene analogs and tested their inhibitory effect on RNA replication in human norovirus (hNV) by means of HG23, a replicon-bearing viral cell line. Initially, authors screened a large number of natural products against NV which resulted into a hit containing stilbene and vinyl moiety. Taking this hit into consideration, various new compounds were synthesized by introducing substituted amides. Many amide derivatives possessed anti-NV activity with  $\text{EC}_{50}$  values within 1 to




Fig. 30. Compound 26 as potent PA–PB1 inhibitor for treatment of the influenza.




Fig. 31. Entrectinib as potent compound for dual inhibition of hNV RdRp and DENV RdRp.

2  $\mu\text{M}$ . However, introduction of piperazine amide resulted in a potent compound **30** with 95.8% inhibition of NV replication. Moreover, compound **30** (Fig. 32) also showed good safety profile with therapeutic index of 41.2 and also showed less toxicity showing the maximum suppressive action on replication of virus [79].

In 2019, Giancotti et al., modified the skeleton of a HuNoV RdRp inhibitor and an inhibitor of Zika virus polymerase to design modified inhibitors for HuNoV RdRp. They synthesized 12 compounds and almost all the compounds showed partial inhibition of HuNoV (Human Noravirus) RdRp activity in *in-vitro* assays. Especially, out of 12, 5 compounds appeared to cause decrease in MNV-induced cytopathic effect with micro molar range  $\text{EC}_{50}$  values. Also, these compounds were not found substantially cytotoxic. These five hits were further subjected to human norovirus replicon assay possessing a geno-group GI. One of the 5 hits, **31** (Fig. 33), having a novel anti-viral scaffold, showed  $\text{EC}_{50}$  values in the low micro molar level in inhibiting HuNoV replication [80].

In 2014, Tarantino et al., evaluated naphthalene di-sulfonate (NAF2), a fragment-molecule derivative from suramin and NF023 (previously reported RdRp inhibitor of norovirus) [81] for their inhibitory action against Norovirus RdRp. This compound showed less RdRp inhibitory activity ( $\text{IC}_{50} = 14 \mu\text{M}$ ) when tested in polymerase inhibition assays. However, this fragment molecule binds to thumb domain which is different binding site when compared to NF023 and suramin. Additionally, another structurally correlated molecule PPNDS (**32**; pyridoxal-5'-phosphate-6-(2'-naphthylazo-6'-nitro-4',8'-disulfonate) tetrasodium salt) was tested for human Norovirus RdRp activity. The compound **32** exhibited  $\text{IC}_{50}$  value of 0.45  $\mu\text{M}$  which is quite closer to suramin ( $0.24 \pm 0.08 \mu\text{M}$ ). Compound **32** also occupied well the thumb domain of hNV-RdRp protein. Various substituents of PPNDS showed many contacts including hydrophobic interactions of nitro group with Leu406, Ile411, Leu443 and Val504. The carbonyl group of pyridoxal ring showed H-bond with Glu168 residue. The sulfonate group at 4th and 8th position on naphthalene ring formed H-bond with Ser410 and Gln414 (Fig. 34). These findings highlighted the significance of binding site (thumb domain) for further development of Norovirus RdRp inhibitors [82].

## 5. Key observations for designing RdRp inhibitors against different viruses

Upon careful investigation of all the small molecule inhibitors reported as RdRp inhibitors for different viruses, some generaliza-




Fig. 32. Modified stilbene analog with piperazine moiety as inhibitor of NV replication.




Fig. 33. Compound 31 as potent HuNoV RdRp inhibitor.




Fig. 34. PPNSD as potent molecule with hNV-RdRp inhibitory activity.

tions can be made. Therefore, here we will highlight the key structural features needed, in each virus type discussed above, to design potent RdRp inhibitors. Firstly, for RdRp inhibitors of BVDV, the most important requirements include H-bond interactions of the scaffold or substituents of the designed hits and Arg264, E265, Arg285, Arg295 and Y674 amino acid residues in the binding pocket. In case of HCV, the most important requirements include H-bond interactions with Asn291, Arg422, Gln446 and Ser476, and hydrophobic interactions with residues like Pro197, Met414, Tyr415, Ile447 and Tyr448 in the binding pocket. While to design RdRp inhibitors for Zika, attention should be given to incorporate pharmacophoric features which could form two H-bond interactions with D535 and D665 amino acid residues as these coordinates with divalent metal ions ( $Mg^{2+}$ ) in the pocket. In chikungunya, analysis of recent RdRp inhibitors suggests that interaction with M2295 at the target site in the RdRp is crucial for the binding and inhibition. In case of dengue, prospective RdRp inhibitors are expected to occupy two sites in the binding pocket. In one site interactions with Cys780, Tyr882 and Met809 are considered crucial while in the other site interactions with Glu510, Gly511, Ser661, and Cys709 are considered vital. To design the RdRp inhibitors for influenza, focus should be laid on the molecules which could form H-bond with Lys376, Glu361 and Arg355, along with  $\pi$ - $\pi$  interactions with His357. In case of norovirus, RdRp inhibitors need to form H-bond interactions with Glu168, Ser410 and Gln414. However, a more important elucidation from these observations is the fact that different RdRp inhibitors are quite selective for the class of virus they are designed due to the different pharmacophoric requirements in each case and therefore, a more focussed and specific approach should be followed to design RdRp inhibitors.

## 6. Conclusion

Although anti-viral research has been continuous, COVID-19 has highlighted a need for a renewed focus on the management and treatment of viral infections. One of the key reasons for this alarming situation is incidence of the novel variants of previously manageable viral classes. The development of anti-viral drugs has become a key task for medicinal chemistry researchers in this past horrible year. And out of all the known protein targets from the viral genome, RdRp remains most sought-after target for the family of RNA-viruses. Expectedly, Molnupiravir, an RdRp inhibitor developed by Merck, has been approved by the FDA for the management of Covid-19 indicating the efficacy of RdRp targeted agents for the management of current pandemic. Consideration of the structure of viral RdRp is crucial for accurate computational as well as knowledge-based drug discovery protocols for pathological conditions like COVID-19. Therefore, the information provided in this compilation could be utilized to increase the library of pre-clinical molecules having potency and efficacy against Covid-19. The main composition of RdRps is quite similar to that of DNA and RNA polymerases, including motifs and sub-domains. Therefore, it is presumed that individually every RdRp inhibitor might possess common structural features or binding conformations that can explain a similar inhibitory effect throughout the RNA virus family. However, our compilation clearly suggests that RdRps belonging to different viruses have different crucial interaction points in their binding pocket and in each case, inhibitors require different pharmacophoric features to have significant affinity. This observation also hints at one of the reasons of why already reported RdRp inhibitors did not fare well against Covid-19. Accordingly, in the current review, the diverse RdRp inhibitors reported in the last few years have been analysed to draw conclusions about the designing of RdRp inhibitors. The structure activity relationship profile and binding conformations of the reported inhibitors are also discussed to elucidate some generalisations for the designing of further potent inhibitors. Overall, the information can be utilized to increase preclinical candidates targeting RdRp for current global pandemic.

## Declaration of Competing Interest

Authors declare no conflict of interest.

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

SP and RKR would like to acknowledge the faculty members of the ISF College of Pharmacy, Moga, Maharishi Markandeshwar (Deemed to University), Mullana, India and Director, CSIR-NEIST, Jorhat for their direction all the times. PKS would like to thank TCSMT for the Post-doctoral fellowship.

## References

- [1] C.E. Cameron, M. Götte, K.D. Raney, *Viral Genome Replication*, Springer, 2009.
- [2] J.-M. Claverie, Viruses take center stage in cellular evolution, *Genome Biol.* 7 (6) (2006) 1–5, doi:10.1186/gb-2006-7-6-110.
- [3] P. Ahlquist, RNA-dependent RNA polymerases, viruses, and RNA silencing, *Science* 296 (5571) (2002) 1270–1273.
- [4] P. Ahlquist, Parallels among positive-strand RNA viruses, reverse-transcribing viruses and double-stranded RNA viruses, *Nat. Rev. Microbiol.* 4 (5) (2006) 371–382.
- [5] S.S. Chakrabarti, U. Kaur, A. Banerjee, U. Ganguly, T. Banerjee, S. Saha, G. Parashar, S. Prasad, S. Chakrabarti, A. Mittal, COVID-19 in India: are biological and environmental factors helping to stem the incidence and severity? *Aging Dis.* 11 (3) (2020) 480–488, doi:10.14336/AD.2020.0402.
- [6] R. Bhatia, S.S. Ganti, R.K. Narang, R.K. Rawal, Strategies and challenges to develop therapeutic candidates against COVID-19 pandemic, *Open Virol J* 14 (1) (2020) 16–21, doi:10.2174/1874357902014010016.
- [7] R. Cannalire, C. Cerchia, A.R. Beccari, F.S. Di Leva, V. Summa, Targeting SARS-CoV-2 proteases and polymerase for COVID-19 treatment: state of the art and future opportunities, *J. Med. Chem.* (2020), doi:10.1021/acs.jmedchem.0c01140.
- [8] S. Choudhary, K. Sharma, P.K. Singh, Von Willebrand factor: a key glycoprotein involved in thrombo-inflammatory complications of COVID-19, *Chem. Biol. Interact.* 348 (2021) 109657, doi:10.1016/j.cbi.2021.109657.
- [9] S. Jensen, A.R. Thomsen, Sensing of RNA viruses: a review of innate immune receptors involved in recognizing RNA virus invasion, *J. Virol.* 86 (6) (2012) 2900–2910.
- [10] J. Díez, M. Ishikawa, M. Kaido, P. Ahlquist, Identification and characterization of a host protein required for efficient template selection in viral RNA replication, *Proc. Natl. Acad. Sci.* 97 (8) (2000) 3913–3918.
- [11] E.C. Holmes, What does virus evolution tell us about virus origins? *J. Virol.* 85 (11) (2011) 5247–5251.
- [12] M. Tonelli, E. Cichero, Fight against H1N1 influenza A virus: recent insights towards the development of druggable compounds, *Curr. Med. Chem.* 23 (18) (2016) 1802–1817.
- [13] P. Singh, S. Pathania, R. Rawal, Exploring RdRp-remdesivir interactions to screen RdRp inhibitors for the management of novel coronavirus 2019-nCoV, *SAR QSAR Environ. Res.* 31 (11) (2020) 857–867.
- [14] R. Bhatia, R.K. Narang, R.K. Rawal, A summary of viral targets and recently released PDB IDs of SARS-CoV-2, *Virology* 478 (2015) 75–85.
- [15] P. Poltronieri, B. Sun, M. Mallardo, RNA viruses: RNA roles in pathogenesis, coreplication and viral load, *Curr. Genomics* 16 (5) (2015) 327–335.
- [16] R. Kumar, S. Mishra, S.K. Maurya, Recent advances in the discovery of potent RNA-dependent RNA-polymerase (RdRp) inhibitors targeting viruses, *RSC Med. Chem.* 12 (3) (2021) 306–320, doi:10.1039/D0MD00318B.
- [17] L. Tian, T. Qiang, C. Liang, X. Ren, M. Jia, J. Zhang, J. Li, M. Wan, X. YuWen, H. Li, RNA-dependent RNA polymerase (RdRp) inhibitors: the current landscape and repurposing for the COVID-19 pandemic, *Eur. J. Med. Chem.* 213 (2021) 113201, doi:10.1016/j.ejmech.2021.113201.
- [18] M. Machitani, M. Yasukawa, J. Nakashima, Y. Furuichi, K. Masutomi, RNA-dependent RNA polymerase, RdRp, a promising therapeutic target for cancer and potentially COVID-19, *Cancer Sci* 111 (11) (2020) 3976–3984.
- [19] K.K.-S. Ng, J.J. Arnold, C.E. Cameron, in: *Structure-function Relationships Among RNA-dependent RNA polymerases, RNA Interference*, Springer, 2008, pp. 137–156.
- [20] J. Černý, B.Č. Bolfíková, J.J. Valdes, L. Grubhoffer, D. Růžek, Evolution of tertiary structure of viral RNA dependent polymerases, *PLoS ONE* 9 (5) (2014) e96070, doi:10.1371/journal.pone.0096070.
- [21] J.-H. Lee, M.S. Chung, K.H. Kim, Structure and function of civicaliviral RNA polymerases, *Viruses* 9 (11) (2017) 329, doi:10.3390/v9110329.
- [22] S.C. Graham, L.P. Sarin, M.W. Bahar, R.A. Myers, D.I. Stuart, D.H. Bamford, J.M. Grimes, The N-terminus of the RNA polymerase from infectious pancreatic necrosis virus is the determinant of genome attachment, *PLoS Pathog.* 7 (6) (2011) e1002085, doi:10.1371/journal.ppat.1002085.
- [23] A.J. te Velthuis, Common and unique features of viral RNA-dependent polymerases, *Cell. Mol. Life Sci.* 71 (22) (2014) 4403–4420.
- [24] G. Lu, P. Gong, A structural view of the RNA-dependent RNA polymerases from the Flavivirus genus, *Virus Res* 234 (2017) 34–43.
- [25] C. Ferrer-Orta, D. Ferrero, N. Verdaguier, RNA-dependent RNA polymerases of picornaviruses: from the structure to regulatory mechanisms, *Viruses* 7 (8) (2015) 4438–4460.
- [26] P. Surana, V. Satchidanandam, D.T. Nair, RNA-dependent RNA polymerase of Japanese encephalitis virus binds the initiator nucleotide GTP to form a mechanistically important pre-initiation state, *Nucleic Acids Res.* 42 (4) (2014) 2758–2773.
- [27] A.M. Collier, O.L. Lyytinen, Y.R. Guo, Y. Toh, M.M. Poranen, Y.J. Tao, Initiation of RNA polymerization and polymerase encapsidation by a small dsRNA virus, *PLoS Pathog* 12 (4) (2016) e1005523, doi:10.1371/journal.ppat.1005523.
- [28] R. Jácome, A. Becerra, S. Ponce de León, A. Lazzano, Structural analysis of monomeric RNA-dependent polymerases: evolutionary and therapeutic implications, *PLoS ONE* 10 (9) (2015) e0139001, doi:10.1371/journal.pone.0139001.
- [29] M. Lukarska, G. Fournier, A. Pflug, P. Resa-Infante, S. Reich, N. Naffakh, S. Cusack, Structural basis of an essential interaction between influenza polymerase and Pol II CTD, *Nature* 541 (7635) (2017) 117–121.
- [30] D. Garriga, C. Ferrer-Orta, J. Querol-Audí, B. Oliva, N. Verdaguier, Role of motif B loop in allosteric regulation of RNA-dependent RNA polymerization activity, *J. Mol. Biol.* 425 (13) (2013) 2279–2287.
- [31] Y. Wang, M. Xiao, J. Chen, W. Zhang, J. Luo, K. Bao, M. Nie, J. Chen, B. Li, Mutational analysis of the GDD sequence motif of classical swine fever virus RNA-dependent RNA polymerases, *Virus Genes* 34 (1) (2007) 63–65.
- [32] X. Yang, E.D. Smidansky, K.R. Maksimchuk, D. Lum, J.L. Welch, J.J. Arnold, C.E. Cameron, D.D. Boehr, Motif D of viral RNA-dependent RNA polymerases determines efficiency and fidelity of nucleotide addition, *Structure* 20 (9) (2012) 1519–1527.
- [33] C. Castro, E.D. Smidansky, J.J. Arnold, K.R. Maksimchuk, I. Moustafa, A. Uchida, M. Götte, W. Konigsberg, C.E. Cameron, Nucleic acid polymerases use a general acid for nucleotidyl transfer, *Nat. Struct. Mol. Biol.* 16 (2) (2009) 212–218.
- [34] R.T. Kidmose, N.N. Vasiliev, A.B. Chetverin, G.R. Andersen, C.R. Knudsen, Structure of the Q $\beta$  replicase, an RNA-dependent RNA polymerase consisting of viral and host proteins, *Proc. Natl. Acad. Sci.* 107 (24) (2010) 10884–10889.
- [35] C. Chu, S. Fan, C. Li, C. Macken, J.H. Kim, M. Hatta, G. Neumann, Y. Kawaoka, Functional analysis of conserved motifs in influenza virus PB1 protein, *PLoS ONE* 7 (5) (2012) e36113, doi:10.1371/journal.pone.0036113.
- [36] C. Ferrer-Orta, A. Arias, C. Escarmís, N. Verdaguier, A comparison of viral RNA-dependent RNA polymerases, *Curr. Opin. Struct. Biol.* 16 (1) (2006) 27–34.
- [37] L. van der Linden, L. Vives-Adrián, B. Selisko, C. Ferrer-Orta, X. Liu, K. Lanke, R. Ulferts, A.M. De Palma, F. Tanchis, N. Goris, The RNA template channel of the RNA-dependent RNA polymerase as a target for development of antiviral therapy of multiple genera within a virus family, *PLoS Pathog.* 11 (3) (2015) e1004733, doi:10.1371/journal.ppat.1004733.
- [38] S.M. McDonald, Y.J. Tao, J.T. Patton, The ins and outs of four-tunneled Reoviridae RNA-dependent RNA polymerases, *Curr. Opin. Struct. Biol.* 19 (6) (2009) 775–782.
- [39] M. Wang, R. Cao, L. Zhang, X. Yang, J. Liu, M. Xu, Z. Shi, Z. Hu, W. Zhong, G. Xiao, Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro, *Cell Res* 30 (3) (2020) 269–271.
- [40] C. Liang, L. Tian, Y. Liu, N. Hui, G. Qiao, H. Li, Z. Shi, Y. Tang, D. Zhang, X. Xie, A promising antiviral candidate drug for the COVID-19 pandemic: a mini-review of remdesivir, *Eur. J. Med. Chem.* 201 (2020) 112527, doi:10.1016/j.ejmech.2020.112527.
- [41] A. Mullard, 2017 FDA drug approvals, *Nat. Rev. Drug Discov.* 17 (2) (2018) 81–86.
- [42] J.B. Westover, A. Mathis, R. Taylor, L. Wandersee, K.W. Bailey, E.J. Sefing, B.T. Hickerson, K.-H. Jung, W.P. Sheridan, B.B. Gowen, Galidesivir limits Rift Valley fever virus infection and disease in Syrian golden hamsters, *Antiviral Res.* 156 (2018) 38–45.
- [43] R.P. Myers, H. Shah, K.W. Burak, C. Cooper, J.J. Feld, An update on the management of chronic hepatitis C: 2015 Consensus guidelines from the Canadian Association for the Study of the Liver, *Can. J. Gastroenterol. Hepatol.* 29 (1) (2015) 19–34.
- [44] F.G. Hayden, N. Shindo, Influenza virus polymerase inhibitors in clinical development, *Curr. Opin. Infect. Dis.* 32 (2) (2019) 176–186.
- [45] R. Ibbá, S. Piras, I. Delogu, R. Loddo, A. Carta, Anti-BVDV Activity Evaluation of Naphthoimidazole Derivatives Compared with Parental Imidazoquinoline Compounds, *Open Medicinal Chem. J.* 14 (1) (2020) 65–70, doi:10.2174/1874104502014010065.
- [46] S. Musiu, Y.P. Castillo, A. Muigg, G. Pürstinger, P. Leyssen, M. Froeyen, J. Neyts, J. Paeshuyse, Quinolincarboxamides Inhibit the Replication of the Bovine Viral Diarrhea Virus by Targeting a Hot Spot for the Inhibition of Pestivirus Replication in the RNA-Dependent RNA Polymerase, *Molecules* 25 (6) (2020) 1283, doi:10.3390/molecules25061283.
- [47] R. Loddo, V. Francesconi, E. Laurini, S. Boccardo, S. Aulic, M. Farneglia, S. Pricl, M. Tonelli, 9-Aminoacridine-based agents impair the bovine viral diarrhoea virus (BVDV) replication targeting the RNA-dependent RNA polymerase (RdRp), *Bioorg. Med. Chem.* 26 (4) (2018) 855–868.
- [48] M.C.S. Santacruz, M. Fabiani, E.F. Castro, L.V. Cavallaro, L.M. Finkielstein, Synthesis, antiviral evaluation and molecular docking studies of N4-aryl substituted/unsubstituted thiosemicarbazones derived from 1-indanones as potent anti-bovine viral diarrhoea virus agents, *Bioorg. Med. Chem.* 25 (15) (2017) 4055–4063.
- [49] S. Musiu, P. Leyssen, M. Froeyen, J.-M. Chezal, J. Neyts, J. Paeshuyse, 3-(imidazo [1, 2-a: 5, 4-b'] dipyridin-2-yl) aniline inhibits pestivirus replication by targeting a hot spot drug binding pocket in the RNA-dependent RNA polymerase, *Antiviral Res.* 129 (2016) 99–103.
- [50] A. Carta, I. Briguglio, S. Piras, P. Corona, G. Boatto, M. Nieddu, P. Giunchedi, M.E. Marongiu, G. Giliberti, F. Iuliano, Quinoline tricyclic derivatives. Design, synthesis and evaluation of the antiviral activity of three new classes of RNA-dependent RNA polymerase inhibitors, *Bioorg. Med. Chem.* 19 (23) (2011) 7070–7084.

- [51] A. Carta, I. Briguglio, S. Piras, P. Corona, R. Ibba, E. Laurini, M. Fermeleglia, S. Pricl, N. Desideri, E. Atzori, A combined in silico/in vitro approach unveils common molecular requirements for efficient BVDV RdRp binding of linear aromatic N-polycyclic systems, *Eur. J. Med. Chem.* 117 (2016) 321–334.
- [52] A. Meguellati, A. Ahmed-Belkacem, A. Nurisso, W. Yi, R. Brillet, N. Berqouch, L. Chavoutier, A. Fortuné, J.-M. Pawlowsky, A. Boumendjel, New pseudodimeric aurones as palm pocket inhibitors of Hepatitis C virus RNA-dependent RNA polymerase, *Eur. J. Med. Chem.* 115 (2016) 217–229.
- [53] M.S. Mohamed, A.I. Sayed, M.A. Khedr, S.H. Soror, Design, synthesis, assessment, and molecular docking of novel pyrrolopyrimidine (7-deazapurine) derivatives as non-nucleoside hepatitis C virus NS5B polymerase inhibitors, *Bioorg. Med. Chem.* 24 (9) (2016) 2146–2157.
- [54] Y. Wei, J. Li, J. Qing, M. Huang, M. Wu, F. Gao, D. Li, Z. Hong, L. Kong, W. Huang, Discovery of novel hepatitis C virus NS5B polymerase inhibitors by combining random forest, multiple e-pharmacophore modeling and docking, *PLoS ONE* 11 (2) (2016) e0148181, doi:10.1371/journal.pone.0148181.
- [55] G. Çakur, İ. Küçükgüzel, R. Guhamazumder, E. Tatar, D. Manvar, A. Basu, B.A. Patel, J. Zia, T.T. Talele, N. Kaushik-Basu, Novel 4-thiazolidinones as non-nucleoside inhibitors of hepatitis C virus NS5B RNA-dependent RNA polymerase, *Arch. Pharm.* 348 (1) (2015) doi:10.1002/ardp.201400247.
- [56] P.J. Therese, D. Manvar, S. Kondepudi, M.B. Battu, D. Sriram, A. Basu, P. Yogeewari, N. Kaushik-Basu, Multiple e-pharmacophore modeling, 3D-QSAR, and high-throughput virtual screening of hepatitis C virus NS5B polymerase inhibitors, *J. Chem. Inf. Model.* 54 (2) (2014) 539–552.
- [57] H.-K. Peng, W.-C. Chen, Y.-T. Lin, C.-K. Tseng, S.-Y. Yang, C.-C. Tzeng, J.-C. Lee, S.-C. Yang, Anti-hepatitis C virus RdRp activity and replication of novel anilinothiazole derivatives, *Antiviral Res.* 100 (1) (2013) 269–275.
- [58] Y. Wang, R. Zhou, Y. Quan, S. Chen, X. Shi, Y. Li, S. Cen, Design, synthesis, and evaluation of novel 4-amino-2-(4-benzylpiperazin-1-yl) methylbenzotriazole compounds as Zika inhibitors, *Bioorg. Med. Chem. Lett.* 30 (4) (2020) 126906, doi:10.1016/j.bmcl.2019.126906.
- [59] A. Pattnaik, N. Palermo, B.R. Sahoo, Z. Yuan, D. Hu, A.S. Annamalai, H.L. Vu, I. Correias, P.K. Prathipati, C.J. Destache, Discovery of a non-nucleoside RNA polymerase inhibitor for blocking Zika virus replication through in silico screening, *Antiviral Res.* 151 (2018) 78–86.
- [60] K. Kovacicova, M.J. van Hemert, Small-molecule inhibitors of chikungunya virus: mechanisms of action and antiviral drug resistance, *Antimicrob. Agents Chemother.* 64 (12) (2020) e01788–20, doi:10.1128/AAC.01788-20.
- [61] J.-s. Yoon, G. Kim, D.B. Jarhad, H.-R. Kim, Y.-S. Shin, S. Qu, P.K. Sahu, H.O. Kim, H.W. Lee, S.B. Wang, Design, synthesis, and anti-RNA virus activity of 6'-fluorinated-aristeromycin analogues, *J. Med. Chem.* 62 (13) (2019) 6346–6362.
- [62] Y. Wada, Y. Orba, M. Sasaki, S. Kobayashi, M.J. Carr, H. Nobori, A. Sato, W.W. Hall, H. Sawa, Discovery of a novel antiviral agent targeting the non-structural protein 4 (nsP4) of chikungunya virus, *Virology* 505 (2017) 102–112.
- [63] A. Khan, M. Khan, S. Saleem, Z. Babar, A. Ali, A.A. Khan, Z. Sardar, F. Hamayun, S.S. Ali, D.-Q. Wei, Phylogenetic analysis and structural perspectives of RNA-dependent RNA-polymerase inhibition from SARS-CoV-2 with natural products, *Interdiscip. Sci. Comput. Life Sci.* 12 (3) (2020) 335–348, doi:10.1039/D0CC03558K.
- [64] M. Beg, F. Athar, Anti-HIV and Anti-HCV drugs are the putative inhibitors of RNA-dependent-RNA polymerase activity of NSP12 of the SARS CoV-2 (COVID-19), *Pharm. Pharmacol. Int. J.* 8 (3) (2020) 163–172.
- [65] P. Procacci, M. Macchiagodena, M. Pagliai, G. Guarnieri, F. Iannone, Interaction of hydroxychloroquine with SARS-CoV2 functional proteins using all-atoms non-equilibrium alchemical simulations, *Chem. Comm.* 56 (62) (2020) 8854–8856.
- [66] C.J. Gordon, E.P. Tchesnokov, J.Y. Feng, D.P. Porter, M. Götte, The antiviral compound remdesivir potently inhibits RNA-dependent RNA polymerase from Middle East respiratory syndrome coronavirus, *J. Biol. Chem.* 295 (15) (2020) 4773–4779.
- [67] Y. Wan, S. Wu, S. Zheng, E. Liang, S. Liu, X. Yao, Q. Zhu, A series of octahydroquinazoline-5-ones as novel inhibitors against dengue virus, *Eur. J. Med. Chem.* 200 (2020) 112318, doi:10.1016/j.ejmech.2020.112318.
- [68] H. Shimizu, A. Saito, J. Mikuni, E.E. Nakayama, H. Koyama, T. Honma, M. Shirouzu, S.-i. Sekine, T. Shioda, Discovery of a small molecule inhibitor targeting dengue virus NS5 RNA-dependent RNA polymerase, *PLoS Negl. Trop. Dis.* 13 (11) (2019) e0007894, doi:10.1371/journal.pntd.0007894.
- [69] S.P. Lim, C.G. Noble, S. Nilar, P.-Y. Shi, F. Yokokawa, Discovery of potent non-nucleoside inhibitors of dengue viral RNA-dependent RNA polymerase from fragment screening and structure-guided design, in: *Dengue and Zika: Control and Antiviral Treatment Strategies*, Springer, 2018, pp. 187–198.
- [70] S. Pelliccia, Y.-H. Wu, A. Coluccia, G. La Regina, C.-K. Tseng, V. Famigliani, D. Masci, J. Hiscott, J.-C. Lee, R. Silvestri, Inhibition of dengue virus replication by novel inhibitors of RNA-dependent RNA polymerase and protease activities, *J. Enzyme Inhib. Med. Chem.* 32 (1) (2017) 1091–1101.
- [71] D. Tarantino, R. Cannalire, E. Mastrangelo, R. Croci, G. Querat, M.L. Barreca, M. Bolognesi, G. Manfroni, V. Cecchetti, M. Milani, Targeting flavivirus RNA dependent RNA polymerase through a pyridobenzothiazole inhibitor, *Antiviral Res.* 134 (2016) 226–235.
- [72] F. Benmansour, C. Eydoux, G. Querat, X. de Lamballerie, B. Canard, K. Alvarez, J.-C. Guillemot, K. Barral, Novel 2-phenyl-5-[(E)-2-(thiophen-2-yl) ethenyl]-1, 3, 4-oxadiazole and 3-phenyl-5-[(E)-2-(thiophen-2-yl) ethenyl]-1, 2, 4-oxadiazole derivatives as dengue virus inhibitors targeting NS5 polymerase, *Eur. J. Med. Chem.* 109 (2016) 146–156.
- [73] D. Manvar, İ. Küçükgüzel, G. Erensoy, E. Tatar, G. Deryabaşoğulları, H. Reddy, T.T. Talele, O. Cevik, N. Kaushik-Basu, Discovery of conjugated thiazolidinone-thiadiazole scaffold as anti-dengue virus polymerase inhibitors, *Biochem. Biophys. Res. Commun.* 469 (3) (2016) 743–747.
- [74] J. Yang, J. Du, C. Huang, T. Wang, L. Huang, S. Yang, L. Li, Discovery of 5-(5-fluoro-1H-pyrrolo [2, 3-b] pyridin-3-yl) pyrazin-2 (1H)-one derivatives as new potent PB2 inhibitors, *Bioorg. Med. Chem. Lett.* 29 (13) (2019) 1609–1613.
- [75] X. Zhao, Y. Wang, Q. Cui, P. Li, L. Wang, Z. Chen, L. Rong, R. Du, A parallel phenotypic versus target-based screening strategy for RNA-dependent RNA polymerase inhibitors of the influenza A virus, *Viruses* 11 (9) (2019) 826, doi:10.3390/v11090826.
- [76] K. Yamada, H. Koyama, K. Hagiwara, A. Ueda, Y. Sasaki, S.-n. Kanesashi, R. Ueno, H.K. Nakamura, K. Kuwata, K. Shimizu, Identification of a novel compound with antiviral activity against influenza A virus depending on PA subunit of viral RNA polymerase, *Microbes Infect* 14 (9) (2012) 740–747.
- [77] S. Lepri, G. Nannetti, G. Muratore, G. Cruciani, R. Ruzziconi, B. Mercorelli, G. Palù, A. Loregian, L. Goracci, Optimization of small-molecule inhibitors of influenza virus polymerase: from thiophene-3-carboxamide to polyamido scaffolds, *J. Med. Chem.* 57 (10) (2014) 4337–4350.
- [78] D. Yi, Q. Li, L. Pang, Y. Wang, Y. Zhang, Z. Duan, C. Liang, S. Cen, Identification of a broad-spectrum viral inhibitor targeting a novel allosteric site in the RNA-dependent RNA polymerases of dengue virus and norovirus, *Front. Microbiol.* 11 (2020) 1440, doi:10.3389/fmicb.2020.01440.
- [79] D.S. Harmalkar, S.-J. Lee, Q. Lu, M.I. Kim, J. Park, H. Lee, M. Park, A. Lee, C. Lee, K. Lee, Identification of novel non-nucleoside vinyl-stilbene analogs as potent norovirus replication inhibitors with a potential host-targeting mechanism, *Eur. J. Med. Chem.* 184 (2019) 111733, doi:10.1016/j.ejmech.2019.111733.
- [80] G. Giancotti, I. Rigo, G. Pasqualetto, M.T. Young, J. Neyts, J. Rocha-Pereira, A. Brancale, S. Ferla, M. Bassetto, A new antiviral scaffold for human norovirus identified with computer-aided approaches on the viral polymerase, *Sci. Rep.* 9 (1) (2019) 1–20, doi:10.1038/s41598-019-54903-7.
- [81] E. Mastrangelo, M. Pezzullo, D. Tarantino, R. Petazzi, F. Germani, D. Kramer, I. Robel, J. Rohayem, M. Bolognesi, M. Milani, Structure-based inhibition of Norovirus RNA-dependent RNA polymerases, *J. Mol. Biol.* 419 (3–4) (2012) 198–210.
- [82] D. Tarantino, M. Pezzullo, E. Mastrangelo, R. Croci, J. Rohayem, I. Robel, M. Bolognesi, M. Milani, Naphthalene-sulfonate inhibitors of human norovirus RNA-dependent RNA-polymerase, *Antiviral Res.* 102 (2014) 23–28.