



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

## RESEARCH LETTER

# Effects of COVID-19 pandemic on physical activity in children and young adults with implanted devices



Anthony J. Mazzella, MD,\* Anil K. Gehi, MD, FHRS,\* Rachel Lampert, MD, FHRS,†  
Scott Buck, MD,‡ Lindsey Rosman, PhD\*

From the \*Division of Cardiology, Department of Medicine, University of North Carolina Hospitals, Chapel Hill, North Carolina, †Division of Cardiology, Department of Medicine, Yale University, New Haven, Connecticut, and ‡Division of Pediatric Cardiology, Department of Pediatrics, University of North Carolina Hospitals, Chapel Hill, North Carolina.

Children and young adults with preexisting cardiovascular disease (CVD) may be disproportionately affected by the collateral health consequences of the coronavirus disease 2019 (COVID-19) pandemic. In addition to a higher risk of morbidity and mortality from COVID-19,<sup>1</sup> young persons with CVD may be more susceptible to alterations in physical activity (PA) and poor health outcomes<sup>2</sup> owing to the unprecedented loss of structured school days, reduced sports participation, increased screen time, and social isolation.<sup>3</sup> However, data on the acute and long-term effects of the pandemic on PA in young persons with established CVD risk remain scarce.

We leveraged data from electronic health records and implanted cardiac devices, which automatically collect continuous PA data, to analyze objective changes in daily PA during distinct phases of the COVID-19 pandemic relative to the same periods 1 year earlier. The study cohort included 261 patients 35 years and younger enrolled in the remote cardiac device monitoring program at the University of North Carolina and 9 affiliated hospitals who received an implanted cardiac monitor, pacemaker, or implantable cardioverter-defibrillator before February 15, 2019, and had continuous PA monitoring from March 15, 2019, to December 31, 2020. From the initial cohort, 174 patients were excluded,


primarily because of incomplete device data (n = 167) or a diagnosis of COVID during study follow-up (n = 5). Linear mixed effects models with daily PA values as the unit of analysis and a random intercept for each patient were used to estimate trajectories of PA during the stay-at-home order in North Carolina (March 15 through May 8, 2020) and reopening phase of the pandemic (May 9 to December 31, 2020) relative to the corresponding periods in 2019. This study was approved by the Institutional Review Board of the University of North Carolina. Statistical analyses were performed using R version 4.0.5 (R Core Team, 2020, Vienna, Austria).

Overall, 87 patients met study inclusion criteria (19 [21.8%] had implanted cardiac monitors, 27 [31.0%] had pacemakers, and 41 [47.12%] had implantable cardioverter-defibrillators). A rapid and significant decrease in PA during the stay-at-home period was observed compared with the corresponding period in 2019 (−25.6 min/d; 95% confidence interval [CI] −22.5 to −28.7 min/d;  $P < .001$ ). PA remained lower after the stay-at-home order was lifted (−11.4 min/d; 95% CI −9.8 to −13.0 min/d;  $P < .001$ ) and after adjustment for age, sex, and race (Figure 1). Nonwhite race and female sex were associated with lower levels of PA during the reopening period (−69.8 minutes; 95% CI −14.7 to −125.4 minutes;  $P = .018$ ).

This study demonstrates that children and young adults with implanted cardiac devices experienced a significant decline in PA during a government-mandated COVID-19 stay-at-home order compared with the same period in 2019. While PA initially increased after the stay-at-home order was lifted, it subsequently decreased as schools reopened for online and later in-person learning in fall 2020. This pattern of prolonged inactivity is concerning given the well-known cardiovascular and psychosocial health benefits of PA in children and young adults.<sup>4</sup> Electrophysiologists are well positioned to use readily available PA data from implanted cardiac devices to counsel young persons on the importance of staying active and preventing clinical deterioration during COVID-19. Findings from this investigation of objective,

**KEYWORDS** Cardiac device; COVID-19; Defibrillator; Pacemaker; Physical activity (Heart Rhythm 2022;19:165–166)

**Funding Sources:** This study was supported by a grant from the National Heart, Lung, and Blood Institute of the National Institutes of Health to Dr Rosman (K23HL141644). **Disclosures:** Dr Gehi receives research funding from Bristol Myers Squibb Foundation, consulting fees from Biosense Webster, and speaker's honoraria from Abbott, Zoll Medical, and Biotronik. Dr Lampert receives research funding and consulting/honoraria from Medtronic and Abbott/St. Jude Medical. Dr Rosman receives consulting fees from Pfizer. The rest of the authors report no conflicts of interest. **Address reprint requests and correspondence:** Dr Anthony J. Mazzella, Division of Cardiology, Department of Medicine, University of North Carolina at Chapel Hill, 160 Dental Circle - CB 7075, Burnett-Womack Bldg, Chapel Hill, NC 27599-7075. E-mail address: [anthony.mazzella@unchealth.unc.edu](mailto:anthony.mazzella@unchealth.unc.edu).



**Figure 1** Comparison of mean activity from 2019 to 2020. Mean physical activity (PA) in minutes per day stratified by year. Linear mixed effects models demonstrate that PA declined significantly during the mandatory stay-at-home order (*blue rectangle*) after adjustment for race, age, sex, and obesity. PA trends associated with the dates of specific government regulated reopenings are also illustrated (*vertical lines*). Summer camps resumed on May 22, 2020; kindergarten through 12th grade schools opened virtually on August 17 and then in-person at full capacity on October 5, 2020. CI = confidence interval; COVID = coronavirus disease.

device-based activity data complement larger survey-based studies of PA in children,<sup>5</sup> emphasizing the need to determine the long-term consequences of public health policies in children and young adults, especially in minority populations, and the importance of developing targeted activity interventions to mitigate potential downstream health consequences.

**References**

1. Kaushik A, Gupta S, Sood M, Sharma S, Verma S. A systematic review of multi-system inflammatory syndrome in children associated with SARS-CoV-2 infection. *Pediatr Infect Dis J* 2020;39:e340–e346.
2. Chaffee BW, Cheng J, Couch ET, Hoefl KS, Halpern-Felsher B. Adolescents' substance use and physical activity before and during the COVID-19 pandemic. *JAMA Pediatr* 2021;175:715–722.
3. Schmidt SCE, Anedda B, Burchartz A, et al. Physical activity and screen time of children and adolescents before and during the COVID-19 lockdown in Germany: a natural experiment. *Sci Rep* 2020;10:21780.
4. Lobelo F, Muth ND, Hanson S, Nemeth BA, Council on Sports Medicine And Fitness; Section on Obesity. Physical activity assessment and counseling in pediatric clinical settings. *Pediatrics* 2020; 145:e20193992.
5. Chambonniere C, Lambert C, Fearnbach N, et al. Effect of the COVID-19 lockdown on physical activity and sedentary behaviors in French children and adolescents: new results from the ONAPS national survey. *Eur J Integr Med* 2021;43:101308.