





Draft Genome Sequence of *Fusarium oxysporum* f. sp. *cubense* Tropical Race 4 from Peru, Obtained by Nanopore and Illumina Hybrid Assembly

Ana M. Leiva,^a  Mathieu Rouard,^b Diana Lopez-Alvarez,^{a,c} Alberto Cenci,^b Catherine Breton,^b Rosalyn Acuña,^d Juan Carlos Rojas,^e Miguel Dita,^f  Wilmer J. Cuellar^a

^aCrops for Nutrition and Health, International Center for Tropical Agriculture (CIAT), Palmira, Colombia

^bBioversity International, Montpellier, France

^cFacultad de Ciencias Agropecuarias, Universidad Nacional de Colombia, Palmira, Colombia

^dServicio Nacional de Sanidad Agraria (SENASA), Lima, Peru

^eInstituto Nacional de Innovación Agraria (INIA), Lima, Peru

^fBioversity International, Cali, Colombia

Ana M. Leiva and Mathieu Rouard contributed equally to this article.

ABSTRACT *Fusarium oxysporum* f. sp. *cubense* tropical race 4 (Foc TR4) is the causal agent of *Fusarium* wilt, a major threat to the banana industry worldwide. Here, we report the genome of a Foc TR4 strain from Peru, sequenced using a combination of Illumina and Oxford Nanopore Technologies.

Fusarium wilt of banana, caused by *Fusarium oxysporum* f. sp. *cubense*, is a devastating fungal disease affecting bananas worldwide. The pathogen population is divided into four races, but tropical race 4 (Foc TR4) is by far the most aggressive, as it attacks several banana types, including Cavendish, which dominates the global banana export economy (1). In the last 6 years, Foc TR4 has spread from Asia into the Middle East and Africa (2), and in 2019, Foc TR4 reached Latin America, in the north of Colombia (3).

In April 2021, banana plants (*Musa acuminata* group AAA, subgroup Cavendish) showing symptoms of *Fusarium* wilt were observed on a farm in Querecotillo, Peru (4°43'54.84"S, 80°33'45.00"W). Diagnostic analyses confirmed the identity of the pathogen as Foc TR4 (4). Pseudostem strands from symptomatic plants were transferred to potato dextrose

TABLE 1 Representative subset of SIX gene homologs detected in isolates of *Fusarium oxysporum* f. sp. *cubense*^a

Race	VCG ^b	BRIP accession code ^c	Presence of SIX gene:									
			1	2	4	6	7	8	9	13		
R1	0123	62895	x ^{d,f}		x ^b	x ^b				x ^a	x ^a	
R2	01214	25609	x ^f							x ^{a,c}	x ^a	
R4	0122	62892	x ^{c,i}						x ^{a3}	x ^a	x ^c	
STR4	0120	44012	x ^g	x ^d	x ^a			x ^a	x ^{a3,b}	x ^a		
TR4	01213	40340	x ^{a,h,i}	x ^a	x ^c	x ^a			x ^{a1,a2}	x ^a	x ^{a,e}	
TR4 ^d			x ^{a,h,i}	x ^a	x ^c	x ^a			x ^{a1}	x ^a	x ^{a,e}	

^a Isolates shown were reported in reference 12, with the addition of the TR4 Peruvian allelic variants. Sequences were searched on the assembly using BLAST matching with high similarity. X denotes the presence of a gene, while the superscript letters correspond to allelic variants of the gene (TR4 SIX1a, h, i; GenBank accession numbers [KX434991](#), [KX434998](#), [KX434999](#), respectively; SIX2a: [KX435000](#); SIX4c: [KX435006](#); SIX6a: SIX8a1: [KX435011](#), [KX435012](#); SIX9a: [KX435015](#); SIX13: [KX435019](#), [KX435023](#)). SIX, secreted in xylem.

^b VCG, vegetative compatibility group.

^c BRIP, Queensland Plant Pathology Herbarium.

^d Peruvian samples.

Editor Antonis Rokas, Vanderbilt University

Copyright © 2022 Leiva et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Miguel Dita, m.dita@cgiar.org, or Wilmer J. Cuellar, w.cuellar@cgiar.org.

The authors declare no conflict of interest.

Received 11 April 2022

Accepted 12 July 2022

Published 8 August 2022




FIG 1 Neighbor-joining phylogenetic tree of 17 *Fusarium oxysporum* f. sp. *cubense* TR4 isolates with 6,462 SNP variants. Support bootstrap values are indicated as a percentage of the 1,000 replicates. The sequences included were from Colombia (SRA accession number [SRR10747097](#), [SRR10125423](#), [SRR10103605](#)), India ([SRR13311628](#)), Indonesia ([SRR10054446](#)), Israel ([SRR10054450](#)), Jordan ([SRR10054448](#)), Laos ([SRR7226878](#)), Lebanon ([SRR7226880](#)), Pakistan ([SRR7226883](#)), Peru ([SRR15514269](#) to [SRR15514272](#)), and the Philippines ([SRR10054447](#)) and were mapped onto strain UK0001 (GenBank accession number [GCA_007994515.1](#)).

agar (PDA) medium and incubated at 25°C. Single-spore isolates from fungal colonies identified as *Fusarium oxysporum* species complex were further purified and used for DNA extraction (5). DNA from 4 samples (PerS1 to PerS4) was extracted using the Illumina DNA prep kit and sequenced using the MiSeq platform (2 × 151 bp). The same DNA sample from PerS4 was further used for sequencing with Oxford Nanopore Technology (FLOW-MIN111, R10.3 chemistry, LSK109 kit) (6). A total of 46,707,802 Illumina and 379,956 Nanopore reads (average length, 2,783 bp) were obtained. The filtered reads (Illumina, 93.60% > Q30) were combined using Unicycler v0.4.8 (7) to make a hybrid genome assembly with a total length of 46,361,425 bp distributed in 115 contigs (G+C content, 47.59%; N_{50} , 1.63 Mbp). The

consensus sequence quality, checked using Qualimap v2.2.1 (8), resulted in an average depth of $19.25\times$ with 6,718,534 reads mapped.

The sequence assembly, mapped using QUAST v5.0.2 (9), showed high contiguity and a total aligned length of 45.9 Mb (94.7% genome fraction) with the highest-quality genome sequence available, strain UK0001 (10). Gene space assessment was performed using BUSCO v5.2.2 (11) (hypocreales odb10), which reported 97.7% completeness (single copy, 97.2%; duplicate, 0.5%; fragmented, 0.5%; missing, 1.8%; $n = 4,494$). The hybrid assembly was used to identify a family of secreted *in xylem* (SIX) genes. The presence or absence of the SIX homologs was checked using BLASTN v2.2.26 (Table 1) to identify the *F. oxysporum* f. sp. *cubense* sequences and matched the expected allelic variants identified in Foc TR4 for SIX1, SIX6, and SIX8 (12). Moreover, the PerS4 reads, combined with previously reported Foc TR4 strains (2, 3, 13–15), were mapped on UK0001 (10) using BWA v0.7.15 (16), and single nucleotide polymorphism (SNP) calling was conducted using GATK v4.1.6 (17). A dissimilarity matrix (simple matching index) and a neighbor-joining phylogenetic tree were subsequently computed using Darwin v6 (18) (Fig. 1). Peruvian samples were clustered together, separated from other strains, including those from Colombia, suggesting independent incursions of Foc TR4 in the Americas (Fig. 1). Default parameters were used for all software unless otherwise specified.

The availability of complete genome sequences and their comparative analysis will contribute to a better understanding of the Foc TR4 population biology, disease epidemiology, and management.

Data availability. The sequence reads have been deposited at GenBank under the BioProject accession number [PRJNA755905](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA755905), and the assembly sequence has been deposited under the GenBank accession number [GCA_021237285.1](https://www.ncbi.nlm.nih.gov/genbank/GCA_021237285.1).

ACKNOWLEDGMENTS

We acknowledge GenLab del Perú S.A.C. for their support with the Illumina sequencing. We thank Viviana Dominguez from the Cassava Crop Protection Team for technical support. Special thanks to the Plant Protection team of SENASA Peru. The genomic analysis was partially realized with the support of MESO@LR and the South Green Platform.

This work was supported in part by grants from STC-CGIAR, Ministerio de Desarrollo Agrario y Riego (MIDAGRI) from Peru, the CGIAR Research Program on Roots, Tubers and Bananas (RTB) and CGIAR's Plant Health (PHI) initiative.

REFERENCES

- Ploetz RC. 2015. Management of Fusarium wilt of banana: a review with special reference to tropical race 4. *Crop Prot* 73:7–15. <https://doi.org/10.1016/j.cropro.2015.01.007>.
- Zheng S-J, García-Bastidas FA, Li X, Zeng L, Bai T, Xu S, Yin K, Li H, Fu G, Yu Y, Yang L, Nguyen HC, Douangboupha B, Khaing AA, Drenth A, Seidl MF, Meijer HJG, Kema GHJ. 2018. New geographical insights of the latest expansion of *Fusarium oxysporum* f. sp. *cubense* tropical race 4 into the Greater Mekong subregion. *Front Plant Sci* 9:457. <https://doi.org/10.3389/fpls.2018.00457>.
- García-Bastidas FA, Quintero-Vargas JC, Ayala-Vasquez M, Schermer T, Seidl MF, Santos-Paiva M, Noguera AM, Aguilera-Galvez C, Wittenberg A, Hofstede R, Sørensen A, Kema GHJ. 2020. First report of Fusarium wilt tropical race 4 in Cavendish bananas caused by *Fusarium odoratissimum* in Colombia. *Plant Dis* 104:994–994. <https://doi.org/10.1094/PDIS-09-19-1922-PDN>.
- Acuña R, Rouard M, Leiva AM, Marques C, Olortegui A, Ureta C, Cabrera-Pintado RM, Rojas JC, Lopez-Alvarez D, Cenci A, Cuellar WJ, Dita M. 29 June 2022. First report of *Fusarium oxysporum* f. sp. *cubense* tropical race 4 causing Fusarium wilt in Cavendish bananas in Peru. *Plant Dis* <https://doi.org/10.1094/PDIS-09-21-1951-PDN>.
- Dita MA, Waalwijk C, Buddenhagen IW, Souza MT, Jr, Kema GHJ. 2010. A molecular diagnostic for tropical race 4 of the banana fusarium wilt pathogen. *Plant Pathol* 59:348–357. <https://doi.org/10.1111/j.1365-3059.2009.02221.x>.
- López-Alvarez D, Leiva AM, Barrantes I, Pardo JM, Dominguez V, Cuellar WJ. 2020. Complete genome sequence of the plant pathogen *Ralstonia solanacearum* strain CIAT-078, isolated in Colombia, obtained using Oxford Nanopore Technology. *Microbiol Resour Announc* 9:e00448-20. <https://doi.org/10.1128/MRA.00448-20>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Okonechnikov K, Conesa A, García-Alcalde F. 2016. Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* 32:292–294. <https://doi.org/10.1093/bioinformatics/btv566>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Warmington RJ, Kay W, Jeffries A, O'Neill P, Farbos A, Moore K, Bebb DP, Studholme DJ. 2019. High-quality draft genome sequence of the causal agent of the current Panama disease epidemic. *Microbiol Resour Announc* 8:e00904-19. <https://doi.org/10.1128/MRA.00904-19>.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>.
- Czislowski E, Fraser-Smith S, Zander M, O'Neill WT, Meldrum RA, Tran-Nguyen LTT, Batley J, Aitken EAB. 2018. Investigation of the diversity of effector genes in the banana pathogen, *Fusarium oxysporum* f. sp. *cubense*, reveals evidence of horizontal gene transfer. *Mol Plant Pathol* 19:1155–1171. <https://doi.org/10.1111/mpp.12594>.
- Thangavelu R, Edwin Raj E, Pushpakanth P, Loganathan M, Uma S. 2021. Draft genome of *Fusarium oxysporum* f. sp. *cubense* strain tropical race-4 infecting Cavendish (AAA) group of banana in India. *Plant Dis* 105:481–483. <https://doi.org/10.1094/PDIS-06-20-1170-A>.
- Maryani N, Lombard L, Poerba YS, Subandiyah S, Crous PW, Kema GHJ. 2019. Phylogeny and genetic diversity of the banana *Fusarium wilt* pathogen

- Fusarium oxysporum* f. sp. *cubense* in the Indonesian centre of origin. *Stud Mycol* 92:155–194. <https://doi.org/10.1016/j.simyco.2018.06.003>.
15. Maymon M, Sela N, Shpatz U, Galpaz N, Freeman S. 2020. The origin and current situation of *Fusarium oxysporum* f. sp. *cubense* tropical race 4 in Israel and the Middle East. *Sci Rep* 10:1590. <https://doi.org/10.1038/s41598-020-58378-9>.
 16. Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26:589–595. <https://doi.org/10.1093/bioinformatics/btp698>.
 17. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytzky A, Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20:1297–1303. <https://doi.org/10.1101/gr.107524.110>.
 18. Perrier X, Jacquemoud-Collet JP. 2006. DARwin. <http://darwin.cirad.fr/>.