

Case Report

Open Access

Giant lymphatic cyst of omentum: a case report

Sanjeev Kumar, Nikhil Agrawal, Rahul Khanna and AK Khanna*

Address: Department of General Surgery, Institute of Medical sciences, Banaras Hindu University, Varanasi, 221005, India

Email: Sanjeev Kumar - ksanjeev79@gmail.com; Nikhil Agrawal - drnkhl@gmail.com; Rahul Khanna - dr_rahul_khanna@rediffmail.com; AK Khanna* - akhannabhu@gmail.com

* Corresponding author

Published: 7 January 2009

Received: 30 September 2008

Cases Journal 2009, 2:23 doi:10.1186/1757-1626-2-23

Accepted: 7 January 2009

This article is available from: <http://www.casesjournal.com/content/2/1/23>

© 2009 Kumar et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Omental cysts are rare abdominal lesions and are difficult to diagnose. Mostly they are detected incidentally during imaging studies performed for unrelated reasons.

Case presentation: Presentation can be both acute and chronic. Acute presentations are usually due to complication in cyst. Imaging is helpful in excluding other causes of lump abdomen.

We encountered a case of giant lymphatic cyst presenting with abdominal swelling, clinically mimicking huge ovarian cyst.

Conclusion: The goal of surgical therapy is complete excision of the cyst, and Omental cysts can be removed without endangering the adjacent bowel.

Case presentation

A 42-year-old female presented to us with progressive distension of abdomen of one year duration. There was no history of anorexia or weight loss. It was associated with mild dull aching pain all over the abdomen. Her menstrual history was unremarkable. On examination, her abdomen was hugely distended. A vague soft, cystic lump was palpable involving whole of the abdomen. Routine blood tests were within normal limits. An abdominal Ultrasound examination was performed, and it showed a giant cystic lesion with internal echogenic particles and septations, extending from epigastrium to pelvis. All internal viscera were shifted postero-laterally. The origin of the lump could not be established. Provisional diagnosis of ovarian cyst was made and patient was advised to undergo CT scan. CECT showed the huge cystic lesion filling the whole peritoneal cavity with homogenous density (Figure 1). The internal genitalia were found to be normal (Figure 2). With these clinical and imaging findings, provisional diagnosis of omental or mesenteric cyst was made.

Patient underwent laparotomy and a huge, soft cystic lesion arising from omentum was found. It was filling the abdominal cavity and has displaced all intraabdominal viscera posteriorly (Figure 3). It was excised. The gross pathologic appearance was of a huge thin walled cyst, with lymph filled cavities. Histopathological examination revealed that lesion was lined by single layered cuboidal cells and filled with lymph. These findings were consistent with lymphatic cyst of omentum.

Discussion

Omental cysts are rare and are mostly derived from lymphatic tissue. They are present in the lesser or greater omentum and are lined by endothelium. Omental cyst occurs in all age groups, but most often presents in children and young adults [1]. Gairdner published the first report of an omental cyst in 1852 [2]. Omental cysts are thought to represent benign proliferations of ectopic lymphatics that lack communication with the normal lymphatic system. Other etiologic theories include (1) failure




Figure 1
CECT scan showing huge intra-peritoneal cyst.

of the embryonic lymph channels to join the venous system, (2) failure of the leaves of the mesentery to fuse, (3) trauma, (4) neoplasm, and (5) degeneration of lymph nodes [3]. Omental cysts can be simple or multiple, unilocular or multilocular, and they may contain hemorrhagic, serous, chylous, or infected fluid. Omental cysts can be discovered as an incidental finding during laparotomy for another condition, or they can manifest as a chronic or acute abdomen. Chronic symptoms include progressive abdominal distension and pain. The mass may be huge, simulating ascites [4]. The most common mode of acute presentation is that of a small-bowel



Figure 2
CECT showing normal internal genitalia of the patient.




Figure 3
Intraoperative photograph showing huge omental lymphatic cyst.

obstruction, which may be associated with intestinal volvulus [5] or infarction, hemorrhage into the cyst, infection, rupture, cystic torsion, and obstruction of the urinary and biliary tract. The imaging modality of choice is abdominal ultrasonography. Ultrasound demonstrates fluid-filled cystic structure, commonly with thin internal septations and sometimes with internal echoes from debris, hemorrhage, or infection. These can be confused with large ovarian cysts in females. Abdominal computed tomography (CT) scanning adds little additional information, although it can reveal that the cyst is not arising from another organ such as the kidney, pancreas, or ovary. The goal of surgical therapy is complete excision of the cyst [4]. Omental cysts can be removed without endangering the adjacent bowel. Prognosis is excellent.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AKK and RK: operating surgeons. NA: collected clinical details including photograph, summarized the case history and prepared first draft. SK: conducted a literature search and prepared final draft. All authors read and approved the final manuscript.

References

1. Walkar AR, Putham TC: **Omental, mesenteric and retroperitoneal cyst. A clinical study of 33 new cases.** *Ann Surg* 1973, 178:13-19.

2. Gairdner WT: **A remarkable cyst in the omentum.** *Trans Path Soc Lond* 1852, **3**:1851.
3. Egozi EI, Ricketts RR: **Mesenteric and omental cysts in children.** *Am Surg* 1997, **63(3)**:287-90.
4. lu Devrim Karaosmanog, Oktar Suna O zhan, cel Cem Yu, zdemir Hakan O: **Huge omental cyst simulating ascites.** *European Journal of Radiology Extra* 2005, **54**:55-57.
5. Lam FCY, Fan TW, Chan KF, Leung YW, Chao NSY: **Omental Lymphangioma with Torsion: Case Report.** *J HK Coll Radiol* 2005, **8**:49-52.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

