

RESEARCH

Open Access

Significance of CO₂ donor on the production of succinic acid by *Actinobacillus succinogenes* ATCC 55618

Wei Zou¹, Li-Wen Zhu¹, Hong-Mei Li¹ and Ya-Jie Tang^{1,2,3,4*}

Abstract

Background: Succinic acid is a building-block chemical which could be used as the precursor of many industrial products. The dissolved CO₂ concentration in the fermentation broth could strongly regulate the metabolic flux of carbon and the activity of phosphoenolpyruvate (PEP) carboxykinase, which are the important committed steps for the biosynthesis of succinic acid by *Actinobacillus succinogenes*. Previous reports showed that succinic acid production could be promoted by regulating the supply of CO₂ donor in the fermentation broth. Therefore, the effects of dissolved CO₂ concentration and MgCO₃ on the fermentation process should be investigated. In this article, we studied the impacts of gaseous CO₂ partial pressure, dissolved CO₂ concentration, and the addition amount of MgCO₃ on succinic acid production by *Actinobacillus succinogenes* ATCC 55618. We also demonstrated that gaseous CO₂ could be removed when MgCO₃ was fully supplied.

Results: An effective CO₂ quantitative mathematical model was developed to calculate the dissolved CO₂ concentration in the fermentation broth. The highest succinic acid production of 61.92 g/L was obtained at 159.22 mM dissolved CO₂ concentration, which was supplied by 40 g/L MgCO₃ at the CO₂ partial pressure of 101.33 kPa. When MgCO₃ was used as the only CO₂ donor, a maximal succinic acid production of 56.1 g/L was obtained, which was just decreased by 7.03% compared with that obtained under the supply of gaseous CO₂ and MgCO₃.

Conclusions: Besides the high dissolved CO₂ concentration, the excessive addition of MgCO₃ was beneficial to promote the succinic acid synthesis. This was the first report investigating the replaceable of gaseous CO₂ in the fermentation of succinic acid. The results obtained in this study may be useful for reducing the cost of succinic acid fermentation process.

Background

Succinic acid, an intermediate in the cycle of tricarboxylic acid (TCA), is one of four-carbon platform chemicals for producing different kinds of petroleum derivatives and biodegradable polymers [1,2]. Succinic acid could be produced by chemical conversion and microbial fermentation [3]. Because of the rising price, the limited reserves of petroleum and the pollution of environment, the oil-based industries had been prompted a movement towards the bio-based chemicals, and the bio-based succinic acid production had drawn the attention from enterprises and research institutes [4,5].

As the end-product of the energy metabolism, succinic acid could be produced by many anaerobic microbes, such as *Actinobacillus succinogenes*, *Anaerobiospirillum succiniciproducens*, *Mannheimia succiniciproducens*, *Escherichia coli*, and other microbes [2,4,6,7]. Especially *A. succinogenes* ATCC 55618, which is a facultative anaerobe isolated from the bovine rumen [8]. In the production of succinic acid by *A. succinogenes*, one of the key factors is the supply of CO₂. A higher concentration of CO₂ could increase the ratio of succinic acid concentration to the other acids production, the ratio of carbon recovery, and the yield of succinic acid [9]. When *A. succinogenes* and *A. succiniciproducens* were used for the production of succinic acid, as a kind of co-substrate of phosphoenolpyruvate (PEP)-carboxykinase in the TCA cycle, CO₂ could promote carbon flow toward the production of succinic

* Correspondence: yajietang@hotmail.com

¹Key Laboratory of Fermentation Engineering (Ministry of Education), Hubei University of Technology, Wuhan 430068, China

Full list of author information is available at the end of the article

acid [10,11]. For the other succinic acid production microorganisms such as *E. coli* and *Mannheimia succiniciprodu-cens*, CO₂ was incorporated into the backbone of three-carbon compound to generate four-carbon oxaloacetate via PEP carboxylase to enhance the production of succinic acid [12,13].

Because of the poor solubility of gaseous CO₂ at 1 atm, many kinds of carbonate and bicarbonate salts were employed as indirect CO₂ donor to enhance the dissolved CO₂ concentration in fermentation broth. MgCO₃ was a preferable carbonate because the addition of MgCO₃ would not lead to a radical change of culture pH, and an increase of Mg²⁺ concentration in fermentation broth showed little negative effect on the metabolism profile and morphology of succinic acid production strain [14]. Some investigators tried to demonstrate the relationship between extra CO₂ donors and succinic acid production [9,15,16]. But there were a few different features in physiological and biochemical characteristics among various kinds of succinic acid producing strains, and the current results were weak in promoting succinic acid production [15,16].

In this study, the dissolved CO₂ concentration and the addition amount of MgCO₃ were quantitatively determined to optimize succinic production by *Actinobacillus succinogenes* ATCC 55618. To calculate the dissolved CO₂ concentration in the fermentation broth, a mathematical model which considers culture pH, temperature, ionic strength, and salt concentration in the fermentation broth were developed. According to the model prediction and experimental verification, this work firstly demonstrated that the supply of gaseous CO₂ had no significant effect on succinic acid production when MgCO₃ was fully supplied.

Methods

Maintenance and preculture of *Actinobacillus succinogenes*

The strain of *A. succinogenes* ATCC 55618 was purchased from American Type Culture Collection (ATCC, Manassas USA), which was maintained in 20% glycerol at -70°C.

The plate was inoculated with the above strain and incubated at 37°C for 2 days. Preculture medium consisted of the following components (g/L): tryptone 17; soya peptone 3; glucose 2.5; NaCl 5; K₂HPO₄ 2.5, and culture pH was adjusted to 7.1-7.5. For the first preculture, 50-mL medium was prepared in a 250-mL anaerobic bottle, and then a colony from a plate culture was inoculated, and followed by 12-hour incubation at 37°C on a rotary shaker at 120 rpm. For the second preculture, 47.5-mL medium was prepared in a 250-mL anaerobic bottle, and inoculated with 2.5-mL first preculture broth, then followed by 12-hour incubation at 37°C on a rotary shaker at 120 rpm.

Fermentation in the stirred-tank bioreactor

The stirred-tank bioreactor used was a 5.0-L (working volume) BioFlo 110 New Brunswick Scientific (NJ, USA) agitated bioreactor with two six-bladed Rushton impellers (5.9-cm i.d.). The lower impeller was 2.5 cm above the reactor bottom, and the vertical distance between two impellers was 8.5 cm. The reactor was aerated through a ring sparger with a pore size of 1.0 mm, which was located 2.2 cm above the reactor bottom. The bioreactor was equipped with probes of pH (Mettler-Toledo GmbH, Switzerland), temperature and foam.

Fermentation medium was composed of (g/L): glucose 100; yeast extract (YE) 16; corn steep liquor (CSL) 12; KH₂PO₄ 3; NaCl 1; MgCl₂·6 H₂O 0.3; and CaCl₂·2 H₂O 0.3. Fermentation was conducted at 37°C and inoculated with 5% (v/v) of the second preculture broth. The rotation speed and external CO₂ gas sparging rate was 200 rpm and 0.1 vvm. The pH was adjusted to 7.5 with 28% (w/w) ammonia solutions.

Four cultures were carried out simultaneously in the stirred-tank bioreactors with homogeneous cell source under well-controlled process conditions but under different culture conditions. The identical cell source and process conditions, other than the experimental condition, made it possible to perform accurate head-to-head comparisons. The results presented here were confirmed to be reproducible in another experiment (data not shown).

Model description and calculation

When a gas mixture of CO₂ and N₂ was supplied into the bioreactor and became liquid-gas phase equilibrium, the CO₂ dissolved concentration in broth at 1 atm could be described by the reduction of Henry's law:

$$C_{\text{CO}_2} = \frac{P_{\text{CO}_2}}{H} \quad (1)$$

Where P_{CO_2} is the CO₂ partial pressure (kPa) in gas mixture which is determined by the mixing ratio of CO₂ and N₂, H is the Henry's constant for CO₂ in the fermentation broth (kPa·L/mol), and C_{CO_2} is the dissolved CO₂ concentration in the fermentation broth (mol/L).

Since a culture medium contains different kinds of salts and organic substances, the solubility of CO₂ was described according to an empirical model suggested by Rischbieter et al. [17], and Weisenberger and Schumpe [18]:

$$\log\left(\frac{H}{H_0}\right) = \log\left(\frac{c_{G,0}}{c_G}\right) = \sum_i (h_i + h_G)c_i + \sum_j (b_n + b_G)c_{n,j} \quad (2)$$

Where H_0 is the Henry's constant for CO₂ in the pure water (kPa·L/mol), c_i is the concentration of ion i (mol/L), $c_{n,j}$ is the concentration of organic substance j (g/L). $c_{G,0}$

and c_G denote the gas solubility in pure water and fermentation broth. h_i indicated ion-specific parameter (L/mol), and the h_i values are: Na^+ : 0.1143; Ca^{2+} : 0.1762; Mg^{2+} : 0.1694; K^+ : 0.0922; Cl^- : 0.0318; H_2PO_4^- : 0.0906; H^+ : 0; OH^- : 0.0839; HCO_3^- : 0.0976; CO_3^{2-} : 0.1423. b_n indicates substance-specific model parameter (m^3/kg), and the b_n for glucose, YE and CSL are: $6.68 \times 10^{-4} \text{ m}^3/\text{kg}$, $7.9 \times 10^{-4} \text{ m}^3/\text{kg}$, and $2.11 \times 10^{-4} \text{ m}^3/\text{kg}$. h_G was estimated by Equation (3) following the suggestions of Weisenberger and Schumpe [18]:

$$h_G = h_{G,0} + h_{G,T}(T - 298.15K) \quad (3)$$

where $h_{G,0}$ and $h_{G,T}$ for CO_2 are -0.0172 L/mol and $-0.338 \times 10^{-3} \text{ L/mol}\cdot\text{K}$, respectively, between 273-313 K [18]. And b_G in Equation (2) was estimated by Equation (4) as suggested by Rischbieter et al. [17]:

$$b_G = b_{G,0} + b_{G,T}(T - 298.15K) \quad (4)$$



where $b_{G,0}$ and $b_{G,T}$ for CO_2 are $-1.86 \times 10^{-4} \text{ m}^3/\text{kg}$ and $0.01 \times 10^{-4} \text{ m}^3/\text{kg}\cdot\text{K}$, respectively, between 283-303 K [17]. Combining Equations (2), (3) and (4), a model of Henry's constant for CO_2 in fermentation medium could be obtained.

$$\log\left(\frac{H}{H_0}\right) = \sum_i (h_i + h_{G,0} + h_{G,T}(T - 298.15K))\epsilon_i + \sum_j (b_j + b_{G,0} + b_{G,T}(T - 298.15K))\epsilon_{nj} \quad (5)$$


In Equation (5), the Henry's constant of CO_2 in the pure water was 4320 kPa L/mol [15]. T is the absolute temperature (K) in culture condition.

After combining all the parameters mentioned above into the Equation (5) and Equation (1), the model used for calculating CO_2 dissolved concentration in broth could be obtained when gaseous CO_2 was used as external CO_2 donor and the result is shown in Figure 1A. There is a correlated linear trend between CO_2 partial pressure and the dissolved CO_2 concentration in fermentation broth. The dissolved CO_2 concentration in the fermentation broth was 5.05, 10.11, 15.16 and 20.22 mM when CO_2 partial pressure was 25.33, 50.66, 75.99 and 101.33 kPa, respectively. And the maximal dissolved CO_2 concentration is 20.22 mM due to the solubility of gaseous CO_2 .

When MgCO_3 was added with the supply of pure gaseous CO_2 at 1 atm, CO_2 , HCO_3^- , and CO_3^{2-} would become in equilibrium in the fermentation broth according to the following equations [15]:



As reported in the previous study [19], the maximum solubility of MgCO_3 in water at 40°C was 139 mM.



According to Equation (1), (5), (6) and (7), the model used for calculating the dissolved CO_2 concentration in the fermentation broth could be obtained when both gas phase CO_2 and MgCO_3 were used as CO_2 donors, and the relationship between the addition amount of MgCO_3 and the dissolved CO_2 concentration under the CO_2 partial pressure of 101.33 kPa is shown in Figure 1B. The CO_2 concentrations in the fermentation broth were 20.22, 54.97, 89.72, 159.22 and 159.22 mM when the addition of MgCO_3 was 0, 2.92, 5.84, 11.68, and 23.35 g/L, respectively. And the maximum dissolved

CO₂ concentration is 159.22 mM due to the solubility of gaseous CO₂ and MgCO₃.

Effect of CO₂ partial pressure

The significance of gaseous CO₂ partial pressure on succinic acid accumulation was studied by setting CO₂ partial pressure at 25.33, 50.66, 75.99 and 101.33 kPa during the whole fermentation process in the stirred-tank bioreactors, which was controlled by adjusting the corresponding mixing ratio of CO₂ and N₂ at 25%, 50%, 75%, 100% (v: v) by gas mix controller (BioFlo110, New Brunswick Scientific NJ, USA), respectively, and the corresponding dissolved CO₂ concentration in the fermentation broth was 5.05, 10.11, 15.16, and 20.22 mM.

Effect of the supply of gaseous CO₂ and the addition of MgCO₃

The maximal dissolved CO₂ concentration in the fermentation broth was 20.22 and 139.00 mM when only gaseous CO₂ and MgCO₃ was supplied, respectively. In order to study the higher dissolved CO₂ concentration on the succinic acid production, the fermentations were conducted by adding MgCO₃ to enhance the dissolved CO₂ concentration. MgCO₃ was added to the broth after a separate sterilization before the inoculation. The effect of the supply of gaseous CO₂ and the addition of MgCO₃ on the fermentation process was studied by adding 2.92, 5.84, 11.68, and 23.35 g/L of MgCO₃ at the CO₂ partial pressure of 101.33 kPa (i.e., 100% CO₂ gas), and its corresponding dissolved CO₂ concentrations in the fermentation broth were 54.97, 89.72, 159.22 and 159.22 mM, respectively. The dissolved CO₂ concentration maintained constant at 159.22 mM even when concentrations higher than 11.68 g/L of MgCO₃ were added at the CO₂ partial pressure of 101.33 kPa. The other culture conditions were the same as the above experiments.

Effect of the addition of higher amount of MgCO₃

Effect of the excessive addition amount of MgCO₃ was studied by adding 30, 40, 50 and 60 g/L of MgCO₃ at the CO₂ partial pressure of 101.33 kPa (i.e., 100% CO₂ gas), and all the corresponding dissolved CO₂ concentration in the fermentation broth was 159.22 mM. The other culture conditions were the same as the above experiments.

Effect of CO₂ donor supply mode

According to the above results and Equation (5), the effect of CO₂ donor supply mode was studied by using two supply modes: 40 g/L MgCO₃ was used as the only CO₂ donor, and 40 g/L MgCO₃ was supplied at the CO₂ partial pressure of 101.33 kPa (i.e., 100% CO₂ gas). The other culture conditions were the same as the above experiments.

Sampling, the determination of succinic acid production


For sampling, about 20-30 mL of broth was taken once from each reactor and the cell growth was monitored by measuring the optical density at 660 nm (OD₆₆₀). At an OD₆₆₀ of 1.0, *A. succinogenes* ATCC 55618 has a concentration of 0.626 g dry cell weight (DCW)/L. For succinic acid determination, 1 mL methanol and 1 mL acetonitrile were added to 1 mL fermentation broth to remove protein and the sample was kept at 4°C overnight. After centrifuging at 11,000 rpm for 30 min, the supernatants were diluted and filtrated through 0.22 μm filter, then analyzed by high-performance liquid chromatography (HPLC, Dionex) using Maisch ReproSil-Pur Basic C18 column. The optimized mobile phase was 5 mM KH₂PO₄ water solution, whose pH was adjusted to 2.8 by H₃PO₄. The column oven temperature was maintained at 40°C and the flow rate was 1 mL/min. The detection wave was 210 nm. Residual sugar level was assayed with phenol-sulfuric acid method [20].

Results and discussion

Effect of CO₂ partial pressure

As one of the direct substrates for the biosynthesis of succinic acid, CO₂ could affect the metabolic flux and the mass distribution of succinic acid [8,21]. The quantitative determination of the dissolved CO₂ concentration in the fermentation broth is beneficial to study the impact of CO₂ partial pressure on the production of succinic acid. Song et al. [15] and Lee et al. [16] reported that succinic acid production could be enhanced by increasing CO₂ partial pressure in the fermentation of *M. succiniciproducens* and *A. succiniciproducens*. Therefore, it was necessary to investigate the effect of CO₂ partial pressure on the accumulation of succinic acid by *A. succinogenes* ATCC 55618.

The effect of CO₂ partial pressure on the succinic acid production is shown in Figure 2A. When the CO₂ partial pressures were 25.33, 50.66, 75.99, and 101.33 kPa, the dissolved CO₂ concentrations in the fermentation broth calculated using Equation (5) was 5.05, 10.11, 15.16, and 20.22 mM, respectively (Table 1). And at the CO₂ partial pressure of 101.33 kPa, the maximal dissolved CO₂ concentration achieved was 20.22 mM, which was the highest dissolved CO₂ concentration when only gaseous CO₂ was supplied. The dissolved CO₂ concentration was increased with the increase of the partial pressure when gaseous CO₂ was used as sole CO₂ donor. The succinic acid productions were 8.84, 10.21, 10.44, and 10.97 g/L as obtained on 48 hour at the CO₂ partial pressure of 25.33, 50.66, 75.99, and 101.33 kPa, respectively, and its corresponding productivities were 0.18, 0.21, 0.22, and 0.23 g/L per hour. This indicated that when gaseous CO₂ was used as the sole CO₂ donor, CO₂ partial pressure showed no significant



effect on the succinic acid accumulation. On the contrary, as reported by Lu et al. [22] and Samuelov et al. [23], a higher available CO₂ concentration could cause higher succinic acid production by increasing the activity of PEP carboxykinase. These indicated that when gaseous CO₂ was used as the sole CO₂ donor, the available dissolved CO₂ concentration was not high enough to increase the production of succinic acid in the fermentation of *A. succinogenes*.

As shown in Figure 2B, the patterns of acetic acid production at various CO₂ partial pressures were similar. The concentrations of other by-products such as formic acid, lactic acid and ethanol were relatively constant at around 5.0, 11.0 and 2.0 g/L, respectively, regardless of the levels of the dissolved CO₂ in the broth.

Figure 2C shows the time profile of residual sugar under various CO₂ partial pressures. The glucose concentration at the CO₂ partial pressure of 101.33 kPa was decreased faster than that at other CO₂ partial pressures during the first 24 hours. The yield of succinic acid against glucose was around 0.21 g succinic acid/g glucose when gaseous CO₂ was used. That means the CO₂ partial pressure showed no significant effect on the succinic acid yield. And there was no significant effect on the cell growth. The OD₆₆₀ was between 6.0 and 6.7 when gaseous CO₂ partial pressure was 25.33, 50.66, 75.99, and 101.33 kPa (Table 1).

Effect of the supply of gaseous CO₂ and the addition of MgCO₃

The maximal dissolved CO₂ concentration is limited by the solubility of gaseous CO₂ when it was supplied as the sole CO₂ donor. In order to investigate the effect of higher dissolved CO₂ concentration on succinic acid production, the fermentations were conducted by adding MgCO₃ at the CO₂ partial pressure of 101.33 kPa (i.e., 100% CO₂ gas) to enhance the dissolved CO₂ concentration in the fermentation broth.


The effect of the supply of gaseous CO₂ and the addition of MgCO₃ on the fermentation process was studied by adding 2.92, 5.84, 11.68, and 23.35 g/L of MgCO₃ at the CO₂ partial pressure of 101.33 kPa (i.e., 100% CO₂ gas), and the corresponding dissolved CO₂ concentrations was 54.97, 89.72, 159.22, and 159.22 mM. Because of the solubility of MgCO₃ and CO₂, the maximal dissolved CO₂ concentration of 159.22 mM was obtained under the addition of 11.68 g/L MgCO₃ with 100% CO₂ gas. Even more than 11.68 g/L of MgCO₃, the dissolved CO₂ concentration maintained constant at 159.22 mM. As shown in Figure 3A, the highest succinic acid productions were 15.26, 15.94, 25.86 and 36.84 g/L as obtained on 48 hour with the addition of 2.92, 5.84, 11.68 and 23.35 g/L MgCO₃, respectively, and its corresponding productivity was 0.32, 0.33, 0.54 and 0.77 g/L per hour,

Table 1 Effect of CO₂ donor on the growth of *A.succinogenes* and succinic acid production parameters

Addition amount of MgCO ₃ (g/L)	Gaseous CO ₂ partial pressure (kPa) ^a	Dissolved CO ₂ concentration (mM)	OD ₆₆₀	Succinic acid productivity (g/L per hour)	Specific succinic acid productivity (g succinic acid/g DCW)	Yield of succinic acid against glucose (g succinic acid/g glucose)
- ^a	25.33	5.05	6.61	0.18	3.50	0.19
-	50.66	10.11	6.69	0.21	3.23	0.22
-	75.99	15.16	6.51	0.22	3.10	0.19
-	101.33	20.22	6.03	0.23	3.57	0.21
2.92	101.33	54.97	10.12	0.32	2.71	0.27
5.84	101.33	89.72	10.27	0.33	2.93	0.26
11.68	101.33	159.22	9.72	0.54	5.13	0.38
23.35	101.33	159.22	10.94	0.77	6.53	0.45
30	101.33	159.22	9.10	0.74	11.04	0.56
40	101.33	159.22	9.62	0.86	13.60	0.60
50	101.33	159.22	9.74	0.85	12.66	0.64
60	101.33	159.22	9.62	0.81	11.44	0.63
40	101.33	159.22	10.31	0.84	12.02	0.58
40	-	139.00	9.65	0.80	11.77	0.54

^a Gas mixture was composed of CO₂ and N₂.

^b - means the CO₂ donor was not add.



respectively. The maximum succinic acid production was increased from 25.86 to 36.84 g/L when the addition amount of MgCO₃ was increased from 11.68 to 23.35 g/L, while the dissolved CO₂ concentration was maintained constant. It can be concluded that the higher dissolved CO₂ concentration was beneficial for the succinic acid biosynthesis. But the dissolved CO₂ concentration was not the only factor affecting succinic acid synthesis; the excessive MgCO₃ also had positive effect. As a kind of neutralization reagent, MgCO₃ could promptly neutralize the organic acid produced during the fermentation process. But when only 11.68 g/L of MgCO₃ was added, the saturated state of MgCO₃ would be lost quickly because of the rapid accumulation of organic acid during the fermentation. And when the addition amounts of MgCO₃ exceeded 11.68 g/L, there will be excessive solid MgCO₃ precipitate. Even if organic acids accumulate, MgCO₃ solution can also keep saturated.

Equation (6) and (7) indicated that the dissolved concentrations of HCO₃⁻, CO₃²⁻ and CO₂ could be enhanced with the addition of MgCO₃ in the fermentation broth. However, MgCO₃ may not be used as CO₃²⁻ donor because there were few reports that CO₃²⁻ could be used directly as substrate by succinic acid producing microorganisms. Although HCO₃⁻ and CO₂ could be used as the co-substrate of PEP carboxylase and improve the production of succinic acid [23], HCO₃⁻ was much less permeable to lipid cell membrane than the uncharged CO₂ molecule because is a kind of polar molecular, and there was no HCO₃⁻ transporter on the membrane of *A. succinogenes* which could deliver HCO₃⁻ from the broth into the cell [24]. So the higher concentration of HCO₃⁻ could not promote the production of succinic acid. And MgCO₃ may be used as indirect CO₂ molecule donor to promote the production of succinic acid in the fermentation process of *A. succinogenes*. On the other hand, when the levels of dissolved CO₂ reached 159.22 mM, there would be insoluble MgCO₃, and that could cause turbid broth. The cells were spread uniformly in the broth, which was helpful to eliminate the cell flocculation and indirectly promoting the succinic acid biosynthesis.

As shown in Figure 3B, the patterns of acetic acid production at the dissolved CO₂ concentration of 54.97 and 89.72 mM were similar. However, when the levels of dissolved CO₂ reached 159.22 mM, the acetic acid production was significantly enhanced. It was distinct from other reports. In the fermentation of *M. succiniciproducens*, the levels of dissolved CO₂ showed little effects on the acetic acid accumulation [15]. The CO₂ concentration has been shown to regulate the levels PEP carboxykinase pathway at high CO₂ levels, and PEP carboxykinase levels rise [23]. However, the enhanced PEP carboxylation may cause higher glucose consumption rate. This effect may cause

more metabolic flow by PEP to pyruvic acid, and further to acetic acid. Meanwhile the production of formic acid, lactic acid and ethanol almost not be improved may be because the raised PEP carboxykinase activity competitively inhibited these key enzymes such as pyruvate formate lyase, lactate dehydrogenase and ethanol dehydrogenase.

The patterns of cell growth with the addition of different concentration of $MgCO_3$ were similar, and the OD_{660} was between 9.7 and 10.9 (Table 1). Figure 3C shows the time profile of residual sugar under various addition amount of $MgCO_3$. When 23.35 g/L of $MgCO_3$ was added, glucose was consumed faster than the other conditions between 12 and 24 hour, which corresponded well to the succinic acid accumulation. The yield of succinic acid against glucose was 0.27, 0.26, 0.38, and 0.45 g succinic acid/g glucose when the addition amount of $MgCO_3$ was 2.92, 5.84, 11.68 and 23.35 g/L, respectively. This indicated that the higher dissolved CO_2 concentration could effectively improve the yield of succinic acid against glucose.

Effect of the addition of higher amount of $MgCO_3$

Effect of the higher addition amount of $MgCO_3$ was studied by adding 30, 40, 50 and 60 g/L of $MgCO_3$ at the CO_2 partial pressure of 101.33 kPa (i.e., 100% CO_2 gas), and all the corresponding dissolved CO_2 concentration in the fermentation broth were 159.22 mM.

As shown in Figure 4A, the pattern of succinic acid production under various addition amount of $MgCO_3$ within the range of investigation was similar. The maximal succinic acid production of 53.55, 61.92, 61.48, and 58.05 g/L was obtained with the addition of 30, 40, 50, and 60 g/L $MgCO_3$, respectively, and its corresponding productivity was 0.74, 0.86, 0.85 and 0.81 g/L per hour. When the addition amount of $MgCO_3$ exceeded 40 g/L, the production and productivity of succinic acid were kept almost constant, but the specific productivity was decreased. This indicated 40 g/L $MgCO_3$ was enough for improving the accumulation of succinic acid. Similarly, Du et al. [25] reported when the addition amount of $MgCO_3$ exceeded 30 g/L, there was no significant change on the production of succinic acid.

The significance of addition amount of $MgCO_3$ on acetic acid accumulation was studied. As shown in Figure 4B, there was no significant effect on the biosynthesis of acetic acid. Similarly, the concentrations of other by-products such as formic acid, lactic acid and ethanol were relatively constant at around 5.0, 11.0 and 2.0 g/L, respectively, regardless of the addition amount of $MgCO_3$.

The cell growth patterns under the addition of $MgCO_3$ were quite similar (Table 1). Figure 4C shows that the time profile of residual sugar under various addition amount of $MgCO_3$. The yield of succinic acid against glucose was 0.56, 0.60, 0.64, and 0.63 g succinic acid/g glucose when the addition amount of $MgCO_3$




Figure 4 Effect of the excessive addition amount of $MgCO_3$ on the succinic acid production. (A) the succinic acid production, (B) the acetic acid production, (C) the sugar consumption.

was 30, 40, 50 and 60 g/L, respectively. It seemed that to obtain a higher yield of succinic acid against glucose, the addition amount of MgCO_3 should no be more than 50 g/L.

Effect of CO_2 donor supply mode

Gaseous CO_2 was widely used as external CO_2 donor and anaerobic environment maintenance agent in succinic acid fermentation process. Calculated from Equation (5, 6, 7), the dissolved CO_2 concentration (139.00 mM) under 40 g/L MgCO_3 was just decreased by 12.76% comparing with that obtained under the addition of 40 g/L MgCO_3 with 100% CO_2 gas. This suggested that the gaseous CO_2 may be removed by the addition of MgCO_3 . In order to testify this proposal, the effect of CO_2 donor supply mode was studied by using two supply modes: 40 g/L MgCO_3 was used alone; 40 g/L MgCO_3 was supplied with 100% CO_2 gas.

As shown in Figure 5A, after 72 h incubation, the production of succinic acid reached 56.14 and 60.38 g/L when 40 g/L MgCO_3 was used as the only CO_2 donor and 40 g/L MgCO_3 was supplied at the CO_2 partial pressure of 101.33 kPa, and the corresponding productivity was 0.80 and 0.84 g/L per hour. The succinic acid production was just decreased by 7.03% without the supply of gaseous CO_2 . As shown in Figure 5B, the acetic acid production was decreased by 17.91% without the supply of gaseous CO_2 . Figure 5C clearly shows the time courses of sugar consumption under different CO_2 supply modes were similar. The yield of succinic acid against glucose was 0.54 g succinic acid/g glucose when 40 g/L MgCO_3 was used alone, and the yield was 0.58 g succinic acid/g glucose when MgCO_3 was supplied with 100% CO_2 . And there was no significant effect on the cell growth whether gaseous CO_2 was used.

Conclusions

In this study, an effective CO_2 quantitative mathematical model was developed to calculate the dissolved CO_2 concentration in the broth during the fermentation of *Actinobacillus succinogenes* ATCC 55618. The model offered a quantitative method for screening the suitable CO_2 donor form and addition amount for the production of succinic acid. There was no significant effect of CO_2 partial pressure on the production of succinic acid when gaseous CO_2 was used as the sole CO_2 donor. But when gaseous CO_2 was used with MgCO_3 , higher amount of MgCO_3 was more effective on promoting the succinic acid synthesis. And the maximum succinic acid production of 61.92 g/L was obtained at 159.22 mM dissolved CO_2 concentration, which was supplied by 40 g/L MgCO_3 with 100% CO_2 gas. And it was concluded that the supply of gaseous CO_2 was not essential when 40 g/L of MgCO_3 was added in the fermentation medium.




Figure 5 Effect of CO_2 donor supply mode on succinic acid production during the *A. succinogenes* fermentation. (A) the succinic acid production, (B) the acetic acid production, (C) the sugar consumption.

This is the first report investigating the replaceable of gaseous CO₂ in the fermentation of succinic acid. The results obtained in this study may be useful for reducing the cost of succinic acid fermentation process.

Acknowledgements

Financial support from the following funding agencies is gratefully acknowledged: the National Natural Science Foundation of China (NSFC, Project Nos. 20976038 and 21176059), the Key Project of Chinese Ministry of Education (Project No. 210132), the Hubei Provincial Natural Science Foundation for Innovative Research Team (Project No. 2008CDA002), the Scientific Research Key Project of Hubei Provincial Department of Education (Project No. Z20101401), the Discipline Leader Project of Wuhan Municipality (Project No. 200951830553), the Key Technology R&D Program of Wuhan Municipality (Project No. 201120822280-2), the Open Project Programs for the Key Laboratory of Fermentation Engineering (Ministry of Education), the National Key Laboratory of Biochemical Engineering (Project No. 2010KF-06), and the State Key Laboratory of Bioreactor Engineering. Ya-Jie Tang also thanks the Chutian Scholar Program (Hubei Provincial Department of Education, China) (2006).

Author details

¹Key Laboratory of Fermentation Engineering (Ministry of Education), Hubei University of Technology, Wuhan 430068, China. ²Laboratory of Synthetic Biology, Shanghai Advanced Research Institute, Chinese Academy of Sciences, Shanghai 201203, China. ³National Key Laboratory of Biochemical Engineering, Institute of Process Engineering, Chinese Academy of Sciences, Beijing 100080, China. ⁴State Key Laboratory of Bioreactor Engineering, East China University of Science and Technology, 130 Meilong Road, Shanghai 200237, China.

Authors' contributions

YJT and WZ designed the experiments, WZ carried out the experimental work, LWZ and WZ analyzed data, WZ drafted the manuscript, HML provided some critical discussions and support for this project. YJT and LWZ critically reviewed and modified the paper. All authors approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 30 June 2011 Accepted: 31 October 2011

Published: 31 October 2011

References

1. Bechthold I, Bretz K, Kabasci S, Kopitzky R, Springer A: **Succinic acid: a new platform chemical for biobased polymers from renewable resources.** *Chem Eng Technol* 2008, **31**(5):647-654.
2. Song H, Lee SY: **Production of succinic acid by bacterial fermentation.** *Enzyme Microb Technol* 2006, **39**:352-361.
3. Sauer M, Porro D, Mattanovich D, Branduardi P: **Microbial production of organic acids: expanding the markets.** *Trends Biotechnol* 2008, **26**(2):100-108.
4. McKinlay JB, Vieille C, Zeikus JG: **Prospects for a bio-based succinate industry.** *Appl Microbiol Biotechnol* 2007, **76**:727-740.
5. Zeikus JG, Jain MK, Elankovan P: **Biotechnology of succinic acid production and markets for derived industrial products.** *Appl Microbiol Biotechnol* 1999, **51**:545-552.
6. Lee KY, Park JM, Kim TY, Yun H, Lee SY: **The genome-scale metabolic network analysis of *Zymomonas mobilis* ZM4 explains physiological features and suggests ethanol and succinic acid production strategies.** *Microb Cell Fact* 2010, **9**:94.
7. Yuzbashev TV, Yuzbasheva EY, Sobolevskaya TI, Laptev IA, Vybornaya TV, Larina AS, Matsui K, Fukui K, Sineoky SP: **Production of succinic acid at low pH by a recombinant strain of the aerobic yeast *Yarrowia lipolytica*.** *Biotechnol Bioeng* 2010, **107**:673-682.
8. Guettler MV, Rumler D, Jain MK: ***Actinobacillus succinogenes* sp. nov., a novel succinic-acid-producing strain from the bovine rumen.** *Int J Syst* 1999, **49**:207-216.

9. Van der Werf MJ, Guettler MV, Jain MK, Zeikus JG: **Environmental and physiological factors affecting the succinate product ratio during carbohydrate fermentation by *Actinobacillus* sp. 130Z.** *Arch Microbiol* 1997, **167**:332-342.
10. McKinlay JB, Zeikus JG, Vieille C: **Insights into *Actinobacillus succinogenes* fermentative metabolism in a chemically defined growth medium.** *Appl Environ Microbiol* 2005, **71**(11):6651-6656.
11. Lee PC, Lee WG, Lee SY, Chang HN: **Succinic acid production with reduced by-product formation in the fermentation of *Anaerobiospirillum succiniciproducens* using glycerol as a carbon source.** *Biotechnol Bioeng* 2001, **72**(1):41-48.
12. Lu S, Eiteman M, Altman E: **Effect of flue gas components on succinate production and CO₂ fixation by metabolically engineered *Escherichia coli*.** *World J Microbiol Biotechnol* 2010, **26**:429-435.
13. Lee SY, Kim JM, Song H, Lee JW, Kim TY, Jang YS: **From genome sequence to integrated bioprocess for succinic acid production by *Mannheimia succiniciproducens*.** *Appl Microbiol Biotechnol* 2008, **79**:11-22.
14. Liu YP, Zheng P, Sun ZH, Ni Y, Dong JJ, Wei P: **Strategies of pH control and glucose-fed batch fermentation for production of succinic acid by *Actinobacillus succinogenes* CGMCC1593.** *J Chem Technol Biotechnol* 2008, **83**:722-729.
15. Song H, Lee JW, Choi S, You JK, Hong WH, Lee SY: **Effects of dissolved CO₂ levels on the growth of *Mannheimia succiniciproducens* and succinic acid production.** *Biotechnol Bioeng* 2007, **98**(6):1296-1304.
16. Lee PC, Lee WG, Kwon S, Lee SY, Chang HN: **Succinic acid production by *Anaerobiospirillum succiniciproducens*: effects of the H₂/CO₂ supply and glucose concentration.** *Enzyme Microb Technol* 1999, **24**:549-554.
17. Rischbieter E, Schumpe A, Wunder V: **Gas solubilities in aqueous solutions of organic substances.** *J Chem Eng Data* 1996, **41**:809-812.
18. Weisenberger S, Schumpe A: **Estimation of gas solubilities in salt solutions at temperatures from 273 K to 363 K.** *AIChE J* 1996, **42**(1):298-300.
19. Mark HF, Othmer DF, Overberger CG, Seaborg GT: *Kirk-Othmer encyclopedia of chemical technology* New York: John Wiley&Sons; 1981.
20. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F: **Colorimetric method for determination of sugars and related substances.** *Anal Chem* 1956, **28**:350-356.
21. Jones RP, Greenfield PF: **Effect of carbon dioxide on yeast growth and fermentation.** *Enzyme Microb Technol* 1982, **4**(4):210-223.
22. Lu S, Eiteman MA, Altman E: **Effect of CO₂ on succinate production in dual-phase *Escherichia coli* fermentations.** *J Biotechnol* 2009, **143**:213-223.
23. Samuelov NS, Lamed R, Lowe R, Zeikus JG: **Influence of CO₂-HCO₃⁻ levels and pH on growth, succinate production, and enzyme activities of *Anaerobiospirillum succiniciproducens*.** *Appl Environ Microbiol* 1991, **57**(10):3013-3019.
24. Badger MR, Price GD: **CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution.** *J Eep Bot* 2003, **54**(383):609-622.
25. Du C, Lin SKC, Koutinas A, Wang R, Dorado P, Webb C: **A wheat biorefining strategy based on solid-state fermentation for fermentative production of succinic acid.** *Bioresour Technol* 2008, **99**:8310-8315.

doi:10.1186/1475-2859-10-87

Cite this article as: Zou et al.: Significance of CO₂ donor on the production of succinic acid by *Actinobacillus succinogenes* ATCC 55618. *Microbial Cell Factories* 2011 **10**:87.