



OPEN

## Shifting and reducing breathing disturbance in patients with very severe obstructive sleep apnea by modified Z-palatoplasty with one-layer closure in one-stage multilevel surgery

Ethan I. Huang<sup>1,2,3</sup>, Yu-Ching Lin<sup>2,3,4,5</sup>, Shu-Yi Huang<sup>3,4,6</sup>, Chin-Kuo Lin<sup>4</sup> & Chieh-Mo Lin<sup>4,7</sup>

Very severe obstructive sleep apnea (OSA) with apnea–hypopnea index (AHI)  $\geq 60$  events/h differs in several areas from OSA with other severities, including having a low-level daytime partial pressure of oxygen and residual on-CPAP (continuous positive airway pressure) AHIs greater than 20/h. Patients with very severe OSA show narrow retroglottal space and confined framework, which is difficult to be enlarged via conventional Uvulopalatopharyngoplasty (UPPP) surgery, resulting in poor response to non-framework surgeries. Our latest report showed efficacy and efficiency for subjects undergoing modified Z-palatoplasty (ZPP) with one-layer closure in a one-stage multilevel surgery. It is unclear whether and how this procedure could help patients with very severe OSA characterized with confined framework. From Mar. 2015 to May 2018, we enrolled 12 patients with very severe OSA receiving one-stage multi-level surgery with modified ZPP with one-layer closure, CO<sub>2</sub> laser partial tongue-base glossectomy, and bilateral septomeatoplasty. Our results show that the surgery reduced AHI from  $73.8 \pm 10.7$  to  $30.8 \pm 23.2$  events/h and achieved a mean AHI reduction of 58.3% ( $p < 0.001$  against 0 reduction or no surgery). The surgery shifted components of the breathing disturbances. It reduced more apnea than hypopnea and might convert some apnea to hypopnea.

Very severe OSA is a distinct subgroup and shows differences in several areas comparing OSA with other severities. In the literature, it refers to OSA with a high AHI or respiratory disturbance index (RDI), ranging from higher than 40<sup>1,2</sup>, 50<sup>3–7</sup>, 60<sup>8–12</sup>, 70<sup>13</sup>, or 100 events/h<sup>14</sup>. Unlike the typical exhibition of hypoxia resulting from repeated cessations of breathing during sleep (see<sup>15</sup> for a review), patients with very severe OSA may have a low level (as 77 mmHg) of diurnal partial pressure of oxygen while they are not in sleep<sup>16</sup>. They may have residual on-CPAP AHIs greater than 20 events/h<sup>17</sup>. Comparing with those with less severity, they have less positional OSA<sup>18</sup>, minimal inspiratory movement of the lateral pharyngeal walls and less maximum cross-sectional area<sup>6</sup>, higher insulin resistance<sup>19</sup>, a high prevalence of hypertension<sup>16</sup>, and about 3 times the episodes of heart block comparing with an unselected group of patients with OSA<sup>20</sup>.

They may comply more with regular use of the CPAP device than those with less severe OSA<sup>21</sup>. However, some refuse using CPAP or cannot benefit from the device. Researchers advocated alternative surgical procedures such as direct skeletal surgery<sup>22</sup>, bariatric surgery<sup>23</sup>, and tracheostomy in patients meeting certain criteria<sup>1,24</sup>. Typically, they have narrow retroglottal space and confined framework<sup>22</sup>, which is difficult to be enlarged via conventional UPPP surgery. These patients may have worse or no response to non-framework surgeries (e.g.,<sup>9,25</sup>).

<sup>1</sup>Department of Otolaryngology, Chang Gung Memorial Hospital, 6 W. Sec. Jiapu Rd., Puzi, Chiayi 61363, Taiwan. <sup>2</sup>School of Medicine, Chang Gung University, Taoyuan, Taiwan. <sup>3</sup>Sleep Center of Chang Gung Memorial Hospital, Chiayi, Taiwan. <sup>4</sup>Division of Pulmonary and Critical Care Medicine, Chang Gung Memorial Hospital, Chiayi, Taiwan. <sup>5</sup>Department of Respiratory Care, Chang Gung University of Science and Technology, Chiayi, Taiwan. <sup>6</sup>Department of Nursing, Chang Gung University of Science and Technology, Chiayi, Taiwan. <sup>7</sup>Graduate Institute of Clinical Medical Sciences, College of Medicine, Chang Gung University, Taoyuan, Taiwan. ✉email: ehuang@alumni.pitt.edu

However, the similar confined framework is commonly seen in some patients with only mild OSA. It hints an opportunity for a non-framework surgery. As a non-framework surgery, modified ZPP with one-layer closure in a one-stage multilevel surgery<sup>26</sup> has revealed efficacy and efficiency as that proposed by Friedman and colleague since 2004<sup>27–29</sup>. It is unclear whether and how this technique could help patients with very severe OSA. Reducing OSA severity may help in its consequences, e.g., hypertension<sup>30</sup>, cognitive deficits<sup>31,32</sup>, and cardiovascular disorders<sup>33–35</sup>. Reducing desaturation might improve coexisting medical conditions. It is unknown how this surgery may improve the severity and desaturation. Smaller body mass index (BMI) and preoperative AHI were reported good predictors for the surgical outcome<sup>36</sup>. Better surgical response has been described in patients with smaller BMI (<30) or smaller AHI (<60)<sup>36</sup>. It is unclear whether they stay good predictors in the category of very severe OSA. Here in this study, we analyzed the component shifts and tested the statistical significance of pre- and post-operative sleep parameters for very severe OSA patients whose AHIs are higher or equal to 60 events/h. We also tested the effectiveness of these two predictors.

## Materials and methods

From Mar. 2015 to May 2018, we enrolled subjects with very severe OSA who met these criteria to the study:

- Age  $\geq 20$  years
- Unsuccessful or refusal of CPAP
- AHI  $\geq 60$  events/h
- Received a one-stage multi-level sleep surgery with the modified ZPP performed with one-layer closure, CO<sub>2</sub> laser partial tongue-base glossectomy, and bilateral septomeatoplasty
- Available preoperative and postoperative polysomnography (PSG) for AHI measurement

We performed preoperative endoscopic evaluation to figure the anatomical stage<sup>37</sup>. Patients were not selected for surgery by any other criteria unlisted above. Epworth Sleepiness Scale (ESS) that comprises 8 4-point scale (0–3) inquiries was used to measure daytime sleepiness, with the total score ranging from 0 to 24. The higher the ESS score, the more that individual's daytime sleepiness in everyday life. Modified ZPP with one-layer closure was carried out as illustrated in our earlier report<sup>26</sup>. Open tongue-base resection was completed with transoral (CO<sub>2</sub>) laser microsurgery for hypopharyngeal obstruction according to the preoperative endoscopic assessment. After the surgery, we cared all patients for in general ward areas with an oximeter monitor. Intravenous Dynastat twice daily was prescribed for 1–3 days. No intravenous or oral narcotics were given to prevent respiratory depression.

We used the percentage of reduction in mean AHI as the primary measure of surgical efficacy<sup>38</sup> to compare with the results across most studies in the literature. It measures the mean change in AHI compared to the mean AHI before surgery. Mean and standard deviation summarized the pre- and post-operative AHI. We performed a paired t-test to examine the change in AHI against no change after the surgery. We examined other associated sleep parameters, including obstructive apnea index (OAI), minimum oxyhemoglobin saturation of pulse oximetry (SpO<sub>2</sub>), mean SpO<sub>2</sub>, desaturation index, and mean desaturation with a paired t-test. Postoperative care and complications were also reported, including suture dehiscence and bleeding. A p-value smaller than 0.05 was deemed significant.

We calculated the correlation coefficient between each of preoperative BMI and AHI vs. AHI reduction to test these two predictors. To clarify the effect of BMI change on AHI reduction, we computed individual BMI change (postoperative BMI–preoperative BMI in PSG records) then calculated the correlation coefficient between individual BMI change and AHI reduction. The statistical significance was tested as  $\alpha = 0.05$ .

The statistical examinations were performed in Matlab 9.4.0.813654 (MathWorks, Natick, Massachusetts, U.S.A.).

**Ethical approval.** The Institutional Review Board (IRB) of Chang Gung Medical Foundation, Taiwan approved the study methods and protocols (IRB number: 201800948B0). We performed the study in accordance with Good Clinical Practice and the applicable laws and regulations. As a retrospective cohort study, the IRB approved the waiver of the participants' consent.


## Results

Ten male and 2 female patients with very severe OSA aged between 25 and 59 years were enrolled to this work. The mean BMI was 28.1 with a standard deviation of 3.3 kg/m<sup>2</sup>. All patients received the 3 procedures in the multilevel sleep surgery listed above. Two and 1 patients underwent routine endoscopic sinus surgery and adenoidectomy, respectively. A PSG followed about 5 months (159  $\pm$  59 days, mean with one standard deviation) after the surgery. Table 1 detailed the individual pre- and post-operative AHI, Friedman anatomic stage, and the surgeries performed.

The mean AHI (with one standard deviation) improved from 73.8  $\pm$  10.7 events/h to 30.8  $\pm$  23.2 after the surgery (Fig. 1). The AHI reduction was 58.3% and was away from 0 ( $p < 0.001$ ). AHI reduction by Friedman anatomical stage was presented in Fig. 2. There were 1, 2, and 9 cases in stage 1, 2, and 3, respectively. Individual AHI change was shown in Fig. 3. One patient (8.3%) displayed normal PSG (AHI < 5 events/h) after the surgery. Three (25%) and 2 (16.7%) improved from very severe to mild and moderate categories, respectively. Four (33%) moved better to the severe level, and 2 (16.7%) remained in the very severe level. Ten out of 12 (83%) of the patients had various improvements on severity grouping. In the components of AHI, the surgery reduced OAI from 44.0  $\pm$  16.8 to 5.8  $\pm$  9.0 events/h,  $p < 0.001$  (Fig. 4A) and improved minimum SpO<sub>2</sub> from 70.3  $\pm$  12.3 to 80.4  $\pm$  7.7%,  $p = 0.0088$  (Fig. 4B). It improved mean SpO<sub>2</sub> from 92.1  $\pm$  4.1 to 95.0  $\pm$  1.2% ( $p = 0.0427$ ), reduced

Case	Age and sex	Preop AHI	Postop AHI	Preop ESS	Postop ESS	Tonsil size	Friedman tongue position	Friedman stage	Preop body mass index	Postop body mass index	Surgeries
1	25M	79.4	29.1	17	16	3	2	1	30	32.2	abce
2	34M	66	0.8	7	8	4	3	2	28.6	29.6	abcd
3	49M	69.8	32.7	12	12	2	3	3	25.8	25.1	abc
4	40M	65.2	12.5	12	5	1	3	3	24.6	24	abc
5	50M	86.5	22	11	11	1	3	3	27.4	25.7	abc
6	31M	85.3	68.6	3	7	2	4	3	32	29.4	abc
7	58M	62.6	77.6	8	2	1	4	3	25.7	23.8	abcd
8	51M	63.1	6.4	11	4	1	4	3	21.8	23.2	abc
9	56F	92.7	13.9	18	12	1	4	3	31.5	30.3	abc
10	59M	60.8	31.5	12	14	1	2	2	27.2	24.3	abc
11	38 M	77.9	39.2	2	1	1	4	3	32.3	31.2	abc
12	58F	76.7	35.2	3	2	1	4	3	30.1	28	abc

**Table 1.** Enrolled subjects ordered by their visit times. Preop: preoperative. Postop: postoperative. Surgeries: surgical procedures in the multilevel surgery. a: modified ZPP with one-layer closure, b: open CO<sub>2</sub>-laser tongue-base resection, c: bilateral septomeatoplasty, d: endoscopic sinusurgery, e: adenoidectomy.




**Figure 1.** Mean and standard deviation of apnea-hypopnea index (AHI) reduced from  $73.8 \pm 10.7$  to  $30.8 \pm 23.2$  events/h after the surgery.

desaturation index from  $65.1 \pm 10.7$  to  $24.1 \pm 17.0$  events/h ( $p < 0.001$ ), and decreased mean desaturation from  $10.8 \pm 5.9$  to  $5.8 \pm 1.9\%$  ( $p = 0.007$ ) (Fig. 5).


There was an event of left inferior tonsil wound hemorrhage in case 5 that resulted in an unplanned return to the operating room. No other major dehiscence or airway complication arose. The mean ESS score declined from  $9.67 \pm 5.23$  to  $7.83 \pm 5.11$ . However, the difference did not reach statistical significance ( $p = 0.39$  from a paired t-test). Scatter plots in Fig. 6 summarize the correlation tests of the two preoperative predictors. There was no statistical correlation between either of these two predictors and AHI reduction ( $r = -0.12$ ,  $p = 0.71$  and  $r = 0.118$ ,  $p = 0.714$ , respectively). Individual changes of BMI and AHI before and after the surgery were illustrated in Fig. 7. Although more patients (9 out of 12) lost weight after the surgery, BMI change was not statistically correlated with AHI reduction ( $r = 0.54$ ,  $p = 0.07$ ).

## Discussion

The results show that the multilevel surgery reduced AHI from 73.8 to 30.8 events/h, resulting in an AHI reduction of 58.3% ( $p < 0.001$ ). It cut desaturation index from 65.1 to 24.1 events/h ( $p < 0.001$ ) and mean desaturation from 10.8 to 5.8% ( $p = 0.007$ ). The surgery improved mean SpO<sub>2</sub> from 92.1 to 95.0% ( $p = 0.0427$ ) and minimum SpO<sub>2</sub> from 70.3 to 80.4% ( $p = 0.0088$ ). These results would help us on decision making with patients with very



**Figure 2.** Reduction of apnea–hypopnea index (AHI) in different Friedman anatomical stage.




**Figure 3.** Individual change of apnea–hypopnea index (AHI) after the surgery. Dotted, dashed, and solid line: anatomical stage 1, 2, and 3, respectively. Very big, big, median, small, and none circle: in the category of very severe, severe, moderate, mild obstructive sleep apnea (OSA), and normal, respectively.

severe OSA with residual on-CPAP AHIs, or have a low level of daytime PaO<sub>2</sub> but refuse the proposed direct skeletal surgery<sup>13</sup>, bariatric surgery<sup>23</sup>, or tracheostomy<sup>24</sup>, or refuse use of CPAP device.

A majority (83%) of the patients improved from the very severe category to milder ones and might reduce its comorbidity (e.g.,<sup>30–35</sup>). The rest (17% or 2 out of 12 patients) remained in the very severe group. The AHI made worse from 62.6 to 77.6 events/h after the surgery in case 7 and a little lessened from 85.4 to 68.6 events/h in case 6 (Table 1). To further understand the change made by the surgery in these 2 patients, we looked into matters of the PSGs before and after the surgery. In case 7 (the subject with worsened AHI after the surgery), the apnea part (i.e., OAI), improved from 38.6 to 31.9 events/h, and his minimum SpO<sub>2</sub> improved from 50 to 84%. In case 6, the OAI reduced from 73 to 1.3 events/h, and his minimum SpO<sub>2</sub> increased from 56 to 71%.

Minimum oxygen (O<sub>2</sub>) saturation has been listed as one of the main sleep-disordered breathing parameters besides AHI (e.g., see<sup>17,29,37,39</sup>) or used as one criterion to classify the severity of OSA (e.g., see<sup>40–42</sup>). So, we analyzed individual changes of minimum SpO<sub>2</sub> in addition to OAI (Fig. 4). The mean OAI reduced from 44.03 to 5.79 events/h with a reduction rate of 86.8%, which showed that the OAI reduction (86.8%) was better than AHI reduction (58.3%). These results showed that the surgery reduced more portion of obstructive apnea than hypopnea. The sum of apneas and hypopneas per hour did not cut as many as the sum of apneas per hour—some



**Figure 4.** Individual change of obstructive apnea index (A) and minimum O<sub>2</sub> saturation (B).  $p < 0.01$  in both panels.

obstructive apneas might become hypopneas after the surgery. The sample size of 12 in this study is small, although it is larger than 9 in Jacobowitz's<sup>43</sup>, 6 in Walker's<sup>40</sup>, 11 in Vilaseca's<sup>9</sup>, and 10 in Mickelson's<sup>44</sup> for patients of very severe OSA. That limits generalizability of the results. It needs future studies with a larger sample size to verify the observation.

To relate the surgical outcome with non-framework sleep surgeries in the literature, we reviewed 253 OSA related articles. Among them, 32 enrolled subjects with very severe OSA. Four provided detailed subject information and allowed us to calculate the mean AHI reduction. In Jacobowitz's report<sup>43</sup>, mean AHI reduction was 77.1/h after UPPP with or without genioglossus advancement, hyoid suspension, or tongue-base radiofrequency treatment ( $n = 9$ ), calculated from their Table 4. The mean AHI reduction was 38.2/h after UPPP ( $n = 6$ ), computed from Table 1 in Walker's study<sup>40</sup>; 14.4/h after UPPP and hyoid advancement with or without mandibular osteotomy with genioglossus advancement ( $n = 11$ ), reckoned from Table 2 in Vilaseca report<sup>9</sup>. The mean RDI reduction was 42.2/h after UPPP and midline glossectomy with or without septomeatoplasty ( $n = 10$ ), calculated from Table 1 in Mickelson's study<sup>44</sup>. Figure 8 presents the comparison of the present study with these reports.


Classical surgical success—AHI  $< 20/h$  and AHI reduction  $> 50%$ , first proposed by Sher et al.<sup>45</sup>—was frequently reported in associated studies with less severity and so might serve as another outcome measure for comparison. The overall success rate in our study was 33% (4/12), which was lower than Jacobowitz's 67% (6/9)<sup>43</sup> but higher than Walker's 17% (1/6)<sup>40</sup>, Vilaseca's 9% (1/11)<sup>9</sup>, and Mickelson's 30% (3/10)<sup>44</sup>.

Some studies recommend postoperative prudence treatment for patients with very severe OSA due to a higher risk of postoperative oxygen desaturation (e.g., Pang, K. P., Siow, J. & Tseng, P.<sup>42</sup>). They usually allocated these patients to the surgical intensive care unit (SICU) after the surgery (e.g., Rotenberg, B.<sup>46</sup>). In the present study, we cared for all patients in general ward areas with an oximeter monitor for 1 to 3 days while they breathe via the mouth because of nasal packing. No immediate or airway complication arose.


Some studies (e.g., Lin, H. S. et al.<sup>36</sup>) disclosed that BMI or preoperative AHI predicts surgical outcome. There was no correlation either between BMI and AHI reduction or between preoperative AHI and AHI reduction in this study. It needs future studies to find a good predictor for patients with very severe OSA.

## Conclusions


Our results show that modified Z-palatoplasty with one-layer closure in the one-stage multilevel surgery achieved a mean AHI and OAI reduction of 58.3% and 86.8% ( $p < 0.001$ ), respectively. It reduced the frequency (desaturation index) and level (mean desaturation) of desaturation and improved mean and minimum oxygen saturation. Analyses show a shift in the components of breathing disturbances. This non-framework surgery reduced more apnea than hypopnea and might convert some apnea to hypopnea.




**Figure 5.** (A) Individual change of mean oxyhemoglobin saturation of pulse oximetry (SpO2) after the surgery. It reveals a ceiling effect. (B) Individual change of desaturation index after the surgery. The surgery reduced the mean desaturation index by 41 events/h ( $p < 0.001$ ). (C) Individual change of mean desaturation after the surgery.



**Figure 6.** Scatter plot and correlation of preoperative body mass index (BMI) versus apnea–hypopnea-index (AHI) reduction (**A**) and preoperative AHI versus AHI reduction (**B**).



**Figure 7.** Scatterplot showed a trend of losing weight after the surgery but no statistic correlation between body-mass-index (BMI) change and apnea–hypopnea-index (AHI) reduction.



**Figure 8.** Comparison of surgical outcome in similar studies. See text for details.

### Data availability

The raw data in the current study are available from the supplementary information.

Received: 24 July 2018; Accepted: 7 April 2021

Published online: 19 April 2021

### References

1. Campanini, A., De Vito, A., Frassinetti, S. & Vicini, C. Role of skin-lined tracheotomy in obstructive sleep apnoea syndrome: personal experience. *Acta Otorhinolaryngol. Ital.* **24**, 68–74 (2004).
2. Haviv, Y. *et al.* A 2-year mean follow-up of oral appliance therapy for severe obstructive sleep apnea: a cohort study. *Oral Dis.* **21**, 386–392. <https://doi.org/10.1111/odi.12291> (2015).
3. Lavie, P., Lavie, L. & Herer, P. All-cause mortality in males with sleep apnoea syndrome: declining mortality rates with age. *Eur. Respir. J.* **25**, 514–520. <https://doi.org/10.1183/09031936.05.00051504> (2005).
4. Veasey, S. C. *et al.* Medical therapy for obstructive sleep apnea: A review by the Medical Therapy for Obstructive Sleep Apnea Task Force of the Standards of Practice Committee of the American Academy of Sleep Medicine. *Sleep* **29**, 1036–1044 (2006).
5. Maimon, N. & Hanly, P. J. Does snoring intensity correlate with the severity of obstructive sleep apnea?. *J. Clin. Sleep Med. (JCSM)* **6**, 475–478 (2010).
6. Brown, E. C. *et al.* Respiratory movement of upper airway tissue in obstructive sleep apnea. *Sleep* **36**, 1069–1076. <https://doi.org/10.5665/sleep.2812> (2013).
7. Olaithe, M. & Bucks, R. S. Executive dysfunction in OSA before and after treatment: A meta-analysis. *Sleep* **36**, 1297–1305. <https://doi.org/10.5665/sleep.2950> (2013).
8. Cohen, R. Obstructive sleep apnea: Oral appliance therapy and severity of condition. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endod.* **85**, 388–392 (1998).
9. Vilaseca, I., Morelló, A., Montserrat, J., Santamaría, J. & Iranzo, A. Usefulness of uvulopalatopharyngoplasty with genioglossus and hyoid advancement in the treatment of obstructive sleep apnea. *Arch. Otolaryngol. Head Neck Surg.* **128**, 435–440. <https://doi.org/10.1001/archotol.128.4.435> (2002).
10. Lee, L.-A. *et al.* Severity of obstructive sleep apnea syndrome and high-sensitivity C-reactive protein reduced after relocation pharyngoplasty. *Otolaryngol. Head Neck Surg.* **144**, 632–638. <https://doi.org/10.1177/0194599810395104> (2011).
11. Lee, L. A. *et al.* Energy types of snoring sounds in patients with obstructive sleep apnea syndrome: A preliminary observation. *PLoS ONE* **7**, e53481. <https://doi.org/10.1371/journal.pone.0053481> (2012).
12. Shiba, T. *et al.* Relationship between severity of obstructive sleep apnea syndrome and retinal nerve fiber layer thickness. *Am. J. Ophthalmol.* **157**, 1202–1208. <https://doi.org/10.1016/j.ajo.2014.01.028> (2014).
13. Huynh, N. T., Emami, E., Helman, J. I. & Chervin, R. D. Interactions between sleep disorders and oral diseases. *Oral Dis.* **20**, 236–245. <https://doi.org/10.1111/odi.12152> (2014).
14. Gay, P. C., Herold, D. L. & Olson, E. J. A randomized, double-blind clinical trial comparing continuous positive airway pressure with a novel bilevel pressure system for treatment of obstructive sleep apnea syndrome. *Sleep* **26**, 864–869. <https://doi.org/10.1093/sleep/26.7.864> (2003).
15. Berry, R., Brooks, R., Gamaldo, C. & American Academy of Sleep, M. *The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications* version 2.2. (American Academy of Sleep Medicine, 2015).
16. Lacedonia, D. *et al.* Characterization of obstructive sleep apnea-hypopnea syndrome (OSA) population by means of cluster analysis. *J. Sleep Res.* **25**, 724–730. <https://doi.org/10.1111/jsr.12429> (2016).
17. Anderson, F. E. *et al.* A randomized crossover efficacy trial of Oral CPAP (Oracle) compared with Nasal CPAP in the management of obstructive sleep apnea. *Sleep* **26**, 721–726. <https://doi.org/10.1093/sleep/26.6.721> (2003).



18. Levendowski, D. J., Seagraves, S., Popovic, D. & Westbrook, P. R. Assessment of a neck-based treatment and monitoring device for positional obstructive sleep apnea. *J. Clin. Sleep Med. (JCSM)* **10**, 863–871. <https://doi.org/10.5664/jcsm.3956> (2014).
19. Otake, K. *et al.* Glucose intolerance in Japanese patients with obstructive sleep apnea. *Intern. Med.* **48**, 1863–1868. <https://doi.org/10.2169/internalmedicine.48.2465> (2009).
20. Becker, H. F., Koehler, U., Stammnitz, A. & Peter, J. H. Heart block in patients with sleep apnoea. *Thorax* **53**(Suppl 3), S29–32 (1998).
21. Yetkin, O., Kunter, E. & Gunen, H. CPAP compliance in patients with obstructive sleep apnea syndrome. *Sleep Breath. SchlafAtmung* **12**, 365–367. <https://doi.org/10.1007/s11325-008-0188-4> (2008).
22. Li, H.-Y., Wang, P.-C., Lee, L.-A., Chen, N.-H. & Fang, T.-J. Prediction of uvulopalatopharyngoplasty outcome: Anatomy-based staging system versus severity-based staging system. *Sleep* **29**, 1537–1541. <https://doi.org/10.1093/sleep/29.12.1537> (2006).
23. Fritscher, L. G., Mottin, C. C., Canani, S. & Chatkin, J. M. Obesity and obstructive sleep apnea-hypopnea syndrome: The impact of bariatric surgery. *Obes. Surg.* **17**, 95–99. <https://doi.org/10.1007/s11695-007-9012-7> (2007).
24. Parati, G., Lombardi, C. & Narkiewicz, K. Sleep apnea: Epidemiology, pathophysiology, and relation to cardiovascular risk. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **293**, R1671–R1683. <https://doi.org/10.1152/ajpregu.00400.2007> (2007).
25. Suh, G. D. Evaluation of open midline glossectomy in the multilevel surgical management of obstructive sleep apnea syndrome. *Otolaryngol. Head Neck Surg.* **148**, 166–171. <https://doi.org/10.1177/0194599812464331> (2013).
26. Huang, E. I., Kuo, C.-L., Chou, Y.-T., Lin, Y.-C. & Huang, S.-Y. Modified Z-palatoplasty with one-layer closure in one-stage multi-level surgery for severe obstructive sleep apnea. *Auris Nasus Larynx* **45**, 791–795. <https://doi.org/10.1016/j.anl.2017.10.002> (2017).
27. Friedman, M., Wilson, M. & Kelley, K. Modification of Z-palatoplasty technique and review of five-year experience. *Oper. Tech. Otolaryngol. Head Neck Surg.* **23**, 30–35. <https://doi.org/10.1016/j.otot.2011.06.004> (2012).
28. Friedman, M. & Hwang, M. S. Z-palatopharyngoplasty. *Oper. Tech. Otolaryngol. Head Neck Surg.* **26**, 90–94. <https://doi.org/10.1016/j.otot.2015.03.008> (2015).
29. Friedman, M., Ibrahim, H. Z., Vidyasagar, R., Pomeranz, J. & Joseph, N. J. Z-palatoplasty (ZPP): a technique for patients without tonsils. *Otolaryngol. Head Neck Surg.* **131**, 89–100. <https://doi.org/10.1016/j.otohns.2004.02.051> (2004).
30. Zimmerman, M. E. & Aloia, M. S. A review of neuroimaging in obstructive sleep apnea. *J. Clin. Sleep Med. (JCSM)* **2**, 461–471 (2006).
31. Cheshire, K., Engleman, H., Deary, I., Shapiro, C. & Douglas, N. J. Factors impairing daytime performance in patients with sleep apnea/hypopnea syndrome. *Arch. Intern. Med.* **152**, 538–541. <https://doi.org/10.1001/archinte.1992.00400150068012> (1992).
32. Kim, H. C. *et al.* Sleep-disordered breathing and neuropsychological deficits. *Am. J. Respir. Crit. Care Med.* **156**, 1813–1819. <https://doi.org/10.1164/ajrccm.156.6.9610026> (1997).
33. Perry, J. C. *et al.* Consequences of subchronic and chronic exposure to intermittent hypoxia and sleep deprivation on cardiovascular risk factors in rats. *Respir. Physiol. Neurobiol.* **156**, 250–258. <https://doi.org/10.1016/j.resp.2006.10.004> (2007).
34. Bananian, S., Lehrman, S. G. & Maguire, G. P. Cardiovascular consequences of sleep-related breathing disorders. *Heart Dis.* **4**, 296–305. <https://doi.org/10.1097/00132580-200209000-00006> (2002).
35. Shahar, E. *et al.* Sleep-disordered breathing and cardiovascular disease. *Am. J. Respir. Crit. Care Med.* **163**, 19–25. <https://doi.org/10.1164/ajrccm.163.1.2001008> (2001).
36. Lin, H. S. *et al.* Transoral robotic surgery for treatment of obstructive sleep apnea: Factors predicting surgical response. *Laryngoscope* **125**, 1013–1020. <https://doi.org/10.1002/lary.24970> (2015).
37. Friedman, M., Ibrahim, H. & Joseph, N. J. Staging of obstructive sleep apnea/hypopnea syndrome: A guide to appropriate treatment. *Laryngoscope* **114**, 454–459. <https://doi.org/10.1097/00005537-200403000-00013> (2004).
38. Caples, S. M. *et al.* Surgical modifications of the upper airway for obstructive sleep apnea in adults: A systematic review and meta-analysis. *Sleep* **33**, 1396–1407 (2010).
39. IP, M. S. M. *et al.* Obstructive sleep apnea is independently associated with insulin resistance. *Am. J. Respir. Crit. Care Med.* **165**, 670–676. <https://doi.org/10.1164/ajrccm.165.5.2103001> (2002).
40. Walker, E. B., Frith, R. W., Harding, D. A. & Cant, B. R. Uvulopalatopharyngoplasty in severe idiopathic obstructive sleep apnoea syndrome. *Thorax* **44**, 205–208. <https://doi.org/10.1136/thx.44.3.205> (1989).
41. Friedman, M. *et al.* The occurrence of sleep-disordered breathing among patients with head and neck cancer. *Laryngoscope* **111**, 1917–1919. <https://doi.org/10.1097/00005537-200111000-00008> (2001).
42. Pang, K. P., Siow, J. & Tseng, P. Safety of multilevel surgery in obstructive sleep apnea: A review of 487 cases. *Arch. Otolaryngol. Head Neck Surg.* **138**, 353–357. <https://doi.org/10.1001/archoto.2012.130> (2012).
43. Jacobowitz, O. Palatal and tongue base surgery for surgical treatment of obstructive sleep apnea: A prospective study. *Otolaryngol. Head Neck Surg.* **135**, 264.e261. <https://doi.org/10.1016/j.otohns.2006.03.029> (2006).
44. Mickelson, S. A. & Rosenthal, L. Midline glossectomy and epiglottidectomy for obstructive sleep apnea syndrome. *Laryngoscope* **107**, 614–619. <https://doi.org/10.1097/00005537-199705000-00011> (1997).
45. Sher, A. E., Schechtman, K. B. & Piccirillo, J. F. The efficacy of surgical modifications of the upper airway in adults with obstructive sleep apnea syndrome. *Sleep* **19**, 156–177 (1996).
46. Rotenberg, B. Early perioperative outcomes after surgery for sleep apnea: A current review of the literature. *Curr. Anesthesiol. Rep.* **4**, 10–18. <https://doi.org/10.1007/s40140-013-0041-6> (2014).

## Author contributions

E.H. designed the study, collected data, read the PSGs, conducted statistical analysis, and wrote the first draft of this manuscript. Y.L. enrolled the patients, helped to interpret and present the data, and made critical comments on this manuscript. S.H., C.K.L. and C.M.L. enrolled the patients, analyzed the data, and revised the manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

**Correspondence** and requests for materials should be addressed to E.I.H.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021