

Adiponectin Upregulates Prolyl-4-Hydroxylase α 1 Expression in Interleukin 6-Stimulated Human Aortic Smooth Muscle Cells by Regulating ERK 1/2 and Sp1

Li Li^{1,2,3}, Ke Zhang^{1,3}, Xiao-Jun Cai^{1,2,3}, Min Feng¹, Yun Zhang¹, Mei Zhang^{1*}

1 The Key Laboratory of Cardiovascular Remodeling and Function Research, Chinese Ministry of Education and Chinese Ministry of Public Health, Shandong University Qilu Hospital, Jinan, Shandong, China, **2** Jinan Central Hospital affiliated to Shandong University, Jinan, Shandong, China

Abstract

Adiponectin is an anti-atherogenic adipokine that inhibits the development of plaque by mechanisms that are not completely understood. Extracellular matrix (ECM) may have a role in the pathogenesis of atherosclerosis. We explored the effect and mechanisms of adiponectin on the synthesis of prolyl-4-hydroxylase (P4H) in interleukin 6 (IL-6)-stimulated human aortic smooth muscle cells (HASMCs). P4H α 1 mRNA level was quantified by RT-PCR, the protein levels of phosphorylated extracellular signal-regulated kinase 1/2 (ERK1/2) and P4H α 1 were quantified by western blot analysis, and activation of specific protein 1 (Sp1) was determined by electrophoretic mobility shift assay and subcellular localization of Sp1 by immunofluorescence analysis. Adiponectin significantly increased P4H α 1 mRNA and protein levels in IL-6-stimulated HASMCs in a dose- and time-dependent manner. As well, ERK1/2 and Sp1 played a crucial role in the effect of adiponectin upregulating P4H α 1 expression in IL-6-stimulated HASMCs. Adiponectin abrogated the effects of IL-6 on collagen III level, which may indicate that P4H α 1 is essential for folding the procollagen polypeptide chains into stabilized collagen. Adiponectin attenuates IL-6-inhibited P4H α 1 synthesis and stabilizes collagen formation in HASMCs through a Sp1-ERK1/2-P4H α 1-dependent pathway.

Citation: Li L, Zhang K, Cai X-J, Feng M, Zhang Y, et al. (2011) Adiponectin Upregulates Prolyl-4-Hydroxylase α 1 Expression in Interleukin 6-Stimulated Human Aortic Smooth Muscle Cells by Regulating ERK 1/2 and Sp1. PLoS ONE 6(7): e22819. doi:10.1371/journal.pone.0022819

Editor: Gangjian Qin, Northwestern University, United States of America

Received: April 14, 2011; **Accepted:** June 29, 2011; **Published:** July 29, 2011

Copyright: © 2011 Li et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the National Basic Research Program of China (973 Program, 2011CB503906), the HI-TECH Technique and Development Program of China (863 Program, 2007AA02Z448), the National Natural Science Foundation of China (No. 30728025, 30970709), the Medical Science and Technology Development Program of Shandong Province (No. 2006GG2202039, 2009HW035). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: spear11@sina.com

³ These authors contributed equally to this work.

Introduction

Experimental results have demonstrated that the pathogenesis of atherosclerosis is based on several mechanisms. Extracellular matrix (ECM) may have a role in the pathogenesis of atherosclerosis [1]. ECM components, especially collagen, are thought to be important in the progression of atherosclerosis. Prolyl-4-hydroxylase (P4H) is a key intracellular enzyme essential for all known types of collagen maturation and secretion [2]. During collagen posttranslational processing, P4H α 1 folds the procollagen polypeptide chains into stable triple helical molecules [3]. Inhibition of P4H can decrease the level of collagen [4]. P4H is regulated by various cytokines, including tumor necrosis factor α , transforming growth factor β , and interleukins (ILs) [5]. Among them, IL-6 is one of the most potent cytokines involved in cardiovascular pathogenesis and actively regulates ECM metabolism [6]. Adiponectin is an adipocyte-specific plasma protein, has anti-inflammatory properties and might regulate ECM metabolism.

The promoter of P4H α 1 possesses several functional enhancer element-binding sites, including specific protein 1 (Sp1), NonO and hnRNP-K sites [7]. Sp1 is one of the first eukaryotic transcription factors to be identified and cloned [8]. It plays an

important role in the transcriptional regulation of ECM metabolism. Previous work has shown that mutations in the Sp1-binding sites of the TbRE are associated with low collagen expression [9]. Although accumulating data have demonstrated that cytokines might decrease P4H α 1 expression through pathways leading to the activation of nuclear proteins, little is known about the defense molecules – responsive cis-elements – of the P4H α 1 transcriptional promoter.

Intracellular signaling transduction pathways activated by cytokines are mitogen-activated protein kinase (MAPK) pathways, of which there are 3 distinct groups: extracellular signal-regulated kinase (ERK), Jun N-terminal kinase, and p38 MAPK. ERK is most associated with the biological effect of inflammation, adiponectin and ECM metabolism. Some studies demonstrated that inflammation and adiponectin are involved in the ERK1/2 pathway. Other studies demonstrated that ECM metabolism is also involved in the pathway [10]. Sp1, as a vital downstream molecule in the MAPK pathway, can be activated by ERK to induce the DNA binding activity [11].

We aimed to investigate whether adiponectin can upregulate P4H α 1 expression in IL-6-stimulated human aortic smooth muscle cells (HASMCs) and to elucidate the corresponding mechanisms to find novel approaches to the management of cardiovascular risk.

Materials and Methods

Materials

HASMCs, cell culture medium and supplements were from Cell Science (Carlsbad, CA, USA). Recombinant human IL-6 was from R&D Systems (Minneapolis, MN, USA). Antibodies against P4H α 1 and Sp1 were from Abcom (Cambridge, MA, USA). Antibodies against phosphor-ERK1/2 and β -actin were from Santa Cruz Biotechnology (Santa Cruz, CA, USA). PD98059 was from Biosource (Camarillo, CA, USA). WP631 was from Alexis Biochemicals (San Diego, CA, USA). All other chemicals were of the highest grade commercially available.

Cell Culture

HASMCs were cultured in smooth muscle cell culture medium up to passage 4. In all experiments, HASMCs were incubated in a humidified incubator at 37°C in a 95% air-5% CO₂ atmosphere until cells reached 80%–90% confluence and were rendered quiescent by serum-free starvation for at least 24 h. HASMCs were stimulated with IL-6 in the presence or absence of 10 multiplicities of infection (MOI) of adiponectin adenovirus (Invitrogen, Carlsbad, CA, USA) for 8 hr. HASMCs were pretreated with 20 μ M of the ERK inhibitor PD98059 or 0.1 μ M of the Sp1 inhibitor WP631 for 1 hr.

RT-PCR Analysis

After stimulation with 20 ng/ml IL-6 for 24 hr, RNA from cultured HASMCs was extracted with use of TriZol (Invitrogen, Carlsbad, CA, USA) following the manufacturer's protocol. Purified RNA was reverse-transcribed into cDNA (RevertAid M-MuLV Reverse Transcriptase, Fermentas UAB, Mainz, Germany). Aliquots of the reverse-transcribed mixture underwent PCR with specific primers as described [12]. Results from quantitative real-time PCR were analyzed by use of the 7500 Real-Time PCR System (Applied Biosystems, Foster City, CA, USA). Results were controlled for real-time efficiency and normalized with those for β -actin.

Western blot analysis

After stimulation with 20 ng/ml IL-6, cell lysates from cultured HASMCs were prepared with use of lysis buffer. Equal amounts of lysates were run on 14% poly-acrylamide gel, transferred onto nitrocellulose membranes (Bio-Rad, Hercules, CA, USA) and incubated at 4°C overnight with the primary antibodies mouse anti- β -actin (Santa Cruz Biotechnology; 1:1000 dilution), goat anti-P4H α 1 (Abcom; 1:250) or rabbit anti-phospho-specific p44/42 MAPK (Thr202 and Tyr204) (Santa Cruz Biotechnology; 1:250). The blots were then incubated with peroxidase-conjugated secondary antibodies (Santa Cruz Biotechnology; 1:2000 dilution) at room temperature for 2 hr, and membranes underwent detection by enhanced chemiluminescence plus reagents (Millipore, Billerica, MA, USA).

Electrophoretic mobility shift assay (EMSA) of Sp1

Briefly, cultured HASMCs were pretreated with WP631 for 1 hr, then stimulated with 20 ng/ml IL-6 for 1 hr. Nuclear extracts were prepared (Active Motif, Carlsbad, CA, USA); the nuclear fraction was transferred into a pre-chilled microcentrifuge tube and stored at -80°C until use. The protein concentration of the cell extract was measured by the Bradford assay (Bio-Rad, Hercules, CA, USA). Activation of Sp1 was determined by EMSA (Roche, Madison, WI, USA) according to the manufacturer's instructions. The generated chemiluminescent signals were recorded on an imaging device.

Immunofluorescence

Subcellular localization of Sp1 was determined by immunofluorescence analysis. Cultured HASMCs were pretreated with the Sp1 inhibitor WP631 or ERK inhibitor PD98059 for 1 hr before stimulation with 20 ng/ml IL-6 for 1 hr. Then cells were fixed with paraformaldehyde. Permeabilized cells were stained by direct immunofluorescence with a FITC-conjugated monoclonal antibody against Sp1 (1:100 dilution). Inclusion-containing cells were visualized under a light fluorescent microscope (Olympus, Japan) and a Bio-Rad MRC 600 confocal laser scanning microscope (Bio-Rad, Hercules, CA).

ELISA

Cultured HASMCs were pretreated with WP631 or PD98059 for 1 hr. After stimulation with 20 ng/ml IL-6 for 24 hr, the levels of type I and III procollagen and collagen secreted from HASMCs were quantified by ELISA (R&D Systems) according to the manufacturer's recommended protocols.

Statistical Analysis

Data are presented as mean \pm SD. Statistical analyses involved ANOVA with subsequent Scheffe *t* test. A $P < 0.05$ was considered statistically significant.

Results

1. Effect of dose and time of IL-6 stimulation on P4H α 1 expression

To explore the effect of IL-6 on the expression of P4H α 1, HASMCs were treated with different doses of recombinant human IL-6 and for different times before being harvested for measurement of P4H α 1 mRNA and protein levels. The expression of P4H α 1 was lowest at 20 ng/ml IL-6 and at 24 hr (Figure 1).

2. Effect of dose and time of adiponectin on P4H α 1 expression

IL-6-stimulated HASMCs were treated with various MOI of adiponectin adenovirus for different times before being harvested for measurement of P4H α 1 mRNA and protein levels. The mRNA and protein expression of P4H α 1 peaked at 10 MOI adiponectin and at 8 hr (Figure 2).

3. Sp1 regulates the effect of adiponectin upregulating P4H α 1 expression in IL-6-stimulated HASMCs

3.1 Effect of adiponectin on Sp1 activation induced by IL-6 in HASMCs. IL-6 significantly upregulated the binding activity of Sp1 in HASMCs as compared with control cells ($P < 0.05$; Figure 3A). Adiponectin alone had no effect on the binding activity of the Sp1 in HASMCs, but it inhibited the IL-6-induced binding activity of Sp1 by 60% ($P < 0.05$). Treatment with WP631, an inhibitor of Sp1, inhibited the IL-6 induced binding activity of Sp1 by 95% ($P < 0.05$). The combination of WP631 and adiponectin abrogated the Sp1 binding activity induced by IL-6 by 90% ($P < 0.05$), whereas adiponectin or WP631 alone did not affect Sp1 binding activity ($P > 0.05$).

3.2 Effect of Adiponectin on Subcellular Localization and Expression of Sp1 Induced by IL-6 in HASMCs. Most unstimulated cells showed diffuse staining in the cytoplasm, as well as weak staining in the nucleus, under normal growth conditions (Fig. 3B). With IL-6 treatment for 1 hr, staining for Sp1 in the nucleus was eight-fold higher than that in control cells ($P < 0.05$). Eight hours after the addition of adiponectin to the cells, the level of nuclear staining was abrogated by 70% ($P < 0.05$). After




Figure 1. Effect of dose and time of IL-6 treatment on P4Hα1 expression. (A and C) mRNA expression of P4Hα1. (B and D) Protein expression of P4Hα1. Data are representative of 3 independent experiments (means±SD). *, $p < 0.05$ vs. con. doi:10.1371/journal.pone.0022819.g001

preincubation with WP631 for 1 hr, the level of nuclear staining was weak ($P < 0.05$); the combination of WP631 and adiponectin also showed weak staining in the nucleus ($P < 0.05$), which confirms the specificity of the Sp1 inhibitor used. However, staining for Sp1 in the nucleus was not influenced by adiponectin or WP631 alone ($P > 0.05$).

3.3 Effect of adiponectin on the expression of P4Hα1 Induced by IL-6 in HASMCs. IL-6 significantly downregulated the level of P4Hα1 mRNA by 65% ($P < 0.05$) and protein by 70% ($P < 0.05$) as compared with control cells (Fig. 3C,D). Eight hours after the addition of adiponectin to the cells, the level of P4Hα1 mRNA was increased 5.8-fold ($P < 0.05$) and that of P4Hα1 protein six-fold ($P < 0.05$). After preincubation with WP631 for 1 hr or the Sp1 inhibitor WP631 and adiponectin combined, the level of P4Hα1 mRNA and protein returned to the level of control cells. However, the expression of P4Hα1 was not influenced by adiponectin or WP631 alone ($P > 0.05$).

4. ERK1/2 leads to adiponectin upregulating P4Hα1 expression in IL-6- mediated HASMCs in an Sp1-dependent manner

4.1 Effect of adiponectin on phosphorylation of ERK induced by IL-6 in HASMCs. IL-6 significantly induced the phosphorylation of ERK in HASMCs as compared with control cells ($P < 0.05$) (Fig. 4A). Eight hours after the addition of adiponectin to the cells, the phosphorylation of ERK induced by IL-6 was abrogated by 70% ($P < 0.05$). After preincubation with PD98059 for 1 hr, the phosphorylation of ERK induced by IL-6 was abrogated by 95% ($P < 0.05$). After preincubation with

PD98059 and adiponectin combined, the phosphorylation of ERK induced by IL-6 was abrogated by 92% ($P < 0.05$). However, the phosphorylation of ERK was not influenced by adiponectin or PD98059 alone ($P > 0.05$).

4.2 Effect of ERK on Sp1 activation induced by IL-6 in HASMCs. IL-6 significantly upregulated the binding activity of Sp1 in HASMCs as compared with control cells ($P < 0.05$) (Fig. 4B). Eight hours after the addition of adiponectin to the cells, the binding activity of Sp1 induced by IL-6 was reduced by 70% ($P < 0.05$). After preincubation with PD98059 for 1 hr, the binding activity of Sp1 induced by IL-6 was reduced by 95% ($P < 0.05$). The combination of PD98059 and adiponectin reduced the IL-6-induced binding activity of Sp1 by 93% ($P < 0.05$). However, the binding activity of Sp1 was not influenced by adiponectin or PD98059 alone ($P > 0.05$).

4.3 Effect of ERK on subcellular localization and expression of Sp1 induced by IL-6 in HASMCs. Most unstimulated cells had diffuse staining in the cytoplasm, as well as weak staining in the nucleus, under normal growth conditions (Fig. 4C). With 1-h treatment with IL-6, staining for Sp1 in the nucleus was increased seven-fold in HASMCs as compared with control cells ($P < 0.05$). Eight hours after the addition of adiponectin, the level of nuclear staining was reduced by 70% ($P < 0.05$). After preincubation with PD98059 for 1 hr, the level of nuclear staining was weak ($P < 0.05$). The combination of PD98059 and adiponectin also produced weak staining in the nucleus ($P < 0.05$). However, staining for Sp1 in the nucleus was not influenced by adiponectin or PD98059 alone ($P > 0.05$).

4.4 Effect of ERK on the expression of P4Hα1 induced by IL-6 in HASMCs. IL-6 significantly downregulated the level of




Figure 2. Effect of dose and time of adiponectin treatment on to P4H α 1 expression. (A and C) mRNA expression of P4H α 1. (B and D) Protein expression of P4H α 1. Data are representative of 3 independent experiments (means \pm SD). *, $p < 0.05$ vs. con; #, $p < 0.05$ vs. IL-6. doi:10.1371/journal.pone.0022819.g002

P4H α 1 mRNA by 65% ($P < 0.05$) and protein by 70% ($P < 0.05$) in HASMCs as compared with control cells (Fig. 4D,E). Eight hours after the addition of adiponectin, the level of P4H α 1 mRNA was increased 6.3-fold ($P < 0.05$) and protein seven-fold ($P < 0.05$). After preincubation with PD98059 for 1 hr or the combination of PD98059 and adiponectin, the level of P4H α 1 mRNA and protein returned to the level of control cells ($P < 0.05$). However, the expression of P4H α 1 was not influenced by adiponectin or PD98059 alone ($P > 0.05$).

5. Effect of adiponectin overexpression on collagen production

The content of procollagen I and III did not change significantly with or without adiponectin or IL-6 treatment (Fig. 5A,B), whereas after the addition of IL-6 to the cells, the content of collagen III decreased significantly in HASMCs as compared with control cells (IL-6: 8.20 ± 0.3 μ g/L vs. control: 12.1 ± 0.1 μ g/L; $P < 0.05$) (Fig. 5D). Eight hours after the addition of adiponectin to the cells, the content of collagen III increased significantly (IL-6: 8.20 ± 0.3 μ g/L vs. IL-6+APN: 10.97 ± 0.2 μ g/L; $P < 0.05$) (Fig. 5D), whereas the content of collagen I did not change significantly in all groups (Fig. 5C).

Discussion

Arterial interstitial collagen confers tensile strength on the fibrous cap and thus determines plaque stability and vulnerability

to rupture. Therefore, the synthesis of collagen in the fibrous cap may be directly responsible for plaque rupture [13]. In this study, we aimed to investigate the role of ECM in the pathogenesis of atherosclerosis and the potential mechanism of adiponectin upregulating P4H α expression in IL-6-stimulated HASMCs. First, we determined that adiponectin can upregulate the expression of P4H α 1 in HASMCs induced by IL-6. Second, we showed that adiponectin upregulating P4H α 1 expression in IL-6-mediated HASMCs depended in part on ERK1/2 activation and Sp1 DNA binding activity. Adiponectin also increased collagen III level in HASMCs alone with upregulated P4H α 1 expression.

Adiponectin is an adipose tissue-secreted protein with anti-inflammatory and anti-atherogenic properties. A proinflammatory state in adipose tissue can reduce secretion of adiponectin and increase that of several cytokines, including TNF α , IL-6, and other proinflammatory molecules [14]. Phosphorylation of GATA2 by Akt increases adipose tissue differentiation and reduces adipose tissue-related inflammation [15]. Other studies have investigated the ratio of TNF to adiponectin and the consequences for inflammation and atherosclerosis. Atherosclerosis was ameliorated in gAd Tg apoE-deficient mice, which was associated with decreased expression of class A scavenger receptor and TNF α [16]. Full-length adiponectin was found to upregulate TNF α and IL-6 protein production differently [17]. In our research, adiponectin attenuated the IL-6-inhibited P4H α 1 synthesis in HASMCs and consequently changed the content of collagen. All




Figure 3. Sp1 regulates the effect of adiponectin upregulating P4Ha1 expression in IL-6-mediated HASMCs. (A) Sp1 DNA binding activity. (B) Subcellular localization of Sp1. (C) mRNA expression of P4Ha1. (D) Protein expression of P4Ha1. *, $p < 0.05$ vs. con; #, $p < 0.05$ vs. IL-6. doi:10.1371/journal.pone.0022819.g003

of these data suggest the significant inverse relationship between level of adiponectin and IL-6 or TNF- α associated with the development of inflammation-related atherosclerosis. Although adiponectin has anti-inflammatory and antiatherogenic properties, it has no effect on normal cells, unless in combination with inflammation. Adiponectin significantly suppressed LPS-induced IL-6 and monocyte chemoattractant protein 1 production and also significantly reduced NF-kappaB activity and IkkappaB-alpha and IKK gene expression as compared with 3T3-L1 induced by LPS alone [18]. Adiponectin may reduce LPS-induced NO production and nitrosative stress and prevent adventitial fibroblasts (AFs) from proliferating, transforming to myofibroblasts, and migrating to the intima, thus worsening atherosclerosis, by inhibiting the AdipoR1-

AMPK-iNOS pathway in AFs [19]. Our present results also show that adiponectin had no effect on P4Ha1 expression, Sp1 DNA binding activity or ERK phosphorylation under basal conditions. With IL-6 stimulation, adiponectin reverted the IL-6-induced abnormal P4Ha1 expression, Sp1 DNA binding activity and ERK phosphorylation to the normal level at least 2 times more than the control. Therefore, adiponectin has anti-inflammatory properties and might regulate inflammatory responses in atherosclerotic lesions.

Several studies have identified that adiponectin might regulate ECM metabolism in atherosclerotic lesions [20-22]. Our previous study of an atherosclerosis mouse model indicates that adiponectin could significantly increase the expression of P4Ha1 in plaque,




Figure 4. ERK1/2 leads to adiponectin upregulating P4H α 1 expression in IL-6-stimulated HASMCs. (A) p-ERK level. (B) Sp1 DNA-binding activity. (C) Subcellular localization of Sp1. (D) mRNA expression of P4H α 1. (E) Protein expression of P4H α 1. *, $p < 0.05$ vs. con; #, $p < 0.05$ vs. IL-6. doi:10.1371/journal.pone.0022819.g004

which may play a major role in the development of the thick fibrous cap of atherosclerotic plaque and prevent plaque rupture [23]. These data suggest that adiponectin might have anti-atherosclerotic properties and regulate ECM metabolism in atherosclerotic lesions, but the potential mechanism was unknown. We further studied the regulation of adiponectin on P4H α 1 expression *in vitro*. The expression of P4H α 1 was mostly induced by IL-6. Adiponectin could upregulate the expression of P4H α 1 in HASMCs induced by IL-6 in a dose- and time-dependent manner, so adiponectin might have anti-inflammatory properties and might regulate ECM metabolism.

To identify signaling pathways in this induction, we studied the activation of Sp1, a known transcriptional factor, that might play an important role in the transcriptional regulation of ECM

metabolism [24–26]. IL-6, sIL-6R, or both were found to decrease both the ratio of Sp1 to Sp3 and DNA binding activities, thus inhibiting COL2A1 transcription [27]. Our results showed that Sp1 DNA binding activity was enhanced on exposure to IL-6. Both the mRNA and protein levels of P4H α 1 were significantly attenuated. To clarify that the effect depended on Sp1, we used the Sp1 inhibitor WP631, which efficiently inhibited Sp1 DNA binding activity. In the presence of the inhibitor, P4H α 1 mRNA and protein levels were significantly enhanced. Adiponectin could counteract all of these effects. Therefore, Sp1 contributed to the adiponectin-induced P4H α 1 expression.

MAPK pathways are important mediators of signal transduction and are involved in multiple intracellular signaling cascades and activated by cytokines [28–31]. Some studies demonstrated




Figure 5. Effect of adiponectin overexpression on collagen production. Content of (A) procollagen I, (B) procollagen III, (C) collagen I, (D) collagen III. Data are representative of 3 independent experiments (means \pm SD). *, $p < 0.05$ vs. con; #, $p < 0.05$ vs. IL-6. doi:10.1371/journal.pone.0022819.g005

that inflammation, adiponectin and ECM metabolism are all involved in the ERK1/2 pathway [32–34]. Sp1, as a vital downstream molecule in the MAPK pathway, can be activated by ERK and induce the Sp1 DNA binding activity [35]. Our results showed phosphorylation of ERK1/2 and Sp1 DNA binding activity enhanced after exposure to IL-6. Both P4H α 1 mRNA and protein levels were significantly attenuated. To clarify that the effect depends on ERK1/2, we used the ERK1/2 inhibitor PD98059, which efficiently inhibited phosphorylation of ERK1/2 and Sp1 DNA binding activity. In the presence of the inhibitor, P4H α 1 mRNA and protein levels were significantly enhanced. Adiponectin could counteract all of these effects. These results supported that adiponectin upregulating P4H α 1 expression in IL-6-mediated HASMCs depended in part on ERK1/2 activation.

The organic matrix of a plaque mainly consists of types I and III collagen. Many studies have fully discussed their content and changes in content in atherosclerotic plaques. The completely processed and cross-linked type III collagen seems to be the major collagen type in atherosclerotic plaques [36]. The density of type III was increased in diabetes mellitus (DM) plaques with a non-significant reduction in type I density in DM as compared with non-DM plaques [37]. The pattern of GPVI-Fc binding was similar to the immunostaining pattern of collagen type III but differed from that of collagen type I, which was more intense in the

cap than in the core [38]. We further explored the effect of adiponectin overexpression on collagen content and found no difference in the content of procollagen I and III on IL-6 stimulation with or without adiponectin treatment; only collagen III content was reversed by adiponectin. Therefore, type III collagen seem to be the major collagen type in atherosclerotic plaques. P4H α 1 is essential for collagen maturation and secretion by folding the procollagen polypeptide chains into stable triple helical molecules, so the change in collagen III content may reflect the activity of P4H α 1 in HASMCs.

In conclusion, we show for the first time that adiponectin attenuates the IL-6-inhibited P4H α 1 synthesis in HASMCs and therefore collagen III content was changed. This effect of adiponectin is through ERK1/2-dependent pathways and by inhibiting Sp1 activation. Our results suggest that adiponectin may play a major role in the development of the thick fibrous cap of atherosclerotic plaque and stabilize plaque by increasing P4H α 1 synthesis in HASMCs stimulated by IL-6.

Author Contributions

Conceived and designed the experiments: LL YZ MZ. Performed the experiments: LL KZ XJC FM. Analyzed the data: LL MZ. Contributed reagents/materials/analysis tools: LL KZ XJC. Wrote the paper: LL.

References

- Libby P (1995) Molecular bases of the acute coronary syndromes. *Circulation* 91: 2844–2850.
- Annunen P, Autio-Harjainen H, Kivirikko KI (1998) The novel type II prolyl 4-hydroxylase is the main enzyme form in chondrocytes and capillary endothelial cells, whereas the type I enzyme predominates in most cells. *J Biol Chem* 273: 5989–5992.
- Kivirikko KI, Pihlajaniemi T (1998) Collagen hydroxylases and the protein disulfide isomerase subunit of prolyl 4-hydroxylases. *Adv Enzymol Relat Areas Mol Biol* 72: 325–398.
- Rocnik EF, Chan BM, Pickering JG (1998) Evidence for a role of collagen synthesis in arterial smooth muscle cell migration. *J Clin Invest* 101: 1889–1898.
- Raveendran M, Senthil D, Utama B, Shen Y, Dudley D, et al. (2004) Cigarette suppresses the expression of P4H α and vascular collagen production. *Biochem Biophys Res Commun* 323: 592–598.
- Valgimigli M, Merli E, Malagutti P, Soukhomovskaia O, Cicchitelli G, et al. (2004) Hydroxyl radical generation, levels of tumor necrosis factor- α , and progression to heart failure after acute myocardial infarction. *J Am Coll Cardiol* 43: 2000–2008.
- Chen L, Shen YH, Wang X, Wang J, Gan Y, et al. (2006) Human Prolyl-4-hydroxylase $\{\alpha\}$ Transcription Is Mediated by Upstream Stimulatory Factors. *J Biol Chem* 281: 10849–10855.
- Jackson SP, Tjian R (1988) O-glycosylation of eukaryotic transcription factors - implications for mechanisms of transcriptional regulation. *Cell* 55: 125–133.
- Milanini-Mongiati J, Pouységur J, Pagès G (2002) Identification of two Sp1 phosphorylation sites for p42/p44 mitogen-activated protein kinases: their implication in vascular endothelial growth factor gene transcription. *J Biol Chem* 277: 20631–20639.
- Rockel JS, Bernier SM, Leask A (2009) Egr-1 inhibits the expression of extracellular matrix genes in chondrocytes by TNF α -induced MEK/ERK signalling. *Arthritis Res Ther* 11: R8.
- Curry JM, Eubank TD, Roberts RD, Wang Y, Pore N, et al. (2008) M-CSF signals through the MAPK/ERK pathway via Sp1 to induce VEGF production and induces angiogenesis in vivo. *PLoS One* 3: e3405.
- Zhang C, Zhang MX, Shen YH, Burks JK, Zhang Y, et al. (2007) TNF- α Suppresses Prolyl-4-Hydroxylase α 1 Expression via the ASK1–JNK–NonO Pathway. *Arterioscler Thromb Vasc Bio* 27: 1760–1767.
- Kivirikko KI, Pihlajaniemi T (1998) Collagen hydroxylases and the protein disulfide isomerase subunit of prolyl 4-hydroxylases. *Adv Enzymol Relat Areas Mol Biol* 72: 325–398.
- Sopasaki VR, Sandqvist M, Gustafson B, Hammarstedt A, Schmelz M, et al. (2004) High local concentrations and effects on differentiation implicate interleukin-6 as a paracrine regulator. *Obes Res* 12: 454–460.
- Menghini R, Marchetti V, Cardellini M, Hribal ML, Mauriello A, et al. (2005) Phosphorylation of GATA2 by Akt increases adipose tissue differentiation and reduces adipose tissue-related inflammation: a novel pathway linking obesity to atherosclerosis. *Circulation* 111: 1946–1953.
- Yamauchi T, Kamon J, Waki H, Imai Y, Shimozawa N, et al. (2003) Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis. *J Biol Chem* 278: 2461–2468.
- Kyriazi E, Tsiotra PC, Boutati E, Ikonomidis I, Fountoulaki K, et al. (2011) Effects of Adiponectin in TNF- α , IL-6, and IL-10 Cytokine Production from Coronary Artery Disease Macrophages. *Horm Metab Res*, [Epub ahead of print].
- Zoico E, Garbin U, Olisio D, Mazzali G, Fratta Pasini AM, et al. (2009) The effects of adiponectin on interleukin-6 and MCP-1 secretion in lipopolysaccharide-treated 3T3-L1 adipocytes: role of the NF- κ B pathway. *Int J Mol Med* 24: 847–851.
- Cai XJ, Chen L, Li L, Feng M, Li X, Zhang K, et al. (2010) Adiponectin inhibits lipopolysaccharide-induced adventitial fibroblast migration and transition to myofibroblasts via AdipoR1-AMPK-iNOS pathway. <http://www.ncbi.nlm.nih.gov/pubmed/218228>.
- Ezure T, Amano S (2007) Adiponectin and leptin up-regulate extracellular matrix production by dermal fibroblasts. *Biofactors* 31: 229–236.
- Kang EH, Lee YJ, Kim TK, Chang CB, Chung JH, Shin K, Lee EY, Lee EB, Song YW (2010) Adiponectin is a potential catabolic mediator in osteoarthritis cartilage. *Arthritis Res Ther* 12: R231.
- Zhang H, Xie H, Zhao Q, Xie GQ, Wu XP, Liao EY, Luo XH (2010) Relationships between serum adiponectin, apelin, leptin, resistin, visfatin levels and bone mineral density, and bone biochemical markers in post-menopausal Chinese women. *J Endocrinol Invest*; 33: 707–711.
- Li L, Cai XJ, Feng M, Rong YY, Zhang Y, et al. (2010) Effect of adiponectin overexpression on stability of preexisting plaques by inducing prolyl-4-hydroxylase expression. *Circ J* 74: 552–559.
- Mukhopadhyay A, Khoo A, Cheong HH, Chan SY, Aalami O, et al. (2007) Targeting of Sp1 transcription factor: a novel therapeutic approach for keloids, an in vitro analysis. *Exp Dermatol* 16: 1023–1031.
- Lee M, Song SU, Ryu JK, Suh JK (2004) Sp1-dependent regulation of the tissue inhibitor of metalloproteinases-1 promoter. *J Cell Biochem* 91: 1260–1268.
- Takahara T, Smart DE, Oakley F, Mann DA (2004) Induction of myofibroblast MMP-9 transcription in three-dimensional collagen I gel cultures: regulation by NF- κ B, AP-1 and Sp1. *Int J Biochem Cell Biol* 36: 353–363.
- Porée B, Kypriotou M, Chadjiachristos C, Beauchef G, Renard E, et al. (2008) Interleukin-6 (IL-6) and/or soluble IL-6 receptor down-regulation of human type II collagen gene expression in articular chondrocytes requires a decrease of Sp1/Sp3 ratio and of the binding activity of both factors to the COL2A1 promoter. *J Biol Chem* 283: 4850–4865.
- Sigala I, Zacharatos P, Toumpanakis D, Michailidou T, Nougssia O, et al. (2011) MAPKs and NF- κ B differentially regulate cytokine expression in the diaphragm in response to resistive breathing: the role of oxidative stress. *Am J Physiol Regul Integr Comp Physiol*. *Am J Physiol Regul Integr Comp Physiol* 300: R1152–1162.
- Kim JY, Kim HJ, Kim SM, Park KR, Jang HJ, et al. (2011) Methylene chloride fraction of the leaves of Thuja orientalis inhibits in vitro inflammatory biomarkers by blocking NF- κ B and p38 MAPK signaling and protects mice from lethal endotoxemia. *J Ethnopharmacol* 133: 687–695.
- Hot A, Zrioual S, Toh ML, Lenief V, Miossec P (2011) IL-17A- versus IL-17F-induced intracellular signal transduction pathways and modulation by IL-17RA and IL-17RC RNA interference in rheumatoid synoviocytes. *Ann Rheum Dis* 70: 341–348.
- Sundararaj KP, Samuvel DJ, Li Y, Sanders JJ, Lopes-Virella MF, et al. (2009) Interleukin-6 released from fibroblasts is essential for upregulation of matrix metalloproteinase-1 expression by u937 macrophages in coculture- crosstalking between fibroblasts and u937 macrophages exposed to high glucose. *J Biol Chem* 284: 567–573.
- Feng M, Cai XJ, Zhang W, Liu XL, Chen L, et al. (2010) Interleukin-6 enhances matrix metalloproteinase-14 expression via the RAF-mitogen-activated protein kinase kinase-extracellular signal-regulated kinase 1/2-activator protein-1 pathway. *Clin Exp Pharmacol Physiol* 37: 162–166.
- Barnes PJ, Adcock IM (1998) Transcription factors and asthma. *Eur Respir J* 12: 221–234.
- Legendre F, Bogdanowicz P, Boumediene K, Pujol JP (2005) Role of interleukin 6(IL-6)/IL-6R-induced signal transducers and activators of transcription and mitogen-activated protein kinase/extracellular. *J Rheumatol* 32: 1307–1316.
- Qureshi HY, Sylvester J, El Mabrouk M, Zafarullah M (2005) TGF- β -induced expression of tissue inhibitor of metalloproteinases-3 gene in chondrocytes is mediated by extracellular signal-regulated kinase pathway and Sp1 transcription factor. *J Cell Physiol* 203: 345–352.
- Bode MK, Mosorin M, Satta J, Risteli L, Juvonen T, et al. (1999) Complete processing of type III collagen in atherosclerotic plaques. *Arterioscler Thromb Vasc Bio* 19: 1506–1511.
- Purushothaman KR, Purushothaman M, Muntner P, Lento PA, O'Connor WN, et al. (2011) Inflammation, neovascularization and intra-plaque hemorrhage are associated with increased reparative collagen content: Implication for plaque progression in diabetic atherosclerosis. *Vasc Med* 16: 103–108.
- Schulz C, Penz S, Hoffmann C, Langer H, Gillitzer A, et al. (2008) Platelet GPVI binds to collagenous structures in the core region of human atherosclerotic plaque and is critical for atheroprotection in vivo. *Basic Res Cardiol* 103: 356–367.