


Environmental toxicants and health adversities: A review on interventions of phytochemicals

Journal of Public Health Research
2023, Vol. 12(2), 1–20
© The Author(s) 2023
DOI: 10.1177/22799036231181226
journals.sagepub.com/home/phj



**Adewale Allen Sokan-Adeaga¹, Micheal Ayodeji Sokan-Adeaga²,
Eniola Deborah Sokan-Adeaga³, Anastasia Nkem Oparaji⁴, Hoseinzadeh Edris⁵,
Esther Oluwabukunola Tella¹, Francis Adeniyi Balogun⁶, Muhammad Aledeh^{7,8} 
and Oluwatosin Emmanuel Amubieya⁹**

Abstract

Toxicity arising from environmental contaminants has attracted global interest in the last few decades, due to the high morbidity and mortality associated with them. Efforts have been made to combat the consequential outcomes of environmental toxicity in humans through traditional remediation techniques and therapeutic measures which have been hampered by one or more limitations. Consequently, this scenario has triggered interest in the medicinal properties of phytochemicals. Thus, this review gives a succinct and in-depth elucidation of the various environmental contaminants and their toxicity effects on humans. It delves into the various classes of phytochemicals and their intervention roles. The study adopted a desk review of existing literatures from scientific reports and peer reviewed articles through triangulation of data sources. “Phytochemicals” are group of secondary metabolites obtained from plants with medicinal properties. These groups of compounds are included but not limited to flavonoids, tannins, saponins, alkaloids, cardenoloids, terpenoids, and phytosteroids. This review corroborates the prophylactic and therapeutics efficacy of these phytochemicals as anti-metastatic, anti-inflammatory, anti-aging, anti-oxidant, anti-microbial and live saving substances with empirical findings from several laboratory, clinical trials and epidemiologic studies. It conclude that given the wide range of medicinal properties of phytochemicals, there is an urgent need for its full optimization in the pharmaceutical industry and future studies should focus on identifying the bioactive molecules in these compounds and its effectiveness against mixer toxicity.

Keywords

Environmental toxicants, phytochemicals, prophylactics, therapeutics, plants, intervention, medicinal properties

Date received: 5 November 2022; accepted: 28 April 2023

¹Department of Environmental Health Science, Faculty of Public Health, College of Medicine, Lead City University, Ibadan, Nigeria

²Department of Community Health and Primary Health Care, Faculty of Clinical Sciences, College of Medicine, University of Lagos, Lagos, Nigeria

³Department of Physiology, Faculty of Basic Medical Sciences, College of Medicine, Ladoke Akintola University of Technology (LAUTECH), Ogbomosho, Oyo State, Nigeria

⁴Department of Nursing Science, Chrisland University, Abeokuta, Ogun State, Nigeria

⁵Incubation and Innovation Center, Saveh University of Medical Sciences, Saveh, Iran

⁶Department of Community Health, Faculty of Public Health, College of Medicine, Lead City University, Ibadan, Nigeria

⁷College of Health, Psychology and Social Care, University of Derby, Derby, United Kingdom

⁸Wiener Gesundheitsverbund, Psychiatric Department, Klinik Donaustadt, Vienna, Austria

⁹Department of Environmental Health Sciences, Faculty of Public Health, University of Ibadan, Ibadan, Nigeria

Corresponding author:

Adewale Allen Sokan-Adeaga, Department of Environmental Health Science, Faculty of Public Health, College of Medicine, Lead City University Ibadan, Toll Gate Area, P.O. Box 30678, Secretariat, Ibadan, Oyo State 200255, Nigeria.

Email: sokanadeaga.adewalleallen@yahoo.com



Introduction

One of the leading causes of diseases, high morbidity and mortality rates in nations across the globe is environmental contamination. According to the World Health Organization (WHO) approximately 7 million deaths per annum is associated with pollution, mostly due to environmental toxicants which are air borne^{1,2}; this casualties is greater than the cumulative death figures recorded from communicable diseases such as malaria, tuberculosis and HIV/AIDS altogether.^{3,4} The greater proportions of these pollution-related deaths take place in developing countries of the world. Exposures to environmental toxicants at early stage in life are primarily precarious.^{5,6} Susceptibility is most pronounced during the first few weeks of the fetal life when the organs primordial are formed. Vulnerability to environmental toxicants during this delicate period can result in perpetual anatomical, behavioral and metabolic alterations. This anomaly can be expressed as short or long-term disorder at any specific point along an individual lifespan from neonate to senescence.⁷ Major acute maladies ascribed to environmental pollution at infancy are diarrheal and pneumonia disease⁸ while the chronic non-communicable diseases associated with early exposure to environmental toxicants include dysfunctional neurobehavioral development, childhood and adult asthma, hypertension, diabetes, cardiovascular diseases, obesity, and cancer as reported by Sly et al.⁹

In some situations, environmental pollution is a crystal-clear phenomenon, while in others, the perception lies greatly in the eyes of the beholder. Air emission and water discharge release man-made pollutants into the environment. The total transformations of pollutants in the atmospheric, soil and hydrologic environments play a critical role in their toxicity and contamination pathways that result in health and ecological consequences.¹⁰ The principal exposure routes to these contaminants are mainly through the lungs, dermal and oral. Many environmental toxicants cause both local and systemic effects. These effects may take place following apparent recovery from acute exposure or as a consequence of repetitive exposures to low dose of contaminants over a longer period. Nevertheless, the manifested health effects from acute or chronic exposure are a function of the chemical involved and the organ it affected. Even for the same contaminant, the degree of toxicity is not the same but varies from one organ to the other.

Copious reports have linked various health challenges and several age-associated diseases to environmental contaminants and chemical exposure.^{11–13} The major means by which humans are exposed to inimical chemicals and their combinations are through occupational and environmental sources.^{14–16} There is no simple “rule” to safeguard or intercede against ailments linked to environmental toxicants exposure. Several contaminants, like heavy metals and persistent organics pollutants (POPs), bioaccumulate

in the human system, and remediation techniques in eliminating these pollutants from the surroundings remain a herculean task and are often expensive. Also, treatments aim at obliterating environmental toxicity from the biological compartment are chelation and cleansing procedures, and corroborative techniques, routinely administered in synergy. The treatment may be very complex and personalized, narrowed to an individual’s distinct needs and requiring the adroitness of an expert, often a combination of pundits.¹⁷ Moreover, numerous environmental contaminants induce signaling pathways that lead to oxidative stress which serve as a precursor to the pathogenesis of several long-term maladies. Consequently, techniques that regulate the impact of toxicants on pathophysiologic processes that leads to disease development and escalation will be of immense public health significance.¹⁸ Emerging findings, obtained from fundamental and epidemiologic studies and clinical trials have proven that “phytochemicals” remain the only viable answer to this menace.^{19–22}

Based on the aforementioned discussion, this review presents background information on environmental toxicants and some of their selected toxicity in animals and humans. It further delves into the literature survey of some phytochemicals popularly found in medicinal plants and presents evidence-based studies to accentuate the therapeutic properties of these compounds in combating and ameliorating several health effects due to environmental toxicity. It is our aspiration that this review will trigger public health interest in the beneficial role of phytochemicals and arouses further research that will lead to the discovery of many phytochemicals from the numerous unexplored medicinal plants across the globe.

Methodology

This write-up adopted a systematic review of information primary from available peer-reviewed articles, scientific reports, and grey literatures (reports from international agencies, academia, government documents, and policy statements). The search for literature was conducted utilizing six electronic databases viz Pub med, Web of Science, Google Scholar, Embase, Science Direct, and Cochrane library, for studies published in English from September, 1962 to September, 2022. Studies focused on environmental toxicants and health adversities, pharmacological properties of phytochemicals, and various laboratories, experimental and clinical studies that discussed the intervention roles of phytochemicals. All papers were reviewed using a predesigned data extraction form.

Overview of environmental toxicants

An environmental toxicant can be viewed as any toxic agent or substance produced by humans or introduced into the environment by human activities. Toxicants come in diverse shapes and sizes, and they may emanate from both

Table 1. Categories of contaminants.

S/N	Types	Examples
1	Organic Contaminants	<ul style="list-style-type: none"> ● Petroleum hydrocarbons (fuels)—benzene, toluene, xylene, polycyclic aromatics, MTBE. ● Chlorinated solvent-trichloroethene, tetrachloroethene, trichloroethane, carbon tetrachloride ● Pesticides—DDT, toxaphene, atrazine, Polychlorinated biphenyls—PCBs ● Coal tar—polycyclic aromatics ● Pharmaceuticals/food additives/cosmetics—drugs ● Pharmaceuticals/food additives/cosmetics—drugs, surfactants, dyes
2	Inorganic Contaminants	<ul style="list-style-type: none"> ● Inorganic “salts”—sodium, calcium, nitrate, sulfate, chloride ● Heavy/trace metals—lead, zinc, cadmium, mercury, arsenic, selenium
3	Radioactive Contaminants	<ul style="list-style-type: none"> ● Solid elements—uranium, strontium, cobalt, plutonium ● Gaseous elements—radon

natural and anthropogenic sources. Thus, toxicants include a wide range of compounds from inorganic substances like metals to numerous organic compounds such as pharmaceutical medicines. Environmental contamination in any possible form has been of primary health interest throughout the globe, because it contributes significantly to the pathophysiology of some human diseases.²³ There are several origins by which chemical toxicants are emitted into the surrounding, although these broadly are grouped into a few classes. This section discusses the diverse kinds of contaminants and the classes they belong.

Types of contaminants

Three primary groups of contaminants exist namely organic, inorganic, and radioactive.²⁴ Consecutively, there are numerous classes of contaminants under each of these groups. The main categories of contaminants are outlined in Table 1. Numerous chemicals are emitted into the surroundings daily. Therefore, during the conduction of site appraisal surveys, it is imperative to specify the type of contaminants being assessed.

Classes of toxicants

Heavy metals. Heavy metals have no broad or standard yardstick of definition. On the basis of setting or assumption, numerous connotations may be ascribed to the term.¹⁷ In chemistry, its definition is based on chemical behavior,²⁵ while for the Physicist; the distinct benchmark for heavy metals definition is atomic number,²⁶ whereas in

metallurgy, it is defined by density.²⁷ The name “Heavy metals” refers to elements with a moderately high density which is lethal even at diminutive concentrations. It is a broad concept that is appropriate to a class of metals and metalloids whose atomic density is more than 4 g/cm^3 , or five times or heavier than water.^{25,28} However, heavy metals are basically classified by chemical characteristics other than by specific gravity; and these include copper (Cu), silver (Ag), iron (Fe), lead (Pb), arsenic (As), mercury (Hg), cadmium (Cd), Zinc (Zn), chromium (Cr), and the platinum group elements.²⁹ They are intrinsically distributed in disproportionate doses in the environment. They either exist in elemental form or are combined in chemical compounds with other elements. Heavy metals are often conveyed over a wide range either as volatile or fine aerosols. This group of metals becomes deleterious to the health of humans when their concentration surpasses the maximum acceptable limits in the surrounding.³⁰ Their virulence in ecosystem depends exclusively on their chemical properties and behaviors.¹⁷

Heavy metal pollution in the ambiance occurs via diverse pathways. Owing to their persistence, they permeate the environment for several years after their initial deposition; and contamination of the environment often takes place through weathering fraction.³¹ They bioaccumulate in soil and crops via natural and man-made processes, and subsequent impacts posed critical problems in the food chain via biomagnification.³² The principal port of entry in the human community is through ingestion and dermal.³³ A common way by which adults is exposed is through industrial activities whereas in juveniles is by ingestion.³⁴ Inappropriate therapy, radiological techniques and cracked thermometers are less consequential means of exposure.^{35,36} Some of the health outcomes of their bioaccumulation in the body are the substitution of essential ions, cell membrane impairment, interference with phosphate ions & sulfhydryl (SH) groups, and competition with vital metabolites for binding sites.³⁷

Agricultural chemicals (pesticides). The word “agricultural chemicals” has been widely substituted by the term “pesticides.” Pesticides can be described as a group of chemicals or agglomeration of chemicals that are widely utilized in farming or in quarantine programs to control, annihilate, or fight off infections or nuisance from the plant; and protect humans from disease vectors. Typical classes of pesticides include fumigants, herbicides, fungicides, rodenticides, insecticides, and plant growth regulators.^{38–40} Pesticides are atypical amidst toxicants because they are applied purposely for the annihilation of certain living things. Ideally, pesticides are expected to be selective, eradicating selected species whilst sparing non-target organism. In reality, most pesticides are unspecific in their action. Taking into consideration the application of pesticides, the gains must be juxtaposed with the threat to environmental safety and mankind




Figure 1. Structures of selected organochloride pesticides.

salubrity. Important factors to be considered are persistence in the surroundings and propensity for bioaccumulation. Susceptibility to these chemicals can be by inhalation, ingestion, or dermal contact. The possible health consequences are governed by the variety of pesticides, the time span and means of vulnerability, and the underlying health conditions of the individual. Inside the living system, pesticides may be metabolized, excreted, or bio-accumulated in the body's fat.^{38,39,41} Numerous inimical health outcomes attributed to chemical pesticides exposure include but not limited to gastrointestinal, carcinogenic, dermatological, endocrine, and neurological effects.⁴²⁻⁴⁴ In addition, inadvertently, purposively, or occupational exposure to pesticides can lead to hospitalization and demise.

- Organochlorine Pesticides:** The chlorinated hydrocarbon insecticides came to limelight in the middle of the 20th century, and include common chemicals like dichlorodiphenyltrichloroethane (DDT), chlordane, heptachlor, dieldrin, aldrin, methoxychlor, toxaphene, mirex, endosulfan, dicofol, lindane, endrin, mirex, and toxaphene. The structures of some familiar organochlorine pesticides are shown in Figure 1. The most popularly known organochlorine pesticide is DDT, whose indiscriminate utilization raised numerous environmental and public health concerns.^{38,45,46} It is ubiquitous and generally conceived to be present in all living beings where it is primarily bioaccumulated in fatty tissue. These groups of pesticides are toxic to the nervous system where they disrupt nerve impulses transmission.




Figure 2. Chemical structures of main organophosphate pesticides.

- Organophosphorus Pesticides (OPs):** are among the popularly used pesticides made up of phosphoric acid esters or thiophosphoric acid esters (Figure 2). They include common pesticides, like parathion, dimethoate, and malathion; many of which are famous for their hormone interference potential.^{43,47,48} Among the OPs, the most widely utilized are glyphosate because of its environmental friendliness over the organochlorines.⁴⁹ These groups of pesticides are linked with disruption of cholinesterase enzymes function,⁴⁹ the decline in insulin production, disturbance of regular cellular metabolism of carbohydrates, lipids and proteins,⁵⁰ genotoxic impacts,⁵¹ and impede the activities of mitochondria, instigating

oxidative stress and impairments of the endocrine and nervous system.⁵⁰

- **Carbamate Pesticides:** These are esters of *N*-methyl (or occasionally *N,N*-dimethyl) carbamic acid (H_2NCOOH). The virulence of this chemical differs based on the alcohol or phenol group. A popularly utilized carbamate insecticide is carbaryl (1-naphthylmethylcarbamate), a far-ranging insecticide (Figure 3). It is generally applied as dust in the agriculture, including home gardens. Carbaryl is not regarded as a persistent compound, due to its rapid hydrolysis. Other carbamate pesticides, such as ziram, aldicarb, and carbofuran have been linked with hormone-disrupting property,^{43,52} probable reproductive




Figure 3. Structure of Carbaryl (insecticide carbamate).

defects,^{52,53} and interference with mitochondrial roles and cellular metabolic actions.⁵⁰

- **Other Classes of Chemical Pesticides:** Botanical insecticides have been in application since ancient times to control insects and are extracts obtained from plants. An example is Nicotine [(S)-3-(1-methyl-2-pyrrolidyl)pyridine], an alkaloid found in several plants, whose insecticidal property was first utilized in 1763. Nicotine is partly toxic dermally and orally. Another botanical insecticide is Pyrethrin (Figure 4), which is an extract obtained from numerous kinds of chrysanthemum, and is among the foremost pesticides utilized by man. It is used at low concentrations and is regarded as non-persistent. Synthetic pyrethroids like permethrin, sumithrin, and fenvalerate, are viewed to be among the innocuous insecticides presently use for agricultural and public health purposes.^{54,55} Nevertheless, there is proof of their potential to exhibit hormone-interference activity,^{43,56,57} and to influence reproductive variables in experimental animals.^{56,58}

Triazines, viz simazine, ametryn, and atrazine, are another group of chemical pesticides. Atrazine is popular among the triazines, and it has been basically linked with




Figure 4. Structure of Pyrethrin.

oxidative stress that leads to endocrine-disrupting consequences and reproductive toxicity,⁵⁹ cell toxicity,^{52,60} and dopaminergic effects.⁶¹ Neonicotinoid pesticides, like thiacloprid, guadipyr, and imidacloprid, are quite new and widely utilized insecticides,⁶² that were popular because of the minimal danger they posed to non-target organism.⁶³ However, research conducted in 2016 denoted that neonicotinoids are capable of accelerating the activity of aromatase, an enzyme involved in breast cancer, and also plays a vital part in its carcinogenesis timeline.⁶⁴

Food additives and contaminants. Chemicals are introduced into food for several reasons: as preservatives with antioxidants, antibacterial, or antifungal features; to modify physical properties, especially for processing; to adjust color; and to modify odor. In totality, food additives have been established to be innocuous and lack long term toxicity. Several of them were introduced at a period when toxicity testing was at its rudimentary; albeit, some of these have been afterward proven to be toxic, such as the type of the inorganic, the most important of which are nitrite and nitrate. Definitely, several of the food additives currently in use globally, lack proper testing. The issue of synergism among these compounds is poorly explored. Moreover, not all toxicants in food are artificial; there are some that are natural.⁶⁵

Solvents. These are commonly found in the workplace although they can still be present in residences. In addition to cutaneous consequences, such as local irritation and defatting; several have systemic toxic impacts, such as impacts on the nerves or, as with benzene, on hematopoietic elements. The well-known solvents fall into the below classes⁶⁶:

- **Glycols and Glycol Ether:** Ethylene and propylene glycols, for instance, in antifreeze give rise to significant exposure in the general population.
- **Aliphatic Hydrocarbons:** like hexane. These may be straight or branched-chain compounds and often occur in mixtures.
- **Aromatic Hydrocarbon:** Benzene is probably of greatest interest in this group; however others, like toluene are also of public health interest.
- **Halogenated Aliphatic Hydrocarbons:** The most popular examples are chloroform, carbon tetrachloride, and methylene dichloride; nevertheless chlorinated ethylenes are also given wider attention.
- **Aliphatic Alcohols:** Popular examples are methanol and ethanol.

Drugs of abuse. Drugs of abuse (Figure 5) can be defined as drugs that either have no medicinal importance or are used at dose concentrations that exceed the dosage needed for treatment. Albeit many of these substances alter only




Figure 5. Chemical structures of the investigated drugs of abuse.

the higher nervous function like reaction time, coordination, and mood; some cause physical addiction and have consequential physical impacts, with fatal overdoses being a periodic phenomenon. This class of drug includes central nervous system (CNS) depressants, viz secobarbital, ethanol, methaqualone (Quaalude); CNS stimulants, like caffeine, nicotine, cocaine, and methamphetamine (speed); opioids, viz meperidine (demerol) and heroin; and hallucinogens like phencyclidine (PCP), lysergic acid diethylamide (LSD), and tetrahydrocannabinol the most active component of marijuana.⁶⁷

Cosmetics. The well-known harmful upshots of modern cosmetics are sporadic allergic reactions and contact dermatitis. Although, the organometallics and aromatic amine dyes or azo employed in ancient times are now obsolete. Bromates and Thioglycerol applied in some cold-wave neutralizers and lotion may be markedly injurious if ingested; ditto ethanol applied as a solvent in perfumes and hair dyes. Applied as instructed, cosmetics have seen to pose a limited risk of systemic poisoning, owing partly to the elimination of hazardous constituents in them and partly to the minute amount absorbed.²⁴

Combustion products. Most air pollutants are from natural and man-made combustion. In terms of human health impacts, the most prominent are the Polycyclic Aromatic Hydrocarbons (PAHs). Albeit they are present in natural products like crude oil and coal, and commonly linked to incomplete oxidation of organic substances. Smoke from coal, wood, tobacco, and oil, for instance, in broiled foods

also emits these contaminants. Due to the carcinogenicity of some of these substances, they have been studied diligently from the position of interactions with DNA, metabolic activation, and other facets of chemical carcinogenesis.²⁴

Toxicity of some selected environmental toxicants

Toxic effects of mercury

Mercury is a popular poisonous metal and its lethal is a frequent cause of acute heavy metal poisoning as reported by the American Association of Poison Control Centers with cases of 3596 in 1997.⁶⁸ Methylmercury is a neurotoxic compound that is liable for microtubule ruination, lipid peroxidation, mitochondrial impairment, and accretion of neurotoxic like glutamate, serotonin, and aspartate.⁶⁹ It is postulated that 8% to 10% of American females possessed inherent mercury levels that may likely trigger neurological abnormalities in the infants they birthed, as enunciated by both the National Academy of Science and Environmental Protection Agency.⁷⁰ Animal studies have affirmed that exposure to toxic mercury caused inimical neurological and behavioral modifications in animals. The primary target organ for mercury is the brain; although it can impair other organs and result in a defect in the muscles, nerves, and kidneys. It leads to derangement of the membrane potential and disrupts intracellular calcium homeostasis. Mercury attaches to free unconjugated thiols as the affinity constants are high.⁶⁹ Mercury vapors can result in asthma, bronchitis, and short-term respiratory difficulties. Mercury contributes significantly to the damaging of tertiary and quaternary protein structure and modifies the cellular properties by binding to the selenohydryl and sulfhydryl groups that react with methyl mercury and disrupt the cellular structure. It also interferes with the transcription and translation process leading to the obliteration of ribosomes and destruction of endoplasmic reticulum and natural killer cells properties. In addition, compromise the cell integrity that results in free radical production.⁶⁸

Toxic effects of lead

Lead metal brings about injury to the cellular component sequel by ionic mechanism and that of oxidative stress. Numerous studies have demonstrated that oxidative stress in biological systems is instigated by the disequilibrium between the free radicals generation and antioxidants production to neutralize the transitional or mend the outcome impairment.^{17,68} The potential of lead metal ions to substitute other bivalent cations like Mg^{2+} , Ca^{2+} , Fe^{2+} and monovalent cations such as Na^+ is primarily responsible for its ionic toxicity effects and resultant impairment of the

metabolic activities in a biological system. Lead toxicity arising from ionic mechanisms is responsible for adverse alterations in diverse biological activities including protein folding, cell binding, apoptosis, intra- and inter-cellular signaling, maturation, protein folding, enzyme regulation, active transport, and neurotransmitter production. Lead has the capacity to replace calcium even at picomolar concentration altering protein kinase C, which modulates memory storage and neural excitation.⁷¹

Toxic effects of arsenic

The toxic impacts of Arsenic can either be short or long-term. Chronic Arsenic Toxicity (CAT) is termed arsenicosis. Numerous reports of CAT in humans center on dermal externalizations due to its particularity in diagnosis. Keratosis and pigmentation are the unique skin aberrations that point to CAT.⁷² Lower exposure levels to arsenic can trigger gastrointestinal disorders, decreased blood cell production, irregular heartbeat, vascular vessel impairment, and prickling sensation at the extremities. Chronic exposure can result in the development of internal cancer, skin lesions, respiratory disease, neurological malignancies, hypertension, vascular disease, diabetes mellitus, and cardiovascular disease.⁷³ Long-term arsenicosis leads to permanent modifications of the essential organs with an attendant high death rate. Despite the enormity of this imaginably deleterious toxicity, there is the absence of an effective remedy for this malady.⁷⁴

Toxicity of persistent organic pollutants (POPs)

The outcomes of acute and chronic exposure to POPs are still ambiguous. However, laboratory probing as well as environmental impact studies in the natural surrounding have stipulated that POPs can lead to immune and reproductive disorders, hormonal disruption, central nervous system (CNS) dysfunction, cancer and developmental anomalies. A few organochlorine chemicals are possibly carcinogenic by aiding tumor development. Polychlorinated Biphenyls (PCBs) [are a group of over 200 anthropogenic and artificial chlorinated and organic chemicals with the same basic chemical structure] are categorized as *probably carcinogenic* to man, while an additional eight of the 12 other POPs pinpoint in the Stockholm Convention are ranked as *possibly carcinogenic* to mankind. The leftover three—endrin, aldrin and dieldrin are grouped by the World Health Organisation as very deleterious (class 1b) owing to their short-term toxicity to laboratory animals. Fetuses and newborns are especially susceptible to POPs exposure because of the transmission of POPs from the mother during the crucial period of gestation. Susceptibility in the course of development has been associated with decreased immunity, CNS damage, developmental anomalies, and cancer/tumor induction or promotion in neonates

Table 2. Classification of POPs according to International Agency for Research on Cancer.

IARC Classification	POPs
Group 1: The agent (mixture) is carcinogenic to humans.	2,3,7,8-Tetrachlorodibenzo-para-dioxin (TCDD)
Group 2A: The agent (mixture) is probably carcinogenic to humans.	Mixtures of polychlorinated biphenyls (PCB)
Group 2B: The agent (mixture) is possibly carcinogenic to humans.	Chlordane DDT Heptachlor Hexachlorobenzene Mirex Toxaphene (mixtures of Polychlorinated camphenes)
Group 3: The agent (mixture or exposure circumstance) is unclassifiable as to carcinogenicity in humans.	Aldrin Dieldrin Endrin Polychlorinated dibenzo-para-dioxins (other than TCDD) Polychlorinated dibenzofuran

Source: http://www.chem.unep.ch/gpa_trial/02health.htm

and juveniles. The likelihood of causing human breast cancer has also been reported.⁷⁵

Table 2 depicted the list of 12 POPs that are humans' carcinogenic agents as highlighted by the International Agency for Research on Cancer during the Stockholm Convention.

Toxicity of polycyclic aromatic hydrocarbons (PAHs)

Each of the PAH compounds exhibits unique health outcomes.^{76,77} Several PAHs are teratogenic, carcinogenic, mutagenic, and immunotoxic to innate beings, such as microbes, animals, and humans.^{76,78,79} The manner of exposure, period, and dosage are crucial variables for determining the virulence of PAHs' toxic outcomes.⁸⁰ Fatal outcomes of PAHs may differ based on variables such as age and underlying health state. Short-term health impacts include regurgitation, eye soreness, disarray, skin irritation, diarrhea, and inflammation.⁷⁸ Anthracene, benzo(a)pyrene, and Naphthalene are explicit skin irritants and skin sensitizers for animals and humans.⁷⁶ Long-term impacts are renal and hepatic derangement, eye cataracts, respiratory distress, pulmonary dysfunction, asthma-like symptoms and compromised immunity.⁷⁷ Naphthalene can bring about the degradation of erythrocytes if inhaled or ingested in large quantities.⁷⁶

Overview of phytochemicals

What are phytochemicals?

The term "phytochemicals" refers to a perplexing number of small molecules from plants, which can be categorized into diverse marked classes based on their biosynthetic origin and structure. On a wider definition, it means plant (Phyto) chemical denoting a large diversity of compounds that are found in plants naturally. A great many bioactive

phytochemicals occur in popularly consumed plant foods and products. Broadly, these compounds can be classified into six main categories based on chemical structures and properties. These groups are carbohydrate, lipids, terpenoids, phenolics, alkaloids, and other nitrogen-containing compounds.⁸¹ Each group can be subdivided based on bio-synthetic origin or biogenesis into different subgroups. Contemporarily, the word "phytochemical" has been employed to differentiate plant chemicals that do not satisfy the traditional meaning of "essential nutrients." Many of these phytochemicals trigger reactions in living organisms, including humans, by expressing both therapeutic and prophylactic features that have been associated with the provision of nutrient for normal cell growth and repair, impede carcinogens and acts as antioxidants, and declines the risk of major non-communicable chronic diseases.^{82,83}

Phytochemicals wield their medicinal prowess by acting synergistically or additively and this nullifies the precarious outcomes linked to the high usage of a mono xenobiotic compound, thus giving the herbal drug(s) a broad spectrum of activity, in addition to diminishing the probability of the pathogens developing resistance or adaptive responses.^{84,85} Some phytochemicals are enunciated in the subsequent sub-section.

Classes of phytochemicals

Flavonoids: are structural derivatives of flavones (see Figure 6). Consisting of conjugated aromatic systems, often bind to sugar(s) as glycosides, and they are phenolic and hydrophilic in nature.⁸⁶ Flavonoids are a vital class of natural products; they are in the group of plant secondary metabolites possessing a polyphenolic structure, and are generally found in vegetables, fruits, and some beverages.⁸⁷ Flavonoids are linked with a wide range of health-enhancing properties and are invaluable constituents in diverse pharmaceutical, nutraceutical, cosmetic and




Figure 6. General structure of flavonoids.

medicinal applications. This is due to their anti-inflammatory, antioxidative, anti-carcinogenic and anti-mutagenic characteristics coupled with their ability to regulate vital cellular enzyme functions. Likewise, they are regarded as powerful inhibitors for numerous enzymes, viz cyclo-oxygenase (COX), xanthine oxidase (XO), phosphoinositide 3-kinase, and lipoxygenase.⁸⁸ Flavonoids have numerous subgroups, such as flavones, isoflavones, flavanones, chalcones, flavanonols, flavonols or catechins, chalcones, and anthocyanins. These subgroups have distinct major sources.⁸⁹ Flavonoids lacking hydroxyl (-OH) on their structure are more potent against microbes vis-à-vis those with -OH, and this corroborates the fact that their target in microbes is the membrane.⁸³

Tannins: these are caustic, pungent phenolic compounds that either coarsen or precipitate or contract proteins. They consist of numerous groups of oligomers and polymers. They can adhere to cellulose, minerals, proteins and starch to form complexes. Tannins are hydrophilic compounds with the exclusion of some higher molecular weight structures. Tannins are generally categorized into two groups namely: hydrolyzable tannins such as ellagitannin, complex tannins, proanthocyanidins, and gallo-tannins; and condensed tannins (Figure 7). These compounds are usually found in both gymnosperm and angiosperm. They have been discovered in 180 families of dicotyledons and 44 families of monocotyledons. Tannins are commonly found in the following families of plant species: Actinidiaceae, Burseraceae, Bixaceae, Anacardiaceae, Dipterocarpaceae, Combretaceae, Grossulariaceae, Ericaceae, Myricaceae for dicotyledons; whereas Typhaceae and Najadaceae for monocotyledons. They exhibit their physiological function by behaving as

antioxidants via free radical scavenging action.^{83,90} thus regulating oxidative stress and averting deteriorating illness. They also hinder tumor growth by activating cell death⁹¹ and impeding the mutagenicity of carcinogens.^{83,92}

Saponins: refer to a group of bioorganic compounds that are found largely amidst plant biodiversity. More precisely, they are naturally occurring glycosides with soapy properties, and as a result, they generate bubbles when agitated in aqueous solutions.⁹³ Structurally saponins possess one or more hydrophilic glycoside sugar moieties (Figure 8) aggregated with a lipophilic triterpene molecule.⁹⁴ Studies depict that saponins exercise biological function and therapeutic effects like hemolytic factor, anti-inflammatory, antifungal, antibacterial, antiviral, anticancer, cytotoxic, molluscicidal and insecticidal action.⁹⁵⁻⁹⁷ The saponin molecule undergoes hydrolysis to give rise to two fragments, an aglycone and a sugar moiety. Isolated amorphous solid saponins have large molecular weight, and contain 27–30 carbon atoms in the non-saccharide fraction (the hydrocarbon skeleton part lacking a sugar chain). This non-saccharide segment referred to as sapogenin, aglycone, or genin. Based on the kind of sapogenin available, the saponins can be segregated into three distinct groups: triterpenoid glycosides, steroid glycosides, and alkaloid glycosides.⁹⁸ The second partition that is, the saccharide moiety—consists of heterogeneity of pentoses or hexoses.⁹³

Alkaloids: These are a large class of naturally existing organic compounds with nitrogen atom or atoms (amino or amido in certain instances) in their structures often located in some cyclic or ring configuration (Figure 9). The presence of nitrogen in these compounds is responsible for their alkalinity. Broadly, based on structure, alkaloids can be classified into groups like quinolones, pyrrolidines, isoquinolines, indoles, tropanes, steroids, pyridines, pyrrolizidines, and trepenoids. Another categorization modality is based on the family of plant species in which these substances are found. A typical example is the opium alkaloids that are found in the opium poppy (*Papaver somniferum*).^{99,100} These two dissimilar classification systems create a disparity between their biological distribution and the chemical varieties of alkaloids because there is no distinctive connection. In its pure form, alkaloids are often odorless, crystalline solids, colorless, although often yellowish and have an astringent taste at times. Presently, over 3000 alkaloids are identified in more than 4000 diverse plant species. Alkaloids are a fascinating class of compounds with a divergent range of functionalities in man and animals. Alkaloids have various physiological outcomes: antimutagenic, local anesthetic, antibacterial, psychotropic, analgesic, hypnotic, antitumor, anti-inflammatory and several others. Popularly known alkaloids are caffeine, nicotine, ephedrine, strychnine, quinine, atropine, and morphine.^{99,100}




Figure 7. Main chemical structures of the tannins.




Figure 8. Generic structures of saponins R=sugar moiety.

Cardenoloids: These are also known as cardiac glycosides and they consist of the most drug-like molecules subjected to numerous examinations and were affirmed to be effective in manufacturing efficacious drugs.^{101,102} These groups are liable for being virulent to livestock and serve as therapy for congestive heart failure. Cardenoloids are steroids capable of exerting a precise forceful effect on the myocardia. A minute quantity can affect a salubrious simulation on an ailing heart. These compounds are beneficial in the treatment of congestive heart failure, flutter, atrial fibrillation, and they behave as emetics and as diuretics.⁸³ They speed the force of contraction in the heart, which is incapable of meeting the rise in oxygen expenditure. As a result, the myocardium acts as a potent pump and is capable of satisfying the need of the vascular system.^{102,103} Cardiac glycosides consist of two distinct categories of compounds that vary in their aglycone structure (Figure 10). They may be C23 or C24 steroids with a primary nucleus of *cyclopentanoperhydro phenanthrene* substituted at C17. Cardenolides possessed a five-membered lactone group in the C17 with α , β -unsaturated γ -lactone ring (butenolide), while the other class, the bufadienolides, was first identified in toad as skin lethal. The C17 substituent has a doubly




Figure 9. General structure of Alkaloids.

unsaturated six-membered lactone ring (α -pyrone). Plants are capable of producing both cardenolides and bufadienolides. A different group is isocardenolides, which possessed a double bond of butenolide ring at position 21 or 22 rather than at position 20. Most clinical interests were pointed to the cardenolides due to their medicinal importance. The two commonly utilized *digitalis inotropes* are Digoxin and digitoxin. Generally, most isocardenolides lack any cardiac activity.¹⁰²

Terpenoids: otherwise called isoprenoids, are a wide and divergent class of naturally existing compounds whose precursors are five carbon isoprene units (Figure 11). Terpenoids distinguished from one another by their basic skeleton and functional groups. Terpenoids are ubiquitous; hence, they occur virtually in all living beings. They add to the scents, color, and flavor of plant's flowers, leaves and fruits. Terpenoids are also crucial for plant growth and development. The terpenoids generated by plants not only safeguard them from insects and predators but also offer




Figure 10. Structure of Cardiac glycosides.




Figure 11. Structure of Terpenoids.

defense from fungal diseases and infections, while in animals, terpenoids serve as precursors of steroids and sterols. Terpenes are simple hydrocarbons, whereas terpenoids are an adjusted class of terpenes with varying functional groups and oxidized methyl groups oscillating at different positions. Terpenoids are categorized into monoterpenes, sesquiterpenes, diterpenes, sesterpenes and triterpenes based on their carbon units. Terpenoids differ in structures and are biologically viable and utilized globally for the therapy of numerous maladies. Several terpenoids impeded diverse human cancer cells and are employed as anticancer medicines such as Taxol and its derivatives. Terpenes and its derivatives are utilized as antimalarial drugs such as artemisinin and associated compounds. They performed various functions viz anti-fungal, anti-protozoan, anti-bacterial, anti-allergens, anti-viral, immune boosters and antineoplastic.^{83,104,105}

Phytosteroids: these are groups of specialized metabolites obtained from plants that are attached to steroid receptor in animals and can activate or suppress downstream receptor-mediated signaling events. Phytosteroids (Figure 12) have varying structures, occasionally very dissimilar from the animal steroids; however, they behave as antagonists, agonists, or habitually exhibit mixed agonist/antagonist properties for steroid receptors.^{106,107} Furthermore, some phytosteroids interrelate with several steroid receptors¹⁰⁸ or hamper steroid metabolizing enzymes,¹⁰⁹ hence exhibiting complicated impacts on the reproductive and hormonal systems. They possess a similar basic cyclic configuration to animal steroids, nevertheless they differ due to different chemical groups attached to the central ring in various positions.⁸³ They are basically




Figure 12. Structure of Phytosterol.

used in the treatment of reproductive ailments viz treatment of sexually transmitted diseases, enhance fertility in women and sex drive in males, and ensure safe parturition. They also behave as sex hormones derivatives and thus they are a possible sources of contraceptives.^{83,98,110} They are also analgesic, anti-microbial, anti-inflammatory, and of use in curing of stomach complications and in reducing serum cholesterol levels.^{83,98} Phytosteroids are marked as powerful inhibitors of macrophage activation, impeding the generation of pro-inflammatory cytokines and LPS-induced fatality and consequently they can act as immunosuppressive agents particularly the phytosterols.^{83,111}

Interventional roles of phytochemicals on various toxicity outcomes

Intervention in oxidative stress, metabolic syndrome and aging

Since time immemorial, it has been known that natural products prolonged the life-span of an organism.¹¹² Epidemiological and laboratory studies suggested these compounds to be potent antioxidants that ameliorate stress-related ailments. Various studies have postulated the efficacy of these compounds in metabolic symptoms and aging.^{113,114} Consequently, comprehensive preclinical assessment on the fundamental pharmacology of these natural compounds may serve as strong scientific basis for clinical implementations. Polyphenols and most especially flavonoids have been proven to offer protection from several age-associated morbidities.¹¹⁵ Numerous studies have shown that augmentation with dietary polyphenols like (-)-epigallocatechin-3-gallate (EGCG) and curcumin can enhance age-related cellular impairment by decreasing production of reactive oxygen species (ROS).¹¹⁶ Conversely, resveratrol and pterostilbene are regarded as outstanding anti-aging chemicals that can regulate oxidative destruction, cell aging, and inflammation; constituents linked with aging¹¹⁷ including flavonoids which is proven to aid anti-aging primarily by regulating metabolic syndrome.¹¹⁸

A few of the proven flavonoids that can handle factors linked to senescence or metabolic syndrome are naringin, hesperidin, and naringenin.¹¹⁹ Importantly, there is an increase in the data bank of preclinical application of phytochemicals in the therapy of diverse conditions attributed to senescence. Nevertheless, certain limitations exist to their applications such as (1) option of experimental models that is clinical pertinent, (2) nebulous elucidated mechanism of action, and (3) appropriate dosage and duration for data interpretation.

As anti-cancer agent

In spite of the fact that significant progress has been made, some tumor still pose poor prognosis and study is still ongoing toward the utilization of innocuous doses of plant-extracted substances. The innovation to treatment strategy due to natural molecules and drugs was instigated by the recognition and application of natural chemotherapeutic substances vis taxanes, anthracyclines, and vinca alkaloids.¹²⁰ Thus, it is reasonable to postulate that compound in foods are presumably to have some protective impacts.

Intriguingly, contemporary research findings suggest an increasing function of polyamines metabolism as an innovative approach against inflammatory ailments. Polyamines are naturally occurring aliphatic compounds, ever-present in all living thing, which interface with nucleotides and proteins and are needed for the growth, viability and specialization of eukaryotic cell.^{114,115,121,122} Based on the aforementioned roles, polyamines metabolism constitutes an intriguing target for anticancer treatment.¹²³ Hence, the application of various natural substances, especially polyphenols, emanating from plants constituents may produce auspicious outcomes in antitumor treatment due to their anti-oxidant properties.¹²⁴ Interestingly, it has been shown that high flavones intake, particularly quercetin and kaempferol, is capable of inducing a significant decrease in serum IL-6 concentration, a popular inflammation-associated cytokine.¹²⁵ Likewise curcumin depicts anti-inflammatory and antioxidant characteristics, and prospective anti-cancer property.¹²⁶ Genistein a soybean isoflavone, has exhibited antitumor property in diverse types of cancer including chronic lymphatic leukemia and neuroblastoma and in diverse organs like ovary, prostate, breast, colon, urinary bladder, stomach, and stomach.¹²⁷ Genistein is regarded as a phytoestrogen because of its structural similarity with mammalian estradiol. Numerous studies have shown the consequential function of isoflavone in the prevention of cancer development in animal models. Indeed, it is proven that dietary soy supplementation mitigate inflammation associated with prostate carcinogenesis.¹²⁸ Another popular anticancer agents possessing both anti-inflammatory and antioxidant feature is lycopene that exhibits treatment actions on numerous tumors.¹²⁹ Specifically, it has been established that lycopene ingestion hinders the development

of cancer.¹³⁰ The recognition of the importance of these natural compounds to the advancement of cancer treatments are well documented nonetheless it is crucial to conduct more meticulous researches and preclinical studies to shed more light on their prospective chemopreventive and antitumor activities.

As therapy for vascular diseases

In this review we shall discuss only the therapeutic impacts of omega 3 polyunsaturated fatty acids (PUFA) and the flavonoid resveratrol due to their prevailing effects on Senescence-Associated Secretory Phenotype (SASP), vascular dysfunction, and cardiovascular disease (CVD).

The effects of omega 3 PUFAs are ascribed to their lower lipid effects which help to decline atherosclerosis development.¹³¹ Omega 3 PUFAs have been demonstrated to mitigate vascular inflammation by down-regulating adhesion molecules and restricting leukocytes binding to the wall of blood vessel.¹³² This latter explicitly sway endothelial-derived nitric oxide production owing to stabilization of lipid rafts like the endothelial cells caveolae, as shown in retinal endothelial cells.¹³³ Nevertheless, the laboratory studies seem to corroborate PUFA's beneficial roles vis-à-vis the clinical proof. Actually, the omega 3 PUFA outcomes on aggrandized endothelial regenerative ability and preservation of vascular endothelial cells homeostasis by virtue of membrane stabilizing potential were marked to have consequential effects on CVD prevention.¹³³ A well-detailed appraisal of the literature recently displayed on Cochrane Database Systematic Review,¹³⁴ encapsulated the findings of numerous randomized clinical trials investigating the impacts of varying doses of PUFA on CVD manifestations. Findings from this survey depicted that higher PUFA intake only moderately lessen risk of coronary heart disease and CVD acute events (i.e. stroke), but in generality has no important influences on its outcomes. A large number of the positive outcomes were linked to modulation of lipid metabolism.⁹¹ Anyway, even a little but notable decline in 10% of morbidity and death for CVD linked to PUFA supplementation remains a momentous clinical outcome.¹³⁵ Conclusively, other phytochemicals with antioxidant properties have been highlighted to contribute a protective action to CVD development and consequently were recommended as significant substances in diet, for example like, β -carotene, curcuma, and others.^{136,137} Specifically, numerous studies underscore the anti-atherogenic role of lycopene in relation with the interdiction of proinflammatory cytokines release.¹³⁸

Role as anti-inflammatory

It has been demonstrated that natural compounds influence numerous pro-inflammatory mediators. Herbal medicines, nutraceuticals, or beneficiary foods with anti-inflammatory

properties, can be utilized as an adjunct to anti-inflammatory drugs resulting in the decline of their consumption, leading to the mitigation of their side effects. A comprehensive review on the anti-inflammatory properties of phytochemicals has been done by several researchers.¹³⁹ Various anti-inflammatory mechanisms have been linked to numerous flavonoids with diverse chemical configurations.¹⁴⁰ Glycosides of apigenin and luteolin are widely distributed flavones.¹⁴¹ Apigenin subdue nitric oxide (NO) and prostaglandin production through hampering of inducible nitric oxide synthase (iNOS) and COX-2, appropriately.¹⁴² Luteolin was also demonstrated to impede chronic inflammation by in vitro co-culture of adipocytes and macrophages and the phosphorylation of Jun N-terminal kinases (JNK) in macrophage.¹⁴³ The most efficacious tumor necrosis factor- α (TNF- α) inhibitors among the flavonoids are luteolin and quercetin.¹⁴⁴ The anti-inflammatory potency of quercetin was clinically assessed in women with rheumatoid arthritis showing an important impact in modulating inflammation and clinical manifestations.¹⁴⁵ In spite of their function in the production of TNF- α , green tea extract has also been shown in contemporary clinical trials to have anti-inflammatory and immunomodulatory properties in autoimmune disease.¹⁴⁶ Consequently, although these compounds cannot replace anti-inflammatory drugs, such as the Disease Modifying Anti-Rheumatic Drugs (DMARDs), but they significantly aid to the decline in their dosage, leading to less costly and innocuous treatment strategy of autoimmune diseases and other inflammation-associated morbidities.

Neurodegenerative diseases prevention

Presently, there is no panacea treatment for neurodegenerative diseases (NDD), and, in a bid to discover new treatment or adjuvant approach for NDDs, numerous natural medicinal plants have attracted interest as prospective neuroprotective agents, and several researches have alluded to the fact that a diet enrich with vegetable products can avert or prolong the initiation of NDD.¹⁴⁷ These attributes may be associated to the presence of polyphenols, a crucial group of phytochemicals that are readily found in vegetables, beverages, fruits, and cereals. In this section we shall discussed the auspicious role of certain phytochemicals in the prevention and treatment of NDD, as illustrated by selected evidence-based studies.

Numerous studies have established that resveratrol possessed important neuroprotective feature both in vitro and in vivo.^{148,149} In vivo, in a mouse model with cerebral amyloid accumulation, orally administered resveratrol lessen microglial activation related to cortical amyloid plaque development.¹⁵⁰ Moreover, chronic dietary resveratrol decrease cognitive damage and perform a neuroprotective role, reducing the amyloid load and decreasing tau hyperphosphorylation in SAMP8 mice, a model of age-associated

Alzheimer's disease (AD).¹⁵¹ Rising evidence has also submitted that resveratrol confers increased advantages to cell and animal models with Parkinson's disease (PD).¹⁵² Curcuminoids comprise three components: curcumin (75%–80%), demethoxycurcumin (15%–20%), and bisdemethoxycurcumin (3%–5%). Curcumin also trigger neuroprotective outcomes via the regulation of pathogenetic oxidative and inflammatory actions both in vitro and in vivo models of AD and PD. In Neuro2a mouse neuroblastoma cells infected with Japanese encephalitis virus, curcumin promotes cell activity by reducing reactive oxygen species (ROS) and hindering proapoptotic signals.¹⁵³ Pretreatment of primary hippocampal cultures with quercetin substantially diminished A β (1-42)-induced cellular toxicity, lipid peroxidation, protein oxidation, and cell death by regulating oxidative stress.¹⁵⁴ More captivatingly, quercetin diminishes extracellular β -amyloidosis, astrogliosis, microgliosis, and tauopathy in the hippocampus and the amygdala and enhances execution of learning and spatial memory burden in old triple AD model mice.¹⁵⁵ Aggregating, the above findings, point to polyphenols as neuroprotective agents. The addicted consumption of dietary polyphenols is validated to impede diverse secondary sources of ROS and proinflammatory cytokines, hence mitigating of NDDs.¹⁵⁶ An advantageous clinical utilization of polyphenols to lessen oxidative impairment in senescence and age-associated ailments may be a potent and auspicious strategy for NDDs therapy.

As powerful antioxidant agents

The first antioxidant molecule identified is ascorbic acid, also known as vitamin C, which is generated in the course of aerobic metabolism, and react rapidly with O₂^{*}, singlet oxygen and ozone (chemically), and H₂O₂ (enzymatically) via ascorbate peroxidase to detoxify their deleterious effects. Aside this, in plants, such acids likewise takes part in the regeneration of carotenoids and vitamin E (tocopherol). The latter can also behave as antioxidant and vital liposoluble redox system, serving as protection against lipid peroxidation.¹⁵⁷ Carotenoids, which are potent antioxidants, are responsible for several coloring in fruits, flowers and leaves; and they are involved in the scavenging of peroxyl radicals through quenching.^{158,159}

Phenolic compounds are one of the most auspicious molecules for advancing salubrity studies. These phytochemicals comprise of a wide diversity of molecules (circa 8000 numerous structures), playing essential functions in the metabolism of the plant,¹⁶⁰ where they are largely distributed. The far-ranging of organic activities of phenolics, one of which is antioxidant and antitumor characteristics, is widely established in different studies.^{157,161} The existence of at least a single phenol ring is vital for such role, with methyl, hydroxyl, or acetyl groups substituting the hydrogen. An elevated antioxidant function has been linked to

the increased number of free hydroxyls and conjugation of side chains to aromatic rings.¹⁵⁸ Terpenoids constitute additional huge family of plant secondary metabolites.¹⁵⁸ In vitro assays depicted that sesquiterpenes, monoterpenes, and diterpenes derived from aromatic plants have significant antioxidant activity.¹⁵⁹ The hypoglycemic and antioxidant property of alkaloid—vindoline, vindolicine, vindolinine, and vindolidine, gotten from *Catharanthus roseus* leaves, have also been highlighted.¹⁶⁰

Therapeutic effects on skin pathologies

According to the European Medicines Agency, about 12 herbal substances, preparations, and mixtures are utilized as traditional herbal medicinal products.¹⁶¹ These 12 herbal substances include—*Foeniculum vulgare*, *Sideritis scardica*, *Valeriana officinalis*, *Echinacea purpurea*, *Hamamelis virginiana*, *Tymus vulgaris*, *Vitis vinifera*, *Eleutherococcus senticosus*, *Melaleuca spp.*, *Calendula officinalis*, *Mentha spp.*, and *Pimpinella anisum*. Their inclusion inside such official list is maintained by scientific evidences denoting their treatment efficacy in several diseases conditions vis basal cell carcinoma, carbuncle, allergy, cellulitis, acne, eczema, impetigo, chickenpox, hives, dermatitis, melanoma, rosacea, vitiligo, wart, psoriasis, impetigo, lupus, measles, blister and squamous cell carcinoma.^{122,157,162–165}

A meta-analysis published show *Vitis vinifera* as an efficacious constituent of medical accessory beneficial in the therapy of local atopic dermatitis.¹⁶⁶ Intriguing outcomes have also been reported in oncological circumstances; *Vitis vinifera* has demonstrate some effectiveness in decreasing radiotherapy-induced dermatitis¹⁶⁷ and in impeding cell multiplication in melanoma and skin non-melanoma cancer,¹⁶⁸ denoting grape seed proanthocyanidin as an apoptosis and autophagy inducer. Limited allergic reactions are enunciated for *Vitis vinifera*.¹⁶⁹ In comprehensive elucidation, local utilization of *Melaleuca alternifolia* derieved oils has been persistently aired to attain a notable amelioration of acne lesions, as reported by numerous autonomous studies.^{162,170} The clinical efficiency of Melaleuca oil is possibly associated to its familiar antibacterial mechanism^{163,171} in addition to anti-inflammatory action.^{164,172} The essential oil from *Melaleuca alternifolia* also depicts antioxidant properties probably efficacious in dermatitis and skin cancers.^{165,173} Therapeutic benefits to skin diseases are suggested for *Agrimoniae herba* (*Agrimonia eupatoria*; in secondary inflammation and superficial lacerations), *Echinacea purpurea* (in modest perfunctory wounds and mild acne), *Soiae oleum* (Glycine max; in benign periodic eczema), *Juglandis folium* (*Juglans regia*; in minor skin irritation), *Matricariae aetheroleum* (in anus and genitals inflammation), *Matricariae fos* (*Matricaria recruta*; in mild skin allergies and sunburns and superficial wounds), *Melaleuca spp* (in insects

bites, mild acne, prickling, lesser skin irritation), *Meliloti herba* (*Melilotus officinalis*; in minor skin inflammation), *Origani dictamnii herba* and *Origani majoranae herba* (*Origanum spp.*; in petty skin irritation and inflammation). Several others phytochemicals are reported in literature with prospective effects on skin, such as anti-senescence property,¹⁷⁴ photoprotection,¹⁷⁵ wound recuperation,¹⁷⁶ and anti-infection.¹⁷⁷

Conclusions

The review critically highlights the various environmental pollutants and their toxicity on living organisms. It also delves into several classes of phytochemicals and their prophylactic and therapeutic interventions in various maladies challenging humanity. This review enunciates that environmental contaminants are generated from both natural and anthropogenic activities, and they become deleterious to the environment and pose a risk to living things when they bioaccumulate above their threshold limits. They are of significant public health interest across the globe, due to their indispensable role in the pathophysiology of several human diseases. Several remediation techniques and treatment measures employed toward mitigating the outcomes effects of environmental toxicity to humans are characterized by one or many limitations; which make the exploitation of phytochemicals a global interest. Phytochemicals are secondary metabolites found in plants and are popular for their numerous pharmacological and medicinal properties. Hence, they exhibit anti-inflammatory, anti-metastatic, anti-oxidant, immune-modulative, nutritive, prophylactic, anti-microbial and life-saving properties. Current researches in literature focus on expanding the database of the efficacy of phytochemicals in reducing or treating various environmental-induced toxicities. However, mechanistic understanding of their pharmacological action needs to be elucidated. Therefore, future research should focus on identifying the bioactive molecule and testing the effectiveness of phytochemicals against mixer toxicity. In addition, there is also need for extensive studies on the optimization and large-scale production of medicinal formulation from these phytochemicals.

Author contributions

AASA conceived the concept. AASA, MASA, and EDSA were involved data sourcing and collation. AASA, MASA, and EDSA wrote the initial draft and revised the manuscript. ANO, HE, EOT, FAB, MA and OEA critically revised the manuscript. All authors read and approved the final version of the manuscript.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Significance for public health

This review article is of high significant impact to public health because it creates public awareness on the pharmacological and medicinal importance of phytochemicals from empirical and evidence-based studies against several maladies associated with environmental toxicity. Environmental toxicants are of public health interest as they contribute to high rate of morbidity and mortality globally. Efforts made to ameliorate their impacts on the environment and humans through remediation techniques, chelation and other cleansing procedures are often limited by complicated techniques. Phytochemicals which are secondary metabolites obtained from plants have wide pharmacological and medicinal properties; which can serve as a panacea to this nagging problem. Since Environmental health is a key part of any comprehensive public health system. This review article helps to enlighten people on environmental exposomes and how they can use natural products from plants to safeguard their health, thus promoting healthier environment and communities.

ORCID iD

Muhammad Aledeh  <https://orcid.org/0000-0002-6831-9005>

References

- World Health Organization. 7 million deaths annual linked to air pollution. Geneva: WHO, http://www.who.int/phe/health_topics/outdoorair/databases/en/ (2022, accessed 20 September 2022).
- World Health Organization. Burden of disease from household air pollution for 2012. Geneva: WHO, http://www.who.int/phe/health_topics/outdoorair/databases/FINAL_HAP_AAP_BoD_24March2014.pdf?ua¼1 (2014, accessed 20 September 2022).
- World Health Organization. Data on the size of the HIV/AIDS epidemic: number of deaths due to HIV/AIDS. Global Health Observatory Data Repository, Geneva, WHO, 2014.
- World Health Organization. Deaths: estimated deaths, data by region. Global Health Observatory Data Repository. Geneva: WHO, <http://apps.who.int/gho/data/view.main.14117?lang¼en> (2014, accessed 9 September 2022).
- Olanrewaju JA, Akinte DO, Sokan-Adeaga AA, et al. A survey on faecal management practices and associated health impacts among residents in selected sub-urban communities in Ibadan, Nigeria. *J Appl Sci* 2022; 22(3): 107–116.
- World Health Organization. Environmental Health Criteria 237. Principles for evaluating health risks in children associated with exposure to chemicals. Geneva: WHO, <http://www.who.int/ipcs/publications/ehc/ehc237.pdf> (2006, Accessed 18 September 2022).
- Grandjean P and Landrigan PJ. Neurobehavioural effects of developmental toxicity. *Lancet Neurol* 2014; 13: 330e8.
- Sokan-Adeaga MA, Sokan-Adeaga AA and Sokan-Adeaga ED. Characterization and antibiotics sensitivity pattern of *Enterobacteriaceae* in Obafemi Awolowo University (OAU) Sewage Oxidation pond. *Glob J Med Res* 2018; 18(1): 7–17.
- Sly PD, Carpenter DO, Van den Berg M, et al. Health consequences of environmental exposures: causal thinking in global environmental epidemiology. *Annals Glob Health* 2016; 82(1): 3–9.
- Allen SAA, Ree AG, Ayodeji SAM, et al. Secondary Inorganic Aerosols: Impacts on the global climate system and Human Health. *Biodiversity Int J* 2019; 3(6): 249–259.
- Delfino RJ, Sioutas C and Malik S. Potential role of ultra-fine particles in associations between airborne particle mass and cardiovascular health. *Environ Health Perspect* 2005; 113(8): 934–946.
- Hennig B, Reiterer G, Majkova Z, et al. Modification of environmental toxicity by nutrients: implications in atherosclerosis. *Cardiovasc Toxicol* 2005; 5(2): 153–160.
- Needham LL, Barr DB, Caudill SP, et al. Concentrations of environmental chemicals associated with neurodevelopmental effects in U.S. Population. *Neurotoxicol* 2017; 26(4): 531–545.
- Olarewaju SO, Sokan-Adeaga AA, Awoaife FB, et al. Assessment of occupational injuries and safety practices among automobile repair artisans in apo and Gudu area of Abuja, Nigeria. *J Epid Soc Nig* 2022; 4(2): 71–86.
- Center for Disease Control and Prevention (CDC). Third National Report on Human Exposure to Environmental Chemicals. Atlanta, GA: Centers for Disease Control and Prevention, <http://www.cdc.gov/exposurereport/> (2005, accessed 20 September 2022)
- Okareh OT, Akin-Brandom T and Sokan-Adeaga AA. Spatial distribution of heavy metals contamination of groundwater in neighborhood communities of Shagamu Industrial Layout, Nigeria. *Glob J Sci Front Res* 2018; 18(2:1): 21–38.
- Sokan-Adeaga AA, Sokan-Adeaga MA, Sokan-Adeaga, et al. Chemobiokinetics, biotoxicity and therapeutic overview of selected heavy metals poisoning: a review. *Biodiversity Int J* 2020; 4(5): 211–222.
- Hennig B, Ettinger AS, Jandacek RJ, et al. Using nutrition for intervention and prevention against environmental chemical toxicity and associated diseases. *Environ Health Perspect* 2007; 115: 493–495.
- Kim C and Park K. Association between phytochemical index and inflammation in Korean adults. *Antioxidants* 2022; 11: 348.
- Dehghani Firouzabadi F, Jayedi A, Asgari E, et al. The association of dietary phytochemical index with metabolic syndrome in adults. *Clin Nutr Res* 2021; 10: 161–171.
- Ghoreishy SM, Aminianfar A, Benisi-Kohansal S, et al. Association between dietary phytochemical index and breast cancer: A case-control study. *Breast Cancer* 2021; 28: 1283–1291.
- Delshad Aghdam S, Siassi F, Nasli Esfahani E, et al. Dietary phytochemical index associated with cardiovascular risk factor in patients with type 1 diabetes mellitus. *BMC Cardiovasc Disord* 2021; 21: 293.
- Zar HJ and Ferkol TW. The global burden of respiratory disease-impact on child health. *Pediatr Pulmonol* 2014; 49(5): 430–434.

24. Hodgson E, Mailman RB and Chambers JE. *Dictionary of toxicology*. 2nd ed. London: Macmillan Reference, 1998. p.162.
25. Hawkes SJ. What is a “heavy metal”? *J Chem Educ* 1997; 74(11): 1374.
26. Gorbachev VM, Zamyatnin YS and Lbov AA. *Nuclear reactions in heavy elements: A data handbook*. Oxford: Pergamon Press, 1980. p.12.
27. Morris CG. *Academic Press Dictionary of Science and Technology*. San Diego: Harcourt Brace Jovanovich, 1992. p.36.
28. Nriagu JO and Pacyna JM. Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature* 1988; 333(6169): 134–139.
29. Khanna S, P, Assessment of heavy metal contamination in different vegetables grown in and around urban areas. *Res J Environ Toxicol* 2011; 5(3): 162–179.
30. Okareh TO, Akin-Brandom T, Sokan-Adeaga AA, et al. Knowledge and perceived health risks associated with heavy metals contamination in groundwater – A case study of Sagamu Local Government Area, Ogun State, Nigeria. *Int J Health Safety Environ* 2020; 6(03): 506–518.
31. Nordberg GF, Fowler BA, Nordberg M, et al. *Handbook on the Toxicology of Metals*. New York, NY: Academic Press, 2005. p.23.
32. Singh J, Upadhyay SK, Pathak RK, et al. Accumulation of heavy metals in soil and paddy crop (*Oryza sativa*), irrigated with water of Ramgarh Lake, Gorakhpur, UP, India. *Toxicol Environ Chem* 2011; 93(3): 462–473.
33. Türkdoğan MK, Kilicel F, Kara K, et al. Heavy metals in soil, vegetables and fruits in the endemic upper gastrointestinal cancer region of Turkey. *Environ Toxicol Pharmacol* 2003; 13(3): 175–179.
34. Ngan V. *Heavy metal toxicity*. Hoboken, NJ: Blackwell Scientific Publications, 2006. p.51.
35. Nriagu JO. Global metal pollution. Poisoning the biosphere? *Environment* 1990; 32(7): 7–33.
36. Smith SR, Jaffe DM and Skinner MA. Case report of metallic mercury injury. *Pediatr Emerg Care* 1997; 13(2): 114–116.
37. Jaishankar M, Mathew BB, Shah MS, et al. Biosorption of few heavy metal ions using agricultural wastes. *J Environ Pollut Hum Health* 2014; 2(1): 1–6.
38. Alewu B and Nosiri C. Pesticides and human health. In: Stoytcheva M. (ed.) *Pesticides in the modern world – effects of pesticides exposure*. London: InTech, 2011, pp.231–250, <http://www.intechopen.com/books/pesticides-in-the-modern-world-effects-of-pesticidesexposure/pesticide-and-human-health>.
39. NSW EPA. What Are Pesticides and How Do They Work?, <http://www.epa.nsw.gov.au/pesticides/pestwhatrhow.htm> (2013, accessed 10 August 2022).
40. Nicolopoulou-Stamati P, Maipas S, Kotampasi C, et al. Chemical Pesticides and human health: the urgent need for a new concept in agriculture. *Public Health Front* 2016; 4: 148.
41. Pirsahab M, Limoe M, Namdari F, et al. Organochlorine pesticides residue in breast milk: a systematic review. *Med J Islam Repub Iran* 2015; 29: 228.
42. Sanborn M, Kerr KJ, Sanin LH, et al. Non-cancer health effects of pesticides. Systematic review and implications for family doctors. *Can Fam Physician* 2007; 53: 1712–1720.
43. Mnif W, Hassine AI, Bouaziz A, et al. Effect of endocrine disruptor pesticides: a review. *Int J Environ Res Public Health* 2011; 8: 2265–2303.
44. Bassil KL, Vakil C, Sanborn M, et al. Cancer health effects of pesticides: systematic review. *Can Fam Physician* 2007; 53: 1704–1711.
45. Turusov V, Rakitsky V and Tomatis L. Dichlorodiphenyltrichloroethane (DDT): ubiquity, persistence, and risks. *Environ Health Perspect* 2002; 110: 125–128.
46. van den Berg H. Global status of DDT and its alternatives for use in vector control to prevent disease. *Environ Health Perspect* 2009; 117: 1656–1663.
47. McKinlay R, Plant JA, Bell JN, et al. Endocrine disrupting pesticides: implications for risk assessment. *Environ Int* 2008; 34: 168–183.
48. Gasnier C, Dumont C, Benachour N, et al. Glyphosate-based herbicides are toxic and endocrine disruptors in human cell lines. *Toxicology* 2009; 262: 184–191.
49. Jaga K and Dharmani C. Sources of exposure to and public health implications of organophosphate pesticides. *Rev Panam Salud Publica* 2003; 14: 171–185.
50. Karami-Mohajeri S and Abdollahi M. Toxic influence of organophosphate, carbamate, and organochlorine pesticides on cellular metabolism of lipids, proteins, and carbohydrates: a systematic review. *Hum Exp Toxicol* 2011; 30(9): 1119–1140.
51. Li D, Huang Q, Lu M, et al. The organophosphate insecticide chlorpyrifos confers its genotoxic effects by inducing DNA damage and cell apoptosis. *Chemosphere* 2015; 135: 387–393.
52. Goad RT, Goad JT, Atieh BH, et al. Carbofuran-induced endocrine disruption in adult male rats. *Toxicol Mech Methods* 2004; 14: 233–239.
53. Jamal F, Haque QS, Singh S, et al. The influence of organophosphate and carbamate on sperm chromatin and reproductive hormones among pesticide sprayers. *Toxicol Ind Health* 2016; 32(8): 1527–1536.
54. Kolaczinski JH and Curtis CF. Chronic illness as a result of low-level exposure to synthetic pyrethroid insecticides: a review of the debate. *Food Chem Toxicol* 2004; 42: 697–706.
55. Ray DE and Fry JR. A reassessment of the neurotoxicity of pyrethroid insecticides. *Pharmacol Ther* 2006; 111: 174–193.
56. Jaensson A, Scott AP, Moore A, et al. Effects of a pyrethroid pesticide on endocrine responses to female odours and reproductive behaviour in male parr of brown trout (*Salmo trutta* L. *Aquat Toxicol* 2007; 81(1): 1–9.
57. Pandey SP and Mohanty B. The neonicotinoid pesticide imidacloprid and the dithiocarbamate fungicide mancozeb disrupt the pituitary-thyroid axis of a wildlife bird. *Chemosphere* 2015; 122: 227–234.
58. Moore A and Waring CP. The effects of a synthetic pyrethroid pesticide on some aspects of reproduction in Atlantic salmon (*Salmo salar* L. *Aquat Toxicol* 2001; 52(1): 1–12.

59. Jin Y, Wang L, Chen G, et al. Exposure of mice to atrazine and its metabolite diaminochlorotriazine elicits oxidative stress and endocrine disruption. *Environ Toxicol Pharmacol* 2014; 37: 782–790.
60. Huang P, Yang J and Song Q. Atrazine affects phosphoprotein and protein expression in MCF-10A human breast epithelial cells. *Int J Mol Sci* 2014; 15: 17806–17826.
61. Ma K, Wu HY, Zhang B, et al. Neurotoxicity effects of atrazine-induced SH-SY5Y human dopaminergic neuroblastoma cells via microglial activation. *Mol Biosyst* 2015; 11: 2915–2924.
62. Goulson D. REVIEW: an overview of the environmental risks posed by neonicotinoid insecticides. *J Appl Ecol* 2013; 50: 977–987.
63. Jeschke P and Nauen R. Neonicotinoids—from zero to hero in insecticide chemistry. *Pest Manag Sci* 2008; 64: 1084–1098.
64. Caron-Beaudoin Denison MS and Sanderson JT. Effects of neonicotinoids on promoter-specific expression and activity of aromatase (CYP19) in human adrenocortical carcinoma (H295R) and primary umbilical vein endothelial (HUVEC) cells. *Toxicol Sci* 2016; 149(1): 134–144.
65. Kotsonis FN, Burdock GA and Flamm WG. *Food toxicology, Casarett and Doull's toxicology: the basic science of poisons*. 6th ed. New York, NY: McGraw-Hill, 2001. pp.1049–1088.
66. Bruckner JV and Warren DA. *Toxic effects of solvents and vapors, Casarett and Doull's Toxicology: the Basic Science of Poisons*. 6th ed. New York, NY: McGraw-Hill, 2001. pp.945–964.
67. Brent J. A review of: medical toxicology. *Clin Toxicol* 2006; 44(3): 355–355.
68. Monisha J, Tenzin T, Naresh A, et al. Toxicity, mechanism and health effects of some heavy metals. *Interdiscip Toxicol* 2014; 7(2): 60–72.
69. Patrick L. Mercury toxicity and antioxidants: Part 1: role of glutathione and alpha-lipoic acid in the treatment of mercury toxicity. *Altern Med Rev* 2002; 7(6): 456–471.
70. Haley BE. Mercury toxicity: genetic susceptibility and synergistic effects. *Med Veritas* 2005; 2(2): 535–542.
71. Flora SJ, Mittal M and Mehta A. Heavy metal induced oxidative stress & its possible reversal by chelation therapy. *Indian J Med Res* 2008; 128: 501–523.
72. Martin S and Griswold W. Human health effects of heavy metals. *Environ Sci Technol Briefs Citizens* 2009; 15: 1–6.
73. Smith AH, Lingas EO and Rahman M. Contamination of drinking-water by arsenic in Bangladesh: a public health emergency. *Bull World Health Organ* 2000; 78(9): 1093–1103.
74. Guha Mazumder DN. Chronic arsenic toxicity & human health. *Indian J Med Res* 2008; 128(4): 436–447.
75. Stober J. Health effects of POPs, Proceedings of the Subregional Awareness Raising Workshop on Persistent Organic Pollutants (POPs) Kranjska Gora, Slovenia, 11-14 May 1998.
76. Rengarajan T, Rajendran P, Nandakumar N, et al. Exposure to polycyclic aromatic hydrocarbons with special focus on cancer. *Asian Pac J Trop Biomed* 2015; 5: 182–189.
77. Abdel-Shafy HI and Mansour MSM. A review on polycyclic aromatic hydrocarbons: source, environmental impact, effect on human health and remediation. *Egypt J Pet* 2016; 25: 107–123.
78. Burchiel SW and Gao J. *Polycyclic aromatic hydrocarbons and the immune system*. In: Vohr H-W (ed.) *Encyclopedia Immunotoxicology*. Berlin: Springer, 2014, pp.1192–1194.
79. Bolden AL, Rochester JR, Schultz K, et al. Polycyclic aromatic hydrocarbons and female reproductive health: a scoping review. *Reprod Toxicol* 2017; 73: 61–74.
80. Rajpara RK, Dudhagara DR, Bhatt JK, et al. Polycyclic aromatic hydrocarbons (pahs) at the Gulf of Kutch, Gujarat, India: occurrence, source apportionment, and toxicity of pahs as an emerging issue. *Mar Pollut Bull* 2017; 119: 231–238.
81. Campos-Vega R and Oomah DB. Chemistry and classification of phytochemicals. In: Tiwari BK, Brunton NP and Brennan CS (eds.) *Handbook of plant food phytochemicals: sources, stability and extraction*. Hoboken: John Wiley & Sons, Ltd, 2013, pp.5–48.
82. Ogunwenmo KO, Idowu OA, Innocent C, et al. Cultivars of *Codiaeum variegatum* (L.) Blume (Euphorbiaceae) show variability in phytochemical and cytological characteristics. *J Biotechnol* 2007; 6: 2400–2405.
83. Ngoci SN, Mwendia CM and Mwaniki CG. Phytochemical and cytotoxicity testing of *Indigofera lupatana* Baker F. *J Anim Plant Sci* 2011; 11(1): 1364–1373.
84. Briskin DP. Medicinal plants and phytomedicines. Linking plant Biochemistry and Physiology to human health. *Plant Physiol* 2000; 124: 507–514.
85. Olila D, Olwa-Odyek and Opuda-Asibo J. Antibacterial and antifungal activities of extracts of *Zanthoxylum chalybeum* and *Warburgia ugandensis*, Ugandan medicinal plants. *J Afri Health Sci* 2001; 1: 66–72.
86. Harborne JB. *Phytochemical methods*. London: Chapman and Hall, 1973. pp.52–114.
87. Ovando C, Hernandez D, Hernandez E, et al. Chemical studies of anthocyanins: a review. *Food Chem* 2009; 113: 859–871.
88. Metodieva D, Kochman A and Karolczak S. Evidence for antiradical and antioxidant properties of four biologically active N,N-diethylaminoethyl ethers of flavanone oximes: a comparison with natural polyphenolic flavonoid rutin action. *Biochem Mol Biol Int* 1997; 41: 1067–1075.
89. Panche AN, Diwan AD and Chandra SR. Flavonoids: an overview. *J Nutr Sci* 2016; 5: e47.
90. Navarro MC, Montilla MP, Cabo MM, et al. Antibacterial, anti-protozoal and antioxidant activity of five plants used in izabal for infectious diseases. *J Phytotherapy Res* 2003; 17: 325–329.
91. Scalbert A, Johnson IT and Saltmarsh M. Polyphenols: Antioxidants and beyond. *Am J Clin Nutr* 2005; 81: 215S–217S.
92. Okuda T. Systematics and health effects of chemically distinct tannins in medicinal plants. *J Phytochemistry* 2005; 66: 2012–2031.
93. El Aziz MMA, Ashour AS and Melad ASG. A review on saponins from medicinal plants: chemistry, isolation, and determination. *J Nanomed Res* 2019; 7(4): 282–288.
94. Vasudeva RN, Sukhendu BG, Pushpalatha B, et al. Triterpenoid saponins: a review on biosynthesis, applications and mechanism of their action. *Int J Pharm Pharm Sci* 2015; 7(1): 24–28.
95. Hassan SM, Haq AU, Byrd JA, et al. Haemolytic and antimicrobial activities of saponin-rich extracts from guar meal. *Food Chem* 2010; 119: 600–605.

96. Sparg SG, Light ME and van Staden J. Biological activities and distribution of plant saponins. *J Ethnopharmacol* 2004; 94(2-3): 219–243.
97. Cheng TC, Lu JF, Wang JS, et al. Antiproliferation effect and apoptosis mechanism of prostate cancer cell PC-3 by flavonoids and saponins prepared from *Gynostemma pentaphyllum*. *J Agric Food Chem* 2011; 59(20): 11319–11329.
98. Hostettmann and Marson A. *Saponins*. Cambridge: Cambridge University Press, 2005. p.21.
99. Kurek J. Introductory Chapter: Alkaloids – their importance in nature and for human life [internet], 2019 Nov 13. <http://dx.doi.org/10.5772/intechopen.85400>
100. Chisholm H. Encyclopedia Britannica. *A dictionary of arts, sciences, literature and general information*. Vol 22. New York, NY: Sagwan Press, 2015. pp.978134014998.
101. Prassas I and Diamandis EP. Novel therapeutic applications of cardiac glycosides. *Reviews* 2008; 7: 926–935.
102. Morsy N. *Cardiac glycosides in medicinal plants. Aromatics and Medicinal Plants – Back to Nature [Internet]*. London: Intechopen Publisher, 2017.
103. Kelly RA. Cardiac glycosides and congestive heart failure. *Am J Cardiol* 1990; 65: E10–E16.
104. Perveen S. *Introductory Chapter: Terpenes and Terpenoids. Terpenes and Terpenoids [internet]*. London: Intechopen Publisher, 2018.
105. Roberts SC. Production and engineering of terpenoids in plant cell culture. *Nat Chem Biol* 2007; 3: 387–395.
106. Lesovaya E, Yemelyanov A, Swart AC, et al. Discovery of compound A - a selective activator of the glucocorticoid receptor with anti-inflammatory and anti-cancer activity. *Oncotarget* 2015; 6(31): 30730–30744.
107. Toh MF, Mendonca E, Eddie SL, et al. Kaempferol exhibits progestogenic effects in ovariectomized rats. *J Steroids Horm Sci* 2014; 5: 136.
108. Pihlajamaa P, Zhang FP, Saarinen L, et al. The phytoestrogen genistein is a tissue-specific androgen receptor modulator. *Endocrinology* 2011; 152: 4395–4405.
109. Blachley JD and Knochel JP. Tobacco chewer's hypokalemia: Licorice Revisited. *N Engl J Med* 1980; 302: 784–785.
110. Edeoga HO, Okwu DE and Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. *Afr J Biotechnol* 2005; 4: 685–688.
111. Soares MBP, Brustolim D, Santos LA, et al. Physalins B, F and G, seco-steroids purified from *Physalis angulata* L., Inhibit lymphocyte function and allogeneic transplant rejection. *Int Immunopharmacol* 2006; 6: 408–414.
112. Dias DA, Urban S and Roessner U. A historical overview of natural products in drug discovery. *Metabolites* 2012; 2(2): 303–336.
113. Patti AM, Al-Rasadi K, Giglio RV, et al. Natural approaches in metabolic syndrome management. *Arch Med Sci* 2018; 14(2): 422–441.
114. Tabatabaei-Malazy O, Larijani B and Abdollahi M. Targeting metabolic disorders by natural products. *J Diabetes Metab Disord* 2015; 14: 57.
115. Truong VL, Jun M and Jeong WS. Role of resveratrol in regulation of cellular defense systems against oxidative stress. *Biofactors* 2018; 44(1): 36–49.
116. Queen B and Tollefsbol T. Polyphenols and aging. *Curr Aging Sci* 2010; 3(1): 34–42.
117. Li YR, Li S and Lin CC. Effect of resveratrol and pterostilbene on aging and longevity. *Biofactors* 2018; 44(1): 69–82.
118. Prasain JK, Carlson SH and Wyss JM. Flavonoids and age-related disease: risk, benefits and critical windows. *Maturitas* 2010; 66(2): 163–171.
119. Alam MA, Subhan N, Rahman MM, et al. Effect of Citrus flavonoids, naringin and naringenin, on metabolic syndrome and their mechanisms of action. *Adv Nutrition* 2014; 5(4): 404–417.
120. Nobili S, Lippi D, Witort E, et al. Natural compounds for cancer treatment and prevention. *Pharmacol Res* 2009; 59(6): 365–378.
121. Hussain T, Tan B, Ren W, et al. Polyamines: therapeutic perspectives in oxidative stress and inflammatory diseases. *Amino Acids* 2017; 49(9): 1457–1468.
122. Forni C, Facchiano F, Bartoli M, et al. Beneficial role of phytochemicals on oxidative stress and Age-Related diseases. *Biomed Res Int* 2019; 2019: 8748253.
123. Lentini A, Tabolacci C, Provenzano B, et al. Phytochemicals and protein–polyamine conjugates by transglutaminase as chemopreventive and chemotherapeutic tools in cancer. *Plant Physiol Biochem* 2010; 48(7): 627–633.
124. Mao XY, Jin MZ, Chen JF, et al. Live or let die: neuroprotective and anti-cancer effects of nutraceutical antioxidants. *Pharmacol Ther* 2018; 183: 137–151.
125. Bobe G, Albert PS, Sansbury LB, et al. Interleukin-6 as a potential indicator for prevention of high-risk adenoma recurrence by dietary flavonols in the polyp prevention trial. *Cancer Prev Res* 2010; 3(6): 764–775.
126. Anand P, Kunnumakkara AB, Newman RA, et al. Bioavailability of curcumin: problems and promises. *Mol Pharm* 2007; 4(6): 807–818.
127. Mukund V, Mukund D, Sharma V, et al. Genistein: its role in metabolic diseases and cancer. *Crit Rev Oncol/Hematol* 2017; 119: 13–22.
128. Lesinski GB, Reville PK, Mace TA, et al. Consumption of soy isoflavone enriched bread in men with prostate cancer is associated with reduced proinflammatory cytokines and immunosuppressive cells. *Cancer Prev Res* 2015; 8(11): 1036–1044.
129. Carini F, David S, Tomasello G, et al. Colorectal cancer: an update on the effects of lycopene on tumor progression and cell proliferation. *J Biol Regul Homeost Agents* 2017; 31: 769–774.
130. Rowles JL, Ranard KM, Smith JW, et al. Increased dietary and circulating lycopene are associated with reduced prostate cancer risk: a systematic review and meta-analysis. *Prostate Cancer Prostatic Dis* 2017; 20(4): 361–377.
131. Wahle KW, Caruso D, Ochoa JJ, et al. Olive oil and modulation of cell signaling in disease prevention. *Lipids* 2004; 39(12): 1223–1231.
132. Baker EJ, Yusof MH, Yaqoob P, et al. Omega-3 fatty acids and leukocyte-endothelium adhesion: novel anti-atherosclerotic actions. *Mol Asp Medicine* 2018; 64: 169–181.
133. Colussi G, Catena C, Novello M, et al. Impact of omega-3 polyunsaturated fatty acids on vascular function and blood

- pressure: relevance for cardiovascular outcomes. *Nutr Metab Cardiovasc Dis* 2017; 27(3): 191–200.
134. Payne C, Wiffen PJ and Martin S. Interventions for fatigue and weight loss in adults with advanced progressive illness. *Cochrane Database Syst Rev* 2012; 1: CD008427.
 135. Kones R, Howell S and Rumana U. N-3 polyunsaturated fatty acids and cardiovascular disease: principles, practices, pitfalls, and promises - a contemporary review. *Med Princ Pract* 2018; 26(6): 497–508.
 136. Gammone MA, Efthymakis K, Pluchinotta FR, et al. Impact of chocolate on the cardiovascular health. *Front Biosci* 2018; 23(3): 852–864.
 137. Zhang H, Liu H, Chen Y, et al. The curcumin-induced vasorelaxation in rat superior mesenteric arteries. *Ann Vasc Surg* 2018; 48: 233–240.
 138. Gammone MA, Pluchinotta FR, Bergante S, et al. Prevention of cardiovascular diseases with carotenoids. *Front Biosci* 2017; 9(1): 165–171.
 139. Howes MJR. Phytochemicals as anti-inflammatory nutraceuticals and phytopharmaceuticals. In: Chatterjee S, Jungraithmayr W and Bagchi D. (eds) *Immunity and inflammation in health and disease: emerging roles of Nutraceuticals and functional foods in immune support*. Elsevier, Cambridge, MA: Academic Press, 2017, pp.363–388.
 140. Gomes A, Fernandes E, Lima J, et al. Molecular mechanisms of anti-inflammatory activity mediated by flavonoids. *Curr Med Chem* 2008; 15(16): 1586–1605.
 141. King HGC. Phenolic compounds of commercial wheat germ. *J Food Sci* 1962; 27(5): 446–454.
 142. Raso GM, Meli R, Di Carlo G, et al. Inhibition of inducible nitric oxide synthase and cyclooxygenase-2 expression by flavonoids in macrophage J774A.1. *Life Sci* 2001; 68(8): 921–931.
 143. Hirai S, Takahashi N, Goto T, et al. Functional food targeting the regulation of obesity-induced inflammatory responses and pathologies. *Mediat Inflammation* 2010; 2010: 1–8.
 144. Xagorari A, Papapetropoulos A and Mauromatis A, M. Luteolin inhibits an endotoxin-stimulated phosphorylation cascade and proinflammatory cytokine production in macrophages. *J Pharmacol Exp Ther* 2001; 296(1): 181–187.
 145. Javadi F, Ahmadzadeh A, Egtesadi S, et al. The effect of quercetin on inflammatory factors and clinical symptoms in women with rheumatoid arthritis: a double-blind, randomized controlled trial. *J Am Coll Nutr* 2017; 36(1): 9–15.
 146. Shamekhi Z, Amani R, Habibagahi Z, et al. A randomized, double-blind, placebo-controlled clinical trial examining the effects of green tea extract on systemic lupus erythematosus disease activity and quality of Life. *Phytother Res* 2017; 31(7): 1063–1071.
 147. Joseph J, Cole G, Head E, et al. Nutrition, brain aging, and neurodegeneration. *J Neurosci* 2009; 29(41): 12795–12801.
 148. Granzotto A and Zatta P. Resveratrol acts not through AntiAggregative pathways but mainly via its scavenging properties against $\text{A}\beta$ and $\text{A}\beta$ -metal complexes toxicity. *PLoS One* 2011; 6(6): e21565.
 149. Yan WJ, Liu RB, Wang LK, et al. Sirt3-mediated autophagy contributes to resveratrol-induced protection against ER stress in HT22 cells. *Front Neurosci* 2018; 12: 116.
 150. Capiralla H, Vingtdoux V, Zhao H, et al. Resveratrol mitigates lipopolysaccharide- and $\text{A}\beta$ -mediated microglial inflammation by inhibiting the TLR4/NF- κ B/STAT signaling cascade. *J Neurochem* 2012; 120(3): 461–472.
 151. Porquet D, Casadesús G, Bayod S, et al. Dietary resveratrol prevents alzheimer's markers and increases life span in samp8. *Age* 2013; 35(5): 1851–1865.
 152. Zhang F, Shi JS, Zhou H, et al. Resveratrol protects dopamine neurons against lipopolysaccharide-induced neurotoxicity through its anti-inflammatory actions. *Mol Pharmacol* 2010; 78(3): 466–477.
 153. Dutta K, Ghosh D and Basu A. Curcumin protects neuronal cells from japanese encephalitis virus-mediated cell death and also inhibits infective viral particle formation by dysregulation of ubiquitin-proteasome system. *J Neuroimmune Pharmacol* 2009; 4(3): 328–337.
 154. Ansari MA, Abdul HM, Joshi G, et al. Protective effect of quercetin in primary neurons against $\text{A}\beta(1-42)$: relevance to Alzheimer's disease. *J Nutr Biochem* 2009; 20(4): 269–275.
 155. Sabogal-Guáqueta AM, Muñoz-Manco JI, Ramírez-Pineda JR, et al. The flavonoid quercetin ameliorates Alzheimer's disease pathology and protects cognitive and emotional function in aged triple transgenic Alzheimer's disease model mice. *Neuropharmacol* 2015; 93: 134–145.
 156. Gardener H and Caunca MR. Mediterranean diet in preventing neurodegenerative diseases. *Curr Nutr Rep* 2018; 7(1): 10–20.
 157. Foyer CH and Noctor G. Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell* 2005; 17(7): 1866–1875.
 158. Jomova K and Valko M. Health protective effects of carotenoids and their interactions with other biological antioxidants. *Eur J Med Chem* 2013; 70: 102–110.
 159. Stahl W and Sies H. Antioxidant activity of carotenoids. *Mol Asp Medicine* 2003; 24(6): 345–351.
 160. Hassan S and Mathesius U. The role of flavonoids in root-rhizosphere signalling: opportunities and challenges for improving plant-microbe interactions. *J Exp Bot* 2012; 63(9): 3429–3444.
 161. European Medicines Agency. Guideline on declaration of herbal substances and herbal preparations ¹ in herbal medicinal products ² /traditional herbal medicinal products, https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-declaration-herbal-substances-herbal-preparations-herbal-medicinal-products/traditional-herbal-medicinal-products-spc_en.pdf (2010, accessed 16 September 2022).
 162. Moran JF, Klucas RV, Grayer RJ, et al. Complexes of iron with phenolic compounds from soybean nodules and other legume tissues: prooxidant and antioxidant properties. *Free Radic Biol Med* 1997; 22(5): 861–870.
 163. Aharoni A, Jongsma MA and Bouwmeester HJ. Volatile science? Metabolic engineering of terpenoids in plants. *Trends Plant Sci* 2005; 10(12): 594–602.

164. Baratta MT, Dorman HJD, Deans SG, et al. Chemical composition, antimicrobial and antioxidative activity of Laurel, sage, rosemary, oregano and coriander essential oils. *J Essent Oil Res* 1998; 10(6): 618–627.
165. Tiong S, Looi C, Hazni H, et al. Antidiabetic and antioxidant properties of alkaloids from *Catharanthus roseus* (L.) G. Don. *Molecules* 2013; 18(8): 9770–9784.
166. Micali G, Paternò V, Cannarella R, et al. Evidence-based treatment of atopic dermatitis with topical moisturizers. *Giornale Italiano di Dermatologia e Venereologia* 2018; 153(3): 396–402.
167. Di Franco R, Sammarco E, Calvanese MG, et al. Preventing the acute skin side effects in patients treated with radiotherapy for breast cancer: the use of corneometry in order to evaluate the protective effect of moisturizing creams. *Radiat Oncol* 2013; 8: 57.
168. Turk S, Malkan UY, Ghasemi M, et al. Growth inhibitory activity of ankaferd hemostat on primary melanoma cells and cell lines. *Sage Open Med* 2017; 5: 2050312116689519.
169. Brito FF, Gimeno PM, Bartolomé B, et al. Vine pollen allergy in areas with a high density of vineyards. *Ann Allergy Asthma Immunol* 2008; 100(6): 596–600.
170. Hammer KA. Treatment of acne with tea tree oil (melaleuca) products: a review of efficacy, tolerability and potential modes of action. *Int J Antimicrob Agents* 2015; 45(2): 106–110.
171. Kwon HH, Yoon JY, Park SY, et al. Comparison of clinical and histological effects between lactobacillus-fermented *chamaecyparis obtusa* and tea tree oil for the treatment of acne: an eight-week double-blind randomized controlled split-face study. *Dermatology* 2014; 229(2): 102–109.
172. Ramage G, Milligan S, Lappin DF, et al. Antifungal, cytotoxic, and immunomodulatory properties of tea tree oil and its derivative components: potential role in management of oral candidosis in cancer patients. *Front Microbiol* 2012; 3: 220.
173. Pazyar N, Yaghoobi R, Bagherani N, et al. A review of applications of tea tree oil in dermatology. *Int J Dermatol* 2013; 52(7): 784–790.
174. Kalyana Sundaram I, Sarangi DD, Sundararajan V, et al. Poly herbal formulation with antielastase and anti-oxidant properties for skin anti-aging. *BMC Complement Altern Med* 2018; 18(1): 33.
175. Prasad R, Singh T and Katiyar SK. Honokiol inhibits ultraviolet radiation-induced immunosuppression through inhibition of ultraviolet-induced inflammation and DNA hypermethylation in mouse skin. *Sci Rep* 2017; 7: 1.
176. Sarandy MM, Novaes RD, Xavier AA, et al. Hydroethanolic extract of *strychnos pseudoquina* accelerates skin wound healing by modulating the oxidative status and microstructural reorganization of scar tissue in experimental type I diabetes. *Biomed Res Int* 2017; 2017: 9538351.
177. Houston DMJ, Robins B, Bugert JJ, et al. In vitro permeation and biological activity of punicalagin and zinc (II) across skin and mucous membranes prone to herpes simplex virus infection. *Eur J Pharm Sci* 2017; 96: 99–106.