

|   |
|---|
| Access this article online  |
| Quick Response Code:  |
|  |
| Website:<br>www.jehp.net  |
| DOI:<br>10.4103/jehp.jehp_1025_20   |

# Improvement of medication error reporting: An applied motivation program in pediatric units

Laleh Massah, Rouhangiz Mohammadi<sup>1</sup>, Mahboobeh Namnabati<sup>2</sup>

## Abstract:

**BACKGROUND:** The rate of reporting medication errors is lower than the actual rate. The role of education, encouragement, and promotion of reporting culture can contribute to the increase of errors reporting. This study was to investigate the effect of motivational program on reporting of medication errors in the pediatric units.

**MATERIALS AND METHODS:** A quasi-experimental study with pre- and post-test design was done through 60 nurses working in the pediatric wards of the selected hospital in Isfahan, Iran in 2018–2019. Nurses participated in the study through the census sampling method. Data were collected using a questionnaire based on eight medication principles completed by the nurses before and after the intervention. The intervention was conducted through a motivational program in two parts of training and positive reinforcement. Data were analyzed through the descriptive statistics (frequency, mean, and standard deviation [SD]) as well as inferential statistics (Wilcoxon test and paired *t*-test).

**RESULTS:** After the intervention, the most medication error reports (60%) were in the principle of right medication (similar name). Furthermore, the total score (SD) of medication error reporting was significantly increased from 7.35 (2.43) before the intervention to 17.04 (5.59) after the intervention ( $t = 8.03, P < 0.001$ ).

**CONCLUSIONS:** Motivational program increased medication error reports. Therefore, the managers can improve medication care and the safety of children through continuous education and encouragement in the hospitals.

## Keywords:

Care, Iran, medication errors, motivation, nurse, pediatrics

Pediatric Department,  
Gharazi Hospital,  
Isfahan, Iran, <sup>1</sup>Pediatric  
Department, Emam  
Hossein Hospital, Isfahan  
University of Medical  
Sciences, Isfahan, Iran,  
<sup>2</sup>Department of Pediatric  
and Neonates, Nursing  
and Midwifery Care  
Research Center, Faculty  
of Nursing and Midwifery,  
Isfahan University of  
Medical Sciences, Isfahan,  
Iran

## Address for correspondence:

Dr. Mahboobeh  
Namnabati,  
Nursing and Midwifery  
Care Research Center,  
Faculty of Nursing  
and Midwifery, Isfahan  
University of Medical  
Sciences, Isfahan, Iran.  
Hezarjarib Blvd., Isfahan,  
Iran.  
E-mail: namnabat@  
nm.mui.ac.ir

Received: 13-08-2020  
Accepted: 12-10-2020  
Published: 31-05-2021

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

and effective communication to prevent medication error and keep the safety of patients.<sup>[2]</sup>

According to the studies, the rate of reporting errors among nurses is lower than the actual rate.<sup>[2,3]</sup> Ghorbanpour Diz *et al.* in a study, conducted in the hospitals in Tehran, showed that only 28% of the 112 nurses who had medication errors over the last year had reported their errors.<sup>[4]</sup> In another study, 91.2% of the participants had reported at least a case of medication error during the last month.<sup>[5]</sup> Previous studies have been conducted on identifying the rate of medication errors. Regarding the rate of

**How to cite this article:** Massah L, Mohammadi R, Namnabati M. Improvement of medication error reporting: An applied motivation program in pediatric units. *J Edu Health Promot* 2021;10:189.

medication errors in hospitalized children, it must be used two or more interventions as a system to develop solutions and involve nurses. In a study indicated that implementing the interventions were reduced about 57% medication error in the pediatric wards.<sup>[6]</sup> In the previous study mentioned to need a medication error reporting program that revealed the overall error rate, the error types and system breakdown points. Regarding a gap between the actual number of medication errors and reporting errors, we need a system has been effective in reporting medication error and improve quality assurance.<sup>[7]</sup>

According above, designing a motivational program to increase nurses' medication error reports may prevent the repetition of similar errors and destructive consequences. There are various interventions to design such a program, one of which is motivating through using techniques such as rewarding and positive reinforcement that can desirably influence the enhancement of the nurses' behavior.<sup>[8]</sup> Motivation is a requirement for the nurse's progress. Motivated nurses can have a better performance.<sup>[9]</sup> Therefore, it is necessary to pay more attention to the motivation of nurses to improve them in reporting attitude. Accordingly, in order to lessen medication errors and their various consequences and increase reporting of errors, the researchers designed a motivational program and evaluated the effect of this program on the rate of medication error reporting in the pediatric wards.

## Materials and Methods

This quasi-experimental study was conducted with pre- and post-test design in one group. It investigated the effect of motivational program on medication error reporting among the nurses working in the pediatric wards of the selected hospital of Isfahan University of Medical Sciences from November 2018 to February 2019 in Isfahan, Iran. The researcher introduced herself and stated the aims and process of the study, invited all of the nurses working in the three internal wards of the hospital who met the inclusion criteria to participate in the study. Census sampling method was performed with 60 nurses in the selected pediatric wards in the hospital. Data collection tool was a two-part researcher-made questionnaire: (1) Demographic information and (2) Medication error reporting questionnaire based on eight medication administration principles. The frequency of medication reporting errors is classified according to eight principles of medication, including 32 questions in eight sections and based on the amount of reporting (0, 1-2, 3-4, and 5 report). The validity of the questionnaire was examined by ten faculty members of the Nursing Faculty of Isfahan University of Medical Sciences, and its reliability was measured using Cronbach's alpha (0.87).

The intervention of this study involved designing a two-part motivational program. The first part was a training program consisting of two sessions on medication errors based on eight medication administration principles (right medication, right dose, right route, right patient, right reason, right time, right recording, and right response), the necessity and importance of medication errors reporting, etc. The second part was the motivational program which was done through collaboration of the head the nurses, educational supervisors, and the hospital authorities with providing positive reinforcement and reward. The nurses who had reported the highest rate of errors were eligible to receive positive reinforcement and reward in the pediatric wards.

The method of the research was such that the questionnaires were first distributed among all nurses in the three wards of the hospital along with giving information about the objectives of the research and then, the intervention was performed for 4 months through the autumn and winter season during which training sessions were held for the nurses advising the importance of medication errors reporting. A set of the same questionnaires was completed by the nurses as posttests. There is a form in the ward for reporting the medication errors that must be filling by the nurse. In collaboration with the patient-safety unit, the accuracy of the submitted reports was verified. Finally, the effect of the motivational program was measured based on the eight principles of medication administration by comparing the mean scores of the research subjects obtained from the medication error reports before and after the intervention. After the intervention, the nurses who were active in medication error reporting were appreciated according to the motivational program.

Data were analyzed using the descriptive (frequency distribution and mean) and analytical (Wilcoxon test and paired *t*-test) statistics. SPSS software (version 16, SPSS Inc., Chicago, IL, USA) was used for the data analysis in this study.

## Ethical considerations

For ethical considerations and confidentiality of the information, the questionnaires were used anonymously. After obtaining the permission of the vice chancellor for research and technology of Isfahan University of Medical Sciences, the researcher referred to the selected hospital. The participant was informed about the purpose of the study and invited to the class. The consent of the nurses was obtained, and the study was approved by the Ethics Committee of Isfahan University of Medical Sciences (IR.MUI.RESEARCH.REC.1397.338).

## Results

The mean (standard deviation) age of nurses was 31.43 (7.44) years with 7.20 (6.21) years of working experience. Most of the nurses were married 34 (56.7%) with bachelor's degree 56 (93.3%) and rotational shiftwork 53 (88.3%). The participating nurses reported that in the past 4 months before the intervention, the rate of the medication errors they committed or witnessed being done by a colleague has been 52 (86.7%), and that they had reported 45 (75%) of the errors. The results of the Wilcoxon test showed that the rate of medication errors reporting was significantly increased in most of the principles after the intervention compared to before the intervention ( $P < 0.05$ ) [Table 1]. In addition, the paired *t*-test showed that the mean of total score for the nurses' medication error reporting was significantly higher in all principles after the intervention than before the intervention ( $P < 0.05$ ). The highest rates were related to the principles of right dose and right patient [Table 2].

## Discussion

In the present study, the total score of medication error reporting was significantly increased after the intervention and the most prevalent medication error was in the principle of right medication related to similar name. It is shown in a study that one of the most important causes of medication errors was similar names and similar packing of the medicines from the viewpoint of nurses.<sup>[10]</sup> Therefore, the familiarity of nurses with similar names and abbreviations of the ward medicines is essential to avoid the errors during the work time.

In the present study, the most frequently reported error in the area of the right dose was due to the incompatibility of the dose with the weight and age of the child. This finding was consistent with the findings of Mozafari and Borji study.<sup>[11]</sup> Wrong dose usually occurs for a variety of reasons, including unfamiliar medicine, lack of knowledge of medications, and incorrect medicine calculations.<sup>[3]</sup>

The findings of the present study showed that there was no significant difference in the rate of medication error reporting for the principle of right route before and after the intervention. Given that this medication error is the most dangerous one and may be seriously harmful to one's health, fear of legal authorities and the difficulties it can cause to one's job continuation may lead the nurses to refuse to report their errors related to this principle. However, different educations and increasing the awareness of nurses and paying more attention to this principle may have reduced this medication error.

Based on the results of the study, the right time of medication administration, the highest rate of errors

belonged to nonadministration of medication in due time. In this regard, Tabatabaee *et al.* showed that the error of medication administration either before or after the due time they had was the highest frequency.<sup>[12]</sup> Similarly, in another study showed 13.3% of oral medications and 18.3% of injectable medications were administered to patients in wrong time. In addition, in this study, the rate of nurses' errors in the administration of medication orders was evaluated and revealed that 60% of nurses did not wash their hands when they prepared medication.<sup>[11]</sup> In the hospital setting of this study, some conditions of the ward such as delay in patient's visit or delays in delivering medication to the ward may sometimes affect the right time of medication administration. Moreover, nurses sometimes give medications to patients sooner or later because of overcrowding or limited number of personnel.

In the present study, investigating each of the principles showed that the motivational program used in this study was effective and educational interventions or the creation of an environment without punishment and encouragement of personnel to report would promote reporting. Consistent with this finding, Farzi *et al.* revealed that the nurses report their mistakes when they feel safe and when they are sure that their report lead to no harmful results for them.<sup>[13]</sup>

The intervention of the present study showed that motivation is a force that can encourage individuals and increase error reporting. The awareness generated through training classes, positive reinforcement with rewards such as score and gift, were strategies used in this motivational program in collaboration with hospital officials. Marquis and Huston refer to positive reinforcement by emphasizing one's desirable behavior and being encouraged for repeating that behavior.<sup>[14]</sup> Similarly, Toloei *et al.* showed motivation factors such as awareness, recognition, and appreciation and encouragement have a great impact on nurses.<sup>[15]</sup> Encouragement and appreciation develop a sense of satisfaction in nurses to the promotion of positive self-esteem, self-actualization, and employee development. Acknowledgment, in addition to being an external reward, can create healthy competition and foster a sense of respect and gratitude in individuals.<sup>[16]</sup>

In addition, interprofessional education program can improve the patient safety. In fact, the solving of medication care problems needs to implement an interprofessional team-based medication as well as improve the safety culture and teamwork climate in the hospital especial in pediatric wards.<sup>[17-19]</sup> In this study, the researcher observed that the nurses were more likely to report an error after passing a course of training and positive reinforcement. The nurses wanted to have these

**Table 1: Frequency distribution and comparison of medication error reporting based on the eight principles of medication administration**

| Eight principles   | Before intervention   |           |         |         | After intervention    |           |           |         | P      |
|--|-----------------------|-----------|---------|---------|-----------------------|-----------|-----------|---------|--------|
|  | Reporting rate, n (%) |           |         |         | Reporting rate, n (%) |           |           |         |        |
|  | Errors number         |           |         |         |                       |           |           |         |        |
|  | 0                     | 1-2       | 3-4     | 5       | 0                     | 1-2       | 3-4       | 5       |        |
| 1. Right medication  |                       |           |         |         |                       |           |           |         |        |
| Similar name   | 46 (76.7)             | 12 (20)   | 2 (3.3) | 0       | 24 (40)               | 34 (56.7) | 2 (3.3)   | 0       | <0.001 |
| Similar shape  | 38 (63.3)             | 18 (30)   | 4 (6.7) | 0       | 25 (41.7)             | 24 (40)   | 10 (16.7) | 1 (1.7) | <0.005 |
| The place of the medication  | 52 (86.7)             | 7 (11.7)  | 1 (1.7) | 0       | 37 (61.7)             | 20 (33.3) | 3 (5)     | 0       | <0.004 |
| Inappropriate and unsafe prescribed medication   | 52 (86.7)             | 7 (11.7)  | 1 (1.7) | 0       | 46 (76.7)             | 11 (18.3) | 2 (3.3)   | 1 (1.7) | <0.046 |
| 2. Right dose  |                       |           |         |         |                       |           |           |         |        |
| Dose noncompliance with the prescription   | 48 (80)               | 10 (16.7) | 2 (3.3) | 0       | 23 (38.3)             | 29 (48.3) | 7 (11.7)  | 1 (1.7) | <0.001 |
| Dose noncompliance with the age and weight of the child  | 42 (70)               | 13 (21.7) | 2 (3.3) | 3 (5)   | 19 (31.1.7)           | 25 (41.7) | 11 (18.3) | 5 (8.3) | <0.001 |
| 3. Right route   |                       |           |         |         |                       |           |           |         |        |
| Oral administration instead of intravenous   | 55 (91.6)             | 4 (6.7)   | 1 (1.7) | 0       | 49 (81.7)             | 9 (15)    | 2 (3.3)   | 0       | <0.11  |
| Intravenous administration Instead of oral administration  | 54 (90)               | 5 (8.3)   | 1 (1.7) | 0       | 51 (85)               | 9 (15)    | 0         | 0       | 59.0   |
| Intramuscular injection instead of intravenous and vice versa  | 55 (91.6)             | 5 (8.3)   | 0       | 0       | 51 (85)               | 9 (15)    | 0         | 0       | 0.25   |
| 4. Right patient   |                       |           |         |         |                       |           |           |         |        |
| Prescribe medicine to another patient  | 46 (76.7)             | 13 (21.7) | 1 (1.7) | 0       | 22 (53.3)             | 23 (38.3) | 4 (6.7)   | 1 (1.7) | <0.001 |
| Lack of attention to the patient identification bracelet   | 52 (86.7)             | 5 (8.3)   | 2 (3.3) | 1 (1.7) | 25 (41.7)             | 26 (43.3) | 7 (11.7)  | 2 (3.3) |        |
| Administration of medicine based on bed number   | 46 (76.7)             | 13 (21.7) | 1 (1.7) | 0       | 32 (53.3)             | 19 (31.7) | 8 (13.3)  | 1 (1.7) |        |
| 5. Right prescription  |                       |           |         |         |                       |           |           |         |        |
| No use of proper solvent   | 49 (81.7)             | 11 (18.3) | 0       | 0       | 40 (66.7)             | 19 (31.7) | 1 (1.7)   | 0       |        |
| Lack of observing proper infusion rate   | 41 (68.3)             | 16 (26.7) | 3 (5)   | 0       | 24 (40)               | 27 (45)   | 8 (13.3)  | 1 (1.7) |        |
| Lack of attention to drug interactions   | 44 (73.3)             | 13 (21.7) | 3 (5)   | 0       | 34 (56.7)             | 23 (38.3) | 3 (5)     | 0       |        |
| Lack of attention to medication precautions  | 50 (83.3)             | 10 (16.7) | 0       | 0       | 45 (75)               | 13 (21.7) | 2 (3.3)   | 0       |        |
| No use of micro if needed  | 52 (86.7)             | 8 (13.3)  | 0       | 0       | 44 (73.3)             | 15 (25)   | 1 (1.7)   | 2.32    |        |
| No use of infusion pump/syringe pump   | 54 (90)               | 6 (10)    | 0       | 0       | 41 (68.3)             | 16 (26.7) | 3 (5)     | 0       |        |
| No hand hygiene  | 46 (76.2)             | 12 (20.2) | 1 (1.7) | 1 (1.7) | 28 (46.7)             | 18 (30)   | 12 (20)   | 2 (3.3) |        |
| No use of alcohol to disinfect the vial  | 49 (81.7)             | 6 (10)    | 2 (3.3) | 3 (5)   | 24 (40)               | 15 (25)   | 18 (30)   | 3 (5)   |        |
| No attention to medication expiration rate   | 53 (88.3)             | 4 (6.7)   | 3 (5)   | 0       | 46 (76.2)             | 12 (20.2) | 2 (3.6)   | 0       |        |
| 6. Right time  |                       |           |         |         |                       |           |           |         |        |
| No observation of the correct time of medication (before or after meals, etc.)   | 45 (75)               | 14 (23.3) | 1 (1.7) | 0       | 39 (65)               | 18 (30)   | 3 (5)     | 0       |        |
| No give of the medicine on due time  | 40 (66.7)             | 20 (33.3) | 0       | 0       | 19 (31.7)             | 31 (60)   | 5 (8.3)   | 0       |        |
| Not giving the drug in proportion to the number of times it is prescribed (e.g., every 8 h every 12 h)                 | 52 (86.7)             | 8 (13.3)  | 0       | 0       | 37 (61.7)             | 22 (36.7) | 1 (1.7)   | 0       |        |
| 7. Right record  |                       |           |         |         |                       |           |           |         |        |
| Noncompliance of the recorded medication with the medication order   | 49 (81.7)             | 10 (16.7) | 1 (1.7) | 0       | 37 (61.7)             | 19 (31.7) | 4 (6.7)   | 0       |        |
| Noncompliance of the recorded medicine shape with the medication order (ampoules instead of tablets or vice versa)     | 52 (86.7)             | 6 (10)    | 2 (3.3) | 0       | 40 (66.7)             | 19 (31.7) | 1 (1.7)   | 0       |        |
| Noncompliance of the medicine dose recorded in the cortex with the medication order                                    | 52 (86.7)             | 7 (11.7)  | 1 (1.7) | 0       | 32 (53.3)             | 25 (41.7) | 3 (5)     | 0       |        |
| Noncompliance of the frequency of the medicine prescribed in the cortex with the medication order (2 or 3 times a day) | 49 (81.7)             | 11 (18.3) | 0       | 0       | 40 (66.7)             | 16 (26.7) | 4 (6.7)   | 0       |        |
| Nonchange of a medicine's new shape in the cortex after being prescribed   | 46 (76.7)             | 14 (23.3) | 0       | 0       | 35 (58.3)             | 24 (40)   | 1 (1.7)   | 0       | 0.03   |
| No record of the prescribed medication   | 46 (76.7)             | 14 (23.3) | 0       | 0       | 25 (41.7)             | 33 (55)   | 2 (3.3)   | 0       | <0.001 |
| No prescription of the recorded medication   | 54 (90)               | 6 (10)    | 0       | 0       | 37 (61.7)             | 1 (31.7)  | 4 (6.7)   | 0       | <0.001 |
| 8. Right response to treatment   |                       |           |         |         |                       |           |           |         |        |
| No assessment of the patient's response to treatment   | 48 (80)               | 9 (15)    | 3 (5)   | 0       | 37 (61.7)             | 15 (25)   | 8 (13.3)  | 0       | 0.18   |

**Table 2: Comparison of the mean total score for nurses' medication error reporting**

| Eight principles            | Before intervention | After intervention | Paired t-test |        |
|-----------------------------|---------------------|--------------------|---------------|--------|
|                             |                     |                    | t-test        | P      |
| Right medication            | 8.33 (1.53)         | 17.92 (1.97)       | 4.08          | <0.001 |
| Right dose                  | 11.11 (2.16)        | 30 (2.56)          | 6.40          | <0.001 |
| Right route                 | 3.33 (1.13)         | 5.74 (1.43)        | 1.40          | 0.16   |
| Right patient               | 7.96 (1.75)         | 21.85 (2.41)       | 6.49          | <0.001 |
| Right prescription          | 7.53 (1.24)         | 17.04 (1.64)       | 6.14          | <0.001 |
| Right time                  | 8.15 (1.23)         | 17.41 (1.70)       | 4.60          | <0.001 |
| Right record                | 6.03 (0.99)         | 15.32 (1.46)       | 5.91          | <0.001 |
| Right response to treatment | 8.33 (2.33)         | 17.22 (3.12)       | 2.46          | 0.02   |
| Total score                 | 7.35 (2.43)         | 17.04 (5.59)       | 8.03          | <0.001 |

classes for interns and medical students. Furthermore, the matron supports and emphasizes that there should not be any fear and anxiety increased by reporting the rate confirmed by statistical results. In this study, there was only one group with a small sample size. Therefore, studies with larger samples are needed to produce more evidence regarding the effects of this motivation program.

## Conclusions

The results of this study showed that the combination of training and motivational program improves reporting of medication errors in nurses. Given that error reporting is known as the basis for maintaining and enhancing patient safety and medication error reporting can be used as an opportunity to learn and prevent the repetition of errors, it is recommended that health care managers increase their quality of medical care and improve the safety of hospitalized children by ongoing training and encouragement of personnel to report errors. It is suggested to use other motivational strategies to improve medication error reporting. Moreover, it will be held the same training course for other personnel in the hospital and professions such as interns and medical students.

## Acknowledgment

We would like to express our gratitude and appreciation to all of the nurses who participated in this study, as well as the managers of the selected hospital of Isfahan University of Medical Sciences who helped us with this research.

## Financial support and sponsorship

This study was financially supported by the Isfahan University of Medical Sciences.

## Conflicts of interest

There are no conflicts of interest.

## References

- Salmani N, Hasanvand S. Evaluation of the frequency and type of medication prescribing errors in the NICU of hospitals in Yazd. *Hayat* 2016;21:53-64.
- Farzi S, Irajpour A, Saghaei M, Ravaghi H. Weak professional interactions as main cause of medication errors in intensive care units in Iran. *Iran Red Crescent Med J* 2017;19:e14946.
- Farzi S, Farzi S, Alimohammadi N, Moladoost A. Medication errors by the intensive care units nurses and the preventive strategies. *J Anesthe Pain* 2015;6:33-45.
- Ghorbanpour Diz M, Mohammad Khan Kermanshahi S, Sedagat M. Evaluation of self-report of medication errors and its barriers in pediatric wards. *Iran J Nurs Pedi Nurs* 2016;3:53-59.
- Farzi S, Saghaei M, Irajpour A, Ravaghi H. The most frequent and important events that threaten patient safety in intensive care units from the perspective of health-care professionals'. *J Res Med Sci* 2018;23:104.
- Alomari A, Sheppard-Law S, Lewis J, Wilson V. Effectiveness of Clinical Nurse's interventions in reducing medication errors in a pediatric ward. *J Clin Nurs* 2020;12:19-21.
- Hartwig SC, Denger SD, Schneider PJ. Severity-indexed, incident report-based medication error-reporting program. *Am J Hosp Pharm* 1991;48:2611-16.
- Saif AA. Educational psychology: Psychology of Learning and Instruction. Tehran: Agah Publishers; 2015. p. 67-70.
- Negarandeh R, Dehghan-Nayeri N, Ghasemi E. Motivating factors among Iranian nurses. *Iran J Nurs Midwifery Res* 2015;20:436-41.
- Farsani L, Rezaei Farsani M, Farokhpour M. The rate, type and factors affecting drug errors from the perspective of nurses working in intensive care and emergency unite of educational hospital of Shahrekord University of medical sciences. *Medi J Develo Strat Medi Edu* 2017;4:61-71.
- Mozafari M, Borji M. Evaluating the rate of nurses errors in administration of medication orders in general departments of hospitals in the City of Ilam. *Iran Nurs Sci Assoc* 2017;12:21-6.
- Tabatabaee S, GHamari M, SHarifi T, Kalhor R, Asadi M. Checking the rate and type of nurses medication errors: A case study in a private hospital. *J Hosp* 2015;14:119-26.
- Farzi S, Farzi S, Taheri S, Ehsani M, Moladoost A. Perspective of nurses toward the patient safety culture in neonatal intensive care units. *IJN* 2017;8:89-94.
- Marquis BL, Huston CJ. Leadership roles and management functions. In: *Nursing Theory and Application* Wolters Kluwer. Philadelphia: Lippincott Williams & Wilkins Inc.; 2020.
- Toloei M, Nayeri Dehghanayeri N, Faghihzadeh S, AslSadooghi Asl A. The nurses motivating factors in relation to patient training. *Hayat* 2006;12:43-51.
- Jodat S, Farajzadeh Z, Saadatjoo SA. A study of job motivation nurses of working in valiasr hospital of Birjand in 2013. *Mod Care Nurs Sci Q Birjand Nurs Midwifery Fac* 2014;10:296-304.
- Irajpour A, Farzi S, Saghaei M, Ravaghi H. Effect of interprofessional education of medication safety program on the medication error of physicians and nurses in the intensive care units. *J Educ Health Promot* 2019;8:196.

18. Cox N, Christensen S, Mayer J, Como N, Tracy M, Sheffield C, Turner K. Implementation of an interprofessional learner team-based medication reconciliation and review in an Internal Medicine-Pediatrics resident continuity clinic. *J Interp Educ Prac* 2020;19:100316.
19. Berry JC, Davis JT, Bartman T, Hafer CC, Lieb LM, Khan N, Brill R. Improved safety culture and teamwork climate are associated with decreases in patient harm and hospital mortality across a hospital program. *J Patient Safety* 2020;16:130-6.