



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

## Can digital contact tracing make up for lost time?



Published Online  
July 16, 2020  
[https://doi.org/10.1016/S2468-2667\(20\)30160-2](https://doi.org/10.1016/S2468-2667(20)30160-2)  
See [Articles](#) page e452

Contact tracing is a fundamental public health intervention, and a mainstay in efforts to control and contain severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus responsible for the COVID-19 pandemic. At the time of writing, the pandemic has caused more than 13 million cases and more than 578 000 deaths.<sup>1</sup> Regions with the most successful containment to date have approached the pandemic with integrated measures that include cohesive leadership, effective communication, physical distancing, wearing of face coverings, improvements in the built environment, promotion of hand hygiene, and support for the staff, supplies, and systems needed to care for patients—with testing and contact tracing as cornerstones of the approach. Despite the emergence of some promising therapies<sup>2</sup> and work towards a future vaccine,<sup>3</sup> basic public health approaches remain the best available prevention and control interventions at this time.

Along with efforts to expand conventional contact tracing programmes, there has been an ongoing debate about the value of digital contact tracing, ranging from issues of privacy, questions about efficacy, lower user adoption rates, and concern from some public health experts that mobile apps might distract resources from the core work of conventional contact tracing. Yet, in the face of ongoing challenges in disease control, the question of whether digital technologies can supplement existing efforts is one that we cannot afford to ignore.

In *The Lancet Public Health*, Mirjam Kretzschmar and colleagues<sup>4</sup> model key steps in SARS-CoV-2 testing and contact tracing across a spectrum of scenarios and identify opportunities to maximise the effectiveness of the process in reducing the effective reproductive number of COVID-19. The study is important as initial large-scale physical distancing policies are relaxed and movement of people increases. Research into how and where to best invest in improving systems of contact tracing is essential, as even those areas with low case burdens will face ongoing transmission events and must be prepared to quell outbreaks as they occur. Not surprisingly, the authors conclude that speed is of the essence in testing and isolating: the study finds that keeping the time between symptom onset and

testing and isolation of an index case at 2 days or less is imperative for success in reducing the reproductive number, and that rapid testing of symptomatic people is at least as important as the efficiency of contact tracing. This study adds to the literature on the role of contact tracing in COVID-19 and highlights the need for adequate testing capacity. The authors also suggest a meaningful contribution to contact tracing from mobile apps, which might minimise notification and tracing delays, although they do not consider a hybrid approach combining conventional and mobile app-based contact tracing.

The authors make several assumptions that might blunt the impact of their findings: they assume that index cases are isolated with no further transmission, yet household transmission has been reported as important even when contact tracing was in place;<sup>5</sup> that all traced contacts, regardless of symptoms, are offered testing, yet capacity to test remains an important challenge in many areas;<sup>6</sup> and that those testing negative (once) do not spread infection, yet this might be an over-simplification of the sensitivity of tests and the dynamics of infectiousness.<sup>7</sup> The importance of these assumptions could be tested in future research and modelling efforts, as could an analysis of a hybrid approach where exposure notification is used to support conventional contact tracing rather than replace it, which seems more likely in practice.

A limitation of the study is the lack of detail on the mobile app technology in the model. While the researchers focus on uptake and speed of notifications—two important parameters—there is a lack of discussion of the efficacy of an app in terms of its detector function (ie, the sensitivity and specificity of an app to determine if a contact event has occurred between two users<sup>8</sup>) and its effector function (ie, the effectiveness in contributing to the desired public health actions by the user, such as entering self-quarantine). The conclusion of the researchers,<sup>4</sup> therefore, that “app-based tracing on its own remains more effective than conventional tracing alone, even with 20% coverage, due to its inherent speed” seems premature without a more nuanced discussion of efficacy and of the potential challenges and harms of digital approaches. This is not to claim that mobile apps are lacking in promise, but they do remain

unproven as a public health intervention. Therefore, as jurisdictions around the world roll out exposure notification apps, there are crucial questions that must be investigated to understand the efficacy of these apps and to make adjustments necessary to build user trust and adoption, if they are to make a contribution to pandemic response.

First, how well do smartphones measure proximity? In other words, what is the effectiveness of the detector function and how many false alarms might be expected for each true contact detected?

Second, how will mobile apps integrate with overall contact tracing programmes? Good contact tracing offers not just an epidemiological intervention—quarantining enough individuals to reduce the reproductive number—but also a human one, that investigates outbreaks and understands linkages, and that recognises and addresses the challenges inherent in quarantine and isolation by providing a variety of supports. In our experience, success at this challenging endeavour requires public health workers as human beings to connect with a person, to build trust on a human level. These vital dynamics are not captured in epidemiological models, nor can we expect that notifications provided by a mobile app will fill the place of the detective work and supportive human interventions at the core of contact tracing.

Third, what factors will encourage users to trust the privacy and security properties of mobile apps? Current adoption rates are low in every jurisdiction where apps have been deployed, with most peaking at download rates of about 20% of the population, and little data available about actual usage levels, which are likely to be lower.<sup>9</sup> Mobile app user behaviour depends on a subtle trust–benefit ratio calculation by users that is challenging to predict in advance.<sup>10</sup> What is behind the public’s decision to use or avoid these apps? Do they have privacy or security concerns or question the benefit of the service? Do they trust public health authorities with their data and do they trust the authorities’ pandemic response?

Fourth, how will mobile apps affect health equity? To be successful in addressing the pandemic, any contact tracing system—conventional or digital—should be evaluated within a health equity framework to avoid perpetuating the deep disparities that the global pandemic has so glaringly exposed.

As contact tracing remains a crucial component of the COVID-19 response, mobile apps offer promise, especially when considering the speed and scale required for tracing to be effective—as highlighted in Kretzschmar and colleagues’ study.<sup>4</sup> However, understanding the potential impact of apps as part of a comprehensive integrated approach requires more evaluation of their use in real life and multidisciplinary engagement of technologists, epidemiologists, public health experts, and the public.

DJW reports grants from the MIT-IBM Watson Artificial Intelligence Laboratory and the US Centers for Disease Control and Prevention. LCI reports grants from The Bill and Melinda Gates Foundation and The National Institutes of Allergy and Infectious Diseases, outside of the submitted work. DJW is co-principal investigator and LCI is senior medical advisor of the Private Automated Contact Tracing initiative, which is a collaboration led by the MIT Computer Science and Artificial Intelligence Laboratory, the MIT Internet Policy Research Initiative, Massachusetts General Hospital Center for Global Health, and MIT Lincoln Laboratory. It includes close collaborators from Boston University, Brown University, Carnegie Mellon University, the MIT Media Lab, and the Weizmann Institute.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

Louise C Ivers, \*Daniel J Weitzner  
weitzner@mit.edu

Center for Global Health, Massachusetts General Hospital, Boston, MA, USA (LCI); Harvard Medical School, Boston, MA, USA (LCI); and Computer Science and Artificial Intelligence Lab, Massachusetts Institute of Technology, Cambridge, MA 02139, USA (DJW)

- 1 Johns Hopkins University of Medicine. COVID-19 dashboard by the Center for Systems Science and Engineering at Johns Hopkins University. <https://coronavirus.jhu.edu/map.html> (accessed July 15, 2020).
- 2 Sanders JM, Monogue ML, Jodlowski TZ, Cutrell JB. Pharmacologic treatments for coronavirus disease 2019 (COVID-19): a review. *JAMA* 2020; **323**: 1824–36.
- 3 Lurie N, Saville, Hatchett R, Halton J. Developing Covid-19 vaccines at pandemic speed. *N Engl J Med* 2020; **382**: 1969–73.
- 4 Kretzschmar ME, Rozhnova G, Bootsma M, van Boven M, van de Wijgert J, Bonten M. Impact of delays on effectiveness of contact tracing strategies for COVID-19: a modelling study. *Lancet Public Health* 2020; published online July 16. [https://doi.org/10.1016/S2468-2667\(20\)30157-2](https://doi.org/10.1016/S2468-2667(20)30157-2).
- 5 Q Bi, Y Wu, S Mei, et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis* 2020; published online April 27. [https://doi.org/10.1016/S1473-3099\(20\)30287-5](https://doi.org/10.1016/S1473-3099(20)30287-5).
- 6 Mervosh S, Fernandez M. Months into virus crisis, US cities still lack testing capacity. *New York Times*. July 6, 2020. <https://www.nytimes.com/2020/07/06/us/coronavirus-test-shortage.html> (accessed July 11, 2020).
- 7 Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in false-negative rate of reverse transcriptase polymerase chain reaction-based SARS-CoV-2 tests by time since exposure. *Ann Intern Med* 2020; published online May 13. <https://doi.org/10.7326/M20-1495>.
- 8 Hatke GF, Montanari M, Appadwedula S, et al. Using Bluetooth Low Energy (BLE) signal strength estimation to facilitate contact tracing for COVID-19. *arXiv* 2020: 2006.15711 (preprint).
- 9 Singer N. Virus-tracing apps are rife with problems. Governments are rushing to fix them. *New York Times*. July 8, 2020. <https://www.nytimes.com/2020/07/08/technology/virus-tracing-apps-privacy.html> (accessed July 12, 2020).
- 10 Shih F, Liccardi I, Weitzner D. Privacy tipping points in smartphones privacy preferences. In Proceedings of the 33rd Annual ACM Conference on Human Factors in Computing Systems. New York, NY: Association for Computing Machinery, 2015: 807–16.