



Natural Killer Cell-Derived Exosomal miR-3607-3p Inhibits Pancreatic Cancer Progression by Targeting IL-26

Hongwei Sun^{1†}, Keqing Shi^{2†}, Kai Qi^{3†}, Hongru Kong¹, Jie Zhang¹, Shengjie Dai¹, Wen Ye¹, Tuo Deng¹, Qiye He^{4,5} and Mengtao Zhou^{6*}

¹ Department of Hepatobiliary Surgery, Key Laboratory of Diagnosis and Treatment of Severe Hepato-Pancreatic Diseases of Zhejiang Province, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China, ² Key Laboratory of Diagnosis and Treatment of Severe Hepato-Pancreatic Diseases of Zhejiang Province, Center of Precision Medicine, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China, ³ Shanghai Institute of Nutrition and Health, Chinese Academy of Sciences, Shanghai, China, ⁴ Singlera Genomics Inc., San Diego, CA, United States, ⁵ Singlera Genomics (Shanghai) Ltd., Shanghai, China, ⁶ Key Laboratory of Diagnosis and Treatment of Severe Hepato-Pancreatic Diseases of Zhejiang Province, Precision Medical Center Laboratory, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China

OPEN ACCESS

Edited by:

José Mordoh,
Leloir Institute Foundation
(FIL), Argentina

Reviewed by:

Mallikarjun Bidarimath,
Cornell University, United States
Bruno Costa-Silva,
Champalimaud Foundation, Portugal

*Correspondence:

Mengtao Zhou
qianazi@yeah.net

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cancer Immunity and Immunotherapy,
a section of the journal
Frontiers in Immunology

Received: 27 May 2019

Accepted: 15 November 2019

Published: 11 December 2019

Citation:

Sun H, Shi K, Qi K, Kong H, Zhang J,
Dai S, Ye W, Deng T, He Q and
Zhou M (2019) Natural Killer
Cell-Derived Exosomal miR-3607-3p
Inhibits Pancreatic Cancer
Progression by Targeting IL-26.
Front. Immunol. 10:2819.
doi: 10.3389/fimmu.2019.02819

Increasing evidences have suggested that natural killer (NK) cells in the tumor microenvironment are involved in the regulation of cancer development. However, the potential biological roles and regulatory mechanisms of NK cells in pancreatic cancer (PC) remain unclear. Co-culture system of NK cells with PC cells is used to test the ability of cancer cell proliferation, migration and invasion both *in vitro* and *in vivo*. And tail vein intravenous transfer was used to test metastasis *in vivo*. Meanwhile, extracellular vesicles (EVs) were separated and examined. Furthermore, reporter assay and Biotin-RNA pull down assay were performed to verify the interaction between molecules. NK cells can inhibit the malignant transformation of co-cultured PC cells both *in vivo* and *in vitro*, which requires miR-3607-3p. miR-3607-3p is found enriched in the EVs of NK cells and transmitted to PC cells, and low level of miR-3607-3p predicts poor prognosis in PC patients. It can also inhibit proliferation, migration and invasion of PC cells *in vitro*. Importantly, IL-26 is found to be a direct target of miR-3607-3p in PC cells. miR-3607-3p enriched in EVs derived from NK cells can inhibit the malignant transformation of PC probably through directly targeting of IL-26.

Keywords: natural killer, pancreatic cancer, extracellular vesicles, miR-3607-3p, IL-26

INTRODUCTION

Pancreatic cancer (PC) is one of the most lethal human malignancies with poor prognosis (1). Due to the lack of early symptoms and diagnose test, PC is usually found at an incurable advanced stage (2, 3). Presently, the treatment of PC mainly relies on surgical resection and chemotherapy (4). Unfortunately, local deterioration and frequent occurrence of distant metastasis results in poor survival rate of PC patients, with an overall 1-year survival rate at 20% and 5-year survival rate <5% (5). Thus, it is critical to explore the mechanism of malignant transformation of PC, identify novel diagnostic markers and therapeutic targets in PC, which might provide new treatment strategy for PC patients.

Studies have shown that tumor cells can conduct a variety of cross-talks within the tumor microenvironment via chemokines, cytokines, and immune cells to regulate the fate of tumor progression (6, 7). Natural killer (NK) cells, a subset of lymphocytes that play an important role in the rapid innate immune response to viral infection and tumor development, are important components in the tumor microenvironment (8, 9). Increasing studies have shown that NK cells play an important role in inhibiting malignant transformation of tumors, however, the detailed mechanisms are still illusive (10–12). Exosomes are cell-derived vesicles (50–100 nm) that can enclose protein, lipid, mRNA, and microRNA (miRNA) (13–15). Multiple studies have shown that a variety of cells can secrete exosomes that might regulate the development of tumor in the tumor microenvironment including NK cells (16–18).

Here we investigated the function of miR-3607-3p, which was enclosed in NK derived EVs, in pancreatic cancer. Our results suggested that miR-3607-3p enriched in EVs of NK cells, could suppress pancreatic cancer development and malignant transformation by directly targeting IL-26.

MATERIALS AND METHODS

Patients and Tissue Samples

Forty primary pancreatic cancer (PC) tissues and 40 normal were collected at the First Affiliated Hospital of Wenzhou Medical University from 01/03/2017 to 03/08/2018. All specimens from resection surgery were frozen and stored in liquid nitrogen. This study was performed with the approval of the Ethic and Research Committees of the First Affiliated Hospital of Wenzhou Medical University and these studies was conducted in accordance with the Declaration of Helsinki Principles. Informed written consent was obtained from all subjects.

Human NK Cell Isolation

Human NK cells were isolated as described before using NK cell isolation kit (Miltenyi Biotech, Bergisch Gladbach, Germany) according to the manufacturer's instruction (19). The PBMC were collected according to the protocols approved by the First Affiliated Hospital of Wenzhou Medical University.

Cell Line

Human PC cells (Mia PaCa-2 and PANC-1) were purchased from the American Type Culture Collection (Rockville, USA). The cells were cultured in RPMI 1640 or DMEM (Gibco) supplemented with 10% fetal bovine serum (FBS), 100 U/mL penicillin, and 100 mg/mL streptomycin at 37°C under 5% CO₂ in a humidified chamber.

Transfection

MiR-3607-3p mimics, inhibitor, and negative control were purchased from GenePharma (Shanghai, China). The transfection was performed using Lipofectamine 2000 (Invitrogen, Waltham, MA USA) following the manufacturer's

instructions. The knockdown efficiency at the mRNA level was assessed using qRT-PCR assay.

CCK-8 Cell Proliferation Assay

PC cells were seeded in 96-well plates at a density of 5×10^3 cells/well. At the indicated time points, viable cells were examined by cell counting kit-8 (CCK-8) according the manufacturer's instructions.

Migration and Invasion Assay

Cell migration and invasion ability were measured using transwell chambers (24-23ll insert, 8 μm, Millipore, Billerica, MA, USA). The chambers were coated with Matrigel Matrix (Sigma, St. Louis, MO). PC cells were seeded at a density of 5×10^4 cells/well in upper chamber with 200 μL serum free medium. The lower chamber was filled with 600 μL of medium containing 10% FBS. After incubation for 48 h, the number of cells passing through the bottom membrane of the chamber was counted under a microscope.

Colony Formation Assay

PC cells transfected with NC-miRNA or miR-3607-3p were seeded in a 24-well plate at a density of 5×10^4 cells/well and cultured for 24 h. Then, the cells were collected and seeded (1,000–1,500 cells/well) in a dish for 10 days. Surviving colonies were fixed, stained with 5% gentian violet (Sigma) and counted. The experiment was carried out in triplicate wells for three times.

Immunoblotting

Total protein was extracted from the PC cells using RIPA buffer (Invitrogen) according to the manufacturer's protocol. Equivalent amounts of proteins from each sample were subjected to SDS-PAGE electrophoresis and then transferred to a PVDF membrane (Millipore), blocked in 5% fat-free milk for 1 h at room temperature, incubated with specific primary antibodies overnight at 4°C with gentle shaking, and followed by detection with enhanced chemiluminescence system (Pierce, Waltham, MA). Primary antibodies were as follows: CD63, TSG101, IL-26, Actin (Abcam, Cambridge, MA).

RNA Isolation and qPCR

Total RNA from PC tissues and cell lines was harvested using the TRIzol reagent (Invitrogen) following the manufacturer's instructions. RNAs were reverse-transcribed into cDNAs using the Prime Script RT reagent Kit (TaKaRa, Dalian, China) according to the manufacturer's protocol. MiR-3607-3p RT primer: GTCGTATCCAGTGCAGGGTCCGAGGTATTTCGCAC TGGATACGACCATCA; U6 (internal control) RT primer: GTCGTATCCAGTGCAGGGTCCGAGGTATTTCGCACTG GATACGACAAAATA TGGAA. SYBR Taq (TaKaRa) was used to measure the expression of miR-3607-3p and IL-26. The primers used were described below: miR-3607-3p Forward: ATGACTGTAAACGCTTTCTG, Reverse: GTGCAGGGTCC GAGGT;

U6 Forward: TGC GG GTGCTCGCTTCGGCAGC, Reverse: GTGCAGGGTCCGAGGT; IL-26, Forward: AAGCAACGAT TCCAGAAGACC; Reverse: AAGTCCTCCACAAAGCGTA TTTT; GAPDH, Forward: ACAACTTTGGTATCGTGGAAGG;

Abbreviations: NK, natural killer; PC, pancreatic cancer; miRNA, microRNA.

Reverse: GCCATCACGCCACAGTTTC. Fold changes of miR-3607-3p and IL-26 were calculated by the equation $2^{-\Delta\Delta Ct}$.

Luciferase Reporter Assays

The dual luciferase reporter assay was performed using Dual-Luciferase Reporter Assay System (Promega, Madison, WI, USA). Briefly, luciferase reporter vector containing the wild-type (WT) and mutant (MT) miR-3607-3p binding site in the 3'-UTR of IL-26 were constructed transiently transfected into MIA PaCa-2 or PANC-1 cells, together with miR-3607-3p mimics or NC using Lipofectamine 2000. Primers used for cloning WT IL-26 3'UTR, Forward: CTACTCGAGACCAAGC CAAGTACATT, Reverse: CTAGCGGCCGCGAAGGAA ACCCAATTTA; Mutant IL-26 3'UTR, Forward: GCCAAG TACATTGATTCTCCCCT, Reverse: AGGGGAGAATCA ATGTACTTGGC. Luciferase activity was measured 48 h later. The renilla luciferase activity were normalized to the firefly luciferase activity in the corresponding well.

Biotin-miRNA Pull-Down Assay

Biotin-miRNA pulldown assay was carried out as described before (20); Briefly, PC cells were washed with cold PBS, harvested by a scraper and treated cell lysis buffer. Supernatant was collected and incubated with biotinylated miR-3607-3p or biotinylated control random RNA for 1 h at 30°C. After incubation, Streptavidin MagneSphere Paramagnetic beads (Promega) was added for 3 h at 4°C. The biotin-miRNA/mRNA complex was eluted and washed for mRNA detection.

Extracellular Vesicle Isolation and Characterization

Extracellular vesicle (EV) isolation and characterization were prepared as described before (21). Purified NK cells were cultured in a serum free medium for 24 h. Cell culture supernatant was centrifuged for 5 min at 500 g and 10 min at 1,500 g to eliminate the cells and debris. EVs were pelleted by ultracentrifuge step at 80,000 g for 120 min and washed with PBS. EVs were evaluated and characterized by flow cytometry and Scanning Electron Microscopy (SEM). Briefly, isolated EVs were put on a copper grid coated with 0.1% Formvar in chloroform. The grids were stained with 1% (vol/vol) uranyl acetate in ddH₂O, and thereafter the EVs were examined immediately by SEM.

In vivo Tumor Formation Assay

Four to six weeks old Male BALB/c nude mice were purchased from SLAC (Shanghai, China). The animal care and experimental protocols were approved by the institutional guidelines of the First Affiliated Hospital of Wenzhou Medical University and by the Animal Care and Use Committee of The First Affiliated Hospital of Wenzhou Medical University. 1×10^7 PC cells were resuspended in 200 μ L PBS medium and were subcutaneously injected into the flank of each nude mouse. The tumors were measured weekly and the tumor volume was calculated following the formula length \times width²/2. The mice were killed at 6 weeks after inoculation.

Statistical Analysis

All results are shown as mean \pm standard deviation (SD) and were analyzed using GraphPad Prism 5 (GraphPad Software, USA) from at least three independent experiments. The differences between groups were analyzed using Two-tail Student's *t*-test or ANOVA; Chi-squared test was used to analyze the frequency of lung metastases; Pearson correlation analysis was used to analyze the correlation between expressions of two genes; Kaplan–Meier's analysis was used for prognostic analyses. Data were considered statistically significant when $p < 0.05$.

RESULTS

NK Cells Co-culture Inhibited Tumor Progression of PC Both *in vitro* and *in vivo*


To test the function of NK cells in the tumor microenvironment, we first cultured NK cells together with pancreatic cancer cell line Mia PaCa-2 and PANC-1. CCK-8 assay indicated that the proliferation of Mia PaCa-2 and PANC-1 cells co-cultured with NK cells was significantly suppressed compared with that of cells cultured alone (Figures 1A,B). Colony formation assay, as shown in Figure 1C, showed that cells co-cultured with NK cells had fewer colonies formed. In line with this, after co-culture with NK cells, Mia PaCa-2 and PANC-1 cells had prominent attenuated cell migration and invasive ability (Figures 1D,E). Furthermore, we employed *in vivo* tumor xenotransplantation mouse model. *In situ* tumor-bearing experiments showed that NK cell co-inoculation could inhibit the tumor growth ability of PANC-1 cells *in vivo* (Figures 1F,G). When intravenously transferred *in vivo* through tail vein, PANC-1 cells co-transferred with NK cells significantly inhibited lung metastasis, as demonstrated by *in vivo* luminescence imaging and H&E staining of the lung tissues (Figures 1H,I). In summary, NK cell can significantly inhibit the proliferation and metastasis of pancreatic cancer cells *in vitro* and *in vivo*.


MiR-3607-3p Is Enriched in EVs of NK Cells

To explore the underlying mechanisms by which NK cells inhibit malignant transformation of pancreatic cancer cells, we speculate that NK cells might act on pancreatic cancer cells through EVs. EVs were isolated and identified from the culture medium of NK cells (Figures 2A–C). In addition, the level of miR-3607-3p in NK cells and its EVs was significantly higher than that in Mia PaCa-2 and PANC-1 (Figure 2D). Furthermore, when co-cultured Mia PaCa-2 and PANC-1 with NK cells or NK cells EVs, we found that the levels of miR-3607-3p were significantly up-regulated in co-cultured Mia PaCa-2 and PANC-1 cells (Figures 2E,F). In summary, the results suggest that NK cells might transmit miR-3607-3p to pancreatic cancer cells via secreted EVs.

MiR-3607-3p Is Down-Regulated in PC and Decreased miR-3607-3p Level Predicts Poor Prognosis in PC Patients

Next, miRNA-3607-3p level was analyzed in different PC cell lines (AsPC-1, PANC-1, Capan-2, CFPAC-1, SW1990 and Mia PaCa-2) and a normal human pancreatic ductal cell control





(hTERT-HPNE). As shown in **Figure 3A**, the level of miR-3607-3p was significantly lower in PC cell lines compared with that in control cell line. Consistently, the level of miR-3607-3p was significantly lower in PC tissues compared with that in normal tissues (**Figure 3B**). In addition, PC patients with lymph node metastasis (LNM⁺) further decreased the expression of miR-3607-3p compared to that in lymph node metastasis free (LNM⁻) patients (**Figure 3C**). Kaplan–Meier’s analysis of the correlation between miR-3607-3p expression and the metastasis-free survival of PC patients indicated that low levels of miR-3607-3p were characterized by worse overall survival and metastasis-free survival rate (**Figures 3D,E**). We also revealed the similar expression pattern of miR-3607-3p in plasma exosomes in PC patients and plasma exosomal miR-3607-3p expression in PC patients with LNM⁺ was significantly lower than that in LNM⁻ PC patients (**Figures 4A,B**). Our results indicate that miR-3607-3p acts as a tumor suppressor in PC and is likely to be involved in tumor metastasis.



MiR-3607-3p Suppresses Proliferation, Migration, and Invasion of PC Cells

To investigate the function of miR-3607-3p in pancreatic cancer cells, miR-3607-3p mimics was transfected to overexpress miR-3607-3p in PC cell line Mia PaCa-2 and PANC-1. As shown

in **Figures 5A–D**, CCK-8 assay and colony formation assays demonstrated that overexpression of miR-3607-3p inhibited the proliferation of Mia PaCa-2 and PANC-1 *in vitro*; Transwell assay results showed that overexpression of miR-3607-3p inhibited the migration and invasion of pancreatic cancer cells *in vitro* (**Figures 5E, 4F**). Interestingly, NK cells extracellular vesicles (NK EVs) treatment was found to be able to inhibit cell viability (**Figures S1A,B**), proliferation (**Figure S1C**), migration (**Figures S1D–F**) and inhibited IL-26 production (**Figure S1G**) in both Mia PaCa-2 and PANC-1 cells. In conclusion, miR-3607-3p inhibits the malignant transformation of pancreatic cancer cells.


IL-26 Is a Direct Target of miR-3607-3p in PC Cells

To explore the mechanism how miR-3607-3p regulates malignant transformation of PC, we performed bioinformatics analysis and Targetscan (http://www.targetscan.org/vert_71/) was used to predict potential downstream targets of miR-3607-3p. IL-26 was predicted as a potential miR-3607-3p target binding to the IL-26 3’UTR (**Figure 6A**). Luciferase reporter vector containing IL-26 3’ UTR (WT) or IL-26 mutated 3’ UTR (MT) was cotransfected with miRNA control or miR-3607-3p mimics into MIA PaCa-2 and PANC-1 cells. The dual luciferase



reporter assay demonstrated that miR-3607-3p bound to the WT IL-26 3'UTR to inhibit the luciferase activity, but not to the mutant IL-26 3'UTR (Figures 5C, 6B). Furthermore, using the biotin-labeled RNA pull-down technique, we found that miR-3607-3p directly interacted with IL-26 mRNA (Figure 5D).

We confirmed that miR-3607-3p inhibited IL-26 expression at both mRNA and protein levels while miR-3607-3p inhibitor enhanced IL-26 mRNA and protein levels in MIA PaCa-2 and PANC-1 cells (Figures 6E–H). Finally, we examined the mRNA levels of IL-26 in 40 pancreatic cancer tissues and




40 normal tissues. IL-26 was highly expressed in pancreatic cancer tissues compared with that in normal tissue (Figure 6I). Consistently, IL-26 mRNA level was significantly higher in tissues from LNM⁺ PC patients than that in tissues from LNM⁻ PC patients (Figure 6J). Correlation analysis showed that there was a significant negative correlation between the expression levels of miR-3607-3p and IL-26 in pancreatic cancer tissues. In summary, IL-26 is a direct target of miR-3607-3p in pancreatic cancer cells.

MiR-3607-3p Was Required for NK Cell-Induced Inhibition of PC Progression

To test whether the inhibition of malignant transformation of PC by NK cells depends on miR-3607-3p, we cultured Mia

PaCa-2 or PANC-1 cells together with NK cells transfected with NC-inhibitor or miR-3607-3p-inhibitor, respectively. The miR-3607-3p levels were confirmed by qPCR (Figure 7A). The results showed that Mia PaCa-2 and PANC-1 cells co-cultured with NK cells transfected with miR-3607-3p-inhibitor had similar levels of cell proliferation, colony formation and cell migration/invasion compared with that of cells co-cultured with only NC-inhibitor (Figures 7B–F). Inhibition of cell proliferation, migration and invasion of Mia PaCa-2 or PANC-1 cells co-cultured with NK cells transfected with NC-inhibitor was significantly restored in cells co-cultured with NK cells transfected with miR-3607-3p-inhibitor (Figures 7B–F). Thus, NK cells inhibited malignant transformation of pancreatic cancer cells at least partially depending on miR-3607-3p.






FIGURE 7 | MiR-3607-3p was required for NK cell-induced inhibition of pancreatic cancer progression. **(A)** qRT-PCR detection of miR-3607-3p in Mia PaCa-2 or PANC-1 co-cultured with only NC-inhibitor (NK cells-/C-inhibitor), NC-inhibitor transfected NK cells (NK cells+/NC-inhibitor) or miR-3607-3p-inhibitor transfected NK cells (NK cells+/miR-3607-3p-inhibitor). **(B,C)** CCK-8 assay showed cell viability of Mia PaCa-2 and PANC-1 cells co-cultured with only NC-inhibitor (NK cells-/NC-inhibitor), NC-inhibitor transfected NK cells (NK cells+/NC-inhibitor) or miR-3607-3p-inhibitor transfected NK cells (NK cells+/miR-3607-3p-inhibitor). **(D)** Colony formation assay showed cell proliferation of Mia PaCa-2 and PANC-1 cells co-cultured with only NC-inhibitor (NK cells-/NC-inhibitor), NC-inhibitor transfected NK cells (NK cells+/NC-inhibitor) or miR-3607-3p-inhibitor transfected NK cells (NK cells+/miR-3607-3p-inhibitor). **(E,F)** Transwell assays showed cell migratory and invasive ability of Mia PaCa-2 and PANC-1 cells co-cultured with only NC-inhibitor (NK cells-/NC-inhibitor), NC-inhibitor transfected NK cells (NK cells+/NC-inhibitor) or miR-3607-3p-inhibitor transfected NK cells (NK cells+/miR-3607-3p-inhibitor). The data represent the mean \pm SD from three independent experiments. $**P < 0.01$; $***P < 0.001$. Two-way ANOVA for **(B,C)**, one-way ANOVA for others.

DISCUSSION

Mounting evidences have shown that exosomal miRNA are involved in cancer biology and clinical applications (14, 22). By using *in vitro* cell co-culture and *in vivo* tumor-bearing experiments, we described a previously un-identified miR-3607-3p which is enriched in EVs derived from NK cells, suppressing PC development and

malignant transformation. Furthermore, we showed that PC patients had low level of miR-3607-3p and decreased miR-3607-3p level could predict poor prognosis. Most importantly, IL-26 was demonstrated as a direct target candidate of miR-3607-3p. Overall, these results help us to further understand the function of miR-3607-3p in PC and provide a potential novel diagnostic and therapeutic target for PC.

NK cells display critical antitumor properties as they respond rapidly to metastatic and malignancies (23). In PC, NK cell activation and function are compromised due to the extensive crosstalk between NK cell and tumor microenvironment, and different strategies are employed to immunomodulate NK cells in anti-tumor therapies (24). Studies have shown that exosomes derived from NK cells exert anti-tumor effect in different cancers, such as melanoma and neuroblastoma (18, 21). Here we showed that co-cultured NK cells inhibited tumor progression of PC both *in vitro* and *in vivo* (Figure 1). However, more effects are needed to distinguish the human NK cells from the endogenous mouse NK cells and to verify the transplanted human NK cells were in the metastasized region of the lung. To understand how NK cell co-culture affects PC progression, we investigated the EVs derived from NK cell. mRNA and microRNAs have been identified in EVs, which could be transmitted to neighboring cells and function in recipient cells (25). Potential target microRNAs were screened and miR-3607-3p was found to be enriched in NK secreted EVs (Figure 2).

miR-3607-3p, a recent defined novel miRNA, has been demonstrated to suppress non-small cell lung cancer development (26). However, other studies have suggested that miR-3607 might promote cancer development and metastasis (27, 28). To explore the function of miR-3607-3p in PC, we examined the levels of miR-3607-3p in PC patients. Our results indicate that miR-3607-3p is down-regulated in PC and decreased miR-3607-3p level predicts poor prognosis in PC patients (Figure 3). Furthermore, the function of miR-3607-3p was confirmed *in vitro* as it could suppress PC cell proliferation, migration and invasion (Figure 4). Using bioinformatics analysis, we predicted IL-26 as a potential miR-3607-3p target and confirmed that using luciferase reporter assay and biotin-miRNA pull-down assay (Figures 5A–D). Consistently, IL-26 expression was much higher in LNM⁺ PC patients and there was a significant negative correlation between the expression levels of miR-3607-3p and IL-26 in pancreatic cancer tissues (Figures 5I–K).

IL-26 is a member of the IL-10 cytokine family and often co-expressed with IL-22 by activated T cells, especially Th17 cells (29). However, the function of IL-26 in human tumor remains largely unknown. Wei You et al. reported that IL-26 expression was enhanced in human gastric cancer cells and enhanced IL-26 facilitated gastric cancer cell proliferation and

survival by regulating STAT1/STAT3 signaling (30). Here we also found increased IL-26 levels in PC patients compared with that in control group. Intriguingly, IL-26 level was further enhanced in LNM⁺ PC patient compared with that in LNM⁻ PC patient (Figures 5I,J), suggesting a role in PC metastasis. However, the downstream signaling of IL-26 in PC is unclear, and whether IL-26 could be used a diagnose marker in PC still need further studies.

CONCLUSIONS

In conclusion, our findings suggest that miR-3607-3p enriched in EVs derived from NK cells inhibits the malignant transformation of PC probably through direct targeting of IL-26, implicating that targeting tumor microenvironment and related molecular cross-talk could be a new strategy of PC therapy.

DATA AVAILABILITY STATEMENT

The datasets analyzed in this article are not publicly available. Requests to access the datasets should be directed to MZ, qianazi@yeah.net.

ETHICS STATEMENT

The animal study was reviewed and approved by Ethic and Research Committees of the First Affiliated Hospital of Wenzhou Medical University.

AUTHOR CONTRIBUTIONS

HS, KS, KQ, HK, JZ, SD, WY, TD, and QH: conception and design. HS, KS, KQ, and HK: collection and assembly of data. HS, and MZ: manuscript writing. HS, KS, KQ, HK, JZ, SD, WY, TD, QH, and MZ: final approval of manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2019.02819/full#supplementary-material>

REFERENCES

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin.* (2011) 61:69–90. doi: 10.3322/caac.20107
- Edwards BK, Brown ML, Wingo PA, Howe HL, Ward E, Ries LA, et al. Annual report to the nation on the status of cancer, 1975-2002, featuring population-based trends in cancer treatment. *J Natl Cancer Inst.* (2005) 97:1407–27. doi: 10.1093/jnci/dji289
- Arslan AA, Helzlsouer KJ, Kooperberg C, Shu XO, Stepnowski E, Bueno-de-Mesquita HB, et al. Anthropometric measures, body mass index, and pancreatic cancer: a pooled analysis from the Pancreatic Cancer Cohort Consortium (PanScan). *Arch Intern Med.* (2010) 170:791–802. doi: 10.1001/archinternmed.2010.63
- Fogel EL, Shahda S, Sandrasegaran K, DeWitt J, Easler JJ, Agarwal DM, et al. A Multidisciplinary approach to pancreas cancer in 2016: a review. *Am J Gastroenterol.* (2017) 112:537–54. doi: 10.1038/ajg.2016.610
- Tuveson DA, Neoptolemos JP. Understanding metastasis in pancreatic cancer: a call for new clinical approaches. *Cell.* (2012) 148:21–3. doi: 10.1016/j.cell.2011.12.021
- Soysal SD, Tzankov A, Muenst SE. Role of the tumor microenvironment in breast cancer. *Pathobiology.* (2015) 82:142–52. doi: 10.1159/000430499
- Yaacoub K, Pedeux R, Tarte K, Guillaudeux T. Role of the tumor microenvironment in regulating apoptosis and cancer progression. *Cancer Lett.* (2016) 378:150–9. doi: 10.1016/j.canlet.2016.05.012

8. Vivier E, Raulet DH, Moretta A, Caligiuri MA, Zitvogel L, Lanier LL, et al. Innate or adaptive immunity? The example of natural killer cells. *Science*. (2011) 331:44–9. doi: 10.1126/science.1198687
9. Jewett A, Kos J, Fong Y, Ko MW, Safaei T, Perisic Nanut M, et al. NK cells shape pancreatic and oral tumor microenvironments; role in inhibition of tumor growth and metastasis. *Semin Cancer Biol*. (2018) 53:178–88. doi: 10.1016/j.semcancer.2018.08.001
10. Yan Y, Steinherz P, Klingemann HG, Dennig D, Childs BH, McGuirk J, et al. Antileukemia activity of a natural killer cell line against human leukemias. *Clin Cancer Res*. (1998) 4:2859–68.
11. Ljunggren HG, Malmberg KJ. Prospects for the use of NK cells in immunotherapy of human cancer. *Nat Rev Immunol*. (2007) 7:329–39. doi: 10.1038/nri2073
12. Cheng M, Zhang J, Jiang W, Chen Y, Tian Z. Natural killer cell lines in tumor immunotherapy. *Front Med*. (2012) 6:56–66. doi: 10.1007/s11684-012-0177-7
13. Thery C. Exosomes: secreted vesicles and intercellular communications. *PLoS Biol Rep*. (2011) 3:15. doi: 10.3410/B3-15
14. Penforis P, Vallabhaneni KC, Whitt J, Pochampally R. Extracellular vesicles as carriers of microRNA, proteins and lipids in tumor microenvironment. *Int J Cancer*. (2016) 138:14–21. doi: 10.1002/ijc.29417
15. Tkach M, Thery C. Communication by extracellular vesicles: where we are and where we need to go. *Cell*. (2016) 164:1226–32. doi: 10.1016/j.cell.2016.01.043
16. Whiteside TL. Tumor-derived exosomes and their role in cancer progression. *Adv Clin Chem*. (2016) 74:103–41. doi: 10.1016/bs.acc.2015.12.005
17. Zheng P, Chen L, Yuan X, Luo Q, Liu Y, Xie G, et al. Exosomal transfer of tumor-associated macrophage-derived miR-21 confers cisplatin resistance in gastric cancer cells. *J Exp Clin Cancer Res*. (2017) 36:53. doi: 10.1186/s13046-017-0528-y
18. Zhu L, Kalimuthu S, Gangadaran P, Oh JM, Lee HW, Baek SH, et al. Exosomes derived from natural killer cells exert therapeutic effect in melanoma. *Theranostics*. (2017) 7:2732–45. doi: 10.7150/thno.18752
19. Zhang LL, Zhang LF, Shi YB. miR-24 inhibited the killing effect of natural killer cells to colorectal cancer cells by downregulating Paxillin. *Biomed Pharmacother*. (2018) 101:257–63. doi: 10.1016/j.biopha.2018.02.024
20. Yamamoto K, Ito S, Hanafusa H, Shimizu K, Ouchida M. Uncovering direct targets of MiR-19a involved in lung cancer progression. *PLoS ONE*. (2015) 10:e0137887. doi: 10.1371/journal.pone.0137887
21. Shoaee-Hassani A, Hamidieh AA, Behfar M, Mohseni R, Mortazavi-Tabatabaei SA, Asgharzadeh S. NK cell-derived exosomes from NK cells previously exposed to neuroblastoma cells augment the antitumor activity of cytokine-activated NK cells. *J Immunother*. (2017) 40:265–76. doi: 10.1097/CJI.0000000000000179
22. Sun Z, Shi K, Yang S, Liu J, Zhou Q, Wang G, et al. Effect of exosomal miRNA on cancer biology and clinical applications. *Mol Cancer*. (2018) 17:147. doi: 10.1186/s12943-018-0897-7
23. Nicholson SE, Keating N, Belz GT. Natural killer cells and anti-tumor immunity. *Mol Immunol*. (2019) 110:40–7. doi: 10.1016/j.molimm.2017.12.002
24. Van Audenaerde JRM, Roeyen G, Darcy PK, Kershaw MH, Peeters M, Smits ELJ. Natural killer cells and their therapeutic role in pancreatic cancer: a systematic review. *Pharmacol Ther*. (2018) 189:31–44. doi: 10.1016/j.pharmthera.2018.04.003
25. Zhang J, Li S, Li L, Li M, Guo C, Yao J, et al. Exosome and exosomal microRNA: trafficking, sorting, and function. *Genomics Proteomics Bioinformatics*. (2015) 13:17–24. doi: 10.1016/j.gpb.2015.02.001
26. Gao P, Wang H, Yu J, Zhang J, Yang Z, Liu M, et al. miR-3607-3p suppresses non-small cell lung cancer (NSCLC) by targeting TGFBR1 and CCNE2. *PLoS Genet*. (2018) 14:e1007790. doi: 10.1371/journal.pgen.1007790
27. Lin Y, Gu Q, Sun Z, Sheng B, Qi C, Liu B, et al. Upregulation of miR-3607 promotes lung adenocarcinoma proliferation by suppressing APC expression. *Biomed Pharmacother*. (2017) 95:497–503. doi: 10.1016/j.biopha.2017.08.052
28. Wu J, Jiang Z, Chen C, Hu Q, Fu Z, Chen J, et al. CircIRAK3 sponges miR-3607 to facilitate breast cancer metastasis. *Cancer Lett*. (2018) 430:179–92. doi: 10.1016/j.canlet.2018.05.033
29. Donnelly RP, Sheikh F, Dickensheets H, Savan R, Young HA, Walter MR. Interleukin-26: an IL-10-related cytokine produced by Th17 cells. *Cytokine Growth Factor Rev*. (2010) 21:393–401. doi: 10.1016/j.cytogfr.2010.09.001
30. You W, Tang Q, Zhang C, Wu J, Gu C, Wu Z, et al. IL-26 promotes the proliferation and survival of human gastric cancer cells by regulating the balance of STAT1 and STAT3 activation. *PLoS ONE*. (2013) 8:e63588. doi: 10.1371/journal.pone.0063588

Conflict of Interest: QH was employed by company Singlera Genomics Inc.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Sun, Shi, Qi, Kong, Zhang, Dai, Ye, Deng, He and Zhou. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.