

Improving Identification of Cognitive Impairment in Fragility Fracture Patients: Impact of Educational Guidelines on Current Practice

Geriatric Orthopaedic Surgery
& Rehabilitation
Volume 11: 1-7
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2151459320935095
journals.sagepub.com/home/gos



Grace E. M. Kennedy, MB, BCh, BAO¹ ,
Parvathy Mohandas, MB, BCh, BAO², Lesley A. Anderson, PhD³,
Maurice Kennedy, PhD⁴, and Denise S. L. Shirley, PhD¹

Abstract

Introduction: Cognitive impairment can hinder a fracture patient's capacity to consent to surgery and negatively impact their postoperative recovery and rehabilitation. National guidelines recommend screening for cognitive impairment upon admission, and the Abbreviated Mental Test Score (AMTS) is a commonly used tool for this. This project aimed to assess current practice regarding documentation of AMTS among frail fracture patients upon admission and to improve AMTS documentation following a simple intervention. **Methods:** Baseline data were obtained by inpatient chart review throughout November to December 2018 in a district general hospital with emergency fracture services. All patients admitted with a fragility hip fracture and patients over 65 years with any fracture were included. National guidelines and baseline results were then distributed among junior doctors. Following an intervention, further data were collected throughout January to February 2019. **Results:** Preintervention, 40 suitable patients (mean age: 82 years) were identified; 9 (22.0%) of whom had an AMTS recorded upon admission. Among the hip fracture subgroup ($n = 25$), 7 (26.9%) had an AMTS recorded. Postintervention, 39 patients (mean age: 80 years) were identified; 15 (38.5%) of whom had an AMTS recorded. Among the hip fracture subgroup ($n = 30$), 11 (36.7%) had an AMTS recorded. Statistical analysis demonstrated a significant improvement in AMTS documentation both among the overall cohort ($P = .001$) and hip fracture patients ($P = .019$). No significant association was found between AMTS documentation and patient age ($P = .566$), grade of admitting doctor ($P = .058$), or prior cognitive/mental health disorder ($P = .256$). **Discussion:** A small yet significant improvement in AMTS documentation among elderly/hip fracture patients was observed following distribution of educational material. Further work should explore the effect of cognitive impairment on outcomes related to orthopedic injuries beyond hip fractures.

Keywords

fracture, delirium, dementia, capacity, rehabilitation

Submitted January 23, 2020. Revised May 8, 2020. Accepted May 19, 2020.

Introduction

Impaired cognitive function is common among elderly fracture patients and can be seen in a spectrum of forms, ranging from chronic dementia, acute delirium, or acute or chronic confusional states. Lower cognitive status upon admission has been previously associated with poorer functional rehabilitation among hip fracture patients.¹ Patients with dementia have been shown to be at higher risk of hip fracture than their cognitively intact counterparts,² while fracture history may be an independent risk factor for developing dementia among patients aged

¹ Ulster Hospital, South Eastern Health and Social Care Trust, Dundonald, Northern Ireland

² Holywell Hospital, Northern Health and Social Care Trust, Antrim, Northern Ireland

³ School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Northern Ireland

⁴ Faculty of Mathematics and Computing, Open University Belfast, Northern Ireland

Corresponding Author:

Grace E. M. Kennedy, Department of Trauma and Orthopaedics, Ulster Hospital, Upper Newtownards Road, Dundonald, Belfast BT16 1RH, Northern Ireland.

Email: gkennedy15@qub.ac.uk



over 65 years.³ Thus, it is possible that an elderly patient may first be identified as having underlying cognitive impairment upon presentation with a fracture.

Both acute delirium and underlying dementia can affect a patient's capacity to consent for surgery, hinder their post-operative recovery, and affect their discharge destination. Peri-operative delirium affects approximately one-third of patients with hip fractures⁴ and has been associated with longer inpatient stays and higher 1-year mortality rates.⁵ Hip fracture patients with prediagnosed dementia have been shown to be significantly more likely to develop delirium postoperatively.⁵

Thus, in 2012, the British Orthopaedic Association Standards for Trauma and Orthopaedics (BOAST) guidelines stated that one should "Assess patient's risk of delirium or dementia by actively looking for cognitive impairment when patients first present with hip fracture and perform regular re-assessment."⁶ More recently, the British Orthopaedic Association extended their guidelines beyond hip fractures, stating that all elderly or frail orthopedic trauma patients should routinely be assessed using a validated delirium assessment tool such as the 4AT.⁷

Numerous scoring systems have been described in acute and chronic cognitive impairment, documentation of which can aid in the detection of cognitive impairment upon admission and for subtle changes in cognition during the inpatient stay. Folstein's Mini-Mental State Examination⁸ is a tool used to estimate the severity of cognitive impairment and to follow the course of cognitive changes in a patient over time. However, the 30-point questionnaire can be somewhat cumbersome and thus many nonspecialists prefer the Abbreviated Mental Test Score (AMTS) as described by Hodkinson.⁹ The AMTS comprises a series of 10 questions; a score of less than 6 is generally regarded to be suggestive of impaired cognition.⁹ However, the AMTS is felt to be too crude compared with other more comprehensive scales to monitor for subtle changes in cognition over time.¹⁰

The AMTS is historically in widespread use throughout the United Kingdom including Northern Ireland. Although the AMTS is also not typically regarded as a validated delirium tool, serial AMTS testing has been shown to be helpful in identifying acute cognitive dysfunction in the elderly population,¹¹ with a decline of 2 or more points having high sensitivity and specificity for diagnosing postoperative delirium.¹² The function of the 4AT, as recently suggested by the British Orthopaedic Association, and the AMTS has been found to be comparable in the hip fracture population.¹³ The AMTS is commonly used in Northern Ireland in the clinical setting¹⁴ and has also been used in numerous studies assessing change in cognitive function among hip fracture patients. For example, Odor et al compared the impact of regional anesthesia on pre- and postoperative AMTS,¹⁵ and Tahir et al used the AMTS on admission to assess baseline cognitive function when screening for postoperative delirium.⁵ Thus, it has been suggested that, due to its brevity and sensitivity, the AMTS is a very useful screening test for elderly trauma patients to assess for impaired cognition of any cause.¹⁶

Table 1. The 10-Point Abbreviated Mental Test Score (AMTS) Template Featured in the Admission Pro Forma Which is in Widespread Use Within the South Eastern Health and Social Care Trust.

Confused: yes/no	
Question	Score
Age in years	
Time of day	
Name of hospital	
Memorize address e.g. 42 West Street	
Recognize person	
Current year	
Name of Queen/Prime Minister	
Date of birth	
Dates of World War II	
Count 20-1	
Recall address—42 West Street	
Score (Total 10)	

Research has shown that while many orthopedic junior doctors are aware of the importance of the use of the AMTS, only a small proportion (approximately 10%) can identify all 10 questions involved.¹⁶ In light of this, and in keeping with BOAST guidelines, an AMTS template (Table 1) has been incorporated into the admission proformas used within Northern Ireland's South Eastern Health and Social Care Trust Trauma & Orthopaedic Unit to serve as an aide memoire. There is also an option to select whether or not the patient appears confused. In cases where the AMTS is not recorded upon admission, this is usually completed during orthogeriatrician review, which may occur several days following admission and possibly after surgery has been completed.

Consideration of cognitive impairment upon admission is of particular relevance in Northern Ireland. Formal diagnoses of both mild cognitive impairment and dementia are routinely made by psychogeriatricians in the community setting. Such psychiatric outpatient clinic letters are not currently accessible on the Northern Ireland Electronic Care Record (NIECR),¹⁷ which is used in the acute hospital setting. Therefore, information available to doctors in the acute setting may be inaccurate or incomplete. Furthermore, recent changes in Northern Ireland legislation, namely the Deprivation of Liberty Safeguards implemented in December 2019, mean that depriving a patient who lacks capacity of their liberty could be an infringement of their human rights. Keeping a patient in hospital for fracture management where they are not free to leave could be considered a deprivation of liberty. Thus, it is important to determine

whether or not a patient has capacity to accept or decline treatment and act in accordance with legislation as necessary.

The aim of this project was to review current practice within the unit regarding documentation of AMTS upon admission among elderly orthopedic trauma patients including hip fracture patients. As an intervention, BOAST guidelines and initial findings were then distributed among junior doctors (foundation year doctors/core surgical trainees) responsible for clerking-in acute fracture patients. The audit loop was then closed to determine the effectiveness of the intervention.

This work was conducted in a district general hospital with emergency trauma and orthopedic services. Current practice within the unit is that after a patient has been accepted for orthopedic admission, they would be clerked-in by a junior member of the surgical team and then subsequently reviewed in-hours by a senior member of the orthogeriatric team. The patient would typically be consented for surgery by a more senior member of the orthopedic team during the post-take ward round. As per Northern Ireland's regional policy, patients deemed to have capacity to consent for surgery sign "Form I" preoperatively. Alternatively, if a patient were deemed not to have capacity but the procedure was believed to be in the patient's best interests, the surgeon signs "Form IV," ideally after consultation with the next of kin and considering any previously expressed wishes. Wherever possible, efforts to treat reversible causes of acute confusion, for example, infection and electrolyte disturbances, are made prior to surgery.

We hypothesized that older patients and those with known cognitive or mental health disorders would be more likely to be screened for cognitive impairment upon admission. We also hypothesized that core trainees would be more aware of the importance of screening for cognitive impairment than foundation year doctors and thus more likely to document the AMTS.

Methods

Data were obtained by means of inpatient chart review and access to the NIECR. Information regarding AMTS upon admission and upon subsequent orthogeriatrician review was recorded, in addition to the nature of consent form used (if applicable) and any history of cognitive impairment/mental health disorders.

Preintervention data were collected throughout November and December 2018. Patients were included if they were admitted to the unit under the care of the Trauma & Orthopaedic team with an acute orthopedic injury and aged >65 years or if they sustained a fragility hip fracture at any age.

The findings from the first cycle of data collection and a copy of BOAST guidelines were distributed in electronic format to all junior doctors in the unit responsible for completing the required admission paperwork. Postintervention data were then collected in a similar manner throughout January and February 2019.

Statistical analysis was performed using Stata 15.0 (StataCorp 2017, Stata Statistical Software: Release 15; StataCorp LLC). Comparisons were undertaken using χ^2 tests for

comparison of categorical variables and *t* tests for continuous variables. For pre- and posttest comparisons of the AMTS, a McNemar χ^2 test was utilized. Multiple logistic regression analyses were used to determine the factors influencing AMTS recording. A *P* value of less than .05 was considered statistically significant.

Results

Patient Demographics

In total, 79 patients were included in the analysis: 40 in the preintervention arm and 39 in the postintervention arm. Preintervention, 40 suitable patients (25 females and 15 males), mean age 81.7 years (range: 65-96), were identified. Of these patients, 25 (62.5%) sustained a hip fracture, mean age 82.1 years (range: 66-94). Other injuries were fractures of the distal femur (*n* = 3), tibia (*n* = 3), ankle (*n* = 2), acetabulum (*n* = 1), humerus (*n* = 1), cervical spine (*n* = 1), and periprosthetic femoral fracture (*n* = 2), as well as prosthetic hip dislocation (*n* = 1) and quadriceps tendon rupture (*n* = 1).

Postintervention, 39 suitable patients (24 females and 15 males), mean age 80.4 years (range: 42-95), were identified. Thirty of these patients (76.9%) sustained a hip fracture, mean age 82.7 years (range: 42-95). Other reasons for admission were fractures of the femoral shaft (*n* = 4), distal femur (*n* = 3), and acetabulum (*n* = 2).

Regarding grade of admitting doctor, 22 patients (55.0%) in the preintervention group and 26 (66.7%) in the postintervention group were admitted by a foundation year doctor. Eighteen patients (45%) in the preintervention group and 12 (30.8%) in the postintervention group were admitted by a core trainee. For 1 patient in the postintervention arm, the grade of admitting doctor was unspecified.

Recording of AMTS

Figure 1 demonstrates the recording of the AMTS for both the overall cohort and the hip fracture subgroup, pre- and postintervention.

Preintervention, only 9 (22.5%) of the 40 patients in the overall cohort and 7 (28.0%) of the 25 patients in the hip fracture subgroup had their AMTS recorded by the admitting doctor (Table 2). Table 2 also shows whether these patients were recorded as being confused or not upon admission. In the multiple regression model, there was no significant association between whether the AMTS was recorded upon admission and patient age (*P* = .566), sex (*P* = .412), type of injury (*P* = .206), confusional status (*P* = .118), or grade of admitting doctor (*P* = .058).

Postintervention, 15 (38.5%) of the 39 patients in the overall cohort and 11 (36.7%) of the 30 patients in the hip fracture subgroup had their AMTS recorded by the admitting doctor (Table 2). In the multiple regression model, there was no significant association between whether the AMTS was recorded upon admission and patient age (*P* = .214), sex (*P* = .120),




Figure 1. The recording of the Abbreviated Mental Test Score (AMTS) for both the overall cohort (A) and the hip fracture subgroup (B) both pre- and postintervention is demonstrated. Note a small but significant increase in the recording of the AMTS has been achieved postintervention for both the cohort overall and the hip fracture subgroup. Red: AMTS not recorded; blue: AMTS recorded.

type of injury ($P = .453$), confusional status ($P = .473$), or grade of admitting doctor ($P = .494$). Statistical analysis demonstrated a significant improvement in AMTS documentation both among the overall cohort ($P = .001$) and the hip fracture patients ($P = .019$).

Nature of Consent Form

Thirty-seven patients underwent operative management in both the pre- and postintervention cohorts. All hip fracture patients underwent operative intervention (preintervention $n = 25$, postintervention $n = 30$). Table 3 outlines the nature of consent form used among the overall cohort and for the hip fracture subgroup pre- and postintervention.

Table 3 also demonstrates the number of patients in each of the groups who had an AMTS recorded upon admission or

upon subsequent orthogeriatrician review (median time to orthogeriatrician review was 1 day following admission for both the pre- and postintervention arms). The mean AMTS for each type of consent form for the overall cohort and hip fracture subgroup, pre- and postintervention, is shown. The mean AMTS was calculated in cases where there was an AMTS score available, either from the admitting doctor's assessment or from the orthogeriatrician's assessment if not completed by the admitting doctor.

Prior Cognitive/Mental Health Disorder

As outlined in Table 4, 15 (37.5%) of the 40 patients in the preintervention cohort and 17 (43.6%) of the 39 patients in the postintervention cohort had evidence of prior cognitive or mental health disorder.

Overall, those with prior cognitive or mental health disorders were not significantly more likely to have their AMTS recorded upon admission ($P = .256$) but were however less likely to be consented for surgery using Form I ($P = .001$). Older patients were also less likely to be consented using Form I than younger patients ($P = .03$). The mean age of those being consented using Form I was 79.6 years (standard deviation: 10.32 years), whereas the mean age of those consented using Form IV was 85.3 years (standard deviation: 5.75 years).

Discussion

The aim of this project was to improve AMTS documentation upon admission among frail fracture patients. Following a simple brief intervention, a statistically significant improvement in AMTS documentation was demonstrated for both the overall elderly fracture patient cohort and the hip fracture subgroup.

The neuropathological hallmarks for Alzheimer's disease are β -amyloid senile plaques and τ -neurofibrillary tangles which accumulate in the parietal and neocortical regions of the brain. Mild cognitive impairment is regarded as a transitional stage between normal aging and Alzheimer disease and has been associated with neurofibrillary tangles in the hippocampus and other medial temporal regions. Introduced into the literature in 1988 by Reisberg and colleagues, Small et al highlighted that mild cognitive impairment is characterized by a cognitive decline without impairment in the ability to carry out activities of daily living.¹⁸ Interestingly, a large proportion of older hip fracture patients without prediagnosed dementia have been found to have evidence of Alzheimer disease pathology on analysis of cerebrospinal fluid biomarkers. This may in fact suggest there is a possibility of preclinical dementia among certain hip fracture patients.¹⁹

The significance of cognitive impairment, resulting from preexisting dementia or acute delirium, is well established among hip fracture patients. A patient's capacity to make decisions regarding their management and functional rehabilitation may be hindered. Delirium has been associated with increasing

Table 2. Documentation of the Abbreviated Mental Test Score (AMTS) and Confusional Status Upon Admission Among the Pre- and Postintervention Cohorts.

	Preintervention group		Postintervention group	
	Cohort overall (N = 40)	Hip fracture subgroup (n = 25)	Cohort overall (n = 39)	Hip fracture subgroup (n = 30)
AMTS recorded	9 (22.5%)	7 (28.0%)	15 (38.5%)	11 (36.7%)
AMTS not recorded	31 (77.5%)	18 (72.0%)	24 (61.5%)	19 (63.3%)
Documented to be confused	6 (15.0%)	4 (16.0%)	8 (20.5%)	6 (20.0%)
Documented to be not confused	21 (52.5%)	14 (56.0%)	13 (33.3%)	12 (40.0%)
Confusional status not recorded	13 (32.5%)	7 (28.0%)	18 (46.1%)	12 (40.0%)

Abbreviations: AMTS, Abbreviated Mental Test Score.

Table 3. Nature of Consent Form Used for Patients Undergoing Operative Intervention.^a

	Preintervention group		Postintervention group	
	Cohort overall undergoing operative management	Hip fracture subgroup undergoing operative management	Cohort overall undergoing operative management	Hip fracture subgroup undergoing operative management
Form I	28 (75.7%)	19 (72.0%)	27 (73.0%)	21 (70%)
No. of patients with AMTS recorded on admission	5	5	7	5
No. of patients with AMTS subsequently recorded by orthogeriatrician	20	14	19	15
Total with AMTS recorded	25	19	26	20
Mean AMTS of recorded values	9.4	9.3	8.2	9.1
Form IV	9 (24.3%)	6 (28.0%)	10 (37.0%)	9 (30%)
No. of patients with AMTS recorded on admission	4	2	6	6
No. of patients with AMTS subsequently recorded by orthogeriatrician	5	4	4	3
Total with AMTS recorded	9	6	10	9
Mean AMTS of recorded values	2.9	3.4	3.7	2.7

Abbreviations: AMTS, Abbreviated Mental Test Score.

^aForm I is used among patients believed to have capacity to make decisions regarding their treatment. Form IV is used among patients felt to lack capacity and therefore the clinician is acting in the patient's best interests.

Table 4. The Incidence of Known Prior Significant Cognitive or Mental Health Disorder Among the Pre- and Postintervention Cohort.

	Preintervention	Postintervention
Cognitive impairment/dementia	8 (20.0%)	9 (23.1%)
Alcohol excess	2 (5.0%)	4 (10.3%)
Depression/bipolar disorder/ anxiety/schizophrenia	5 (12.5%)	3 (7.7%)
Learning disability	0	1 (2.6%)
Total	15 (37.5%)	17 (43.6%)

age, female sex, underlying dementia, poorer general physical health, chest or urinary infections, and increased length of hospital stay.^{4,5} Inadequate analgesia has also been shown to negatively affect a patient's AMTS, and regional analgesia has been

found to reduce the risk of AMTS deterioration perioperatively.¹⁵ Following hip fracture, patients with impaired cognition are less likely to return to their own homes or have successful rehabilitation.¹ Furthermore, hip fracture patients with delirium have been found to experience increased length of hospital stay and increased 1-year mortality,⁵ and delirium superimposed on dementia is a strong predictor of functional dependence, institutionalization, and mortality.²⁰

Hip fracture surgery has been associated with a higher incidence of delirium than elective orthopedic surgery.²¹ However, the literature is lacking regarding the role of cognitive impairment in the recovery and rehabilitation regarding other orthopedic injuries, as studied in this cohort. We suggest that cognitive impairment may be of particular relevance among those with injuries affecting weight-bearing status (eg, ankle fractures) or use of mobility aids (eg, distal radial fractures).

Challenges can also arise when managing patients with delirium on an acute trauma ward, and antipsychotics are often prescribed in elderly patients. However, current evidence does not support the use of antipsychotics for the prevention of delirium, and antipsychotics have not been shown to improve delirium duration or severity.²² From our current results, it is unclear if the AMTS specifically helps with the management of delirium in the acute orthopedic patient. However, a low AMTS upon admission may highlight that a patient has a degree of cognitive impairment and is thus more likely to experience perioperative delirium.

Interestingly, we found that older fracture patients were not significantly more likely to be screened for cognitive impairment upon admission than younger patients. We suggest that if an elderly patient has an established dementia diagnosis, the admitting doctor may not see it necessary to record the AMTS. It should also be remembered that the AMTS value achieved upon admission cannot be assumed to represent the patient's best cognitive function, as their cognition may be affected for various reasons, including pain, unfamiliar environment, and time of day.

To our knowledge, this is the first reported study of associations between the AMTS and type of consent form used in Northern Ireland. Although the numbers were too small to make meaningful statistical comparisons, the mean AMTS for both the pre- and postintervention arms were greater than 8/10 among those consented using Form I and below 4/10 among those consented using Form IV (Table 3). While it is important to remember that level of cognition and capacity are difficult to define numerically, and capacity is regarded as time- and question-specific, we suggest that further work is warranted to investigate whether a cutoff score exists at which the clinician can have reasonable confidence in saying that a patient likely lacks capacity.

None of the patients had an AMTS recalculated between the time of initial AMTS recording and their surgery in cases where there was a delay of a few days in getting to theater. This may be significant as patients previously deemed to lack capacity may have improved with the correction of reversible factors; similarly, patients deemed to have capacity upon admission may have developed delirium. None of the patients had their AMTS recalculated postoperatively or prior to discharge. Therefore, we were not able to objectively determine whether undergoing surgery was associated with a deterioration in the AMTS.

Disappointingly, only a small improvement in the number of patients having their AMTS recorded upon admission was seen following education and distribution of BOAST guidelines. We suggest that barriers to compliance with BOAST guidelines include competing demands from more clinically urgent tasks (particularly during out-of-hours work), the assumption that the AMTS will be subsequently recorded by the orthogeriatric team, and possible lack of awareness due to rotational/shift-based work seen among junior doctors.

Future work should explore the effectiveness, or otherwise, of highlighting the clinical relevance of BOAST guidelines during both the unit induction sessions with each rotation of

junior doctors and weekly junior doctor teaching sessions. A focus group to explore trainee understanding of the importance of screening for cognitive impairment may also help to identify further barriers to compliance and means by which these may be overcome.

It can also be suggested that while the AMTS score upon admission with a fracture may not be entirely representative of baseline cognitive function, it plays an important role in encouraging the trauma team to screen for dementia and/or delirium and perhaps refer onward where appropriate, as well as identifying patients who may be at heightened risk of perioperative delirium. The AMTS may also serve as part of a preoperative risk assessment tool for cognitive impairment in the fracture clinic setting among patients with other orthopedic injuries for whom the decision is made to operate at outpatient review.

We are mindful that this project was commenced prior to the update on the British Orthopaedic Association guidelines suggesting the use of a validated delirium assessment tool.⁷ These guidelines have more recently suggested the use of the 4AT, which has been shown to be both a sensitive and specific method of screening for delirium in hospitalized older people.²³ However, in the United Kingdom, and in our trauma unit, the AMTS has historically been used. We suggest that an area for further work would be to compare AMTS and 4AT scores upon admission and postoperatively.

Conclusion

In conclusion, a small but significant improvement in compliance with BOAST guidelines was seen following distribution of BOAST guidelines in written format. Further work should explore means by which compliance with these guidelines can be improved and also the effect of cognitive impairment on outcomes related to injuries beyond hip fractures.

Authors' Note

Approval obtained from the South Eastern Health and Social Care Trust audit department.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Grace E. M. Kennedy  <https://orcid.org/0000-0002-1693-6052>

References

1. Heruti RJ, Lusky A, Barell V, Ohry A, Adunsky A. Cognitive status at admission: does it affect the rehabilitation outcome of elderly patients with hip fracture? *Arch Phys Med Rehabil.* 1999; 80(4):432-436.

2. Friedman SM, Menzies IB, Bukata SV, Mendelson DA, Kates SL. Dementia and hip fractures: development of a pathogenic framework for understanding and studying risk. *Geriatr Orthop Surg Rehabil.* 2010;1(2):52-62.
3. Tsai CH, Chuang CS, Hung CH, et al. Fracture as an independent risk factor of dementia: a nationwide population-based cohort study. *Medicine (Baltimore).* 2014;93(26):e188.
4. Smith TO, Cooper A, Peryer G, Griffiths R, Fox C, Cross J. Factors predicting incidence of post-operative delirium in older people following hip fracture surgery: a systematic review and meta-analysis. *Int J Geriatr Psychiatry.* 2017;32(4):386-396.
5. Tahir M, Malik SS, Ahmed U, Kozdryk J, Naqvi SH, Malik A. Risk factors for onset of delirium after neck of femur fracture surgery: a prospective observational study. *SICOT J.* 2018;4:27.
6. BOAST-I Version 2 Guidelines. Published 2012. Accessed December 22, 2019. <https://www.boa.ac.uk/wp-content/uploads/2014/12/BOAST-1.pdf>
7. BOAST. The care of the older or frail orthopaedic trauma patient. Published 2019. Accessed December 22, 2019. <https://www.boa.ac.uk/standards-guidance/boasts.html>
8. Folstein MF, Folstein SE, McHugh PR. Mini-mental state: a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12:189-198.
9. Hodkinson HM. Evaluation of a mental test score for assessment of mental impairment in the elderly. *Age Ageing.* 1972;1(4):233-238.
10. Sheehan B. Assessment scales in dementia. *Ther Adv Neurol Disord.* 2012;5(6):349-358.
11. Jitapunkul S, Pillay I, Ebrahim S. Delirium in newly admitted elderly patients: a prospective study. *Q J Med.* 1992;83(300):307-314.
12. Ní Chonchubhair A, Valacio R, Kelly J, O'Keefe S. Use of the abbreviated mental test to detect postoperative delirium in elderly people. *Br J Anaesth.* 1995;75(4):481-2.
13. Lindsay WA, Moppett IK. Interchangeability of AMT4 and AMTS in a hip fracture population. *Anaesthesia.* 2019;74(11):1477-1478.
14. Beringer TRO, Clarke J, Elliott JRM, Marsh DR, Heyburn G, Steele IC. Outcome following proximal femoral fracture in Northern Ireland. *Ulster Med J.* 2006;75(3):200-206.
15. Odor PM, Chis Ster I, Wilkinson I, et al. Effect of admission fascia iliaca compartment blocks on postoperative abbreviated mental test scores in elderly fractured neck of femur patients: a retrospective cohort study. *BMC Anesthesiol.* 2017 January 5; 17(1):2.
16. Jain NP, Guyver PM, McCarthy P, et al. Use of the abbreviated mental test score by junior doctors on patients with fractured neck of femur. *Arch Orthop Trauma Surg.* 2008;128(2):235-8.
17. NI Electronic Care Record 2012. Accessed June 09, 2020 . <http://www.ehealthandcare.hscni.net/niecr/niecr.aspx>
18. Small GW, Kepe V, Ercoli LM, et al. PET of brain amyloid and tau in mild cognitive impairment. *N Engl J Med.* 2006;355(25):2652-2663.
19. Oh ES, Blennow K, Bigelow GE, et al. - β 42 and in without. *PLoS One.* 2018;13(9):e0204695.
20. Morandi A, Davis D, Fick DM, et al. Delirium superimposed on dementia strongly predicts worse outcomes in older rehabilitation inpatients. *J Am Med Dir Assoc.* 2014;15(5):349-354.
21. Bruce AJ, Ritchie CW, Blizard R, Lai R, Raven P. The incidence of delirium associated with orthopedic surgery: a meta-analytic review. *Int Psychogeriatr.* 2007;19(2):197-214.
22. Neufeld KJ, Yue J, Robinson TN, Inouye SK, Needham DM. Antipsychotic medication for prevention and treatment of delirium in hospitalized adults: a systematic review and meta-analysis. *J Am Geriatr Soc.* 2016;64(4):705-714.
23. Bellelli G, Morandi A, Davis DHJ, et al. Validation of the 4AT, a new instrument for rapid delirium screening: a study in 234 hospitalised older people. *Age Ageing.* 2014;43(4):496-502.